表面引发的 DNA 杂交链式反应的石英晶体微天平研究

王萍 葛志磊 裴昊 王丽华 樊春海*

(中国科学院上海应用物理研究所物理生物学实验室 上海 201800)

摘要 通过石英晶体振荡技术研究了杂交链式反应这种核酸扩增的方法. 石英晶体微天平可以表征在晶体和溶液的界面上的 DNA 层, 并获得粘性穿透深度这一重要参数. 根据石英晶体表面吸附质量和振荡频率之间的关系, 我们测量了表面引发的杂交链式反应的动力学过程, 并获得界面层的粘度、剪切模量等参数. 这一工作为研究固液界面上核酸反应过程, 特别是杂交链式反应的机制提供了新的途径.

关键词 杂交链式反应; 石英晶体微天平; DNA; 粘度; 界面; 穿透深度

Quartz Crystal Microbalance Studies on Surface-Initiated DNA Hybridization Chain Reaction

Wang, Ping Ge, Zhilei Pei, Hao Wang, Lihua Fan, Chunhai*

(Laboratory of Physical Biology, Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800)

Abstract The hybridization chain reaction (HCR), a type of nucleic acid amplification method, was studied by a quartz crystal resonator technique. We probed the properties of the boundary DNA layer with a thickness of the viscous penetration depth at the crystal-solution interface. According to the relationship between the change of resonant frequency and the mass adsorbed on the crystal surface, the kinetics of surface-initiated HCR was measured, as well as the viscosity and the shear modulus of H1/H2 solution with different concentrations near a solid substrate. This work provides a new method for studying nucleic acid reactions at solid surface and mechanistic insight for HCR-based amplification and biodetection.

Keywords hybridization chain reaction; quartz crystal microbalance; DNA; viscosity; interface; penetration depth

1 Introduction

Hybridization chain reaction (HCR) is a type of nucleic acid amplification reactions and this class of mechanisms suggests the possibility of constructing biosensors solely from unmodified single-stranded DNA.^[1-3] When metastable DNA hairpin probes meet an initiator molecule, they polymerize non-covalently to a long linear chain. The driving force for this reaction is the formation of inter-strand base pairs between previously unpaired bases in the polymerizing hairpins. Figure 1 shows the reaction pathway involving three species of DNA strands: hairpin probes H1, H2 and the initiator. The DNA hairpin structure is a good reservoir of hybridization energy that can be used as a fuel. Pierce and Dirks first proposed the concept of HCR and provided a mechanistic study of the free-running process in solution.^[4] They report a multiplexed fluorescent *in situ* hybridization method based on orthogonal amplification with HCR,^[1] small conditional RNAs that undergo HCR to induce cell death via an innate immune response.^[3] More recently, surface-initiated HCR was employed as amplifier for biomolecule detection.^[5-7]

Classical methods such as chromophore, fluorophore, or enzyme tags required for detection can affect the interac-

Figure 1 Reaction pathway of HCR occurring at the gold surface

tion between molecules,^[8-22] and nonspecific adsorption of analysts to a well modified surface can lead to random orientation of binding sites and possible denaturation of substrate.^[23] These issues present formidable challenges to accurate quantitation using the techniques that can be used to study macromolecules at the solid-solution interface. Since the solid-solution interface is sandwiched by two condensed phases, many techniques that are widely used for studying phenomena at solid-vapor interfaces are difficult to be applied. Quartz crystal microbalance (QCM) is a powerful tool used in solid-liquid interface which develop

* E-mail: fchh@sinap.ac.cn

Received August 3, 2012; published September 5, 2012.

Project supported by the National Natural Science Foundation of China (No. 21075128) and Shanghai Municipal Commission for Science and Technology (No. 10QA1408200).

项目受国家自然科学基金(No. 21075128)和上海市科委(No. 10QA1408200)资助.

a label-free, time-efficient, quantitative assay format that allows for multifunctional substrates, multilayer adsorptions, and nonrigid biological recognition.^[24-26]

Quartz crystal resonator is a surface-sensitive technique that relies on measurements of the response curves of a quartz crystal oscillator under a frequency sweep of the acoustic shear mode excitation applied across the crystal.^[27-30] The propagation of the excitation in the medium with which the crystal is in contact directly affects the frequency and its higher harmonics. The shifts of the resonant peaks and the changes in the width of the peaks are related to the mass deposition onto the crystal surface and the viscoelastic properties of the deposited mass. This technique is very sensitive for detecting small changes in the mass deposition on the surface and in the coupling between the deposited entities and the environment. Therefore it has been increasingly used in studies of the adsorption of macromolecules on various substrates.^[31-33] When a quartz crystal is immersed in liquid, the acoustic excitation from the crystal decays exponentially in the liquid, the technique thus probes a region next to the surface with a depth comparable to a so-called viscous penetration depth.^[29] QCM technique has been widely used in biology and chemistry. For example, it was employed to explore the supported lipid bilayers (SLBs) formation on various substrates.^[31,34-37] Structure changes and mechanical properties of high molecule polymers were studied using QCM-D technique in the boundary layer.^[23-28] Besides, antibody/antigen reaction or other nanomaterials with 3D structures and multilayers were studied by QCM-D technique.^[38-40] In this work, we conducted a mechanistic study on surface-initiated HCR by using quartz crystal oscillator technique. We monitored the kinetics of HCR in different conditions and the mechanical properties of adsorbed DNA film near the boundary layer.

2 Results and discussion

In this study, a freshly cleaned quartz crystal with a fundamental resonant frequency of 5 MHz was installed in the liquid cell. Gold films evaporated on the surfaces of the quartz crystals as the electrodes were used as the adsorption substrates. The initiator DNA were immobilized on the gold electrode by thiol group. After a solution with H1/H2 hairpin (consisting two nucleic acid hairpin species) was added to the cell, we observed the frequency and dissipation changes gradually in real time. HCR is a simple and easy-control system which is suitable to explore the relation between the mechanical properties in boundary layer and the length or modified density of DNA double helix. It may have implications to improve our understanding of biomolecule reactions at the solid-liquid interface that is widely used in gene chips.

2.1 Probe concentration dependence of HCR

The key to a HCR system is the storage of potential energy in short loops protected by long stems.^[1,2,4] Amplification of the initiator recognition event continues until the supply of H1 or H2 is exhausted. We changed the concentration of H1/H2 solution that was flown through the liquid cell, while keeping the initiator DNA concentration at 100 nmol/L. Figure 2 (a) and (b) displays

the measured resonant frequency shift (Δf) and the dissipation factor changes (ΔD) of the quartz crystal in solutions of H1/H2 with different concentrations. Both Δf and ΔD changed rapidly as the solution was flown through the liquid cell. As the concentration of H1/H2 increased, Δf and ΔD changed more rapidly. That means the velocity of this reaction is more rapidly and the H1/H2 molecule in HCR are not orderly one by one reaction as described in Figure 1, but a random process which speeds was affected by the reactant concentration. However, the reaction was finished after 30 min even with the highest concentration and Δf and ΔD reached a fixed value. As a result, higher concentration of the H1/H2 solution results in faster speed of HCR.

Figure 2 Experiment of probe concentration dependence of HCR. (a) Change of resonant frequency as a function of time. (b) Change of dissipation factor as a function of time. (c) Fitting curve of the kinetics of HCR

2.2 Initiator concentration dependence of the hybridization chain reaction

The average molecule weight of the resulting polymers is inversely related to the initiator concentration.^[1,3,4] We changed the concentration of the initiator on the surface of crystal via changing the concentration of the initiator solution, then flew 1 mL H1/H2 hairpin solution of the same concentration (200 nmol/L) each time. As shown in Figure 3 (a) and (b), the resonant frequency (Δf) decrease and the dissipation factor (ΔD) increase which demonstrate there are more and more H1/H2 bind to the initiator and become a longer polymer. After about 20000 s, the Δf and ΔD reached a fix value until the H1/H2 solution is exhausted. Added the buffer solution again and found that the curves kept in the stable value, which illustrates the

Figure 3 Experiment of the initiator concentration dependence of HCR. (a) Change of resonant frequency as a function of time. (b) Change of dissipation factor as a function of time. (c) Fitting curve of the kinetics of HCR

DNA double strands were chemically adsorbed on the gold-coated crystal surface and no H1/H2 hairpin remain in solution. Considering the reaction time is approximately the same even in different initiator concentration, the reaction velocity increases as the initiator concentration increase and we obtained the polymers of higher molecule weight with lower initiator.

2.3 Kinetics of the hybridization chain reaction

Many biological reactions occur at the liquid-solid interface rather than in liquid solution, especially in various biochips and nanomaterials appear these years.^[41-45] However, the kinetic and mechanical properties of molecules interactions in boundary layer can not be readily deduced from that in solution.^[40] First, the molecules modified on the surface reduce their translational and rotational freedom, and results in a preferred orientation of the interaction partners to each other compare to the normal of the surface; second, lateral and rotational diffusion of the modified molecule is much slower than in solution.

The mass of H1/H2 react and bind on the surface of gold-coated quartz crystal is related to the changes of resonant frequency. There is a linear relationship between the frequency response of an oscillating piezoelectric crystal and mass deposited on that crystal. This is mathematically expressed by the Sauerbrey equation^[46]:

$$\Delta f = C \cdot \Delta m \quad (1)$$

Where Δf is the change of resonant frequency of quartz crystal, C is the sensitivity factor, Δm is the change in mass. If we describe the HCR simply as follows:

$H1 + H2 \xrightarrow{k} H1 \cdot H2$, the kinetics of hybridization chain reaction can be shown as Equation 2, finally, we can gain the relation between the adsorbed mass of H1/H2 and the resonant frequency as Equation 3.

$$k = \frac{\Delta C_{H1 \cdot H2}}{\Delta T} \quad (2)$$

$$k = \frac{\Delta f \cdot \Delta T}{C \cdot V} \quad (3)$$

Where ΔT is the reaction time of HCR, $\Delta C_{H1 \cdot H2}$ is the concentration of $H1 \cdot H2$, V is the volume of the liquid cell. Because V and ΔT is about the same in this reaction, we can finally obtain this relation: $k \sim \Delta f \sim C \cdot \Delta m$. As shown in Figure 2 (c), the kinetics curve is nearly linear when the concentration is higher than 100 nmol/L: $k = 27.6 + 0.318 \Delta C_{H1 \cdot H2}$, we can conclude that if the H1/H2 concentration is higher, the rate of HCR will be higher simultaneously. Whereas in Figure 3 (c), the fitting curve is: $k = -105.2 \exp(-\Delta C_{H1 \cdot H2}/128.9) + 111.0$. As the initiator concentration grows, the curve is non-linear. When the concentration reaches a high threshold, the kinetics of HCR turns to be stable. This issue can be explained by the boundary effects which is a simple geometric consideration. If the initiator DNA can be treated approximately as rounded balls, a flat rigid surface in contact with a layer of initiator would result in void spaces between the initiator and the surface. Thus, when

the surface of the substrate is close packing, there is no big change of the reaction kinetics even the initiator concentration increases.

2.4 Shear modulus and viscosity of the film near the liquid-solid interface

As Sauerbrey equation holds for mass loading that is rigid with low dissipation but has been shown to overestimate mass adsorption in liquid loading, viscoelastic samples, and no rigid recognition and binding of biological molecules.^[24,25,34,47,48] Careful consideration of the reaction monofilm and liquid loading has shown that Δf and ΔD relate not only to adsorbed mass but also to the shear modulus and viscosity of the contact liquid, it is important to use a viscoelastic model such as Voigt-based model to quantify the changes in mass and effective viscoelastic properties of the DNA film in hybridization chain reaction.^[29] As seen in Equations (4) and (5), where ρ_{sol} and η_{sol} are the density and the viscosity of the solution; h_f and ρ_f is the thickness and the density of the adsorbed film; $m_q = 0.85$; $\omega = 2\pi f = 2\pi f_0 = 10\pi \times 10^6$

$$\Delta f = -\frac{1}{2\pi m_q} \left[\sqrt{\frac{\rho_{\text{sol}} \eta_{\text{sol}} \omega}{2}} + (h_f \rho_f \omega - h_f \rho_{\text{sol}} \omega \frac{\eta_f \eta_{\text{sol}} \omega^3}{\mu_f^2 + \eta_f^2 \omega^2}) \right] \quad (4)$$

$$\Delta D = \frac{1}{\pi f m_q} \left[\sqrt{\frac{\rho_{\text{sol}} \eta_{\text{sol}} \omega}{2}} + (h_f \rho_{\text{sol}} \omega \frac{\mu_f \eta_{\text{sol}} \omega}{\mu_f^2 + \eta_f^2 \omega^2}) \right] \quad (5)$$

This expression has been used in determining the viscoelastic properties of homogeneous solutions in contact with a quartz resonator.^[27,29,30,49] However, it should be noted that the materials parameters contained in these expressions are those for the boundary layer of the solutions adjacent to the crystal-solution interface and may differ from those for a bulk liquid. The viscosity (η) and the shear modulus (μ) of HCR boundary layers were subsequently calculated using Eqs. 4 and 5 based on the measured Δf and ΔD . Figure 4 shows the changes of shear modulus (μ) and viscosity (η) in H1/H2 solution concentration dependence experiment. When the concentration increases, the shear modulus and the viscosity decrease because there are more H1 and H2 react and form to longer DNA double helix. The longer this DNA molecule, the smaller the shear modulus and the viscosity, which indicates the DNA molecule is like a brush modified on the crystal surface which the longer brush is softer. Figure 5 shows the changes of shear modulus and viscosity in initiator concentration dependence experiment. When the concentration of the initiator solution increases, the density of HCR chain near the boundary increases. When the initiator concentration is above about 100 nmol/L, the viscosity and the shear modulus decrease slowly. We consider that the steric hindrance of the initiator molecule adsorbed on the crystal surface obstruct more H1/H2 molecule form to a longer chain, that cause the viscosity and the shear modulus decrease not so much like it in Figure 4. However, both the

Figure 4 Change of shear modulus and viscosity as a function of concentration. (a~b) Experiment of probe concentration dependence of HCR. (c~d) Experiment of the initiator concentration dependence of HCR

chain length and the density of chain can make μ and η decrease, but the effect of chain density is much less than that of chain length. These results are consistent to our previous results which is the scaled polymer molecules studied by QCM.^[25]

3 Conclusion

We have studied the hybridization chain reaction using quartz crystal resonator technique. The kinetics of this reaction was obtained in different conditions. First, the resonant frequency and the dissipation factor increase as the H1/H2 solution concentration was increased; Second, as the initiator concentration increases, the reaction velocity of HCR increases rapidly when the initiator concentration is below 100 nmol/L, when the initiator concentration is above 100 nmol/L, the reaction velocity of HCR is approximately keep constant. In addition, the mechanical properties of HCR on the liquid-solid interface are illustrated. The viscosity and shear modulus of the H1/H2 film adsorbed on the crystal surface decrease as the density or the concentration increase, but the effect of density is smaller than that of concentration to a certain extent because of the steric hindrance.

4 Experimental section

DNA oligonucleotides were purchased from Invitrogen and used as received.^[4] For the HCR system shown in Figure 1, concentrated DNA solutions were prepared in buffer that was later diluted to reaction conditions: 1 mol/L NaCl, 50 mmol/L Na₂HPO₄, pH 7.5. To ensure that H1 and H2 form hairpin monomers, the strands were snap-cooled before use (heat at 95 °C for 3 min and then cool to room temperature for at least 30 min).

In the experiments, the gold-coated quartz crystal with a fundamental resonant frequency of 5 MHz was bought from Q-Sense. Prior to use, the gold electrode surface were thoroughly cleaned with piranha solution [$V(\text{H}_2\text{SO}_4) : V(\text{H}_2\text{O}_2) = 3 : 1$], then rinsed with Milli-Q water, and blown dry with a stream of dry nitrogen gas. A droplet of 100 μL initiator DNA solution was incubated with gold for approximately 16 h in 4 °C, the gold surface was washed with Milli-Q water, then immersed into the mercaptohexanol (MCH) solution (1 mmol/L) for 30 min, it was washed with Milli-Q water again, and blown it dry with nitrogen gas. The liquid cell was initially filled with pure water; the frequency shift and the change of the dissipation factor of the crystal were measured in air and in pure water to check against the standards. After that, buffer solution was added to the cell and then the H1/H2 solution (consisting two nucleic acid hairpin species) with different conditions was flown through the cell. When the equilibrium of the solution was clearly reached, flown buffer solution again to check whether the HCR film was bind to the gold surface chemically. The temperature of the liquid cell was controlled at 24 °C throughout the experiment.

References

- [1] Choi, H. M. T.; Chang, J. Y.; Trinh, L. A.; Padilla, J. E.; Fraser, S. E.; Pierce, N. A. *Nat. Biotechnol.* **2010**, *28*, 1208.
- [2] Venkataraman, S.; Dirks, R. M.; Rothmund, P. W. K.; Winfree, E.;

- Pierce, N. A. *Nat. Nanotechnol.* **2007**, *2*, 490.
- [3] Venkataraman, S.; Dirks, R. M.; Ueda, C. T.; Pierce, N. A. *Proc. Natl. Acad. Sci.* **2010**, *107*, 16777.
- [4] Dirks, R. M.; Pierce, N. A. *Proc. Natl. Acad. Sci.* **2004**, *101*, 15275.
- [5] Green, S. J.; Lubrich, D.; Turberfield, A. J. *Biophys. J.* **2006**, *91*, 2966.
- [6] Lubrich, D.; Green, S. J.; Turberfield, A. J. *J. Am. Chem. Soc.* **2009**, *131*, 2422.
- [7] Turberfield, A. J.; Mitchell, J. C.; Yurke, B.; Mills, A. P., Jr.; Blakey, M. I.; Simmel, F. C. *Phys. Rev. Lett.* **2003**, *90*, 118102.
- [8] Niu, S.; Jiang, Y.; Zhang, S. *Chem. Commun.* **2010**, *46*, 3089.
- [9] Zhang, C. Y.; Yeh, H. C.; Kuroki, M. T.; Wang, T. H. *Nat. Mater.* **2005**, *4*, 826.
- [10] Bath, J.; Green, S. J.; Allen, K. E.; Turberfield, A. J. *Small* **2009**, *5*, 1513.
- [11] McKee, M. L.; Milnes, P. J.; Bath, J.; Stulz, E.; Turberfield, A. J.; O'Reilly, R. K. *Angew. Chem. Int. Ed.* **2010**, *49*, 7948.
- [12] Muscat, R. A.; Bath, J.; Turberfield, A. J. *Nano Lett.* **2011**, *11*, 982.
- [13] Turberfield, A. J. *Nat. Chem.* **2011**, *3*, 580.
- [14] Boeneman, K.; Mei, B. C.; Dennis, A. M.; Bao, G.; Deschamps, J. R.; Mattoussi, H.; Medintz, I. L. *J. Am. Chem. Soc.* **2009**, *131*, 3828.
- [15] Song, S.; Liang, Z.; Zhang, J.; Wang, L.; Li, G.; Fan, C. *Angew. Chem., Int. Ed.* **2009**, *48*, 8670.
- [16] Zhang, J.; Song, S. P.; Wang, L. H.; Pan, D.; Fan, C. H. *Nat. Protoc.* **2007**, *2*, 2888.
- [17] Liu, G.; Wan, Y.; Zou, Z.; Ren, S.; Fan, C. *Chin. J. Anal. Chem.* **2011**, *39*, 953 (in Chinese).
- [18] Yan, J.; Hu, M.; Li, D.; He, Y.; Zhao, R.; Jiang, X.; Song, S.; Wang, L.; Fan, C. *Nano Res.* **2008**, *1*, 490.
- [19] Song, S.; Liang, Z.; Zhang, J.; Wang, L.; Li, G.; Fan, C. *Angew. Chem., Int. Ed.* **2009**, *48*, 8670.
- [20] Yan, J.; Song, S.; Li, B.; Zhang, Q.; Huang, Q.; Zhang, H.; Fan, C. *Small* **2010**, *6*, 2520.
- [21] Fan, C.; Wang, S.; Hong, J. W.; Bazan, G. C.; Plaxco, K. W.; Heeger, A. J. *Proc. Natl. Acad. Sci.* **2003**, *100*, 6297.
- [22] He, S.; Song, B.; Li, D.; Zhu, C.; Qi, W.; Wen, Y.; Wang, L.; Song, S.; Fan, H.; Fan, C. *Adv. Funct. Mater.* **2010**, *20*, 453.
- [23] Gerdon, A. E.; Wright, D. W.; Cliffel, D. E. *Anal. Chem.* **2004**, *77*, 304.
- [24] Wang, P.; Fang, J.; Hou, Y.; Du, X.; Zhu, D.-M. *J. Phys. Chem. C* **2008**, *113*, 729.
- [25] Wang, P.; Fang, J.; Qin, S.; Kang, Y.; Zhu, D.-M. *J. Phys. Chem. C* **2009**, *113*, 13793.
- [26] Wu, B.; Wu, K.; Wang, P.; Zhu, D.-M. *J. Phys. Chem. C* **2006**, *111*, 1131.
- [27] Rodahl, M.; Kasemo, B. *Sens. Actuators, A* **1996**, *54*, 448.
- [28] Johannsmann, D.; Reviakine, I.; Rojas, E.; Gallego, M. *Anal. Chem.* **2008**, *80*, 8891.
- [29] Voinova, M. V.; Rodahl, M.; Jonson, M.; Kasemo, B. *Phys. Scr.* **1999**, *59*, 391.
- [30] Kanazawa, K. K.; Gordon, J. G. *Anal. Chem.* **1985**, *57*, 1770.
- [31] Hook, F.; Kasemo, B.; Nylander, T.; Fant, C.; Sott, K.; Elwing, H. *Anal. Chem.* **2001**, *73*, 5796.
- [32] Kitano, K.; Inoue, Y.; Matsuno, R.; Takai, M.; Ishihara, K. *Colloids Surf., B* **2009**, *74*, 350.
- [33] Amstad, E.; Zurcher, S.; Mashaghi, A.; Wong, J. Y.; Textor, M.; Reimhult, E. *Small* **2009**, *5*, 1334.
- [34] Cho, N.-J.; Frank, C. W.; Kasemo, B.; Hook, F. *Nat. Protoc.* **2010**, *5*, 1096.
- [35] Edvardsson, M.; Svedhem, S.; Wang, G.; Richter, R.; Rodahl, M.; Kasemo, B. *Anal. Chem.* **2008**, *81*, 349.
- [36] Granéli, A.; Rydström, J.; Kasemo, B.; Höök, F. *Langmuir* **2003**, *19*, 842.
- [37] Reimhult, E.; Zäch, M.; Höök, F.; Kasemo, B. *Langmuir* **2006**, *22*, 3313.
- [38] Höök, F.; Rodahl, M.; Brzezinski, P.; Kasemo, B. *Langmuir* **1998**, *14*, 729.
- [39] Arlett, J. L.; Myers, E. B.; Roukes, M. L. *Nat. Nanotechnol.* **2011**, *6*, 203.
- [40] Gavutis, M.; Lata, S.; Piehler, J. *Nat. Protoc.* **2006**, *1*, 2091.
- [41] Thierry, B.; Jasieniak, M.; de Smet, L. C. P. M.; Vasilev, K.; Griesser, H. J. *Langmuir* **2008**, *24*, 10187.
- [42] Braun, T.; Ghatkesar, M. K.; Backmann, N.; Grange, W.; Boulanger, P.; Letellier, L.; Lang, H.-P.; Bietsch, A.; Gerber, C.; Hegner, M. *Nat. Nanotechnol.* **2009**, *4*, 179.
- [43] Federici, S.; Oliviero, G.; Hamad-Schifferli, K.; Bergese, P.

- Nanoscale* **2010**, 2, 2570.
- [44] Choi, S.; Yang, Y.; Chae, J. *Biosens. Bioelectron.* **2008**, 24, 893.
- [45] Pei, H.; Lu, N.; Wen, Y.; Song, S.; Liu, Y.; Yan, H.; Fan, C. *Adv. Mater.* **2010**, 22, 4754.
- [46] Sauerbrey, G. *Z Phys.* **1959**, 155, 206.
- [47] Gerdon, A. E.; Wright, D. W.; Clifffel, D. E. *Anal. Chem.* **2005**, 77, 304.
- [48] Zhu, D.-M.; Wu, K.; Wu, B.; Wang, P.; Fang, J.; Hou, Y.; Zhang, G. *J. Phys. Chem. C* **2007**, 111, 18679.
- [49] Rodahl, M.; Hook, F.; Krozer, A.; Brzezinski, P.; Kasemo, B. *Rev. Sci. Instrum.* **1995**, 66, 3924.

(Qin, X.)