氯化亚铁催化的醛与伯胺和仲胺的氧化酰胺化反应

李远明 贾凡 马丽娜 李志平*

(中国人民大学化学系 北京 100872)

摘要 酰胺化反应是合成含有酰胺键的天然产物、材料以及药物分子中一类非常重要的化学反应. 发展高原子经济性和环境友好的新型酰胺键的合成方法仍然是该领域具有重要研究意义和价值的研究内容. 报道了在氯化亚铁作为催化剂和叔丁基过氧化氢作为氧化剂的条件下, 实现了芳香醛和脂肪醛分别与伯胺和仲胺的直接酰胺化反应, 获得了较高的收率. 进一步研究发现, 该反应体系对氮杂环化合物也有较好的适用性.

关键词 铁催化; 氧化; 酰胺化; 醛; 胺

FeCl₂-Catalyzed Oxidative Amidation of Aldehydes with Primary and Secondary Amines

Li, Yuanming Jia, Fan Ma, Lina Li, Zhiping*

(Department of Chemistry, Renmin University of China, Beijing 100872)

Abstract The amidation reaction is an important process in both academia and industry for the synthesis of natural products, functional polymers, and pharmaceuticals. The development of new amidation methods for constructing such important class of organic compounds is still attracting considerable interests. We herein reported a general oxidative amidation of aldehydes with a variety of free primary or secondary amines using inexpensive iron(II) chloride as catalyst and *tert*-butyl hydroperoxide (TBHP) as oxidant. The present amidation reactions afford the corresponding secondary and tertiary amides in moderate to excellent yields. In addition, the amidations of benzimidazole and pyrazole were also achieved by the developed method.

Keywords iron catalysis; oxidative; amidation; aldehyde; amine

1 Introduction

The amide bond widely exists in natural products, polymers, and pharmaceuticals.^[1] Historically, chemical reactions for the formation of amide bond are among the most classical transformations in organic chemistry. The most well-established routes toward amides synthesis are the reactions of amines with carboxylic acids (or their derivatives),^[2] the Schmit reaction,^[3] the Beckmann rearrangement reaction,^[4] and modified Staudiger reaction. Among the emerging amide formation reactions,^[6] the direct cross-dehydrogenative coupling (CDC) of aldehydes with amines has evolved into one of the most promising methods with potential industrial applications.^[7]

Three strategies have been developed for amide bond formation using aldehydes and primary or secondary amines as starting materials (Scheme 1). The aldehydes are activated through acyl bromide,^[8] acyl cyanide,^[9] nitrile,^[10] or active ester^[11] as intermediate to couple with amines (Strategy I). Amines can also be transformed into *N*-chloroamines^[12] and anionic amido complex^[13] which play as active intermediates to react with aldehydes (Strategy II). Following the pioneering work of Yoshida,^[14] sev-

eral transition metal (Pd,^[15] Rh,^[16] Cu,^[17] Fe,^[18] Zn^[19]) catalyzed amidation reactions have been reported to date (Strategy III). Despite the rapid progress have been made in amidation of aldehydes with amines, there are still some shortcomings. Some of the catalysts reported are either expensive or difficult to handle. The pre-activation of amines is still inefficient from the synthetic point of view. As such, the efficient and easily-handled catalytic systems bearing broader substrate scope are in great demand.

Scheme 1 Types of transformation of aldehyde with amine

Recently, we reported iron-catalyzed oxidative ami-

* E-mail: zhipingli@ruc.edu.cn; Tel.: 010-62514226; Fax: 010-62516444

Received April 27, 2015; published June 5, 2015.

Supporting information for this article is available free of charge via the Internet at <http://sioc-journal.cn>.

Project supported by the Fundamental Research Funds for the Central Universities and the Research Funds of Renmin University of China (No. 10XNL017).

项目受中国人民大学科学研究基金(中央高校基本科研业务费专项资金资助)项目成果(No. 10XNL017)资助.

dation of tertiary Amines.^[20] Based on previous results, we herein reported a general method for the formation of secondary and tertiary amides from aldehydes and free amines in iron(II) chloride catalyzed oxidative system.

2 Result and Discussion

2.1 Optimization of the reaction conditions

Reaction was conducted by using benzaldehyde **1a** and benzylamine **2a** as the model for the screening of reaction conditions (Table 1). As anticipated, the desired amide **3a** was detected in the iron-catalyzed oxidative system, albeit in trace amount (Entry 1). By increasing the amount of **1a** and TBHP, 56% and 71% yields of the product **3a** were obtained respectively by using 2.5 mol% FeCl₂ as catalyst (Entries 2 and 3). However, other iron salts and first-row transition metal salts provided worse results (Entries 4~11). To our delight, a 90% yield of the desired product **3a** was achieved when 5 equiv. of **1a** were used (Entry 12). Notably, in the absence of catalyst, **3a** was obtained in only 18% yield (Entry 13). By varying oxidants (Entries 14, 15), the results indicated that TBHP was the best choice of oxidant.

Table 1 Optimization of the reaction conditions^a

Entry	1a /equiv.	[Oxidant]/equiv.	Cat.	Yield ^b /%
1	1	TBHP (1.0)	FeCl ₂	<5
2	3	TBHP (1.0)	FeCl ₂	56
3	3	TBHP (2.0)	FeCl ₂	71
4	3	TBHP (2.0)	FeCl ₃	19
5	3	TBHP (2.0)	FeBr ₂	11
6	3	TBHP (2.0)	Fe(OAc) ₂	<5
7	3	TBHP (2.0)	FeSO ₄ ·7H ₂ O	<5
8	3	TBHP (2.0)	Fe ₂ (CO) ₉	<5
9	3	TBHP (2.0)	CuBr ₂	46
10	3	TBHP (2.0)	CuSO ₄ ·5H ₂ O	50
11	3	TBHP (2.0)	CoCl ₂	50
12	5	TBHP (2.0)	FeCl ₂	90
13	5	TBHP (2.0)	—	18
14	5	H ₂ O ₂ (2.0)	FeCl ₂	0
15	5	O ₂ (1 atm)	FeCl ₂	0

^a Conditions: **2a** (0.5 mmol), catalyst (0.0125 mmol), CH₃CN (3.0 mL), 85 °C, 1 h; ^b Reported yields were based on **2a** and determined by ¹H NMR using an internal standard.

2.2 The reaction scope of aromatic aldehydes with primary amines

With the optimized conditions in hand, we next investigated the scope and generality of such transformation with various aromatic aldehydes and primary amines (Scheme 2). The representative electron-donating or electron-withdrawing groups on phenyl ring of benzaldehyde sub-

strates produced good yields of secondary amides (**3a**~**3c**, **3e**). However, with nitro group, a diminished yield was obtained (**3d**). Various 4-substituted functional groups on the phenyl ring of anilines are well-suited for this transformation (**3f**, **3j**). Due to the steric effect of 2,6-dimethylaniline, the corresponding amide **3k** was obtained in 53% yield. To our satisfaction, the reaction also works with acetic hydrazide, providing the desired amide **3l** in 52% yield.

Conditions: **1** (2.5 mmol), **2** (0.5 mmol), FeCl₂ (0.0125 mmol), CH₃CN (3.0 mL), 85 °C, 1 h; ^b Reported yields were based on **2** and determined by ¹H NMR using an internal standard; isolated yields were given in parentheses.

Scheme 2 The reaction of aromatic aldehydes with primary amines

2.3 The reaction scope of aromatic aldehydes with secondary amines

Subsequently, a variety of secondary amines were investigated under the developed reaction conditions (Scheme 3). Under our optimized condition, a range of tertiary

Conditions: **1** (2.5 mmol), **2** (0.5 mmol), FeCl₂ (0.0125 mmol), CH₃CN (3.0 mL), 85 °C, 1 h; ^b Reported yields were based on **2** and determined by ¹H NMR using an internal standard; isolated yields were given in parentheses.

Scheme 3 The reaction of aromatic aldehydes with secondary amines

amides were obtained in good yields using secondary amines (**4a**~**4e**). The formation of more challenging *N*-phenyl tertiary amides (**4f**~**4h**) were also obtained in moderate yields under the present catalytic oxidative system, which was not previously accessible through *N*-phenyl anilines.^[17b,18] Gratifyingly, 2-thiophenecarboxaldehyde was also compatible, providing the desired amide **4i** efficiently. It should be noted that other reported oxidative amidation of aldehydes with secondary amines generally required longer reaction time and special class of substrates.^[15]

2.4 The reaction scope of aliphatic aldehydes with amines

Next, we explored the scope of aliphatic aldehydes (Scheme 4). In contrast to aromatic aldehydes, the reactions of aliphatic aldehydes with amines were messy under the standard reaction conditions, probably due to the undesired oxidation of amine.^[20a] To our satisfactory, the slow addition of amines using syringe pump partially solved the problem and the moderate yields of the desired amides (**5a**~**5f**) were obtained.

Conditions: **1** (2.5 mmol), **2** (0.5 mmol), FeCl₂ (0.0125 mmol), CH₃CN (3.0 mL), 85 °C, 1 h; ^b Reported yields were based on **2** and determined by ¹H NMR using an internal standard; isolated yields were given in parentheses.

Scheme 4 The reaction scope of aliphatic aldehydes with amines

2.5 The reactions of aldehydes with other amines

Furthermore, when methoxymethylamine hydrochloride **6a** was tested, *N*-methoxy-*N*-methylamide **7a** was obtained in 54% yield by using calcium carbonate as base (Eq. 1). Gratifyingly, the present reaction conditions were also suitable for the reactions of benzaldehyde with benzimidazole **6b** and pyrazole **6c**, resulted in 81% and 85% yields of the corresponding products (**7b**, **7c**) respectively (Eqs. 2 and 3).

2.6 The mechanistic studies

When benzoic acid **6d** instead of benzaldehyde **1a** reacted with benzylamine **2a**, the desired amide **3a** was not detected by ¹H NMR under the optimized reaction condi-

tions (Eq. 4). The result excluded that aldehydes were oxidized to carboxylic acids and followed by the reaction with amines to give amides.

In addition, imine **6e** could not be transformed into the desired amide **3a** under the optimized reaction conditions (Eq. 5). The result indicated that imine does not likely play as a reaction intermediate.^[21]

Recently, FeSO₄·7H₂O catalyzed the synthesis of amides from aldehydes and amine hydrochloride salts was reported by Chen.^[18] It is worth noting that **3a** was not detected in our reaction condition using FeSO₄·7H₂O as catalyst (Table 1, Entry 7). To probe the possible reasons, NaCl was added into the reactions by the use of FeSO₄·7H₂O as catalyst. Interestingly, a 36% yield of **3a** was obtained, while NaCl itself could not promote the desired amidation. These results suggested that both iron and chloride ions might play co-effects on the present amide formation, which needs further studies to elucidate the exact reasons. The comparison experiments were summarized in Table 2.

Table 2 The effects of iron sources

Cat.	Additive	Yield of 3a
FeCl ₂	—	90% (79%)
FeSO ₄ ·7H ₂ O	—	Trace
FeSO ₄ ·7H ₂ O	NaCl (5 mol%)	36%
—	NaCl (5 mol%)	trace

Accordingly, a simplified mechanism for the present oxidative amidation was proposed (Scheme 5). Free amine undergoes nucleophilic attack to aldehyde to give a carbi-

noamine intermediate, which is subsequently oxidized to the amide by iron/TBHP.

Scheme 5 A proposed mechanism

3 Conclusions

In summary, we developed a general protocol for the oxidative amidation of aldehydes with a variety of primary and secondary amines by simple iron salt as catalyst. The amidations of benzimidazole and pyrazole were also achieved, which demonstrated the advantages of the method. Further explorations directed towards synthetic applications are under progressing in our lab.

References

- [1] (a) Li, L.; Zhou, B.; Ye, L. *Chin. J. Org. Chem.* **2015**, 35, 655. (李龙, 周波, 叶龙武, 有机化学, **2015**, 35, 655.); (b) Pattabiraman, V. R.; Bode, J. W. *Nature* **2011**, 480, 471; (c) Allen, C. L.; Williams, J. M. J. *Chem. Soc. Rev.* **2011**, 40, 3405.
- [2] (a) Valeur, E.; Bradley, M. *Chem. Soc. Rev.* **2009**, 38, 606; (b) Montalbetti, C. A. G. N.; Falque, V. *Tetrahedron* **2005**, 61, 10827.
- [3] Lang, S.; Murphy, J. A. *Chem. Soc. Rev.* **2006**, 35, 146.
- [4] (a) Qin, C.; Zhou, W.; Chen, F.; Ou, Y.; Jiao, N. *Angew. Chem., Int. Ed.* **2011**, 50, 12595; (b) Li, Q.; Yan, L.; Xia, D.; Shen, Y. *Chin. J. Org. Chem.* **2011**, 31, 2034. (李倩, 严罗一, 夏定, 申永存, 有机化学, **2011**, 31, 2034.); (c) Owston, N. A.; Parker, A. J.; Williams, J. M. J. *Org. Lett.* **2007**, 9, 3599; (d) Mao, D.; Lu, G.; Chen, Q.; Xie, Z.; Zhang, Y. *Acta Chim. Sinica* **2001**, 59, 1139. (毛东森, 卢冠忠, 陈庆龄, 谢在库, 张玉贤, 化学学报, **2001**, 59, 1139.)
- [5] (a) Köhn, M.; Breinbauer, R. *Angew. Chem. Int. Ed.* **2004**, 43, 3106; (b) Damkaci, F.; DeShong, P. J. *Am. Chem. Soc.* **2003**, 125, 4408; (c) Saxon, E.; Bertozzi, C. R. *Science* **2000**, 287, 2007.
- [6] (a) Chen, J.; Mao, J.; Zheng, Y. *Chin. J. Org. Chem.* **2015**, 35, 672. (陈洁, 毛金成, 郑洋, 有机化学, **2015**, 35, 672.); (b) Wang, H.; Yang, L.; Xiao, Y.; Mao, P.; Zhu, X.; Yuan, J.; Qu, L. *Chin. J. Org. Chem.* **2014**, 34, 355. (王宏雁, 杨亮茹, 肖咏梅, 毛璞, 朱晓亮, 袁金伟, 屈凌波, 有机化学, **2014**, 34, 355.); (c) Martinelli, J. R.; Clark, T. P.; Watson, D. A.; Munday, R. H.; Buchwald, S. L. *Angew. Chem., Int. Ed.* **2007**, 46, 8460; (d) Cho, S. H.; Yoo, E. J.; Bae, I.; Chang, S. J. *Am. Chem. Soc.* **2005**, 127, 16046.
- [7] (a) Ding, Y.; Zhang, X.; Zhang, D.; Chen, Y.; Wu, Z.; Wang, P.; Xue, W.; Song, B.; Yang, S. *Tetrahedron Lett.* **2015**, 56, 831; (b) Ghosh, S. C.; Ngiam, J. S. Y.; Chai, C. L. L.; Seayad, A. M.; Dang, T. T.; Chen, A. *Adv. Synth. Catal.* **2012**, 354, 1407; (c) Ekoue-Kovi, K.; Wolf, C. *Chem. Eur. J.* **2008**, 14, 6302.
- [8] Markó, I. E.; Mekhelfia, A. *Tetrahedron Lett.* **1990**, 31, 7237.
- [9] Gilman, N. W. *J. Chem. Soc. D* **1971**, 733.
- [10] Shie, J.-J.; Fang, J.-M. *J. Org. Chem.* **2003**, 68, 1158.
- [11] (a) De Sarkar, S.; Studer, A. *Org. Lett.* **2010**, 12, 1992; (b) Vora, H. U.; Rovis, T. *J. Am. Chem. Soc.* **2007**, 129, 13796; (c) Tan, B.; Toda, N.; Barbas, C. F. *Angew. Chem., Int. Ed.* **2012**, 51, 12538; (d) Yao, H.; Tang, Y.; Yamamoto, K. *Tetrahedron Lett.* **2012**, 53, 5094; (e) Ji, M.; Lim, S.; Jang, H.-Y. *RSC Adv.* **2014**, 4, 28225.
- [12] (a) Vanjari, R.; Guntreddi, T.; Singh, K. N. *Green Chem.* **2014**, 16, 351; (b) Cadoni, R.; Porcheddu, A.; Giacomelli, G.; De Luca, L. *Org. Lett.* **2012**, 14, 5014; (c) Porcheddu, A.; Luca, L. D. *Adv. Synth. Catal.* **2012**, 354, 2949.
- [13] (a) Wang, J.; Li, J.; Xu, F.; Shen, Q. *Adv. Synth. Catal.* **2009**, 351, 1363; (b) Seo, S.; Marks, T. J. *Org. Lett.* **2008**, 10, 317; (c) Seo, S.; Marks, T. J. *Org. Lett.* **2007**, 10, 317; (d) Li, J.; Xu, F.; Zhang, Y.; Shen, Q. *J. Org. Chem.* **2009**, 74, 2575.
- [14] Tamaru, Y.; Yamada, Y.; Yoshida, Z. *Synthesis* **1983**, 474.
- [15] Ekoue-Kovi, K.; Wolf, C. *Org. Lett.* **2007**, 9, 3429.
- [16] (a) Dai, W.; Liu, Y.; Tong, T.; Li, X.; Luo, F. *Chin. J. Catal.* **2014**, 35, 1012; (b) Tillack, A.; Rudloff, I.; Beller, M. *Eur. J. Org. Chem.* **2001**, 2001, 523.
- [17] (a) Yoo, W.-J.; Li, C.-J. *J. Am. Chem. Soc.* **2006**, 128, 13064; (b) Ghosh, S. C.; Ngiam, J. S. Y.; Seayad, A. M.; Tuan, D. T.; Chai, C. L. L.; Chen, A. *J. Org. Chem.* **2012**, 77, 8007; (c) Suresh Kumar, A.; Thulasiram, B.; Bala Laxmi, S.; Rawat, V. S.; Sreedhar, B. *Tetrahedron* **2014**, 70, 6059.
- [18] Ghosh, S. C.; Ngiam, J. S. Y.; Chai, C. L. L.; Seayad, A. M.; Dang, T. T.; Chen, A. *Adv. Synth. Catal.* **2012**, 354, 1407.
- [19] Zhang, M.; Wu, X.-F. *Tetrahedron Lett.* **2013**, 54, 1059.
- [20] (a) Jia, F.; Li, Z. *Org. Chem. Front.* **2014**, 1, 194; (b) Li, Y.; Jia, F.; Li, Z. *Chem. Eur. J.* **2013**, 19, 82; (c) Li, Y.; Ma, L.; Jia, F.; Li, Z. *J. Org. Chem.* **2013**, 78, 5638; (d) Li, Y.; Ma, L.; Li, Z. *Chin. J. Org. Chem.* **2013**, 33, 704. (李远明, 马丽娜, 李志平, 有机化学, **2013**, 33, 704.); (e) Ma, L.; Li, Y.; Li, Z. *Sci. China Chem.* **2015**, 58, 1310.
- [21] (a) Leung, C. H.; Voutchkova, A. M.; Crabtree, R. H.; Balcells, D.; Eisenstein, O. *Green Chem.* **2007**, 9, 976; (b) Gao, J.; Wang, G.-W. *J. Org. Chem.* **2008**, 73, 2955.

(Cheng, F.)