可见光促进的双键三氟甲基化反应研究

苟宝权 杨超* 张磊 夏吾炯*

(哈尔滨工业大学化工与化学学院 城市水资源和环境国家重点实验室 哈尔滨 150080)

摘要 报道了一种可见光促进的双键直接三氟甲基化的方法。研究中利用 $\text{Ru}(\text{bpy})_3\text{Cl}_2$ 为催化剂, Togni's 试剂为三氟甲基来源, 在可见光促进下实现了含羰基烯烃的三氟甲基化。该反应具有方法简便, 条件温和, 可以获得较高的反应收率, 同时避免了有毒试剂的使用。

关键词 可见光; 烯烃; 三氟甲基化; 光催化剂; 操作简便

Visible-Light Induced Trifluoromethylation of Internal Olefinic C—H Bonds through Photoredox Catalysis

Gou, Baoquan Yang, Chao* Zhang, Lei Xia, Wujiong*

(School of Chemistry and Chemical Engineering, State Key Lab of Urban Water Resource and Environment, Harbin Institute of Technology, Harbin 150080)

Abstract An approach for direct trifluoromethylation of internal olefins of α -oxoketene dithioacetals has been achieved by using $\text{Ru}(\text{bpy})_3\text{Cl}_2$ as photocatalyst and Togni's reagent as trifluoromethylating agent under irradiation with visible light. Under a nitrogen atmosphere, a mixture of α -oxoketene dithioacetal (0.1 mmol), Togni's reagent (0.15 mmol), $\text{Ru}(\text{bpy})_3\text{Cl}_2$ (0.005 mmol), and Na_2CO_3 (0.3 mmol) in DMSO (1 mL) was stirred at room temperature for 72 h under 5 W Blue LEDs, which led to the trifluoromethylated products in 40%~90% yield. This protocol provides an efficient and easy access to prepare trifluoromethylated dithioalkyl α -oxoketene acetals under mild conditions, and is highlighted by its operational simplicity and avoiding using toxic reagent. Furthermore, the gram-scale reaction of **1a** suggested the potential application of this protocol in organic synthesis.

Keywords visible light; olefin; trifluoromethylation; photocatalyst; operational simplicity

1 Introduction

The trifluoromethylation of organic molecules is of significant importance in the field of pharmaceuticals, agrochemicals, and advanced materials^[1] owing to the incorporation of CF_3 moiety into compounds resulted in enhanced properties such as lipophilicity, metabolic stability, bioavailability, and binding selectivity.^[2] Among which, for example, Panomifene^[3] exhibits antie-strogenic and tumorinhibiting activities superior to those of tamoxifen when used for the clinical treatment of breast cancer. Therefore a variety of methods have been reported to form C— CF_3 bonds,^[4] including forming C— CF_3 bonds on C=O,^[5] C=N,^[6] or C=C,^[7,8] with cross-coupling the trifluoromethyl reagents and prefunctionalized aromatics,^[9] or other methods.^[10] The trifluoromethylated olefins are generally derived from the reaction of prefunctionalized olefins with trifluoromethylating reagents,^[11] as well as adding trifluoromethylating reagents to terminal alkynes,^[12] or from other indirect routes.^[13] Owing to its low reactivity, direct olefinic C—H trifluoromethylation

still remains as an interesting topic and a few methods were developed up to now by transition-metal-catalyzed^[14] and metal free^[15] means, wherein quinones,^[16a,16b] uracil,^[16c] and terminal olefins^[16d~16h] were utilized as starting materials.^[17,18] Hence, introducing new protocols for trifluoromethylation of olefinic C—H bonds are still demanded and continue to draw the attention of the synthetic community.

Literatures were documented that α -oxoketene dithioacetals, strongly electron-donating moieties and easily cleaved, possess unique features when serving as versatile reagents^[19] in the synthesis of heterocycles^[20] and aromatic compounds,^[21] as well as odorless thiol equivalent compounds.^[22] Therefore its preparation and transformation has attracted particular interests from the realm of organic synthesis and among which the trifluoromethylation reaction has been realized by metal-free or Cu^{II} -catalyzed conditions.^[7,8,23]

Recently, visible-light-mediated radical functionalization of olefins to introduce the trifluoromethyl group has been shown to be a feasible and efficient method.^[16g,16h]

* E-mail: xyang@hit.edu.cn (C. Y.); xiaoj@hit.edu.cn (W. X.)

Received July 15, 2016; published August 17, 2016.

Supporting information for this article is available free of charge via the Internet at <http://sioc-journal.cn>.

Project supported by the National Natural Science Foundation of China (Nos. 21272047, 21372055 and 21472030), State Key Laboratory of Urban Water Resource and Environment (No. 2015DX01) and the Fundamental Research Funds for the Central Universities (No. HIT.BRETIV.201310).

项目受国家自然科学基金(Nos. 21272047, 21372055 and 21472030)、城市水资源与水环境国家重点实验室(No. 2015DX01)和中央高校基础研究(No. HIT.BRETIV.201310)资助。

Yet trifluoromethylation of the internal olefin remains an elusive goal. As α -oxoketene dithioacetals are polar functionalized alkenes, we therefore envisioned that internal olefins can be tuned to become highly polarized by attaching to such strongly electron-donating group, and those modified internal olefins can be utilized as backbones to prepare multi-substituted trifluoromethylated olefins under photocatalytic conditions. Herein, we report a conceptually new approach to the trifluoromethylation of α -oxoketene dithioacetals via visible light photoredox catalysis (Scheme 1).

Scheme 1 Trifluoromethylation of internal olefinic C—H bonds

2 Results and Discussion

We commenced our investigation with α -oxoketene dithioacetal **1a** as a model substrate. Treating **1a** with Togni's reagent **4a**, *fac*-[Ir(ppy)₃] and K₂CO₃ in DMSO, the trifluoromethylated α -oxoketene dithioacetal was

Table 1 Screening of reaction conditions^a

Entry	PC (mol%)	Base	4a/equiv.	Solvent	Yield ^b /%
1	5a (5.0)	Na ₂ CO ₃	3	DMSO	70 ^c
2	5a (5.0)	Na ₂ CO ₃	3	DMSO	0 ^d
3	5b (5.0)	Na ₂ CO ₃	3	DMSO	78
4	5c (5.0)	Na ₂ CO ₃	3	DMSO	87 ^c ,92
5	5c (5.0)	Na ₂ CO ₃	3	DMSO	58 ^e
6	5c (5.0)	Na ₂ CO ₃	2	DMSO	91
7	5c (5.0)	Na₂CO₃	1.5	DMSO	87^c,92
8	5c (5.0)	Na ₂ CO ₃	1.5	CH ₃ OH	58
9	5c (5.0)	Na ₂ CO ₃	1.5	DMF	62
10	5c (5.0)	Na ₂ CO ₃	1.5	CH ₂ Cl ₂	41
11	5c (5.0)	Na ₂ CO ₃	1.5	CH ₃ CN	trace
12	5c (5.0)	Na ₂ CO ₃	1.5	toluene	49
13	5c (5.0)	—	1.5	DMSO	90
14	5c (5.0)	KH ₂ PO ₄	1.5	DMSO	72
15	5c (5.0)	KF	1.5	DMSO	90
16	5c (5.0)	DBU	1.5	DMSO	76
17	5c (1.0)	Na ₂ CO ₃	1.5	DMSO	70
18	5c (2.0)	Na ₂ CO ₃	1.5	DMSO	81

^a Conditions: **1a** (0.1 mmol), N₂ atmosphere. ^b The yield was determined by gas chromatography. ^c Isolated Yield. ^d No light. ^e Under air.

obtained in good yield within 39 h under 5 W Blue LEDs at room temperature (Table 1, Entry 1). With the promising initial result, we continued to optimize the reaction conditions. First, different photocatalysts were evaluated (Table 1, Entries 3 and 4). We were pleased to find that Ru(bpy)₃Cl₂ was more effective to the reaction. Next, we examined the effect of other parameters on the reaction, including solvent, base, the stoichiometry of Togni's reagent and atmosphere (Table 1, Entries 5 ~ 18). In addition, control experiment showed that the visible light was required for the transformation (Table 1, Entry 2).

With the optimized conditions in hand, a variety of α -oxoketene dithioacetals were prepared and then subjected to the reaction conditions. The results were listed in Scheme 2. Substrates bearing different groups, including methoxy, chloro, bromo, and fluoro groups at *para*, *ortho* or *meta* position of aromatic ring, were tolerant on the reaction conditions to provide corresponding products

^a Conditions: **1** (0.1 mmol), Togni's reagent (0.15 mmol), Ru(bpy)₃Cl₂ (5% mmol), Na₂CO₃ (3 mmol), in DMSO, N₂ atmosphere. ^b Isolated yield.

Scheme 2 Trifluoromethylation of α -oxoketene dithioacetals^{a,b}

(Scheme 2, **2a**~**2n**). When the aromatic ring connecting to carbonyl group was changed to methyl, the substrate was still suitable for the reaction, giving the desired ketene dithioacetals **2o** in 58% yield. By changing the dithioalkyl moiety to $-\text{S}(\text{CH}_2)_3\text{S}-$, we noticed that the corresponding substrate exhibited lower reactivity to form the target product **2s** (40%). And to our surprise, the acyclic dimethylthio analogue of **1** reacted smoothly with Togni's reagent to afford product **2r** in 90% yield.

On comparison with previous documents,^[8,23] visible-light photoredox catalytic reactions showed more extensive substrate scope, better efficiency and mild reaction condition. Moreover, the trifluoromethylation of 2-thienoyl and furoyl substrates could undergo successfully to afford both mono- and di-trifluoromethylation products (Scheme 2, **2p**, **2p'**, and **2q**, **2q'**) under standard conditions. Based on that, we anticipated controlling the scale of Togni's reagent should produce different ratios of mono- and di-trifluoromethylation products. As showed in Table 2, when the Togni's reagent increased to 3 equiv., the di-trifluoromethylation products could be obtained as the single or major products (Table 2, Entries 3 and 6).

Table 2 mono- and di-trifluoromethylation of the heterocyclic substrates^{a,b}

Entry	R	Togni's reagent/equiv.	2 Yield ^b /%	2' Yield ^b /%
1	O	1.0	43	13
2	O	1.5	27	29
3	O	3.0	0	72
4	S	1.0	75	10
5	S	1.5	58	29
6	S	3.0	40	44

^a Conditions: **1** (0.1 mmol), Ru(bpy)₃Cl₂ (5% mmol), Na₂CO₃ (3 mmol), in DMSO, N₂ atmosphere. ^b Isolated yield.

In addition, the scope was extended to the bromination reaction by changing the Togni's reagent to CBr₄. The reaction proceeded smoothly and the brominated α -oxoketene dithioacetals was obtained in 79% isolated yield (Scheme 3).

Scheme 3 Bromination of **1a**

A plausible reaction pathway for the trifluoromethylation of α -oxoketene dithioacetals was

proposed as shown in Scheme 4. Once activated by visible light, Ru(bpy)₃²⁺ reached to the excited state Ru(bpy)₃^{2+*} which then would readily be oxidized by Togni's reagent **4a** to produce Ru(bpy)₃³⁺, while CF₃ radical was released simultaneously. In succession, addition of CF₃ radical to substrate **1** delivered the trifluoromethylated secondary carbon radical intermediate **A**. Afterwards, the single-electron oxidation of **A** occurred to form the carbon center cation intermediate **B**, which then produce the desired product **2** by deprotonation.

Scheme 4 Proposed possible mechanism for the trifluoromethylation of α -oxoketene dithioacetals

During our investigation, to test the potential application of this protocol, a gram-scale reaction of **1a** was carried out under standard reaction conditions and the desired product was obtained in 82% yield (Eq. 1).

3 Conclusion

In conclusion, we have developed a visible-light-induced trifluoromethylation of α -oxoketene dithioacetals under mild conditions. This approach features not only in exhibiting high efficiency but also in expanding the substrate scope that is difficult to access with known methods.

References

- [1] Kirsch, P. *Modern Fluoro-Organic Chemistry*, Wiley-VCH, Weinheim, **2004**.
- [2] (a) Müller, K.; Faeh, C.; Diederich, F. *Science* **2007**, *317*, 1881. (b) O'Hagan, D. *Chem. Soc. Rev.* **2008**, *37*, 308. (c) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. *Chem. Soc. Rev.* **2008**, *37*, 320. (d) Furuya, T.; Kamlet, A. S.; Ritter, T. *Nature* **2011**, *473*, 470.
- [3] Liu, Y. X.; Shimizu, M.; Hiyama, T. *Angew. Chem., Int. Ed.* **2004**, *43*, 879.

- [4] (a) Liang, T.; Neumann, C. N.; Ritter, T. *Angew. Chem., Int. Ed.* **2013**, *52*, 8214. (b) Charpentier, J.; Früh, N.; Togni, A. *Chem. Rev.* **2015**, *115*, 650. (c) Alonso, C.; Marigorta, E. M.; Rubiales, G.; Palacios, F. *Chem. Rev.* **2015**, *115*, 1847. (d) Zeng, T.; Xuan, J.; Chen, J.; Lu, L.; Xiao, W. *Imag. Sci. Photochem.* **2014**, *32*, 415. (e) Peng, R.; Yang, B.; Chen, Z.; Xu, L.; Wang, S. *Chin. J. Org. Chem.* **2014**, *34*, 980. (彭蕊, 杨彬森, 陈志敏, 徐力, 王少华, 有机化学, **2014**, *34*, 980.) (f) Wang, G.; He, X.; Dai, J.; Xu, H. *Chin. J. Org. Chem.* **2014**, *34*, 837. (王光祖, 赫侠平, 戴建军, 许华建, 有机化学, **2014**, *34*, 837.) (g) Zhang, Q.; Jin, C.; Zhang, Y. *Chin. J. Org. Chem.* **2014**, *34*, 662. (张霁, 金传飞, 张英俊, 有机化学, **2014**, *34*, 662.) (h) He, Z.; Huang, Y.; Francis, V. *Acta Chim. Sinica* **2013**, *71*, 700. (何展荣, 黄毅勇, Francis, V. 化学学报, **2013**, *71*, 700.)
- [5] Sanhueza, I. A.; Bonney, K. J.; Nielsen, M. C.; Schoenebeck, F. *J. Org. Chem.* **2013**, *78*, 7749.
- [6] Zhang, Y. Q.; Liu, J. D.; Xu, H. *Org. Biomol. Chem.* **2013**, *11*, 6242.
- [7] For selected recent reports on addition of CF₃ reagents to alkenes, see: (a) Zhu, R.; Buchwald, S. L. *J. Am. Chem. Soc.* **2012**, *134*, 12462. (b) Egami, H.; Shimizu, R.; Kawamura, S.; Sodeoka, M. *Angew. Chem., Int. Ed.* **2013**, *52*, 4000. (c) Deb, A.; Manna, S.; Modak, A.; Patra, T.; Maity, S.; Maiti, D. *Angew. Chem. Int. Ed.* **2013**, *52*, 9747. (d) Wang, X.; Ye, Y. X.; Zhang, S. N.; Feng, J. J.; Xu, Y.; Zhang, Y.; Wang, J. B. *J. Am. Chem. Soc.* **2011**, *133*, 16410.
- [8] Mao, Z. F.; Huang, F.; Yu, H. F.; Chen, J. P.; Yu, Z. K.; Xu, Z. Q. *Chem. Eur. J.* **2014**, *20*, 3439.
- [9] For selected recent reports on the trifluoromethylation of aromatics, see: (a) Ye, Y. D.; Sanford, M. S. *J. Am. Chem. Soc.* **2012**, *134*, 9034. (b) Zeng, Y. W.; Zhang, L. J.; Zhao, Y. C.; Ni, C. F.; Zhao, J. W.; Hu, J. B. *J. Am. Chem. Soc.* **2013**, *135*, 2955. (c) Wang, X.; Xu, Y.; Mo, F. Y.; Ji, G.; Qiu, D.; Feng, J. J.; Ye, Y. X.; Zhang, S. N.; Zhang, Y.; Wang, J. B. *J. Am. Chem. Soc.* **2013**, *135*, 10330. (d) Hafner, A.; Bräse, S. *Angew. Chem., Int. Ed.* **2012**, *51*, 3713.
- [10] For selected recent examples, see: (a) Hu, M. Y.; Ni, C. F.; Hu, J. B. *J. Am. Chem. Soc.* **2012**, *134*, 15257. (b) Novk, P.; Lishchynskyi, A.; Grushin, V. V. *J. Am. Chem. Soc.* **2012**, *134*, 16167. (c) Shimizu, R.; Egami, H.; Hamashima, Y.; Sodeoka, M. *Angew. Chem., Int. Ed.* **2012**, *51*, 4577. (d) Niedermann, K.; Früh, N.; Senn, R.; Czarniecki, B.; Verel, R.; Togni, A. *Angew. Chem., Int. Ed.* **2012**, *51*, 6511. (e) Li, Z.; Gevorgyan, V. *Angew. Chem., Int. Ed.* **2012**, *51*, 1225.
- [11] For Selected recent reports, see: (a) He, Z.; Luo, T.; Hu, M. Y.; Cao, Y. J.; Hu, J. B. *Angew. Chem., Int. Ed.* **2012**, *51*, 3944. (b) Li, Y.; Wu, L. P.; Neumann, H.; Beller, M. *Chem. Commun.* **2013**, *49*, 2628.
- [12] Janson, P. G.; Ghoneim, I.; Ilchenko, N. O.; Szab, K. J. *Org. Lett.* **2012**, *14*, 2882.
- [13] Omote, M.; Tanaka, M.; Ikeda, A.; Nomura, S.; Tarui, A.; Sato, K.; Ando, A. *Org. Lett.* **2012**, *14*, 2286.
- [14] For selected recent examples of transition-metal-catalyzed (hetero)arene C-H Trifluoromethylation, see: (a) Zhang, X.-G.; Dai, H.-X.; Wasa, M.; Yu, J.-Q. *J. Am. Chem. Soc.* **2012**, *134*, 11948. (b) Seo, S.; Taylor, J. B.; Greaney, M. F. *Chem. Commun.* **2013**, *49*, 6385. (c) Chu, L. L.; Qing, F.-L. *J. Am. Chem. Soc.* **2012**, *134*, 1298. (d) Mejia, E.; Togni, A. *ACS Catal.* **2012**, *2*, 521. (e) Ji, Y. L.; Brueckl, T.; Baxter, R. D.; Fujiwara, Y.; Seiple, I. B.; Su, S.; Blackmond, D. G.; Baran, P. S. *Proc. Natl. Acad. Sci. USA* **2011**, *108*, 14411. (f) Ye, Y. D.; Lee, S. H.; Sanford, M. S. *Org. Lett.* **2011**, *13*, 5464. (g) Mu, X.; Chen, S. J.; Zhen, X. L.; Liu, G. S. *Chem. Eur. J.* **2011**, *17*, 6039. (h) Wang, X. S.; Truesdale, L.; Yu, J.-Q. *J. Am. Chem. Soc.* **2010**, *132*, 3648.
- [15] (a) Yang, Y. D.; Iwamoto, K.; Tokunaga, E.; Shibata, N. *Chem. Commun.* **2013**, *49*, 5510. (b) Wu, X. Y.; Chu, L. L.; Qing, F.-L. *Tetrahedron Lett.* **2013**, *54*, 249.
- [16] (a) Ilchenko, N. O.; Janson, P. G.; Szab, K. J. *Chem. Commun.* **2013**, *49*, 6614. (b) Wang, X.; Ye, Y. X.; Ji, G. J.; Xu, Y.; Zhang, S. N.; Feng, J. J.; Zhang, Y.; Wang, J. B. *Org. Lett.* **2013**, *15*, 3730. (c) Uraguchi, D.; Yamamoto, K.; Ohtsuka, Y.; Tokuhisa, K.; Yamakawa, T. *Appl. Catal. A* **2008**, *342*, 137. (d) Feng, C.; Loh, T.-P. *Chem. Sci.* **2012**, *3*, 3458. (e) Feng, C.; Loh, T.-P. *Angew. Chem., Int. Ed.* **2013**, *52*, 12414. (f) Besset, T.; Cahard, D.; Pannecoucke, X. *J. Org. Chem.* **2014**, *79*, 413. (g) Jiang, H.; Huang, C. M.; Guo, J. J.; Zeng, C. Q.; Zhang, Y.; Yu, S. Y. *Chem. Eur. J.* **2012**, *18*, 15158. (h) Iqbal, N.; Choi, S.; Kim, E.; Cho, E. J. *J. Org. Chem.* **2012**, *77*, 11383. (i) Yasu, Y.; Koike, T.; Akita, M. *Angew. Chem., Int. Ed.* **2012**, *51*, 9567.
- [17] (a) Yu, H. F.; Yu, Z. K. *Angew. Chem., Int. Ed.* **2009**, *48*, 2929. (b) Yu, H. F.; Jin, W. W.; Sun, C. L.; Chen, J. P.; Du, W. M.; He, S. B.; Yu, Z. K. *Angew. Chem., Int. Ed.* **2010**, *49*, 5792. (c) Pan, L.; Bi, X. H.; Liu, Q. *Chem. Soc. Rev.* **2013**, *42*, 1251.
- [18] Wang, L. D.; He, W.; Yu, Z. K. *Chem. Soc. Rev.* **2013**, *42*, 599.
- [19] Junjappa, H.; Ila, H.; Asokan, C. V. *Tetrahedron* **1990**, *46*, 5423.
- [20] Selected recent reports: (a) Fu, Z. Q.; Wang, M.; Ma, Y. H.; Liu, Q.; Liu, J. *J. Org. Chem.* **2008**, *73*, 7625. (b) Misra, N. C.; Panda, K.; Ila, H.; Junjappa, H. *J. Org. Chem.* **2007**, *72*, 1246. (c) Rao, H. S. P.; Sivakumar, S. *J. Org. Chem.* **2006**, *71*, 8715.
- [21] Selected recent reports: (a) Hu, J. L.; Zhang, Q.; Yuan, H. J.; Liu, Q. *J. Org. Chem.* **2008**, *73*, 2442. (b) Bi, X. H.; Dong, D. W.; Liu, Q.; Pan, W.; Zhao, L.; Li, B. *J. Am. Chem. Soc.* **2005**, *127*, 4578. (c) Sundaram, G. S. M.; Venkatesh, C.; Kumar, U. K. S.; Ila, H.; Junjappa, H. *J. Org. Chem.* **2004**, *69*, 5760.
- [22] Selected recent reports: (a) Yu, H. F.; Dong, D. W.; Ouyang, Y.; Wang, Y.; Liu, Q. *Synlett* **2007**, 151. (b) Dong, D. W.; Ouyang, Y.; Yu, H. F.; Liu, Q.; Liu, J.; Wang, M.; Zhu, J. *J. Org. Chem.* **2005**, *70*, 4535.
- [23] Xu, C.; Liu, J. X.; Ming, W. B.; Liu, Y. J.; Liu, J.; Wang, M.; Liu, Q. *Chem. Eur. J.* **2013**, *19*, 9104.

(Zhao, X.)