室温下叔丁醇钾促进烷基芳基硫醚的合成与机理研究

郭芳杰 何雨轩 王景芸 孙京* 周明东*

(辽宁石油化工大学化学与材料科学学院 抚顺 113001)

摘要 研究了在无金属参与条件下芳基溴化物与烷基硫醇的 C—S 交叉偶联反应。研究表明在室温、KO^tBu 强碱作用下, 各种缺电子的芳基溴化物与烷基硫醇均可以顺利地发生 C—S 交叉偶联反应, 高产率地得到了一系列烷基芳基硫醚类产物。本研究同时对非金属参与的偶联反应机理进行了探讨。

关键词 C—S 交叉偶联; 芳基溴化物; 机理; 烷基芳基硫醚

Potassium *tert*-Butoxide Promoted Formation of Alkyl Aryl Thioethers at Room Temperature: Synthesis and Mechanism

Guo, Fangjie He, Yuxuan Wang, Jingyun Sun, Jing* Zhou, Mingdong*

(School of Chemistry and Materials Science, Liaoning Shihua University, Fushun 113001)

Abstract The C—S cross-coupling of aryl halides with alkyl thiols under transition metal-free conditions was investigated. Good to excellent yields can be obtained for a variety of electron-poor aryl halides with alkyl thiols in the presence of KO^tBu even at room temperature. The mechanisms for transition metal-free coupling reactions are discussed.

Keywords C—S cross-coupling; aryl halides; mechanism; alkyl aryl thioethers

Aryl sulfides are important synthetic intermediates which can be used in biological or pharmaceutical industry.^[1] Catalytic cross-coupling of aryl halides with thiols proved to be one of very effective method for the construction of C—S bond, wherein palladium^[2] and copper^[3] salts or their complexes have been commonly used as effective catalysts under strong basic conditions. It is generally believed that the transition metal catalyzed coupling of aryl halides with thiols firstly undergoes the oxidative addition of M(0) with ArX, then the ligand exchanges of Ar—M—X with RSH, and finally the reductive elimination of Ar—M—SR to form the desired Ar—SR coupling product.^[4] Interestingly, it has been found that the reactivity for the coupling of electron rich or poor aryl halides are quite different. In the presence of a strong base, the coupling of electron-poor aryl halides can be facilitated even in the absence of a transition metal catalyst under a comparatively high temperature (> 80 °C).^[5] For these electron-poor aryl halides, the reaction is assumed to proceed via S_NAr mechanism. Despite significant efforts have been made in this transformation, it is

surprising that so far most reported processes are mainly focused on the coupling of aryl halides and aryl thiols.^[6] However, the coupling of aryl halides with alkyl thiols is unfortunately still not well-established, even though alkyl aryl thioethers are also important synthetic intermediates. Based on this consideration, we were interested in investigating the C—S cross-coupling of aryl halides with alkyl thiols under transition metal-free conditions. Surprisingly, it was found that the reaction could be smoothly proceeded even at room temperature. It is known that the S_NAr reaction normally proceeds at high temperatures. Such a phenomenon motivated us to make further studies on the mechanism of this reaction. Therefore, we wish to report our recent findings on this work herein.

1 Result and discussion

The C—S cross-coupling of electron-poor 4-bromoacetophenone (**1a**) and butanethiol (**2a**) was initially explored to examine the reactivity. Different solvents were

* Corresponding authors. E-mail: sunjing@lnpu.edu.cn, mingdong.zhou@lnpu.edu.cn

Received December 2, 2016; revised January 14, 2017; published online February 17, 2017.

Project supported by the National Natural Science Foundation of China (No. 21101085), the Natural Science Foundation of Liaoning Province (No. 2015020196), the Fushun Science & Technology Program (No. FSKJHT 201423), the Liaoning Excellent Talents Program in University (No. LJQ2012031), and the Talent Scientific Research Fund of Liaoning Shihua University (No. 2016XJJ-006).

国家自然科学基金(No. 21101085)、辽宁省自然科学基金(No. 2015020196)、抚顺市科技厅(No. FSKJHT 201423)、辽宁省高等学校优秀人才支持计划(No. LJQ2012031)、辽宁石油化工大学引进人才科研启动基金(No. 2016XJJ-006)资助项目。

firstly applied using 1.5 equiv. of KO^tBu at 100 °C under nitrogen protection. The reaction was not able to proceed in toluene, but moderate yield was achieved in tetrahydrofuran (THF) (Table 1, Entries 1 and 2). To our delight, excellent yield could be obtained when carrying out the reaction in pyridine, 1,2-dimethoxyethane (DME), dimethyl sulfoxide (DMSO) and *N,N*-dimethylformamide (DMF) (Entries 3~6). To our surprise, further studies indicated that excellent yields of **3aa** could also be maintained when decreasing the temperature from 100 °C to room temperature (Entries 6~9). To further determine the reactivity, various bases were studied at room temperature. The results indicated that the replacement of KO^tBu with NaO^tBu, KOH or NaOH also afforded excellent product yields at room temperature (Entries 10~12), while K₂CO₃ led to a loweryield (Entry 13) and Na₂CO₃ proved to be not quite effective (Entry 14). Decreasing of the amount of KO^tBu to 1.0 equiv., the product yield also decreased (Entry 15).

Table 1 C—S cross-coupling of 4-bromoacetophenone and butanethiol^a

Entry	Solvent	Temperature/°C	Base (equiv.)	Yield ^b /%
1	Toluene	100	KO ^t Bu (1.5)	0
2	THF	100	KO ^t Bu (1.5)	52
3	Pyridine	100	KO ^t Bu (1.5)	>99
4	DME	100	KO ^t Bu (1.5)	>99
5	DMSO	100	KO ^t Bu (1.5)	>99
6	DMF	100	KO ^t Bu (1.5)	>99
7	DMF	80	KO ^t Bu (1.5)	>99
8	DMF	50	KO ^t Bu (1.5)	>99
9	DMF	r.t.	KO ^t Bu (1.5)	>99 (95 ^c)
10	DMF	r.t.	NaO ^t Bu (1.5)	>99
11	DMF	r.t.	NaOH (1.5)	>99
12	DMF	r.t.	KOH (1.5)	>99
13	DMF	r.t.	K ₂ CO ₃ (1.5)	84
14	DMF	r.t.	Na ₂ CO ₃ (1.5)	8
15	DMF	r.t.	KO ^t Bu (1.0)	83
16	DMF	r.t.	KO ^t Bu (2.0)	>99

^a Reaction conditions: 4-bromoacetophenone (1 mmol), butanethiol (1.1 mmol), base in 3 mL of solvent for 1 h under N₂ protection; ^b GC yields; ^c Isolated yields.

Based on the above studies, a condition **A** (1 mmol of 4-bromoacetophenone, 1.1 mmol of butanethiol, 1.5 equiv. of KO^tBu in 3 mL of DMF at room temperature for 1 h under N₂ protection) was applied as a standard condition for further study on the substrate scope of this reaction (Table 2). It turned out that reactions between butanethiol and all examined electron-poor aryl bromides can be facilitated very well at room temperature, leading to the corresponding cross-coupling products in good yields. Excellent yields can be obtained for the coupling of 4-bromoacetophenone with heptanthiol, 2-methyl-2-propanethiol or cyclohexylthiol (**3aa**~**3ad**). A comparatively lower yield can be achieved

when using bulky alkylthiols such as 3-methyl-2-butanethiol (**3ae**). For the sake of comparison, the coupling of 4-bromoacetophenone with aryl thiol such as 4-methylbenzenethiol and 4-chlorobenzenethiol was also conducted, and no obvious difference on activity was observed as compared to alkyl thiols (**3af**, **3ag**). Additionally, excellent yields were also obtained for the coupling of various CF₃ or NO₂ substituted aryl bromide with butanethiol (**3ba**~**3fa**). However, this reaction did not occur when the substrates bearing with F, H or Me group (**3ga**~**3ia**). Obviously, the electronic effect of substituted aryl bromides had a significant impact on this transformation.

Table 2 C—S cross-coupling of aryl bromides with thiols^{a,b}

3aa , 95%	3ab , 94%	
3ac , 84%	3ad , 78%	3ae , 69%
3af , 92%	3ag , 91%	
3ba , 93%	3ca , 95%	3da , 95%
3ea , 91%	3fa , 92%	3ga , 0%
3ha , 0%	3ia , 0%	

^a Reaction conditions: aryl halide (1 mmol), thiol (1.1 mmol), KO^tBu (1.5 equiv.) in 3 mL of DMF at room temperature for 1 h under nitrogen atmosphere; ^b Isolated yields.

2 Mechanistic considerations

As mentioned previously, the S_NAr reaction normally occurs at high temperatures. In our case, the coupling reaction could be smoothly facilitated at room temperature.^[5] Additionally, Liu *et al.*^[5d] also found that the coupling of

1-chloro-4-nitrobenzene with 1-octanethiol in the presence of 3 equiv. of KOH in PEG-600 (solvent) could proceed at room temperature, in which 90% of octyl *p*-nitrobenzyl thioether was obtained. Nevertheless, the mechanism was not discussed in this work. The main concern is whether C—S coupling of electron-poor aryl halides with thiols can proceed via a S_NAr mechanism at room temperature. Therefore, the mechanism needs to be carefully examined. First, if the reaction proceeds via an aryne mechanism, two isomers (*para* and *meta*-substituted thioethers) should be observed in a roughly 1 : 1 molar ratio when using *para*-substituted aryl halides as substrates.^[5b,7] However, only one isomer could be detected for all examined electron-poor aryl bromides. Therefore, the aryne mechanism can be ruled out. A mechanism proceeding via a benzene radical intermediate induced by the KO^tBu/DMF super-base system should not be ignored. KO^tBu or NaO^tBu has been found to be able to promote the direct C—H arylation of unactivated benzene with aryl halides, which has been independently described by the group of Shi, Kong/Lei, and Shirakawa/Hayashi.^[8–10] Moreover, Yan *et al.*^[11] have also reported a series of KO^tBu/DMF promoted radical coupling reactions. To further explore the mechanism, radical trapping experiments were carried out using 1~3 equiv. of tetramethylpiperidine *N*-oxide (TEMPO) or 2,6-di-*tert*-butyl-4-methylphenol (BHT) as the radical scavengers in the presence of 1.5 equiv. of KO^tBu in DMF at room temperature. However, the reaction could not be significantly hindered by both scavengers. Thus, a benzene radical based mechanism appears unlikely. Therefore, the C—S coupling using electron-poor aryl halides might proceed via a S_NAr mechanism, even at room temperature, which is in accordance to the reported mechanism.^[5b]

3 Conclusions

C—S cross-coupling of aryl bromides with alkyl thiols under transition metal-free conditions is described. The examined coupling of electron-poor aryl halides with alkylthiols proved to be highly effective at room temperature under transition metal-free conditions. The reaction is applicable for aryl bromides bearing various electron-withdrawing groups. The reaction is also applicable for various linear, bulky and cyclic alkyl thiols. Reaction mechanism studies indicate that the reaction likely undergoes a S_NAr mechanism, even at room temperature. This reaction protocol provides an efficient and cost-effective method for the synthesis of alkyl aryl thioethers under mild conditions.

4 Experimental

4.1 General method

All reactions were carried out under a nitrogen atmosphere with standard Schlenk techniques. All solvents were purified and dried according to standard methods prior to use. The GC-MS spectra were determined by a Thermo Fisher ISQ GC-MS; NMR spectra were recorded on a Bruker Ascend HD 500 spectrometer; IR spectra were rec-

orded on a Nicolet 6700 FT-IR spectrometer; The melting points of compounds were obtained on a Shang Hai Shen Guang WRS-3 melting point instrument; HRMS (ESI) determinations were carried out on a Bruker Daltonics MicrOTOF II spectrometer.

4.2 General procedure for the coupling reaction of aryl bromides with thiols

In a typical procedure, aryl bromide (1.0 mmol), thiols (1.1 mmol) and KO^tBu (168 mg, 1.5 mmol) were added to a Schlenk tube under an atmosphere of nitrogen. DMF (3 mL) was then injected under vigorous stirring 1 h. After the reaction, the mixture was quenched by addition of distilled water. The organic phase was then separated, and the aqueous phase was extracted with ethyl acetate for three times. The combined organic layers were dried with MgSO₄, and the sample was conducted for GC or GC-MS analysis. The sample solution was further evaporated under vacuum to remove the solvent. The obtained crude product was purified by column chromatography (eluent, petroleum) on silica gel to afford the desired thioethers.

1-(4-(Butylthio)phenyl)ethanone (**3aa**):^[12] Colorless liquid, 198 mg, 95% yield. ¹H NMR (500 MHz, CDCl₃) δ : 7.85 (d, *J*=8.5 Hz, 2H), 7.29 (d, *J*=8.5 Hz, 2H), 2.98 (t, *J*=7.0 Hz, 2H), 2.55 (s, 3H), 1.71~1.65 (m, 2H), 1.51~1.44 (m, 2H), 0.94 (t, *J*=7.0 Hz, 3H).

1-(4-(Heptylthio)phenyl)ethanone (**3ab**):^[13] Colorless liquid, 207 mg, 94% yield. ¹H NMR (500 MHz, CDCl₃) δ : 7.86 (d, *J*=8.5 Hz, 2H), 7.30 (d, *J*=9.0 Hz, 2H), 2.99 (t, *J*=7.5 Hz, 2H), 2.57 (s, 3H), 1.73~1.66 (m, 2H), 1.46~1.43 (m, 2H), 1.34~1.26 (m, 6H), 0.89 (t, *J*=7.0 Hz, 3H).

1-(4-((3-Methylbutan-2-yl)thio)phenyl)ethanone (**3ac**): Colorless liquid, 162 mg, 84% yield. ¹H NMR (500 MHz, CDCl₃) δ : 7.85 (d, *J*=8.5 Hz, 2H), 7.35 (d, *J*=8.5 Hz, 2H), 3.39~3.34 (m, 1H), 2.55 (s, 3H), 1.98~1.92 (m, 1H), 1.30 (d, *J*=7.0 Hz, 3H), 1.04~1.00 (m, 6H); ¹³C NMR (125 MHz, CDCl₃) δ : 196.9, 144.3, 134.0, 128.6, 128.1, 48.1, 32.5, 26.3, 19.9, 18.4, 16.9; IR (neat) ν : 2966, 2940, 2874, 1680, 1589, 1552, 1394, 1350, 1262, 1181, 1097, 957, 817, 762, 601 cm⁻¹; HRMS (EI) calcd for C₁₃H₁₈SO: 222.1078, found 222.1080.

1-(4-(Cyclohexylthio)phenyl)ethanone (**3ad**):^[4e,14] White solid, 161 mg, 78% yield. m.p. 67.4~67.6 °C (lit.^[14] 66~68 °C); ¹H NMR (500 MHz, CDCl₃) δ : 7.85 (d, *J*=8.5 Hz, 2H), 7.35 (d, *J*=8.5 Hz, 2H), 3.33~3.30 (m, 1H), 2.56 (s, 3H), 2.09~1.95 (m, 4H), 1.80~1.77 (m, 4H), 1.66~1.59 (m, 2H).

1-(4-(*tert*-Butylthio)phenyl)ethanone (**3ae**):^[15] Colorless liquid, 124 mg, 69% yield. ¹H NMR (500 MHz, CDCl₃) δ : 7.91 (d, *J*=8.5 Hz, 2H), 7.62 (d, *J*=8.5 Hz, 2H), 2.60 (s, 3H), 1.32 (s, 9H).

1-(4-(*p*-Tolylthio)phenyl)ethanone (**3af**):^[6b] Yellow solid, 197 mg, 92% yield. m.p. 94.1~94.2 °C (lit.^[6b] 89~92 °C); ¹H NMR (500 MHz, CDCl₃) δ : 7.83~7.79 (m, 2H), 7.42 (d, *J*=8.5 Hz, 2H), 7.23 (d, *J*=8.0 Hz, 2H), 7.16 (d, *J*=8.0 Hz, 2H), 2.54 (s, 3H), 2.39 (s, 3H).

1-(4-((4-Chlorophenyl)thio)phenyl)ethanone (**3ag**):^[16]

White solid, 212 mg, 91% yield. m.p. 56.0~56.7 °C (lit.^[16] 40~42 °C); ¹H NMR (500 MHz, CDCl₃) δ: 7.84 (d, *J*=9.0 Hz, 2H), 7.41~7.35 (m, 4H), 7.22 (d, *J*=8.5 Hz, 2H), 2.55 (s, 3H).

Butyl(2-(trifluoromethyl)phenyl)sulfane (**3ba**): Colorless liquid, 217 mg, 93% yield. ¹H NMR (500 MHz, CDCl₃) δ: 7.63 (d, *J*=7.5 Hz, 1H), 7.46~7.44 (m, 2H), 7.25~7.23 (m, 1H), 2.97 (t, *J*=7.5 Hz, 2H), 1.69~1.63 (m, 2H), 1.49~1.41 (m, 2H), 0.93 (t, *J*=9.0 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ: 137.1, 131.8, 129.9, 126.7, 125.2, 33.6, 30.8, 21.9, 13.5; IR (neat) ν: 2959, 2931, 2869, 1596, 1568, 1471, 1433, 1310, 1255, 1171, 1126, 1023, 760 cm⁻¹; HRMS (EI) calcd for C₁₁H₁₃F₃S: 234.0690, found 234.0683.

Butyl(3-(trifluoromethyl)phenyl)sulfane (**3ca**):^[17] Colorless liquid, 223 mg, 95% yield. ¹H NMR (500 MHz, CDCl₃) δ: 7.52 (s, 1H), 7.47~7.45 (m, 1H), 7.39~7.37 (m, 2H), 2.96 (t, *J*=9.5 Hz, 2H), 1.69~1.60 (m, 2H), 1.49~1.41 (m, 2H), 0.94 (t, *J*=9.5 Hz, 3H).

Butyl(4-(trifluoromethyl)phenyl)sulfane (**3da**):^[18] Colorless liquid, 223 mg, 95% yield. ¹H NMR (500 MHz, CDCl₃) δ: 7.50 (d, *J*=8.5 Hz, 2H), 7.34 (d, *J*=8.5 Hz, 2H), 2.96 (t, *J*=7.0 Hz, 2H), 1.69~1.63 (m, 2H), 1.51~1.45 (m, 2H), 0.94 (t, *J*=7.5 Hz, 3H).

Butyl(2-nitrophenyl)sulfane (**3ea**):^[12] Colorless liquid, 192 mg, 91% yield. ¹H NMR (500 MHz, CDCl₃) δ: 8.17~8.15 (m, 1H), 7.54~7.51 (m, 1H), 7.41 (d, *J*=8.0 Hz, 1H), 7.22~7.19 (m, 1H), 2.94 (t, *J*=7.5 Hz, 2H), 1.72~1.67 (m, 2H), 1.53~1.47 (m, 2H), 0.95 (t, *J*=7.5 Hz, 3H).

Butyl(4-nitrophenyl)sulfane (**3fa**):^[12] Colorless liquid, 194 mg, 92% yield. ¹H NMR (500 MHz, CDCl₃) δ: 8.11 (d, *J*=9.0 Hz, 2H), 7.31 (d, *J*=9.0 Hz, 2H), 3.01 (t, *J*=8.0 Hz, 2H), 1.73~1.66 (m, 2H), 1.53~1.47 (m, 2H), 0.96 (t, *J*=7.5 Hz, 3H).

Supporting Information ¹H NMR and ¹³C NMR spectra. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Yao, H.; Richardson, D. E. *J. Am. Chem. Soc.* **2003**, *125*, 6211.
(b) Qin, Z.; Kastrati, I.; Chandrasena, R. E. P.; Liu, H.; Yao, P.; Petukhov, P. A.; Bolton, J. L.; Thatcher, G. R. J. *J. Med. Chem.* **2007**, *50*, 2682.
(c) Bagley, M. C.; Davis, T.; Dix, M. C.; Rokicki, M.; Kipling, D. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 5107.
(d) Bryan, C. S.; Braunger, J. A.; Lautens, M. *Angew. Chem. Int. Ed.* **2009**, *121*, 7198.
(e) Beletskaya, I. P.; Ananikov, V. P. *Chem. Rev.* **2011**, *111*, 1596.
(f) Ubale, A. U.; Bhute, M. V.; Malpe, G. P.; Raut, P. P.; Chipade, K. S.; Ibrahim, S. G. *J. Sadui Chem. Soc.* **2016**, *20*, 227.
(g) Zhou, Y. B.; Gao, F.; Zhao, Y.; Lu, J. J. *J. Sadui Chem. Soc.* **2014**, *18*, 589.
(h) Zaafarany, I.; Boller, H. J. *J. Sadui Chem. Soc.* **2010**, *14*, 183.
(i) An, Y. N.; Li, J. X.; Li, M.; Li, C. S.; Yang, S. R. *Chin. J. Org. Chem.* **2017**, *37*, 720 (in Chinese).
(安艳妮, 李建晓, 李蒙, 李春生, 杨少容, 有机化学, **2017**, *37*, 720.)
- [2] (a) Byeun, A.; Baek, K.; Han, M. S.; Lee, S. *Tetrahedron Lett.* **2013**, *54*, 6712.
(b) Alvaroa, E.; Hartwig, J. F. *J. Am. Chem. Soc.* **2009**, *131*, 7858.
(c) Li, J. X.; Li, C. S.; Yang, S. R.; An, Y. N.; Wu, W. Q.; Jiang, H. *J. Org. Chem.* **2016**, *81*, 2875.
(d) Li, J. X.; Li, C. S.; Yang, S. R.; An, Y. N.; Wu, W. Q.; Jiang, H. *J. Org. Chem.* **2016**, *81*, 7771.
- [3] (a) Luo, F.; Pan, C.; Li, L. P.; Chen, F.; Cheng, J. *Chem. Commun.* **2011**, *47*, 5304.
(b) Mallick, S.; Rana, S.; Parida, K. *Dalton. Trans.* **2011**, *40*, 9169.
- [4] (a) Zhang, J.; Medley, C. M.; Krause, J. A.; Guan, H. *Organometallics* **2010**, *29*, 6393.
(b) Venkanna, G. T.; Arman, H. D.; Tonzetich, Z. J. *ACS Catal.* **2014**, *4*, 2941.
(c) Jammi, S.; Barua, P.; Rout, L.; Saha, P.; Punniyamurthy, T. *Tetrahedron Lett.* **2008**, *49*, 1484.
(d) Baldovino-Pantaleón, O.; Hernández-Ortega, S.; Morales-Morales, D. *Adv. Syn. Catal.* **2006**, *348*, 236.
(e) Guan, P.; Cao, C.; Liu, Y.; Li, Y.; He, P.; Chen, Q.; Liu, G.; Shi, Y. *Tetrahedron Lett.* **2012**, *53*, 5987.
- [5] (a) Zhang, Y. G.; Ngeow, K. C.; Ying, J. Y. *Org. Lett.* **2007**, *9*, 3495.
(b) Yuan, Y.; Thomé, I.; Kim, S. H.; Chen, D.; Beyer, A.; Bonnamour, J.; Zuidema, E.; Chang, S.; Bolm, C. *Adv. Synth. Catal.* **2010**, *352*, 2892.
(c) Cano, R.; Ramón, D. J.; Yus, M. *J. Org. Chem.* **2011**, *76*, 654.
(d) Duan, Z.; Ranjit, S.; Liu, X. *Org. Lett.* **2010**, *12*, 2430.
- [6] (a) Shi, Y.; Cai, Z.; Guan, P.; Pang, G. *Synth. Lett.* **2011**, *22*, 2090.
(b) Fu, C. F.; Liu, Y. H.; Peng, S. M.; Liu, S. T. *Tetrahedron* **2010**, *66*, 2119.
(c) Ma, D.; Geng, Q.; Zhang, H.; Jiang, Y. *Angew. Chem., Int. Ed.* **2010**, *49*, 1291.
(d) Xu, R.; Wan, J. P.; Mao, H.; Pan, Y. *J. Am. Chem. Soc.* **2010**, *132*, 15531.
(e) Wu, J. R.; Lin, C. H.; Lee, C. F. *Chem. Commun.* **2009**, *29*, 4450.
(f) Iglesias, M. J.; Prieto, A.; Nicasio, M. C. *Adv. Synth. Catal.* **2010**, *352*, 949.
- [7] (a) Bradshaw, J. S.; Chen, E. Y.; Hales, R. H.; South, J. A. *J. Org. Chem.* **1972**, *37*, 2051.
(b) March, J. *Advanced Organic Chemistry*, Vol. 4, John Wiley & Sons, Inc, New York, **1985**, p. 576.
(c) Varala, R.; Ramu, E.; Alam, M. M.; Adapa, S. R. *Chem. Lett.* **2004**, *33*, 1614.
- [8] Sun, C.; Li, H.; Yu, D.; Yu, M.; Zhou, X.; Lu, X.; Huang, K.; Zheng, S.; Li, B.; Shi, Z. *Nat. Chem.* **2010**, *2*, 044.
- [9] Liu, W.; Cao, H.; Zhang, H.; Zhang, H.; Chung, K. H.; He, C.; Wang, H.; Kong, F. Y.; Lei, A. *J. Am. Chem. Soc.* **2010**, *132*, 16737.
- [10] Shirakawa, E.; Itoh, K.; Higashino, T.; Hayashi, T. *J. Am. Chem. Soc.* **2010**, *132*, 15537.
- [11] (a) Wei, W. T.; Liu, Y.; Ye, L. M.; Lei, R. H.; Zhang, X. J.; Yan, M. *Org. Biomol. Chem.* **2015**, *13*, 817.
(b) Chen, Y. Y.; Zhang, N. N.; Ye, L. M.; Chen, J. H.; Shun, X.; Zhang, X. J.; Yan, M. *RSC Adv.* **2015**, *5*, 48046.
- [12] Qiao, Z.; Wei, J.; Jiang, X. *Org. Lett.* **2014**, *16*, 1212.
- [13] Xu, J. A.; Wei, T. Z.; Xia, J. K.; Zhang, Q. H.; Wu, H. S. *Chirality* **2004**, *16*, 341.
- [14] Zhang, H.; Cao, W. G.; Ma, D. W. *Synth. Commun.* **2007**, *37*, 25.
- [15] Martyn, J.; Soren, L. B.; Mogens, B. N. *J. Org. Chem.* **2013**, *78*, 4348.
- [16] Gorczynski, M. J.; Leal, R. M.; Mooberry, S. L.; Bushweller, J. H.; Brown, M. L. *Bioorg. Med. Chem.* **2004**, *12*, 1029.
- [17] Jean, M.; Renault, J.; Weghe, P. V. D.; Asao, N. *Tetrahedron Lett.* **2010**, *51*, 378.
- [18] Taniguchi, N.; Onami, T. *J. Org. Chem.* **2004**, *69*, 915.

(Li, L.; Lu, Z.)