I₂/CuO 催化的 3-乙酰基香豆素和硫脲一锅反应合成 3-(2-氨基-4-噻唑基)香豆素衍生物

赵 芳 胡 洋 李 巧 胡胜利*

(湖北师范大学化学化工学院 黄石 435002)

摘要 香豆素和噻唑是两类具有广泛的生物活性和药理作用的杂环化合物, 合成含噻唑基的香豆素衍生物在药物化学上具有重要的意义. 本研究发展了一种方便高效地通过 3-乙酰基香豆素和硫脲/*N*-取代硫脲在 I₂/CuO 催化作用下一锅反应合成 3-(2-氨基-4-噻唑基)香豆素衍生物的方法. 所有目标化合物通过核磁共振、红外光谱、质谱分析证实.

关键词 香豆素; 2-氨基噻唑; 一锅反应; I₂/CuO

I₂/CuO-Catalyzed One-Pot Synthesis of 3-(2-Amino-4-thiazolyl) Coumarin Derivatives from 3-Acetylcoumarins and Thiourea

Zhao, Fang Hu, Yang Li, Qiao Hu, Shengli*

(College of Chemistry and Chemical Engineering, Hubei Normal University, Huangshi 435002)

Abstract Coumarin and thiazole are two classes of important heterocycles with many biological activities and pharmacological effects. It is very meaningful to synthesize thiazolyl coumarin derivatives in medicinal chemistry area. In this study, a facile and efficient one-pot process was developed to synthesize substituted aminothiazoles from 3-acetylcoumarins and thiourea/*N*-substituted thioureas under the media of I₂/CuO. All target molecules were characterized by NMR, IR spectra and MS.

Keywords coumarin; 2-aminothiazole; one-pot reaction; I₂/CuO

1 Introduction

Coumarins play a key role in the medicinal area because of their structural diversity and pharmaceutical properties. Various synthetic coumarin derivatives constitute one of the most widely used groups of anti-human immunodeficiency virus,^[1] anti-inflam-matory,^[2,3] antioxidant^[4] and anticancer agents.^[5~7] Further, 2-aminothiazoles have been employed in the preparation of different important drugs required for the treatment of allergies, hypertension, inflammation, bacterial, and human immunodeficiency virus (HIV) infections.^[8~11] Therefore, it is very meaningful to synthesize 2-amino-4-thiazolyl coumarin derivatives in medicinal chemistry area.

Numerous methods were reported for the preparation of 2-amino-4-thiazolyl coumarin derivatives. In general, Hantzsch thiazole synthesis, involving the reaction of 3-(2-bromoacetyl) coumarin and various thioureas, is the most widely used methods.^[12~21] In recent years, much attention

has been paid to one-pot synthesis,^[22~26] which is a strategy to improve the efficiency of a chemical reaction because of avoiding a separation process and purification of the intermediates. The use of molecular iodine in organic synthesis has been known for a long time.^[27~29] Recently, Zhu and co-workers^[30] developed a I₂/CuO-catalyzed tandem cyclization strategy involving the direct use of aromatic methyl ketones and thiourea for one-pot synthesis of substituted 2-aminothiazole. Although 3-acetylcoumarins also belong to methyl ketone, to the best of our knowledge, the direct use of 3-acetylcoumarin and thiourea for one-pot synthesis of 2-amino-4-thiazolyl coumarin has not been reported. Herein, we reported a facile, efficient method for synthesis of 3-(2-amino-4-thiazolyl) coumarins from 3-acetylcoumarins and thioureas.

2 Results and discussion

To explore optimized reaction condition, 3-acetylcoumarin (**1a**) was first used as the model substrate to react

* Corresponding author. E-mail: hushengli168@126.com

Received August 23, 2019; revised October 19, 2019; published online November 7, 2019.

Project supported by the Science Technology Foundation for Creative Research Group of Hubei Department of Education (No. T201311) and the Foundation of Hubei Provincial Ministry of Education (No. D20112507).

湖北省高校优秀中青年科技创新团队(No. T201311)和湖北省教育厅(No. D20112507)资助项目.

with thiourea in different solvents for the synthesis of desired product 3-(2-aminothiazol-4-yl)coumarin (**3a**). As shown in Eq. 1 and Table 1, the product **3a** was successfully obtained in 72%~85% yields using ethanol, methanol, propanol and butanol as solvents under the media of I_2/CuO (Table 1, Entries 1~4). However, **3a** was not obtained when tertbutanol (*t*-BuOH), acetonitrile (MeCN), tetrahydrofuran (THF), toluene or dimethylformamide (DMF) was used as solvent (Entries 5~9). It was obvious that EtOH was the optimal solvent in the reaction. Further, the molar ratio of substrate, I_2 and CuO was also optimized. When the molar ratio of $n(1a)/n(2a)/n(CuO)/n(I_2)$ was varied from 1.0 : 1.0 : 1.0 : 1.0 to 1.0 : 1.1 : 1.1 (Entries 10~12), the product **3a** was afforded in 86%~88% yields in ethanol at reflux. Therefore, $n(1a) : n(2a) : n(I_2) : n(CuO) = 1.0 : 1.0 : 1.1 : 1.1$ is the optimal condition. Finally, the effect of reaction temperature on the yield of the product **3a** was also examined (Entry 13). The experiment result indicates that 78 °C is the optimal temperature.

Table 1 Effect of solvent, temperature and molar ratio of $n(1a)/n(2a)/n(CuO)/n(I_2)$ on the yield of **3a**

Entry	Solvent	<i>t</i> /°C	$n(1a) : n(2a) : n(CuO) : n(I_2)$	Yield ^a /%
1	EtOH	78	1.0 : 1.0 : 1.0 : 1.0	85
2	MeOH	65	1.0 : 1.0 : 1.0 : 1.0	82
3	PrOH	97	1.0 : 1.0 : 1.0 : 1.0	78
4	BuOH	118	1.0 : 1.0 : 1.0 : 1.0	72
5	<i>t</i> -BuOH	83	1.0 : 1.0 : 1.0 : 1.0	0 ^b
6	MeCN	82	1.0 : 1.0 : 1.0 : 1.0	0 ^b
7	THF	66	1.0 : 1.0 : 1.0 : 1.0	0 ^b
8	Toluene	111	1.0 : 1.0 : 1.0 : 1.0	0 ^b
9	DMF	90	1.0 : 1.0 : 1.0 : 1.0	0 ^b
10	EtOH	78	1.0 : 1.0 : 1.1 : 1.1	88
11	EtOH	78	1.0 : 1.0 : 1.5 : 1.5	86
12	EtOH	78	1.0 : 1.1 : 1.1 : 1.1	87
13	EtOH	65	1.0 : 1.0 : 1.1 : 1.1	75

^a Isolated yield. ^b No reaction.

After the optimization of reaction conditions, the reactions of different acetylcoumarins (**1**) with thiourea and *N*-substituted thioureas (Eq. 2) were examined to establish the generality of the present transformation. The results are listed in Table 2. The condensation of different 3-acetylcoumarins including electron-donating groups (such as Me, NEt_3) or electron-withdrawing groups (such as OMe, Cl, Br and NO_2) on their phenyl ring with thiourea gave the expected products 3-(2-amino-4-thiazolyl) coumarins **3b**~**3h** in good to excellent yields (80%~89%). Furthermore,

it was found that the reaction of *N*-substituted thioureas (such as *N*-methylthiourea and *N*-phenylthiourea) and acetylcoumarins could afford the 2-aminothiazole derivatives **3i**~**3x** in 58%~85% yields. The treatment of **1a** with thiourea, *N*-methylthiourea and *N*-phenylthiourea gave the desired products **3a**, **3i** and **3q** in 88%, 84% and 66% yields, respectively. The slightly lower yields might be attributed to the steric hindrance of *N*-methylthiourea and *N*-phenylthiourea.

Table 2 Synthesis of **3** from 3-acetylcoumarins and thiourea

Compd.	R ¹	R ²	R ³	R ⁴	R ⁵	Yield ^a /%
3a	H	H	H	H	H	88
3b	H	Me	H	H	H	82
3c	H	H	H	OMe	H	89
3d	H	Cl	H	H	H	85
3e	H	Br	H	H	H	83
3f	H	NO_2	H	H	H	81
3g	H	H	NEt_3	H	H	80
3h		Benzo	H	H	H	82
3i	H	H	H	H	Me	84
3j	H	Me	H	H	Me	76
3k	H	H	H	OMe	Me	85
3l	H	Cl	H	H	Me	82
3m	H	Br	H	H	Me	79
3n	H	NO_2	H	H	Me	74
3o	H	H	NEt_3	H	Me	75
3p		Benzo	H	H	Me	77
3q	H	H	H	H	Ph	66
3r	H	Me	H	H	Ph	63
3s	H	H	H	OMe	Ph	68
3t	H	Cl	H	H	Ph	65
3u	H	Br	H	H	Ph	62
3v	H	NO_2	H	H	Ph	58
3w	H	H	NEt_3	H	Ph	60
3x		Benzo	H	H	Ph	63

^a Isolated yield.

Based on previous report,^[30] a proposed mechanism for this reaction is given in Scheme 1. Initially, 3-acetyl-coumarins (**1**) undergoes the formation of intermediate 3-(2-iodoacetyl)coumarin in the media of I_2 and CuO, then, 3-(2-iodoacetyl)coumarins reacted with thiourea giving the corresponding 3-(2-amino-4-thiazolyl) coumarins.

Scheme 1 Proposed mechanism for the synthesis of 2-aminothiazolyl coumarins (**3**)

3 Conclusions

A facile, efficient one-pot synthesis of 3-(2-amino-4-thiazolyl) coumarins from 3-acetylcoumarins and thiourea under the media of I_2/CuO in refluxing ethanol was developed. Further application of this methodology for the synthesis of various coumarin-based heterocyclic compounds is under way in our laboratory.

4 Experimental section

4.1 Instruments and reagents

NMR spectra were measured on a VXR 300 instrument. Chemical shifts are reported relative to the internal standard of tetramethylsilane (TMS). IR spectra were obtained as KBr pellet samples on a Nicolet 5700 FTIR spectrometer. Mass spectra were measured on an LCQ Advantage MAX (ESI). The elemental analyses were measured on a Vario EL-III instrument CHNSO elemental analyzer. 3-Acetylcoumarins (**1**) were synthesized according to literature method.^[31] Other reagents were obtained from commercial suppliers and used without further purification.

4.2 General procedure for the synthesis of 3-(2-amino-4-thiazolyl) coumarin derivatives **3a**~**3x**

A mixture of 3-acetylcoumarin (**1**) (1.0 mmol), iodine (279.2 mg, 1.1 mmol), and CuO (88 mg, 1.1 mmol) in anhydrous ethanol (30 mL) was heated at reflux. After the reaction (1~5 h, monitored by TLC), thiourea (**2**) (1.0 mmol) was added and the mixture was refluxed for 1 h. After that, the solvent was removed under reduce pressure, and 50 mL of water was added to the residue. The mixture was extracted with EtOAc three times (50 mL×3). The extract was washed with 10% $Na_2S_2O_3$ and then a small amount of 5% NaOH solution, dried over anhydrous Na_2SO_4 and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel using petroleum ether/EtOAc as the eluent to afford the corresponding purified products **3**.

3-(2-Aminothiazol-4-yl)-2H-chromen-2-one (**3a**): Yel-

low solid; m.p. 224~225 °C (lit.^[12] 225~227 °C); 1H NMR (300 MHz, DMSO- d_6) δ : 8.51 (s, 1H), 7.83 (d, J =7.4 Hz, 1H), 7.61 (t, J =7.4 Hz, 1H), 7.51 (s, 1H), 7.45~7.35 (m, 2H), 7.15 (br, 2H); ^{13}C NMR (75 MHz, DMSO- d_6) δ : 167.4, 158.8, 152.2, 143.3, 138.0, 131.4, 128.8, 124.5, 120.5, 119.3, 115.7, 108.6; IR (KBr) ν_{max} : 3388, 1691, 1639, 1531, 1442, 1374, 1091, 751 cm^{-1} ; ESI-MS m/z : 244.89 $[M+H]^+$.

3-(2-Aminothiazol-4-yl)-6-methyl-2H-chromen-2-one (**3b**): Yellow solid, m.p. 235~237 °C; 1H NMR (300 MHz, $CDCl_3$) δ : 8.39 (s, 1H), 7.70 (s, 1H), 7.18~7.23 (m, 3H), 4.97 (s, 2H, NH_2), 2.34 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ : 166.2, 151.0, 143.9, 138.8, 134.2, 132.4, 127.9, 120.7, 119.3, 116.5, 116.0, 111.0, 20.8; IR (KBr) ν_{max} : 3395, 1630, 1535, 1378, 1262, 1091, 1024, 805 cm^{-1} ; ESI-MS m/z : 259.05 $[M+H]^+$. Anal. calcd for $C_{13}H_{10}N_2O_2S$: C 60.45, H 3.90, N 10.85; found C 60.32, H 3.79, N 10.71.

3-(2-Aminothiazol-4-yl)-8-methoxy-2H-chromen-2-one (**3c**): Yellow solid, m.p. 256~258 °C (lit.^[18] 256~258 °C); 1H NMR (300 MHz, $CDCl_3$) δ : 8.47 (s, 1H), 7.37~7.51 (m, 1H), 7.30~7.36 (m, 1H), 7.29~7.30 (m, 2H), 7.18 (s, 2H), 3.93 (s, 3H); ^{13}C NMR (75 MHz, $CDCl_3$) δ : 167.4, 158.5, 146.2, 143.2, 138.3, 133.7, 131.6, 124.6, 120.6, 119.9, 113.7, 108.8, 56.0; IR (KBr) ν_{max} : 3452, 2962, 1631, 1391, 1262, 1094, 1026, 804 cm^{-1} ; ESI-MS m/z : 274.89 $[M+H]^+$.

3-(2-Aminothiazol-4-yl)-6-chloro-2H-chromen-2-one (**3d**): Yellow solid, m.p. 204~205 °C (lit.^[18] 206~208 °C); 1H NMR (300 MHz, DMSO- d_6) δ : 8.45 (s, 1H), 7.98 (d, J =2.4 Hz, 1H), 7.45~7.64 (m, 3H), 7.18 (s, 2H, NH_2); ^{13}C NMR (75 MHz, DMSO- d_6) δ : 167.9, 158.8, 151.2, 143.5, 137.1, 131.3, 128.9, 128.0, 121.8, 121.2, 118.2, 110.0; IR (KBr) ν_{max} : 3444, 1639, 1372, 1086, 708 cm^{-1} ; ESI-MS m/z : 278.84 $[M+H]^+$.

3-(2-Aminothiazol-4-yl)-6-bromo-2H-chromen-2-one (**3e**): Yellow solid, m.p. 208~209 °C (lit.^[18] 210~212 °C); 1H NMR (300 MHz, $CDCl_3$) δ : 8.40 (s, 1H), 7.77~7.79 (s, 1H), 7.67~7.70 (m, 1H), 7.71~7.74 (m, 1H), 7.56~7.59 (m, 1H), 5.08 (s, 2H, NH_2); ^{13}C NMR (75 MHz, $CDCl_3$) δ : 166.4, 159.0, 154.1, 151.6, 145.9, 137.1, 133.8, 132.2, 130.3, 118.0, 117.1, 112.0; IR (KBr) ν_{max} : 3422, 1721, 1625, 1541, 1363, 1196, 1094, 823 cm^{-1} ; ESI-MS m/z : 323.05 $[M+H]^+$.

3-(2-Aminothiazol-4-yl)-6-nitro-2H-chromen-2-one (**3f**): Yellow solid, m.p. 181~182 °C; 1H NMR (300 MHz, DMSO- d_6) δ : 8.52 (d, J =2.7 Hz, 1H), 8.18~8.22 (m, 3H), 7.48 (s, 2H), 7.34 (d, J =9.0 Hz, 1H); ^{13}C NMR (75 MHz, DMSO- d_6) δ : 169.7, 157.5, 156.8, 148.3, 143.7, 142.6, 126.9, 124.7, 123.5, 118.9, 117.8, 114.5; IR (KBr) ν_{max} : 3426, 1737, 1616, 1526, 1344, 1093, 841 cm^{-1} ; ESI-MS m/z : 289.14 $[M]^+$. Anal. calcd for $C_{12}H_7N_3O_4S$: C 49.83, H 2.44, N 14.53; found C 49.76, H 2.37, N 14.47.

3-(2-Aminothiazol-4-yl)-7-(diethylamino)-2H-chromen-2-one (**3g**): Yellow solid, m.p. 198~199 °C (lit.^[32] 199~201 °C); 1H NMR (300 MHz, $CDCl_3$) δ : 8.37 (s, 1H), 7.59 (s, 1H), 7.27~7.35 (m, 1H), 6.35 (d, J =8.9 Hz, 1H), 6.51 (s, 1H), 5.03 (s, 2H, NH_2), 3.42 (q, J =7.2 Hz, 4H, CH_2),

1.22 (q, $J=7.2$ Hz, 6H, CH₃); ¹³C NMR (75 MHz, CDCl₃) δ : 166.1, 160.7, 155.7, 150.6, 144.9, 139.6, 129.3, 114.5, 109.2, 109.0, 107.9, 97.1, 44.8, 12.5; IR (KBr) $\nu_{\max}$: 3438, 2972, 1704, 1623, 1526, 1410, 1238, 1125, 770 cm⁻¹; ESI-MS m/z : 316.15 [M+H]⁺.

2-(2-Aminothiazol-4-yl)-3H-benzo[f]chromen-3-one (**3h**): Yellow solid, m.p. 286~287 °C (lit.^[18] 287~289 °C); ¹H NMR (300 MHz, CDCl₃) δ : 9.33 (s, 1H), 8.44~8.47 (m, 1H), 7.90~7.98 (m, 2H), 7.84 (s, 1H), 7.68~7.73 (m, 1H), 7.47~7.60 (m, 2H), 5.06 (s, 2H); ¹³C NMR (75 MHz, CDCl₃) δ : 162.3, 159.8, 156.2, 153.2, 143.5, 134.7, 132.5, 130.4, 129.5, 128.9, 128.1, 126.0, 122.1, 119.9, 116.6, 111.1; IR (KBr) $\nu_{\max}$: 3442, 2972, 1694, 1638, 1526, 1373, 1238, 1086, 786 cm⁻¹; ESI-MS m/z : 259.05 [M+H]⁺.

3-(2-(Methylamino)thiazol-4-yl)-2H-chromen-2-one (**3i**): Yellow solid, m.p. 170~171 °C; ¹H NMR (300 MHz, CDCl₃) δ : 8.54 (s, 1H), 7.78 (s, 1H), 7.51~7.60 (m, 2H), 7.29~7.36 (m, 2H), 5.24 (s, 1H), 3.05 (d, $J=4.8$ Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ : 169.4, 159.7, 152.8, 144.2, 138.5, 131.1, 128.1, 124.5, 121.1, 119.7, 116.3, 109.4, 32.2; IR (KBr) $\nu_{\max}$: 3443, 2127, 1638, 1378, 1084, 471 cm⁻¹; ESI-MS m/z : 259.01 [M+H]⁺. Anal. calcd for C₁₃H₁₀N₂O₂S: C 60.45, H 3.90, N 10.85; found C 60.23, H 3.78, N 10.67.

6-Methyl-3-(2-(methylamino)thiazol-4-yl)-2H-chromen-2-one (**3j**): Yellow solid, m.p. 129~130 °C; ¹H NMR (300 MHz, CDCl₃) δ : 8.49 (s, 1H), 7.76 (s, 1H), 7.30~7.37 (m, 3H), 5.33 (s, 1H), 3.06 (d, $J=4.5$ Hz, 3H), 2.42 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ : 169.5, 159.9, 151.0, 143.8, 138.7, 134.1, 132.3, 128.0, 120.6, 119.4, 116.0, 109.0, 32.3, 20.8; IR (KBr) $\nu_{\max}$: 3444, 1638, 1086, 476 cm⁻¹; ESI-MS m/z : 273.01 [M+H]⁺. Anal. calcd for C₁₄H₁₂N₂O₂S: C 61.75, H 4.44, N 10.29; found C 61.87, H 4.24, N 10.01.

8-Methoxy-3-(2-(methylamino)thiazol-4-yl)-2H-chromen-2-one (**3k**): Yellow solid, m.p. 178~179 °C; ¹H NMR (300 MHz, CDCl₃) δ : 8.50 (s, 1H), 7.77 (s, 1H), 7.14~7.23 (m, 2H), 7.05 (d, $J=7.4$ Hz, 1H), 5.32 (s, 1H), 3.40 (s, 3H), 3.04 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ : 169.5, 159.1, 146.8, 144.2, 142.4, 138.6, 124.2, 121.3, 120.3, 119.7, 113.0, 109.5, 56.2, 32.2; IR (KBr) $\nu_{\max}$: 3443, 1721, 1639, 1469, 1269, 1094, 713 cm⁻¹; ESI-MS m/z : 289.09 [M+H]⁺. Anal. calcd for C₁₄H₁₂N₂O₃S: C 58.32, H 4.20, N 9.72; found C 58.20, H 4.07, N 9.59.

6-Chloro-3-(2-(methylamino)thiazol-4-yl)-2H-chromen-2-one (**3l**): Yellow solid, m.p. 172~173 °C; ¹H NMR (300 MHz, CDCl₃) δ : 8.45 (s, 1H), 7.78 (s, 1H), 7.55~7.56 (m, 1H), 7.45~7.46 (m, 1H), 7.30~7.43 (m, 1H), 5.19 (s, 1H), 3.06 (d, $J=5.0$ Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ : 169.4, 159.1, 151.1, 143.8, 137.1, 134.3, 130.9, 127.2, 122.0, 120.8, 117.7, 110.2, 32.2; IR (KBr) $\nu_{\max}$: 3440, 1732, 1689, 1557, 1470, 1364, 1081, 773 cm⁻¹; ESI-MS m/z : 292.87 [M+H]⁺. Anal. calcd for C₁₃H₉ClN₂O₂S: C 53.34, H 3.10, N 9.57; found C 53.18, H 3.01, N 9.36.

6-Bromo-3-(2-(methylamino)thiazol-4-yl)-2H-chromen-

2-one (**3m**): Yellow solid, m.p. 185~186 °C; ¹H NMR (300 MHz, CDCl₃) δ : 8.40~8.45 (d, $J=12.8$ Hz, 1H), 7.70~7.79 (m, 3H), 7.56~7.59 (m, 1H), 5.15 (s, 1H), 3.06 (d, $J=5.0$ Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ : 169.4, 154.1, 151.6, 145.9, 137.0, 132.2, 130.2, 125.4, 121.4, 118.4, 118.0, 110.3, 32.1; IR (KBr) $\nu_{\max}$: 3440, 1722, 1669, 1551, 1476, 1402, 1356, 1201, 1081, 822, 774 cm⁻¹; ESI-MS m/z : 336.99 [M+H]⁺. Anal. calcd for C₁₃H₉BrN₂O₂S: C 46.31, H 2.69, N 8.31; found C 46.24, H 2.58, N 8.16.

3-(2-(Methylamino)thiazol-4-yl)-6-nitro-2H-chromen-2-one (**3n**): Yellow solid, m.p. 146~147 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ : 8.50~8.59 (m, 2H), 8.33~8.37 (m, 1H), 7.79 (s, 1H), 7.44~7.47 (m, 1H), 5.15 (s, 1H), 3.07 (d, $J=5.0$ Hz, 3H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ : 169.4, 158.1, 156.0, 144.2, 143.3, 136.6, 125.5, 123.7, 122.9, 119.9, 117.3, 111.1, 32.1; IR (KBr) $\nu_{\max}$: 3408, 1742, 1720, 1620, 1583, 1476, 1342, 1213, 1092, 839, 746 cm⁻¹; ESI-MS m/z : 303.00 [M]⁺. Anal. calcd for C₁₃H₉N₃O₄S: C 51.48, H 2.99, N 13.85; found C 51.40, H 2.89, N 13.76.

7-Diethylamino-3-(2-(methylamino)thiazol-4-yl)-2H-chromen-2-one (**3o**): Yellow solid, m.p. 123~124 °C; ¹H NMR (300 MHz, CDCl₃) δ : 8.41 (s, 1H), 7.58 (s, 1H), 7.36 (d, 1H), 6.61~6.62 (m, 1H), 6.52~6.59 (m, 1H), 5.21 (s, 1H), 3.42 (q, $J=7.1$ Hz, 4H), 3.03 (d, $J=4.7$ Hz, 3H), 1.22 (t, $J=7.0$ Hz, 6H); ¹³C NMR (75 MHz, CDCl₃) δ : 169.4, 160.8, 155.6, 150.5, 145.1, 139.5, 129.3, 114.6, 109.1, 109.0, 106.1, 97.0, 44.8, 32.2, 12.5; IR (KBr) $\nu_{\max}$: 3442, 1712, 1623, 1403, 1235, 1125, 771 cm⁻¹; ESI-MS m/z : 330.13 [M+H]⁺. Anal. calcd for C₁₇H₁₉N₃O₂S: C 61.98, H 5.81, N 12.76; found C 61.81, H 5.69, N 12.64.

2-(2-(Methylamino)thiazol-4-yl)-3H-benzo[f]chromen-3-one (**3p**): Yellow solid, m.p. 200~201 °C; ¹H NMR (300 MHz, CDCl₃) δ : 9.35 (s, 1H), 8.43 (d, $J=8.4$ Hz, 1H), 7.89~7.91 (m, 2H), 7.83 (s, 1H), 7.67~7.72 (m, 1H), 7.47~7.57 (m, 2H), 5.28 (s, 1H), 3.09 (d, 3H, $J=4.8$ Hz); ¹³C NMR (75 MHz, CDCl₃) δ : 169.6, 159.8, 152.3, 144.8, 134.5, 132.4, 130.4, 129.4, 128.9, 128.0, 126.0, 122.1, 120.2, 116.6, 114.0, 109.3, 32.3; IR (KBr) $\nu_{\max}$: 3441, 1632, 1554, 1378, 1086, 810 cm⁻¹; ESI-MS m/z : 309.14 [M+H]⁺. Anal. calcd for C₁₇H₁₂N₂O₂S: C 66.22, H 3.92, N 9.08; found C 66.11, H 3.76, N 9.01.

3-(2-(Phenylamino)thiazol-4-yl)-2H-chromen-2-one (**3q**): Yellow solid, m.p. 187~189 °C (lit.^[18] 188~190 °C); ¹H NMR (300 MHz, DMSO-*d*₆) δ : 10.36 (s, 1H), 8.70 (s, 1H), 7.97 (d, $J=6.9$ Hz, 1H), 7.75~7.79 (m, 2H), 7.62~7.67 (m, 1H), 7.36~7.48 (m, 5H), 7.01 (t, $J=7.3$ Hz, 1H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ : 162.5, 158.8, 152.3, 143.6, 141.0, 138.6, 131.7, 129.1, 129.0, 124.7, 121.5, 120.3, 119.3, 117.1, 115.9, 109.9; IR (KBr) $\nu_{\max}$: 3442, 2921, 1606, 1543, 1087, 740 cm⁻¹; ESI-MS m/z : 321.17 [M+H]⁺.

6-Methyl-3-(2-(phenylamino)thiazol-4-yl)-2H-chromen-2-one (**3r**): Yellow solid, m.p. 123~124 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ : 10.36 (s, 1H), 8.64 (s, 1H), 7.75~7.78 (m, 4H), 7.34~7.44 (m, 4H), 7.01 (t, $J=7.32$

Hz, 1H), 2.41 (s, 3H, CH₃); ¹³C NMR (75 MHz, DMSO-*d*₆) δ: 162.5, 160.0, 150.5, 143.7, 141.0, 138.6, 134.0, 132.6, 129.1, 128.6, 121.4, 120.2, 119.0, 117.1, 115.6, 109.8, 20.3; IR (KBr) ν_{max}: 3436, 1704, 1604, 1543, 1497, 1440, 1313, 1094, 744 cm⁻¹; ESI-MS *m/z*: 335.08 [M+H]⁺. Anal. calcd for C₁₉H₁₄N₂O₂S: C 68.24, H 4.22, N 8.38; found C 68.12, H 4.13, N 8.26.

8-Methoxy-3-(2-(phenylamino)thiazol-4-yl)-2H-chromen-2-one (**3s**): Yellow solid, m.p. 169~170 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ: 10.36 (s, 1H), 8.66 (s, 1H), 7.74~7.80 (m, 3H), 7.49~7.51 (m, 1H), 7.32~7.41 (m, 4H), 7.00 (t, *J*=7.3 Hz, 1H), 3.94 (s, 3H, OCH₃-H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ: 162.5, 158.5, 147.3, 146.2, 143.5, 141.6, 140.9, 138.8, 129.1, 124.6, 121.4, 120.4, 119.8, 117.1, 113.8, 110.0, 56.1; IR (KBr) ν_{max}: 3436, 1639, 1266, 1097, 804 cm⁻¹; ESI-MS *m/z*: 350.99 [M+H]⁺. Anal. calcd for C₁₉H₁₄N₂O₃S: C 65.13, H 4.03, N 7.99; found C 65.01, H 3.97, N 7.81.

6-Chloro-3-(2-(phenylamino)thiazol-4-yl)-2H-chromen-2-one (**3t**): Yellow solid, m.p. 215~216 °C (lit.^[18] 215~217 °C); ¹H NMR (300 MHz, DMSO-*d*₆) δ: 10.36 (s, 1H), 8.60 (s, 1H), 8.07~8.15 (m, 2H), 7.67~7.81 (m, 3H), 7.63~7.67 (m, 1H), 7.47~7.51 (m, 1H), 7.39 (d, *J*=7.8 Hz, 1H), 7.02 (t, *J*=7.2 Hz, 1H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ: 162.5, 158.4, 150.9, 143.4, 140.9, 137.3, 133.8, 131.0, 129.1, 128.4, 121.4, 120.8, 119.5, 117.8, 117.1, 110.7; IR (KBr) ν_{max}: 3441, 1721, 1612, 1548, 1076, 745 cm⁻¹; ESI-MS *m/z*: 354.92 [M+H]⁺.

6-Bromo-3-(2-(phenylamino)thiazol-4-yl)-2H-chromen-2-one (**3u**): Yellow solid, m.p. 229~231 °C (lit.^[18] 229~231 °C); ¹H NMR (300 MHz, DMSO-*d*₆) δ: 10.36 (s, 1H), 8.60 (s, 1H), 8.30 (s, 1H), 7.75~7.81 (m, 3H), 7.36~7.44 (m, 3H), 7.18 (s, 1H), 7.00~7.02 (m, 1H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ: 162.5, 158.0, 153.6, 145.6, 143.4, 140.9, 137.2, 136.6, 132.5, 129.1, 125.4, 121.4, 120.0, 118.4, 117.1, 116.3; IR (KBr) ν_{max}: 3443, 1717, 1605, 1545, 1094, 741 cm⁻¹; ESI-MS *m/z*: 398.96 [M+H]⁺.

6-Nitro-3-(2-(phenylamino)thiazol-4-yl)-2H-chromen-2-one (**3v**): Yellow solid; m.p. 190~191 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ: 10.13 (s, 1H), 8.34 (br, 1H), 7.85 (br, 1H), 7.69~7.70 (m, 1H), 7.39~7.44 (m, 3H), 7.26~7.28 (m, 4H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ: 176.6, 166.9, 138.5, 137.0, 131.6, 131.5, 130.1, 129.7, 129.6, 129.3, 128.6, 126.2, 124.3, 122.4, 114.8, 103.8; IR (KBr) ν_{max}: 3425, 1613, 1526, 1384, 1263, 599 cm⁻¹; ESI-MS *m/z*: 365.99 [M+H]⁺. Anal. calcd for C₁₈H₁₁N₃O₄S: C 59.17, H 3.03, N 11.50; found C 59.10, H 2.96, N 11.42.

7-(Diethylamino)-3-(2-(phenylamino)thiazol-4-yl)-2H-chromen-2-one (**3w**): Yellow solid, m.p. 198~199 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ: 10.27 (s, 1H), 8.52 (s, 1H), 7.74 (d, *J*=10.8 Hz, 2H), 7.66 (d, *J*=8.8 Hz, 1H), 7.55 (s, 1H), 7.35~7.40 (m, 2H), 6.96~7.01 (m, 1H), 6.75~6.78 (m, 1H), 6.60 (s, 1H), 3.46 (q, *J*=7.2 Hz, 4H, CH₂), 1.45 (t, *J*=7.2 Hz, 6H, CH₃); ¹³C NMR (75 MHz, DMSO-*d*₆) δ: 162.2, 159.9, 158.3, 155.3, 152.9, 147.6, 144.6, 139.6, 132.4, 129.8, 122.5, 121.2, 116.9, 114.9, 110.1, 107.5, 44.4, 12.3; IR (KBr) ν_{max}: 3438, 1694, 1616, 1508,

1124, 1079, 744 cm⁻¹; ESI-MS *m/z*: 392.01 [M+H]⁺. Anal. calcd for C₂₂H₂₁N₃O₂S: C 67.50, H 5.41, N 10.73; found C 67.43, H 5.28, N 10.61.

2-(2-(Phenylamino)thiazol-4-yl)-3H-benzo[f]chromen-3-one (**3x**): Yellow solid, m.p. 201~202 °C; ¹H NMR (300 MHz, DMSO-*d*₆) δ: 10.43 (s, 1H), 9.37 (s, 1H), 8.44 (d, *J*=8.4 Hz, 1H), 8.22 (d, *J*=9.0 Hz, 1H), 8.11 (d, *J*=8.1 Hz, 1H), 7.78~7.90 (m, 4H), 7.64~7.71 (m, 2H), 7.45 (t, *J*=7.8 Hz, 2H), 7.05 (t, *J*=7.2 Hz, 1H); ¹³C NMR (75 MHz, DMSO-*d*₆) δ: 162.8, 158.8, 151.9, 143.8, 141.0, 133.9, 133.0, 130.1, 129.2, 129.1, 128.7, 128.6, 126.2, 121.6, 121.5, 119.5, 117.2, 116.5, 113.2, 109.9; IR (KBr) ν_{max}: 3441, 1632, 1554, 1378, 1086, 810, 778 cm⁻¹; ESI-MS *m/z*: 371.09 [M+H]⁺.

References

- [1] Spino, C.; Dodier, M. Sotheeswaran, S. *Bioorg. Med. Chem. Lett.* **1998**, 8, 3475.
- [2] Witaicenis, A.; Seito, L. N.; Di Stasi, L. C. *Chem. Biol. Interact.* **2010**, 186, 211.
- [3] Arora, R. K.; Kaur, N.; Bansal, Y.; Bansal, G. *Acta Pharm. Sin. B* **2014**, 4, 368.
- [4] Basile, A.; Sorbo, S.; Spadaro, V. *Molecules* **2009**, 14, 939.
- [5] Huang, X.; Liu, J. *Chin. J. Org. Chem.* **2018**, 38, 1233 (in Chinese). (黄新炜, 刘建利, 有机化学, **2018**, 38, 1233.)
- [6] Sun, H.; Sun, H.; Liu, M.; Huang, W.; Yang, G. *Chin. J. Org. Chem.* **2018**, 38, 2067 (in Chinese). (孙海燕, 孙宏顺, 刘明珍, 黄伟, 杨光富, 有机化学, **2018**, 38, 2067.)
- [7] Fang, Z. X.; Liu, J. Y.; Qiao, Y. H. *Chin. J. Org. Chem.* **2018**, 38, 1985 (in Chinese). (房智兴, 刘巨艳, 乔艳红, 有机化学, **2018**, 38, 1985.)
- [8] Kim, J.; Moon, Y.; Ham, S. W. *Bull. Korean Chem. Soc.* **2011**, 32, 2893.
- [9] Ding, C.; Pan, Y.; Yin, X.; Tan, C. *Chin. J. Org. Chem.* **2019**, 39, 2099 (in Chinese). (丁成荣, 潘亚运, 殷许, 谭成侠, 有机化学, **2019**, 39, 2099.)
- [10] Xu, J.; Ma, L.; Liu, X. B. *Chin. J. Org. Chem.* **2018**, 38, 1680 (in Chinese). (徐姣, 马玲, 刘秀波, 有机化学, **2018**, 38, 1680.)
- [11] Zhang, Q.; Wang, Y.; Zhao, Y.; Ma, C.; Xu, X.; Gu, W.; Yang, Y.; Wang, S. *Chin. J. Org. Chem.* **2019**, 39, 2616 (in Chinese). (张强健, 王芸芸, 赵雨珣, 马崇慧, 徐徐, 谷文, 杨益琴, 王石发, 有机化学, **2019**, 39, 2616.)
- [12] Koelsch, C. F. *J. Am. Chem. Soc.* **1950**, 72, 2993.
- [13] Vardhan, V. A.; Rao, V. R. *Indian J. Chem., Sect. B* **1997**, 36, 1085.
- [14] Ramanna, S.; Rao, V. R.; Kumari, T. S.; Rao, T. V. P. *Phosphorus Sulfur Relat. Elem.* **1995**, 107, 197.
- [15] Czerney, P.; Hartmann, H. *J. Prakt. Chem.* **1983**, 325, 551.
- [16] Ramagiri, R. K.; Vedula, R. R.; Thupurani, M. K. *Phosphorus sulfur Relat. Elem.* **2015**, 190, 1393.
- [17] Zhuravel, I. O.; Kovalenko, S. M.; Vlasov, S. V.; Chernykh, V. P. *Molecules* **2005**, 10, 444.
- [18] Ramesh, G.; Janardhan, B.; Rajitha, B. *Res. Chem. Intermed.* **2015**, 41, 8099.
- [19] Chunduru, V. S. R.; Rao, R. V. *J. Chem. Res.* **2010**, 34, 50.
- [20] Saeed, A.; Ejaz, S. A.; Shehzad, M.; Hassan, S.; al-Rashida, M.; Lecka, J.; Sévigny, J.; Iqbal, J. *RSC Adv.* **2016**, 6, 21026.
- [21] Hande, P. E.; Samui, A. B.; Kulkarni, P. S. *Sens. Actuators, B* **2017**, 246, 597.

- [22] Patil, N. T.; Shinde, V. S.; Gajula, B. *Org. Biomol. Chem.* **2012**, *10*, 211.
- [23] Fan, L.; Yao, C.; Shu, M. *Heterocycl. Commun.* **2016**, *22*, 63.
- [24] Jadhvar, S. C.; Kasraliker, H. M.; Goswami, S. V.; Chakrawar, A. V.; Bhusare, S. R. *Res. Chem. Intermed.* **2017**, *43*, 7211.
- [25] Jannati, S.; Esmacili, A. A. *Res. Chem. Intermed.* **2017**, *43*, 6817.
- [26] Reddy, K. M. K.; Sadik, S. M.; Saichaithanya, N.; Peddanna, K.; Reddy, N. B.; Sravya, G.; Grigory V, Z.; Reddy, C. S. *Res. Chem. Intermed.* **2017**, *43*, 7087.
- [27] Kim, K. M.; Ryu, E. K. *Tetrahedron Lett.* **1996**, *37*, 1441.
- [28] More, S. V.; Sastry, M. N. V.; Wang, C. C.; Yao, C. F. *Tetrahedron Lett.* **2005**, *46*, 6345.
- [29] Behmadi, H.; Roshani, M.; Saadati, S. M. *Chin. Chem. Lett.* **2009**, *20*, 5.
- [30] Zhu, Y. P.; Yuan, J. J.; Zhao, Q.; Lian, M.; Gao, Q. H.; Liu, M. C.; Yang, Y.; Wu, A. X. *Tetrahedron* **2012**, *68*, 173.
- [31] Bogdal, D. J. *Chem. Res., Synop.* **1998**, 468.
- [32] Lin, W.; Cao, X.; Yuan, L. Ding, Y. *Chem. Eur. J.* **2010**, *16*, 6454.

(Cheng, F.)