双官能团硫脲催化吲哚-3-酮对 β,γ -不饱和酮酯的不对称 Michael 加成反应研究

刘耀宗^{*a} 许鹏飞^b 马建军^a 李晓明^a 梁睿渊^a 滕志君^a

(^a 甘肃银光聚银化工有限公司 甘肃白银 730900)

(^b 兰州大学化学化工学院 功能有机分子化学国家重点实验室 兰州 730000)

摘要 通过双官能团硫脲催化的吲哚-3-酮对不饱和酮酯的不对称 Michael 加成反应研究, 以很高的产率、很好的对映选择性和非常好的非对映选择性得到了手性的 2-取代吲哚-3-酮类化合物, 对吲哚类化合物的手性合成具有积极意义。

关键词 不对称催化; 双官能团硫脲; Michael 加成; 吲哚-3-酮; 不饱和酮酯

Bifunctional Thioureas Catalyzed Asymmetric Michael Addition of 1-Acetylindolin-3-ones to β,γ -Unsaturated α -Keto Esters

Liu, Yaozong^{*a} Xu, Pengfei^b Ma, Jianjun^a Li, Xiaoming^a

Liang, Ruiyuan^a Teng, Zhijun^a

(^a Gansu Yinguang Juyin Chemical Co., Ltd., Baiyin, Gansu 730900)

(^b State Key Laboratory of Applied Organic Chemistry, College of Chemistry and Chemical Engineering, Lanzhou University, Lanzhou 730000)

Abstract The asymmetric Michael addition of 1-acetylindolin-3-ones to β,γ -unsaturated α -keto esters catalyzed by bifunctional thioureas has been developed. Enantio-pure 2-substituted indolin-3-one derivatives were obtained easily in excellent yields (up to 99%) with good diastereoselectivity (up to 10 : 1) and excellent enantioselectivities (up to 99.5%), which would be useful for the chiral synthesis of indole-related compounds.

Keywords asymmetric catalysis; bifunctional thiourea; Michael addition; 1-acetylindolin-3-one; β,γ -unsaturated α -keto ester

1 Introduction

The catalytic asymmetric Michael addition reaction is one of the most important organic reactions for the carbon-carbon bond-formation in organic synthesis. Remarkable progress has been made in the organocatalytic version of this reaction in recent years.^[1,2] The development of novel and effective transformations of this reaction is still under exploring. Compared to other Michael acceptors, the reactions using β,γ -unsaturated α -keto esters were very limited.^[3] So the enantioselective version of β,γ -unsaturated α -keto esters should be explored due to the fact that the adducts can be functionalized readily to corresponding amino acid esters or α -hydroxy acid esters.^[4]

2-Substituted indolin-3-ones exist widely in biologically active natural products and pharmaceutical compounds, and have been used as versatile intermediates in organic and natural-product synthesis.^[5] Although a lot of prepara-

tive methods have been developed,^[6] the enantioselective synthesis of these structures is much less explored.^[7] Then the development of an enantioselective, metal-free and highly effective approach for the asymmetric synthesis of enantio-pure 2-substituted indolin-3-one derivatives appeared to be of great significance. As a part of our continuing interest in exploring the preparation of 2-substituted indolin-3-ones,^[7e-7f] bifunctional thioureas were found to be efficient catalysts for the asymmetric Michael addition of indolin-3-ones with β,γ -unsaturated α -keto esters, which provides a direct access to optically pure 2-substituted indolin-3-ones.

2 Results and discussion

The reaction of 1-acetylindolin-3-one (**2a**) with β,γ -unsaturated α -keto ester **3a** catalyzed by thiourea-tertiary amine catalyst **1a** (10 mol%) derivated from quinine in

* Corresponding author. E-mail: 1048632722@qq.com

Received November 23, 2019; revised January 4, 2020; published online January 15, 2020.

dichloromethane (DCM) was first investigated.^[8] To our delight, the desired product **4a** was obtained in 91% yield with 6 : 1 *dr* and 98% *ee* (Table 1, Entry 1). Then different bifunctional thiourea catalysts **1a**~**1f** were screened under the same reaction conditions (Table 1, Entries 1~6). All thiourea-tertiary amine catalysts gave excellent yields and enantioselectivities, while the thiourea-primary amine catalyst **1e** and thiourea-secondary amine catalyst **1f** gave disappointing results. The catalyst **1a** provided the best *dr* with excellent yield and *ee* value, therefore, catalyst **1a** was chosen to further optimize the reaction.

Table 1 Optimization of the reaction conditions^a

Entry	Cat.	Solvent	<i>t</i> /h	Yield ^b /%	<i>dr</i> ^c	<i>ee</i> ^d /%
1	1a	DCM	6	91	6 : 1	98
2	1b	DCM	6	95	4 : 1	−98
3	1c	DCM	6	87	3 : 1	−99
4	1d	DCM	6	94	5 : 1	96
5	1e	DCM	24	<20	1 : 1	44
6	1f	DCM	12	89	4 : 1	−60
7	1a	CHCl ₃	6	92	5 : 1	97
8	1a	Toluene	6	82	6 : 1	97
9	1a	THF	24	93	6 : 1	98
10	1a	CH ₃ CN	24	95	6 : 1	96
11	1a	Acetone	24	84	6 : 1	98
12 ^e	1a	DCM	10	93	6 : 1	98.5
13 ^f	1a	DCM	12	95	6 : 1	99

^a The reaction was performed using **2a** (0.2 mmol), **3a** (0.3 mmol), catalyst **1** (10 mol%) in 0.4 mL of solvent. ^b Isolated yields. ^c Determined by ¹H NMR spectroscopic analysis. ^d Determined by HPLC on a chiral stationary phase. ^e Reaction at 0 °C. ^f Reaction at −10 °C.

Then, different organic solvents with 10% catalyst **1a** were screened, all the solvents led to excellent enantiomeric excess values, and CH₂Cl₂ was identified as the most ideal solvent which afforded the best result (Table 1, Entry 1). When the reaction temperature was decreased from room temperature to 0 °C, a little higher *ee* value of 98.5%

was obtained (Table 1, Entry 12). When the reaction temperature was reduced to −10 °C, the best result was obtained in 95% yield with 6 : 1 *dr* and 99% *ee* (Table 1, Entry 13).

Under the optimized reaction conditions, a wide range of β,γ -unsaturated α -keto esters **3** and 1-acetylindolin-3-ones **2** were investigated in this asymmetric Michael addition reaction. The results are summarized in Table 2. Good *dr*, excellent yields and *ee* values were obtained for the desired products. Structural variation of keto esters could be well tolerated (Table 2). β,γ -Unsaturated α -keto esters bearing either electron-withdrawing (Table 2, Entries 2, 4, 5, 7, 8) or electron-donating (Table 2, Entries 3, 6, 9, 10) groups on the γ -phenyl rings were found to proceed well. These transformations gave the corresponding products **4b**~**4j** in 86%~99% yields with high levels of enantioselectivity (92%~99.5% *ee*). The position and electronic properties of substituents on aromatic rings of α -keto esters had limited influence on the stereochemical outcome. Heteroaromatic 2-furyl substituted keto ester **3k** was also effectively engaged in the reaction under the optimized reaction conditions (85% yield, 4 : 1 *dr*, 99% *ee*; Table 2, Entry 11). Unfortunately, cyclohexyl substituted keto ester **3l** (Table 2, Entry 12) proceeded very slowly under the optimized reaction conditions and even at room temperature. Substituted 1-acetyl-indolin-3-ones containing an electron-dona-

Table 2 Catalytic asymmetric Michael addition of 1-acetylindolin-3-ones to β,γ -unsaturated α -keto esters^a

Entry	R ¹	R ²	<i>t</i> /h	Yield ^b /% (4)	<i>dr</i> ^c	<i>ee</i> ^d /%
1	H	Ph	12	95 (4a)	6 : 1	99
2	H	2-ClC ₆ H ₄	6	95 (4b)	9 : 1	97
3	H	2-MeOC ₆ H ₄	6	99 (4c)	9 : 1	92
4	H	2,4-Cl ₂ C ₆ H ₃	12	99 (4d)	10 : 1	99
5	H	3-ClC ₆ H ₄	12	89 (4e)	7 : 1	98
6	H	3,4-(CH ₃) ₂ C ₆ H ₃	12	93 (4f)	6 : 1	92
7	H	4-ClC ₆ H ₄	4	94 (4g)	7 : 1	99
8	H	4-BrC ₆ H ₄	4	94 (4h)	7 : 1	99.5
9	H	4-MeC ₆ H ₄	12	94 (4i)	6 : 1	98
10	H	4-MeOC ₆ H ₄	24	86 (4j)	6 : 1	92
11	H	2-Furyl	24	85 (4k)	4 : 1	99
12	H	Cyclohexyl	24	Trace	ND ^e	ND
13	5-Br	Ph	4	98 (4l)	8 : 1	98
14	5-Cl	Ph	4	88 (4m)	6 : 1	98
15	5-Me	Ph	12	89 (4n)	6 : 1	98
16	7-Me	Ph	4	86 (4o)	9 : 1	90

^a The reaction was performed using **2** (0.2 mmol), **3** (0.3 mmol), catalyst **1a** (10 mol%) in 0.4 mL of solvent. ^b Isolated yields. ^c Determined by ¹H NMR spectroscopic analysis. ^d Determined by HPLC on a chiral stationary phase. ^e Reaction at 0 °C. ^f Reaction at −10 °C.

ting group or an electron- withdrawing group were also tested in the reaction, and satisfying results were obtained (Table 2, Entries 13~16). With this protocol, highly enantio-enriched 2-substituted 1-acetylindolin-3-ones could be easily obtained.

The absolute configuration of the product was determined by the crystals of compound **5a** (CCDC 842611) bearing a chlorine atom, which was obtained from the reduction of **4e**. As shown in Figure 1, the newly formed stereogenic centers in the Michael adducts were confirmed as (1*S*,11*R*).

Figure 1 X-Ray crystal structure of **5a**

On the basis of the observed reactivity and experimental results of the Michael addition reactions, it was proposed that the reaction proceeds via a dual activation model. As shown in Figure 2, a thiourea moiety of the catalyst **1a** interacts through hydrogen bonding with two adjacent carbonyl groups of the keto ester and enhances their electrophilicity while the tertiary amine deprotonates an acidic proton of 1-acetylindolin-3-one, generating a ternary complex. This proposed dual activation model might be responsible for the stereochemical outcome of the Michael adducts.

Figure 2 Proposed transition state model in the Michael addition

3 Conclusions

In summary, a bifunctional thiourea-tertiary amine catalyzed asymmetric Michael addition of 1-acetylindolin-3-ones to β,γ -unsaturated α -keto esters was developed in excellent yields, good diastereoselectivities and excellent

enantioselectivities, providing a practical method to synthesize highly enantiopure 2-substituted indolin-3-ones. Further investigation and application of this methodology are ongoing.

4 Experimental section

4.1 Representative procedure for the catalytic asymmetric Michael addition reaction

To a solution of β,γ -unsaturated α -keto ester **3** (0.3 mmol, 1.5 equiv.) and catalyst **1a** (12 mg, 0.02 mmol, 0.1 equiv.) in dry CH_2Cl_2 (0.4 mL) cooled at -10°C was added 1-acetylindolin-3-one **2** (0.2 mmol, 1.0 equiv.). The solution was stirred at -10°C until **2** was consumed and monitored by thin-layer chromatography (TLC). The reaction mixture was concentrated *in vacuo* and purified by flash column chromatography (petroleum ether/ ethyl acetate as eluent) to afford the product.

(*R*)-Ethyl 4-((*S*)-1-acetyl-3-oxoindolin-2-yl)-2-oxo-4-phenylbutanoate (**4a**): colorless oil, 72 mg (95% yield, 99% *ee*). $[\alpha]_{\text{D}}^{20} -169$ (*c* 1.0, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) δ : 8.16 (s, 1H), 7.40~7.34 (m, 2H), 6.96~6.94 (m, 6H), 4.59 (s, 1H), 4.39~4.34 (m, 3H), 4.03 (s, 1H), 3.43 (s, 1H), 2.64 (s, 3H), 1.40 (t, *J*=7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 198.0, 193.2, 168.5, 160.2, 153.2, 136.6, 135.3, 128.1, 127.8, 127.5, 125.1, 123.6, 122.6, 118.2, 66.1, 62.6, 41.8, 38.6, 23.9, 13.8; IR (KBr) ν : 3402.0, 2983.5, 1726.5, 1675.7, 1606.3, 1464.2, 1380.3, 1282.5, 1068.8, 761.7 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{22}\text{NO}_5$ $[\text{M}+\text{H}]^+$ 380.1492, found 380.1483. The enantiomeric excess was determined by HPLC with an AS-H column (hexane/*i*-PrOH, *V*: *V*=70:30), 1.0 mL/min; minor enantiomer t_{R} =15.0 min, major enantiomer t_{R} =21.4 min.

(*R*)-Ethyl 4-((*S*)-1-acetyl-3-oxoindolin-2-yl)-4-(2-chlorophenyl)-2-oxobutanoate (**4b**): colorless oil, 78 mg (95% yield, 97% *ee*). $[\alpha]_{\text{D}}^{20} -205$ (*c* 1.0, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) δ : 8.21 (s, 1H), 7.46 (d, *J*=8.0 Hz, 1H), 7.40 (t, *J*=8.0 Hz, 1H), 7.13 (dd, *J*=2.0, 8.0 Hz, 1H), 7.03~6.87 (m, 4H), 4.79 (s, 1H), 4.66 (s, 1H), 4.40~4.32 (m, 3H), 3.34 (s, 1H), 2.66 (s, 3H), 1.40 (t, *J*=6.8 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 198.6, 192.9, 169.1, 160.2, 153.5, 137.0, 134.7, 133.8, 129.5, 128.9, 128.7, 126.5, 125.2, 123.8, 122.8, 118.4, 66.3, 62.8, 39.2, 37.2, 24.0, 14.0; IR (KBr) ν : 3339.8, 2983.2, 1726.7, 1680.0, 1606.3, 1465.8, 1378.9, 1277.2, 1040.6, 759.3 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{21}\text{ClNO}_5$ $[\text{M}+\text{H}]^+$ 414.1103, found 414.1109. The enantiomeric excess was determined by HPLC with an AD-H column (hexane/*i*-PrOH, *V*: *V*=90:10), 1.0 mL/min; major enantiomer t_{R} =33.3 min, minor enantiomer t_{R} =38.5 min.

(*R*)-Ethyl 4-((*S*)-1-acetyl-3-oxoindolin-2-yl)-4-(2-methoxyphenyl)-2-oxobutanoate (**4c**): colorless oil, 81 mg (99% yield, 92% *ee*). $[\alpha]_{\text{D}}^{20} -423$ (*c* 1.0, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) δ : 8.18 (s, 1H), 7.44 (d, *J*=7.6 Hz, 1H), 7.38~7.34 (m, 1H), 6.96~6.89 (m, 3H), 6.61~6.56 (m, 2H), 4.80~4.75 (m, 1H), 4.62 (s, 1H), 4.46~4.34 (m,

3H), 3.71 (s, 3H), 3.26 (d, $J=19.6$ Hz, 1H), 2.66 (s, 3H), 1.41 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 198.8, 193.7, 169.2, 160.4, 157.0, 153.6, 136.6, 128.6, 128.3, 125.1, 124.3, 123.3, 122.6, 120.2, 117.8, 110.0, 66.1, 62.7, 55.1, 38.5, 33.2, 23.7, 14.0; IR (KBr) ν : 3342.3, 2935.7, 1726.0, 1676.4, 1464.0, 1380.8, 1283.1, 1245.0, 1077.8, 1025.9, 757.6 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{24}\text{NO}_6$ $[\text{M}+\text{H}]^+$ 410.1598, found 410.1589. The enantiomeric excess was determined by HPLC with an AS-H column (hexane/*i*-PrOH, $V:V=90:10$), 1.0 mL/min; minor enantiomer $t_{\text{R}}=24.0$ min, major enantiomer $t_{\text{R}}=27.5$ min.

(*R*)-Ethyl 4-((*S*)-1-acetyl-3-oxoindolin-2-yl)-4-(2,4-dichlorophenyl)-2-oxobutanoate (**4d**): colorless oil, 88 mg (99% yield, 99% *ee*). $[\alpha]_{\text{D}}^{20} -213$ (c 1.0, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) δ : 8.21 (s, 1H), 7.48~7.44 (m, 2H), 7.16 (d, $J=2.4$ Hz, 1H), 7.02 (t, $J=7.6$ Hz, 1H), 6.96 (d, $J=8.8$ Hz, 1H), 6.88 (dd, $J=2.4$ Hz, 8.8 Hz, 1H), 4.74 (s, 1H), 4.66 (s, 1H), 4.40~4.32 (m, 3H), 3.31 (d, $J=5.2$ Hz, 1H), 2.65 (s, 3H), 1.40 (t, $J=6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 198.4, 192.6, 168.9, 160.1, 153.4, 137.2, 135.4, 133.8, 132.6, 129.2, 126.8, 124.9, 124.0, 122.8, 118.5, 66.2, 62.8, 39.2, 36.8, 24.0, 13.9; IR (KBr) ν : 3404.6, 2983.8, 1729.1, 1681.5, 1605.3, 1466.3, 1379.7, 1275.2, 1050.3, 761.7 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{20}\text{Cl}_2\text{NO}_5$ $[\text{M}+\text{H}]^+$ 448.0713, found 448.0724. The enantiomeric excess was determined by HPLC with an AS-H column (hexane/*i*-PrOH, $V:V=70:30$), 1.0 mL/min; major enantiomer $t_{\text{R}}=12.6$ min, minor enantiomer $t_{\text{R}}=16.4$ min.

(*R*)-Ethyl 4-((*S*)-1-acetyl-3-oxoindolin-2-yl)-4-(3-chlorophenyl)-2-oxobutanoate (**4e**): colorless oil, 73 mg (89% yield, 98% *ee*). $[\alpha]_{\text{D}}^{20} -139$ (c 1.0, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) δ : 8.18 (s, 1H), 7.46~7.40 (m, 2H), 7.02~6.91 (m, 4H), 6.86 (d, $J=7.2$ Hz, 1H), 4.61 (s, 1H), 4.40~4.35 (m, 3H), 4.03 (s, 1H), 3.43 (s, 1H), 2.63 (s, 3H), 1.41 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 197.8, 192.8, 168.4, 160.2, 153.0, 137.6, 136.9, 133.8, 129.1, 128.3, 127.8, 126.5, 125.1, 124.0, 122.9, 118.2, 66.0, 62.8, 41.5, 38.5, 24.0, 13.9; IR (KBr) ν : 3403.4, 2984.2, 1723.8, 1676.5, 1603.6, 1465.6, 1379.9, 1282.8, 1077.6, 1043.7, 761.2 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{21}\text{ClNO}_5$ $[\text{M}+\text{H}]^+$ 414.1103, found 414.1104. The enantiomeric excess was determined by HPLC with an AS-H column (hexane/*i*-PrOH, $V:V=70:30$), 1.0 mL/min; minor enantiomer $t_{\text{R}}=14.8$ min, major enantiomer $t_{\text{R}}=21.4$ min.

(*R*)-Ethyl 4-((*S*)-1-acetyl-3-oxoindolin-2-yl)-4-(3,4-dimethylphenyl)-2-oxobutanoate (**4f**): colorless oil, 76 mg (93% yield, 92% *ee*). $[\alpha]_{\text{D}}^{20} -117$ (c 1.0, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) δ : 8.18 (s, 1H), 7.44~7.35 (m, 2H), 6.95 (t, $J=7.6$ Hz, 1H), 6.74~6.66 (m, 3H), 4.57 (s, 1H), 4.40~4.34 (m, 3H), 3.96 (s, 1H), 3.36 (d, $J=18.0$ Hz, 1H), 2.62 (s, 3H), 2.00 (s, 3H), 1.98 (s, 3H), 1.40 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 198.3, 193.4, 168.5, 160.3, 153.3, 136.5, 136.0, 135.7, 132.7, 129.3, 129.1, 125.5, 125.2, 123.6, 122.6, 118.2, 66.2, 62.6, 41.4, 38.9, 23.9, 19.3, 19.0, 13.9; IR (KBr) ν : 3406.7,

2980.2, 1725.2, 1676.8, 1606.2, 1464.5, 1381.2, 1282.4, 1087.9, 1047.1, 762.5 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{26}\text{NO}_5$ $[\text{M}+\text{H}]^+$ 408.1815, found 408.1815. The enantiomeric excess was determined by HPLC with an AS-H column (hexane/*i*-PrOH, $V:V=70:30$), 1.0 mL/min; minor enantiomer $t_{\text{R}}=10.4$ min, major enantiomer $t_{\text{R}}=15.2$ min.

(*R*)-Ethyl 4-((*S*)-1-acetyl-3-oxoindolin-2-yl)-4-(4-chlorophenyl)-2-oxobutanoate (**4g**): colorless oil, 78 mg (94% yield, 99% *ee*). $[\alpha]_{\text{D}}^{20} -143$ (c 1.0, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) δ : 8.20 (s, 1H), 7.44 (t, $J=8.0$ Hz, 2H), 7.03~6.97 (m, 3H), 6.91 (d, $J=8.0$ Hz, 2H), 4.61 (s, 1H), 4.40~4.35 (m, 3H), 4.03 (s, 1H), 3.41 (s, 1H), 2.65 (s, 3H), 1.40 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 197.9, 192.9, 168.5, 160.3, 153.3, 137.1, 134.0, 133.6, 129.5, 128.1, 125.0, 124.0, 122.9, 118.2, 66.0, 62.8, 41.2, 38.7, 24.1, 13.9; IR (KBr) ν : 3441.4, 2984.2, 1729.0, 1677.0, 1606.5, 1465.7, 1380.3, 1282.6, 1086.6, 1043.8, 761.2 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{21}\text{ClNO}_5$ $[\text{M}+\text{H}]^+$ 414.1103, found 414.1111. The enantiomeric excess was determined by HPLC with an AD-H column (hexane/*i*-PrOH, $V:V=70:30$), 1.0 mL/min; major enantiomer $t_{\text{R}}=15.3$ min, minor enantiomer $t_{\text{R}}=18.9$ min.

(*R*)-Ethyl 4-((*S*)-1-acetyl-3-oxoindolin-2-yl)-4-(4-bromophenyl)-2-oxobutanoate (**4h**): colorless oil, 86 mg (94% yield, 99.5% *ee*). $[\alpha]_{\text{D}}^{20} -113$ (c 1.0, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) δ : 8.18 (s, 1H), 7.44 (t, $J=7.6$ Hz, 2H), 7.13 (d, $J=8.8$ Hz, 2H), 7.02 (t, $J=7.6$ Hz, 1H), 6.85 (d, $J=8.8$ Hz, 2H), 4.61 (s, 1H), 4.40~4.35 (m, 3H), 4.04 (s, 1H), 3.43 (s, 1H), 2.63 (s, 3H), 1.40 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 197.9, 192.9, 168.5, 160.3, 153.2, 137.1, 134.7, 131.1, 129.9, 125.0, 124.0, 122.9, 121.7, 121.0, 118.2, 66.0, 62.8, 41.4, 38.7, 24.2, 13.9; IR (KBr) ν : 3405.4, 2983.8, 1723.5, 1676.2, 1605.1, 1465.2, 1380.4, 1282.8, 1078.1, 1042.6, 760.9 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{21}\text{BrNO}_5$ $[\text{M}+\text{H}]^+$ 458.0598, found 458.0606. The enantiomeric excess was determined by HPLC with an AD-H column (hexane/*i*-PrOH, $V:V=70:30$), 1.0 mL/min; major enantiomer $t_{\text{R}}=14.4$ min, minor enantiomer $t_{\text{R}}=18.2$ min.

(*R*)-Ethyl 4-((*S*)-1-acetyl-3-oxoindolin-2-yl)-2-oxo-4-*p*-tolylbutanoate (**4i**): colorless oil, 74 mg (94% yield, 98% *ee*). $[\alpha]_{\text{D}}^{20} -149$ (c 1.0, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) δ : 8.18 (s, 1H), 7.44~7.36 (m, 2H), 6.96 (t, $J=7.6$ Hz, 1H), 6.85~6.79 (m, 4H), 4.59 (s, 1H), 4.40~4.34 (m, 3H), 4.01 (s, 1H), 3.38 (d, $J=16.4$ Hz, 1H), 2.63 (s, 3H), 2.09 (s, 3H), 1.42 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 198.2, 193.4, 168.6, 160.3, 153.3, 137.2, 136.7, 132.3, 128.6, 128.0, 125.1, 123.7, 122.7, 118.2, 66.2, 62.7, 41.5, 38.9, 24.0, 20.7, 13.9; IR (KBr) ν : 3404.4, 2983.0, 1726.9, 1676.1, 1464.5, 1380.1, 1282.2, 1043.9, 761.5 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{24}\text{NO}_5$ $[\text{M}+\text{H}]^+$ 394.1649, found 394.1655. The enantiomeric excess was determined by HPLC with an AS-H column (hexane/*i*-PrOH, $V:V=70:30$), 1.0 mL/min; minor enantiomer $t_{\text{R}}=11.6$ min, major enantiomer $t_{\text{R}}=14.9$ min.

(*R*)-Ethyl 4-((*S*)-1-acetyl-3-oxoindolin-2-yl)-4-(4-meth-

oxyphenyl)-2-oxobutanoate (**4j**): colorless oil, 70 mg (86% yield, 92% *ee*). $[\alpha]_{\text{D}}^{20} -116$ (*c* 1.0, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) δ : 8.21 (s, 1H), 7.45~7.39 (m, 2H), 6.98 (t, $J=7.6$ Hz, 1H), 6.88 (t, $J=8.8$ Hz, 1H), 6.53 (t, $J=8.8$ Hz, 1H), 4.58 (s, 1H), 4.40~4.35 (m, 3H), 4.00 (s, 1H), 3.61 (s, 3H), 3.42 (d, $J=8.4$ Hz, 1H), 2.65 (s, 3H), 1.40 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 198.4, 193.5, 168.8, 160.4, 158.9, 136.8, 131.0, 129.3, 127.4, 125.2, 123.9, 122.8, 118.2, 114.6, 113.4, 66.4, 62.8, 55.1, 41.2, 39.0, 24.0, 14.0; IR (KBr) ν : 3400.9, 2932.5, 1725.2, 1675.0, 1464.4, 1380.5, 1252.9, 1040.9, 762.0 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{24}\text{NO}_6$ $[\text{M} + \text{H}]^+$ 410.1598, found 410.1587. The enantiomeric excess was determined by HPLC with an AS-H column (hexane/*i*-PrOH, $V: V=70:30$), 1.0 mL/min; minor enantiomer $t_{\text{R}}=18.6$ min, major enantiomer $t_{\text{R}}=24.9$ min.

(*S*)-Ethyl 4-((*S*)-1-acetyl-3-oxoindolin-2-yl)-4-(furan-2-yl)-2-oxobutanoate (**4k**): colorless oil, 63 mg (85% yield, 99% *ee*). $[\alpha]_{\text{D}}^{20} -104$ (*c* 0.5, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) δ : 8.30 (s, 1H), 7.55~7.49 (m, 2H), 7.07 (t, $J=8.0$ Hz, 1H), 6.96 (s, 1H), 5.97~5.94 (m, 2H), 4.58 (s, 1H), 4.38 (dd, $J=7.2$, 14.4 Hz, 2H), 4.20 (s, 2H), 3.42 (d, $J=15.6$ Hz, 1H), 2.56 (s, 3H), 1.41 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 197.4, 192.6, 168.4, 160.1, 153.2, 149.9, 141.9, 136.8, 124.6, 123.8, 123.1, 118.4, 109.8, 108.2, 65.2, 62.8, 37.6, 36.0, 24.0, 13.9; IR (KBr) ν : 3396.2, 2983.6, 1726.2, 1676.8, 1604.9, 1466.8, 1379.8, 1292.2, 1070.7, 760.8 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{20}\text{NO}_6$ $[\text{M} + \text{H}]^+$ 370.1285, found 370.1291. The enantiomeric excess was determined by HPLC with an AS-H column (hexane/*i*-PrOH, $V: V=70:30$), 1.0 mL/min; minor enantiomer $t_{\text{R}}=15.7$ min, major enantiomer $t_{\text{R}}=23.0$ min.

(*R*)-Ethyl 4-((*S*)-1-acetyl-5-bromo-3-oxoindolin-2-yl)-2-oxo-4-phenylbutanoate (**4l**): colorless oil, 90 mg (98% yield, 98% *ee*). $[\alpha]_{\text{D}}^{20} -212$ (*c* 1.0, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) δ : 8.10 (s, 1H), 7.52 (s, 1H), 7.45 (dd, $J=2.0$, 8.8 Hz, 1H), 7.05~7.03 (m, 3H), 6.96~6.94 (m, 2H), 4.63 (s, 1H), 4.42~4.36 (m, 3H), 4.01 (s, 1H), 3.40 (d, $J=20.0$ Hz, 1H), 2.67 (s, 3H), 1.41 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 196.8, 193.5, 168.7, 160.2, 152.1, 139.2, 135.0, 129.0, 129.0, 128.2, 128.0, 126.7, 125.3, 119.9, 116.8, 66.4, 62.9, 42.0, 38.6, 23.9, 14.0; IR (KBr) ν : 3419.8, 2982.8, 1723.3, 1677.7, 1459.5, 1375.7, 1287.0, 1264.8, 1068.2, 1043.4, 701.3 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{21}\text{BrNO}_5$ $[\text{M} + \text{H}]^+$ 458.0598, found 458.0605. The enantiomeric excess was determined by HPLC with an AD-H column (hexane/*i*-PrOH, $V: V=70:30$), 1.0 mL/min; major enantiomer $t_{\text{R}}=11.4$ min, minor enantiomer $t_{\text{R}}=15.0$ min.

(*R*)-Ethyl 4-((*S*)-1-acetyl-5-chloro-3-oxoindolin-2-yl)-2-oxo-4-phenylbutanoate (**4m**): colorless oil, 73 mg (88% yield, 98% *ee*). $[\alpha]_{\text{D}}^{20} -229$ (*c* 1.0, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) δ : 8.15 (s, 1H), 7.36 (s, 1H), 7.31 (dd, $J=2.0$, 8.8 Hz, 1H), 7.04~7.02 (m, 3H), 6.96~6.93 (m, 2H), 4.64 (s, 1H), 4.41~4.36 (m, 3H), 4.01 (s, 1H), 3.40 (d, $J=19.2$ Hz, 1H), 2.67 (s, 3H), 1.41 (t, $J=7.2$ Hz, 3H);

^{13}C NMR (100 MHz, CDCl_3) δ : 197.0, 193.4, 168.6, 160.2, 151.8, 136.4, 135.0, 129.4, 128.2, 128.1, 127.9, 126.2, 122.1, 119.5, 116.8, 66.5, 62.8, 42.0, 38.5, 23.8, 13.9; IR (KBr) ν : 3440.6, 2984.3, 1726.4, 1678.8, 1605.5, 1463.3, 1377.3, 1286.6, 1263.5, 1068.9, 1044.7, 835.6, 701.7 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{21}\text{ClNO}_5$ $[\text{M} + \text{H}]^+$ 414.1103, found 414.1113. The enantiomeric excess was determined by HPLC with an AS-H column (hexane/*i*-PrOH, $V: V=70:30$), 1.0 mL/min; major enantiomer $t_{\text{R}}=10.2$ min, minor enantiomer $t_{\text{R}}=13.4$ min.

(*R*)-Ethyl 4-((*S*)-1-acetyl-5-methyl-3-oxoindolin-2-yl)-2-oxo-4-phenylbutanoate (**4n**): colorless oil, 70 mg (89% yield, 98% *ee*). $[\alpha]_{\text{D}}^{20} -197$ (*c* 1.0, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) δ : 8.07 (s, 1H), 7.20 (d, $J=8.0$ Hz, 2H), 7.01~6.95 (m, 5H), 4.58 (s, 1H), 4.40~4.35 (m, 3H), 4.02 (s, 1H), 3.42 (s, 1H), 2.64 (s, 3H), 2.22 (s, 3H), 1.40 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 198.1, 193.4, 168.4, 160.4, 151.6, 137.9, 135.4, 133.7, 128.2, 128.0, 127.6, 125.2, 122.3, 118.0, 66.3, 62.7, 41.9, 38.8, 23.9, 20.5, 13.9; IR (KBr) ν : 3397.2, 2926.3, 1728.0, 1673.0, 1487.7, 1378.8, 1282.5, 1068.5, 1043.0, 704.2 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{24}\text{NO}_5$ $[\text{M} + \text{H}]^+$ 394.1649, found 394.1659. The enantiomeric excess was determined by HPLC with an AS-H column (hexane/*i*-PrOH, $V: V=70:30$), 1.0 mL/min; minor enantiomer $t_{\text{R}}=15.5$ min, major enantiomer $t_{\text{R}}=19.9$ min.

(*R*)-Ethyl 4-((*S*)-1-acetyl-7-methyl-3-oxoindolin-2-yl)-2-oxo-4-phenylbutanoate (**4o**): colorless oil, 68 mg (86% yield, 90% *ee*). $[\alpha]_{\text{D}}^{20} -291$ (*c* 1.0, CH_2Cl_2); ^1H NMR (400 MHz, CDCl_3) δ : 7.28 (d, $J=7.2$ Hz, 1H), 7.16 (d, $J=7.2$ Hz, 1H), 6.96~6.93 (m, 6H), 4.62 (d, $J=8.8$ Hz, 1H), 4.38 (dd, $J=7.2$, 14.4 Hz, 2H), 4.23 (dd, $J=10.4$, 20.0 Hz, 1H), 3.35 (dd, $J=3.6$, 20.0 Hz, 1H), 2.56 (s, 3H), 2.07 (s, 3H), 1.40 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 199.0, 193.4, 168.8, 160.3, 153.0, 138.5, 135.7, 130.6, 128.4, 128.1, 127.7, 127.6, 125.3, 120.2, 68.0, 62.8, 42.7, 38.5, 23.2, 20.2, 14.0; IR (KBr) ν : 3442.0, 2981.8, 2927.7, 1726.6, 1713.4, 1686.6, 1590.6, 1453.7, 1375.2, 1274.2, 1038.2, 1044.2, 701.0 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{24}\text{NO}_5$ $[\text{M} + \text{H}]^+$ 394.1649, found 394.1638. The enantiomeric excess was determined by HPLC with an AD-H column (hexane/*i*-PrOH, $V: V=70:30$), 1.0 mL/min; major enantiomer $t_{\text{R}}=11.1$ min, minor enantiomer $t_{\text{R}}=15.8$ min.

4.2 Representative procedure for the synthesis of compound 5

To a solution of **4e** (0.1 mmol, 1.0 equiv.) in 1.0 mL of tetrahydrofuran (THF) cooled to 0 °C were added 0.125 mL of concentrated AcOH and 11 mg (0.2 mmol, 2.0 equiv.) of NaCNBH_3 . The reaction mixture was warmed up to room temperature. After **4e** was consumed, 3 mL of brine was added and the pH was adjusted to 7 with saturated NaHCO_3 solution. The aqueous layer was extracted with ethyl acetate (10 mL $\times$ 3) and the combined organic layers were dried over Na_2SO_4 . After evaporation of the solvents under vacuum, the residue was purified by flash

column chromatography to obtain (4*R*)-ethyl 4-((*S*)-1-acetyl-3-oxoindolin-2-yl)-4-(3-chlorophenyl)-2-hydroxybutanoate (**5**) (**5a** : **5b** = 1 : 1, 75% yield). Colorless oil. ^1H NMR (400 MHz, CDCl_3) δ : 8.17 (s, 2H), 7.51 (s, 1H), 7.46~7.40 (m, 2H), 7.04~6.84 (m, 10H), 4.68 (s, 2H), 4.55 (s, 1H), 4.31~4.23 (m, 2H), 4.21~4.11 (m, 1H), 4.06~3.95 (m, 2H), 3.83~3.67 (m, 2H), 3.16 (s, 1H), 2.96 (s, 2H), 2.77~2.71 (m, 1H), 2.51 (s, 3H), 2.49 (s, 3H), 2.38~2.27 (m, 2H), 1.33 (t, $J=7.2$ Hz, 3H), 1.27 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 198.0, 174.7, 174.5, 168.8, 168.1, 152.8, 138.6, 136.8, 133.9, 133.6, 129.2, 128.9, 128.8, 128.5, 127.6, 127.6, 127.0, 126.8, 123.9, 123.0, 118.3, 68.5, 68.5, 68.0, 67.0, 62.0, 42.9, 42.5, 33.9, 33.0, 24.8, 24.0, 14.1, 14.0; IR (KBr) ν : 3413.9, 2933.3, 1717.4, 1673.7, 1466.9, 1383.3, 1295.2, 1210.3, 1108.3, 1024.6, 761.5 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{23}\text{ClNO}_5$ [$\text{M}+\text{H}$] $^+$ 416.1259, found 416.1248.

Supporting Information Crystallography data, copies of NMR and HRMS as well as chiral HPLC chromatograms of the prepared active 2-substituted 1-acetylindolin-3-ones. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Harutyunyan, S. R.; Hartog, T. D.; Geurts, K.; Minnaard, A. J.; Feringa, B. L. *Chem. Rev.* **2008**, *108*, 2824.
(b) Dalko, P. I.; Moisan, L. *Angew. Chem., Int. Ed.* **2004**, *43*, 5138.
(c) Li, N.; Xi, G.-H.; Wu, Q.-H.; Liu, W.-H.; Ma, J.-J.; Wang, C. *Chin. J. Org. Chem.* **2009**, *29*, 1018 (in Chinese).
(李宁, 郝国宏, 吴秋华, 刘伟华, 马晶军, 王春, 有机化学, **2009**, *29*, 1018.)
- [2] (a) Sulzer-Mossé, S.; Alexakis, A. *Chem. Commun.* **2007**, 3123.
(b) Tsogoeva, S. B. *Eur. J. Org. Chem.* **2007**, 1701.
(c) Guan, X.-Y.; Yang, L.-P.; Hu, W.-H. *Angew. Chem., Int. Ed.* **2010**, *49*, 2190.
(d) Li, M.-F.; Guo, X.; Jin, W.-F.; Zheng, Q.; Liu, S.-Y.; Hu, W.-H. *Chem. Commun.* **2016**, 52, 2736.
(e) Jia, Y.-B.; Zhang, R.; Xie, B.; Chen, G.-S.; Liu, T.-S.; Chen, Y.-X. *J. Guangzhou Univ. (Nat. Sci. Ed.)* **2009**, *13*, 47 (in Chinese).
(贾永兵, 张睿, 谢彬, 陈国术, 刘天穗, 陈亿新, 广州大学学报 (自然科学版), **2009**, *13*, 47.)
- [3] (a) Zhou, L.; Lin, L.-L.; Wang, W.-T.; Ji, J.; Liu, X.-H.; Feng, X.-M. *Chem. Commun.* **2010**, 46, 3601.
(b) Herrera, R. P.; Monge, D.; Zamora, E. M.; Fernández, R.; Las-saletta, J. M. *Org. Lett.* **2007**, *11*, 3303.
(c) Liu, S.-S.; Xu, Z.-H.; Wang, X.; Zhu, H.-R.; Wang, M.-C. *J. Org. Chem.* **2019**, *84*, 13881.
(d) Zhang, M.-L.; Wu, Z.-J.; Zhao, J.-Q.; Luo, Y.; Xu, X.-Y.; Zhang, X.-M.; Yuan, W.-C. *Org. Lett.* **2016**, *18*, 5110.
(e) Zhi, Y.-L.; Huang, J.-H.; Liu, N.; Lu, T.; Dou, X.-W. *Org. Lett.* **2017**, *19*, 2378.
- [4] (a) Rueping, M.; Nachtsheim, B.; Moreth, S. A.; Bolte, M. *Angew. Chem., Int. Ed.* **2008**, *47*, 593.
(b) Yang, W.; Pan, X.-J.; Yang, D.-Q. *Chin. J. Org. Chem.* **2015**, *35*, 1216 (in Chinese).
(杨文, 潘雪静, 杨定乔, 有机化学, **2015**, *35*, 1216.)
- [5] (a) Baran, P. S.; Corey, E. J. *J. Am. Chem. Soc.* **2002**, *124*, 7904.
(b) Karadeolian, A.; Kerr, M. A. *Angew. Chem., Int. Ed.* **2010**, *49*, 1133.
(c) Higuchi, K.; Sato, Y.; Tsuchimochi, M.; Sugiura, K.; Hatori, M.; Kawasaki, T. *Org. Lett.* **2009**, *11*, 197.
- [6] (a) Witkop, B. J. *Am. Chem. Soc.* **1950**, *72*, 614.
(b) Lednicer, D. L.; Emmert, D. E. *J. Heterocycl. Chem.* **1970**, *7*, 575.
(c) Liu, Y.; McWhorter, W. W. *J. Org. Chem.* **2003**, *68*, 2618.
(d) Buller, M. J.; Cook, T. G.; Kobayashi, Y. *Heterocycles* **2007**, *72*, 163.
(e) Higuchi, K.; Sato, Y.; Kojima, S.; Tsuchimochi, M.; Sugiura, K.; Hatori, M.; Kawasaki, T. *Tetrahedron* **2010**, *66*, 1236.
- [7] (a) Higuchi, K.; Masuda, K.; Koseki, T.; Hatori, M.; Sakamoto, M.; Kawasaki, T. *Heterocycles* **2007**, *73*, 641.
(b) Yin, Q.; You, S.-L. *Chem. Sci.* **2011**, *2*, 1344.
(c) Sun, W.-S.; Hong, L.; Wang, R. *Chem.-Eur. J.* **2011**, *17*, 6030.
(d) Rueping, M.; Raja, S.; Nunez, A. *Adv. Synth. Catal.* **2011**, *353*, 563.
(e) Liu, Y.-Z.; Cheng, R.-L.; Xu, P.-F. *J. Org. Chem.* **2011**, *76*, 2884.
(f) Liu, Y.-Z.; Zhang, J.; Xu, P.-F.; Luo, Y.-C. *J. Org. Chem.* **2011**, *76*, 7551.
(g) Jin, C.-Y.; Wang, Y.; Liu, Y.-Z.; Shen, C.; Xu, P.-F. *J. Org. Chem.* **2012**, *77*, 11307.
(f) Zhao, Y.-L.; Wang, Y.; Cao, J.; Liang, Y.-M.; Xu, P.-F. *Org. Lett.* **2014**, *16*, 2438.
(g) Chen, S.-R.; Wang, Y.-M.; Zhou, Z.-H. *J. Org. Chem.* **2016**, *81*, 11432.
- [8] (a) Sigman, M. S.; Vachal, P.; Jacobsen, E. *Angew. Chem., Int. Ed.* **2000**, *39*, 1279.
(b) Schreiner, P. R. *Chem. Soc. Rev.* **2003**, *32*, 289.
(c) Zuend, S. J.; Matthew, P. C.; Mathieu, P. L.; Jacobsen, E. N. *Nature* **2009**, *461*, 968.
(d) Zhang, C.-H.; Yang, J.; Zhou, W.-Q.; Tan, Q.-Y.; Yang, Z.; He, L.; Zhang, M. *Org. Lett.* **2019**, *21*, 8620.
(e) Zhang, Y.-P.; You, Y.; Zhao, J.-Q.; Zhang, X.-M.; Xu, X.-Y.; Yuan, W.-C. *J. Org. Chem.* **2019**, *84*, 7984.
(f) Li, X.-L.; Kong, X.-W.; Yang, S.; Meng, M.; Zhan, X.-Y.; Zeng, M.; Fang, X.-Q. *Org. Lett.* **2019**, *21*, 1979.
(g) Lu, N.; Fang, Y.-H.; Gao, Y.; Wei, Z.-L.; Cao, J.-G.; Liang, D.-P.; Lin, Y.-J.; Duan, H.-F. *J. Org. Chem.* **2018**, *83*, 793.
(h) Ma, Z.-W.; Liu, X.-F.; Liu, J.-T.; Tao, J.-C. *Chin. J. Org. Chem.* **2018**, *38*, 183 (in Chinese).
(马志伟, 刘晓峰, 刘俊桃, 陶京朝, 有机化学, **2018**, *38*, 183.)

(Zhao, C.)