碘促进苯乙酮与 2-(芳基乙烯基)苯胺反应合成 2-酰基喹啉

杜兴鹏 张玺 侯飞 刘小军 王喜存* 权正军*

(西北师范大学化学化工学院 兰州 730070)

摘要 报道了碘存在下 2-(芳基乙烯基)苯胺和苯乙酮通过环化反应得到 2-酰基喹啉的合成方法, 该法反应条件温和、无需金属催化剂的存在, 底物适用范围广, 产率良好或适中, 为多样化 2-酰基喹啉衍生物的合成提供了一种有效途径。

关键词 2-(芳基乙烯基)苯胺; 苯乙酮; 2-酰基喹啉; 碘催化

Iodine-Mediated Synthesis of 2-Acylquinoline from Acetophenone and 2-(Arylvinyl)aniline

Du, Xingpeng Zhang, Xi Hou, Fei Liu, Xiaojun

Wang, Xicun* Quan, Zhengjun*

(College of Chemistry and Chemical Engineering, Northwest Normal University, Lanzhou 730070)

Abstract The cyclization reaction between 2-(arylvinyl)aniline and acetophenone in presence of iodine was reported, which offered an efficient and cost effective strategy to generate 2-acylquinoline derivatives under mild and metal-free conditions.

Keywords 2-(arylvinyl)aniline; acetophenone; 2-acylquinoline; metal-free

1 Introduction

The quinoline skeleton is a common structural unit mainly found in various natural compounds and has excellent pharmacological activity and biological activity, especially in anticancer drugs.^[1] Furthermore, in terms of organic synthesis, the quinoline ring is of great importance as a guiding group and some common ligands.^[2] Because of its prominence in the field of chemistry, it is still a challenging research hotspot to develop convenient, efficient and green synthetic methods for the preparation of quinoline and its derivatives.^[3]

Based on the efforts made in quinoline, many classical cases have been developed in the past decades, such as the synthesis of doebner-vonmiller,^[4] combes^[5] and povarov-reaction^[6]. Recently, Li *et al.*^[7] reported that the highly substituted quinoline product was obtained by oxidative tandem cyclization of α -(*N*-arylamino)carbonyl compounds with aryl olefins under the action of copper. Jiang group^[8] reported a new approach that 2,3-disubstituted quinoline was obtained by Pd-catalyzed oxidative cyclization of *o*-vinylanilines and alkynes containing molecular oxygen. Ma *et al.*^[9] have developed the ruthenium-cata-

lyzed construction of quinoline from 2-alkenylaniline and sulfoxonium ylide. Although many synthetic methods of quinoline have been reported, it is deficient because raw material preparation is relatively complicated^[10]. Recently, Wu group^[11] reported the non-metal promoted synthesis of quinoline derivatives from alkenes and aryl amines in the presence of iodine. The development of non-metals promotes the synthesis of quinoline rings with great potential. These cases need to be further enriched and expanded. Our group is interested in the preparation of some *N*-heterocycle derivatives using 2-(arylvinyl)aniline.^[12] Owing to its potential ability to synthesize heterocyclic compounds,^[13] we have developed an important substructure of 2-acylquinoline route which directly obtained from 2-(arylvinyl)aniline and acetophenone mediated by iodine.

2 Results and discussion

Initially, the reaction conditions of 4-methylacetophenone (**1a**) and 2-(1-phenylvinyl) aniline (**2a**) were screened in dimethyl sulfoxide (DMSO) at 80 °C (Table 1). When 2,3-dichloro-5,6-dicyanobenzoquinone (DDQ), K₂S₂O₈, and H₂O₂ were used as oxidants, the target product was not obtained (Entries 1~3). Subsequently, when 0.5 equiv. of

* Corresponding authors. E-mail: quanzhengjun@hotmail.com; wangxicun@nwnu.edu.cn

Received November 24, 2019; revised January 9, 2020; published online January 15, 2020.

Project supported by the National Natural Science Foundation of China (No. 21562036).

国家自然科学基金(No. 21562036)资助项目.

I_2 was used as the oxidant, the target product **3a** was achieved in 47% yield (Entry 5). Then the amount of I_2 was explored, and it was found that when I_2 increased to 1.5 or 2 equiv., the target product **3a** can be obtained in yields of 76% and 80% (Entries 7, 8). Taking into account the economic benefits and green policy, 1.5 equiv. of I_2 was chosen as the best quantity (Entry 7). The screening of temperature suggested that 80 °C was the best choice for the transformation. Next, some reaction solvents were screened. It was found that DMSO was the most excellent solvent. The reaction did not occur smoothly in other solvents, such as CH_3CN , 1,4-dioxane, *N,N*-dimethylformamide (DMF) (Entries 11~13). When bases were added into the reaction system, the target product was not obtained (Entries 14~17). Finally, when DMSO (2 equiv.) was used as the additive and CH_3CN as solvent, the target product was not obtained under standard conditions. At the same time, other iodine sources were optimized. It was found that the target product could not be obtained (Entries 20~22). Thus, the optimal conditions were got. When DMSO was used as solvent, I_2 was used as an oxidant, the compound **3a** was isolated in yield of 76% by reacting for 12 h at 80 °C (Table 1, Entry 7).

Table 1 Optimized reaction conditions^a

Entry	Oxidant (equiv.)	<i>T</i> /°C	Solvent	Base	Yield ^b
1	DDQ (1.0)	80	DMSO	—	Trace
2	K ₂ S ₂ O ₈ (1.0)	80	DMSO	—	Trace
3	H ₂ O ₂ (1.0)	80	DMSO	—	Trace
4	I_2 (0.2)	80	DMSO	—	Trace
5	I_2 (0.5)	80	DMSO	—	47%
6	I_2 (1.0)	80	DMSO	—	63%
7	I_2 (1.5)	80	DMSO	—	76%
8	I_2 (2.0)	80	DMSO	—	80%
9	I_2 (1.5)	60	DMSO	—	67%
10	I_2 (1.5)	40	DMSO	—	52%
11	I_2 (1.5)	80	Dioxane	—	Trace
12	I_2 (1.5)	80	DMF	—	Trace
13	I_2 (1.5)	80	CH_3CN	—	Trace
14	I_2 (1.5)	80	DMSO	Cs ₂ CO ₃	Trace
15	I_2 (1.5)	80	DMSO	KOH	Trace
16	I_2 (1.5)	80	DMSO	K ₂ CO ₃	Trace
17	I_2 (1.5)	80	DMSO	<i>t</i> -BuONa	Trace
18 ^c	I_2 (1.5)	80	CH_3CN	—	N.R.
19	I_2 (1.5)	100	DMSO	—	78%
20	KI (1.5)	80	DMSO	—	NR
21	NH ₄ I (1.5)	80	DMSO	—	NR
22	NaI (1.5)	80	DMSO	—	NR

^a Reaction conditions: **1a** (0.12 mmol), **2a** (0.1 mmol), I_2 (1.5 equiv.), DMSO (2 mL), 80 °C, 12 h. ^b Isolated yields. ^c DMSO (2 equiv.), CH_3CN (2 mL).

With the optimized conditions in hand, the scope of the

reaction was explored (Table 2). The result shows that a wide range of acetophenones and 2-(arylvinyl)anilines are effective substrates in this system. When the para-substituted acetophenone is electron-donating group or electron-withdrawing group, the corresponding target product can be obtained in a superior yield. Fortunately, the substrate that contains electron-rich group [$N(CH_3)_2$] or electron-withdrawing group (NO_2) can obtain the target product (**3l** and **3k**) in 70% and 61% yields. Similarly, *ortho*- and *meta*-substituted acetophenones have also successfully provided quinoline products with excellent yields. It is noteworthy that 1-naphthylethyl ketone and 2-acetylfuran also obtain the target products in moderate yields (**3m**, **3n**). Under the standard reaction conditions, the substrate containing OH is compatible, and the target product (**3x**) is obtained in 53% yield. For 2-(1-arylvinyl)aniline, the electron-donating group (CH_3 , OMe, Et) (**3c**, **3p**, **3q**) or electron-withdrawing group (Cl) (**3y**~**3ab**) can obtain the target production in moderate to good yield. All in all, the position of substituents does not affect the yield. When 2-(1-arylvinyl)aniline is disubstituted, the target products can also be obtained in moderate yields (**3ad**~**3ag**). Due to some target products contain halogens and other active sites, we can continue to further studies^[14], such as Heck and Suzuki coupling reactions and so on.

To verify the possible mechanisms, a series of controlled experiments have been done (Scheme 1). Firstly, aryl methyl ketone (**1a**, 1.0 mmol) and I_2 (1.5 mmol) were heated in DMSO at 80 °C to provide phenylglyoxal **1b** and corresponding hydrate **1c** (Scheme 1a). When acetophenone was replaced by iodoacetophenone **1d** under the standard conditions, the target product **3c** was obtained by reacting with **2a**, and the yield was 78% (Scheme 1b). These results confirm that benzoyl iodide and phenylglyoxal are intermediates of the reaction. Next, intermediate **1c** also obtained the target product in 80% yield under standard conditions (Scheme 1c), and the target product was not obtained without I_2 . What's more, the free radical inhibitor 2,2,6,6-tetramethylpiperidinooxy (TEMPO) was added to obtain the target product (**3c**) in 67% yield under standard conditions, which proved that the reaction was not a free radical process (Scheme 1d). In addition, **3c** was obtained in good yield under argon atmosphere (Scheme 1e). This result shows that the oxidation process is carried out under the promotion of iodine. In summary, I_2 plays a vital role in this reaction.

Based on our experimental studies and related literature^[15] reported, we proposed a reasonable reaction mechanism in Scheme 2. Acetophenone is converted to phenylglyoxal by iodization/Kornblum oxidation sequence, meanwhile HI and DMS are released. The obtained aldehyde **1b** reacts with amine **2a** to obtain the corresponding imine **B**. Intermediate **B** reacts with HI to form intermediate **C**, then intermediate **C** is cyclized intramolecularly to form intermediate **D**. Finally, the intermediate **D** is deprotonated to obtain **E**. By aromatization of **E** under the action of iodine obtain the final product **3d**.

Table 2 Scope of 2-acylquinolines of vinyl anilines with acetophenone

 3a, 76%	 3b, 74%
 3c, R' = H, 82%	 3d, R' = 4-CH ₃ , 75%
 3e, R' = 3-CH ₃ , 73%	 3f, R' = 4-OCH ₃ , 79%
 3g, R' = 4-Br, 71%	 3h, R' = 3-Br, 70%
 3i, R' = 2-Br, 67%	 3j, R' = 4-I, 66%
 3k, R' = 4-NO ₂ , 61%	 3l, R' = 4-N(CH ₃) ₂ , 70%
 3m, 56%	 3n, 67%
 3o, 73%	
 3p, 76%	 3q, R'' = H, 78%
 3r, R'' = 4-CH ₃ , 80%	 3s, R'' = 3-CH ₃ , 76%
 3t, R'' = 4-Br, 79%	 3u, R'' = 3-Br, 74%
 3v, R'' = 2-Br, 66%	 3w, R'' = 4-I, 63%
 3x, R'' = 4-OH, 53%	 3y, 69%
 3z, 64%	 3aa, 60%
 3ab, 63%	
 3ac, 52%	 3ad, 66%
 3ae, 59%	 3af, 53%
 3ag, 64%

Scheme 1 Control experimental

Scheme 2 Proposed mechanism

3 Conclusions

In summary, we have developed an iodine-mediated strategy for the synthesis of aryl (4-arylquinolin-2-yl) ketone. In this reaction system, without metal participation and additional additives, the quinoline derivatives are synthesized easily, efficiently, and have good substrates tolerance. This method not only enriches the non-metal libraries involved in the synthesis of quinoline derivatives but also explores the cyclization potential of vinyl aniline.

4 Experimental

4.1 General information

Unless otherwise noted, all commercially available compounds were used as provided without further purification. ^1H NMR and ^{13}C NMR analyses were performed on a Varian Mercury plus-400 and Agilent 600 MHz DD2 instruments with CDCl_3 and $\text{DMSO}-d_6$ as solvent and tetramethylsilane (TMS) as the internal standard. Flash chromatography was performed using 200~300 mesh silica gel with the indicated solvent system according to standard techniques. Analytical thin-layer chromatography (TLC) was performed on pre-coated, glass-backed silica gel plates. Melting points were measured with an XT-4 apparatus. High-resolution mass spectra (HRMS) (ESI) were obtained with a 6502B Q-TOF mass spectrometer. Column chromatography was performed generally on silica gel (200~300 mesh) and TLC analyses were conducted on silica gel GF254 plates. All reagents were used directly without any further purification unless otherwise specified.

4.2 General procedure

A sealed tube was charged with acetophenone (**1a**) (0.12 mmol) and iodine (0.15 mmol) at room temperature, and DMSO (2 mL) was added. The resulting mixture was stirred at 80 °C after disappearance of the reactant (monitored by TLC), then 4-methyl-2-(1-phenylvinyl)aniline (**2a**) (0.1 mmol) was added at 80 °C for another 12 h. After the reaction completed, 40 mL of water was added to the mixture, and then the mixture was extracted with EtOAc (40 mL $\times$ 3). The extract was washed with 10% $\text{Na}_2\text{S}_2\text{O}_3$ solution, dried over anhydrous Na_2SO_4 and evaporated. The residue was purified by column chromatography on silica gel [$V(\text{petroleum ether}) : V(\text{EtOAc}) = 20 : 1$] to yield the desired product **3c** as yellow solid (26.5 mg, 82% yield).

(4-Phenylquinolin-2-yl)(*p*-tolyl)methanone (**3a**): 24.6 mg, 76% yield. Yellow solid, m.p. 92~94 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.27 (d, $J=8.4$ Hz, 1H), 8.18 (d, $J=8.4$ Hz, 2H), 8.02 (s, 1H), 8.01 (d, $J=8.4$ Hz, 1H), 7.80~7.76 (m, 1H), 7.62~7.58 (m, 1H), 7.58~7.49 (m, 5H), 7.33 (d, $J=7.8$ Hz, 2H), 2.46 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 193.55, 154.62, 149.55, 147.33, 143.98, 137.70, 133.56, 131.61, 130.91, 129.80, 129.58, 128.91, 128.64, 128.62, 128.29, 127.36, 125.78, 120.96, 21.78; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{18}\text{NO}$ [$\text{M}+\text{H}$] $^+$ 324.1383, found 324.1384.

(4-Phenylquinolin-2-yl)(*m*-tolyl)methanone (**3b**): 23.9 mg, 74% yield. Yellow solid, m.p. 72~74 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.27 (d, $J=8.4$ Hz, 1H), 8.05~7.99 (m, 4H), 7.80~7.77 (m, 1H), 7.63~7.59 (m, 1H), 7.58~7.51 (m, 5H), 7.44 (d, $J=7.2$ Hz, 1H), 7.40 (t, $J=7.8$ Hz, 1H), 2.45 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.23, 154.47, 149.59, 147.34, 137.95, 137.68, 136.15, 133.89, 131.68, 130.94, 129.82, 129.58, 128.84, 128.64, 128.34, 128.00, 127.40, 125.79, 120.92, 21.41; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{18}\text{NO}$ [$\text{M}+\text{H}$] $^+$ 324.1383, found 324.1386.

(6-Methyl-4-phenylquinolin-2-yl)(phenyl)methanone (**3c**): 26.5 mg, 82% yield. Yellow solid, m.p. 133~135 °C (lit.^[11d] 125~128 °C); ^1H NMR (600 MHz, CDCl_3) δ : 8.26 (d, $J=7.8$ Hz, 2H), 8.15 (d, $J=8.4$ Hz, 1H), 8.02 (s, 1H), 7.74 (s, 1H), 7.64~7.60 (m, 2H), 7.56 (d, $J=4.2$ Hz, 4H), 7.52 (t, $J=7.8$ Hz, 3H), 2.51 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 193.89, 153.41, 148.78, 145.92, 138.79, 137.90, 136.34, 132.90, 132.13, 131.44, 130.66, 129.54, 128.61, 128.50, 128.08, 127.44, 124.50, 121.06, 22.00; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{18}\text{NO}$ [$\text{M}+\text{H}$] $^+$ 324.1383, found 324.1385.

(6-Methyl-4-phenylquinolin-2-yl)(*p*-tolyl)methanone (**3d**): 25.3 mg, 75% yield. Yellow solid, m.p. 124~126 °C (lit.^[11d] 113~115 °C); ^1H NMR (600 MHz, CDCl_3) δ : 8.18 (d, $J=7.8$ Hz, 2H), 8.16 (d, $J=9.0$ Hz, 1H), 7.99 (s, 1H), 7.74 (s, 1H), 7.61 (dd, $J=8.4$, 1.8 Hz, 1H), 7.56~7.49 (m, 5H), 7.32 (d, $J=7.8$ Hz, 2H), 2.51 (s, 3H), 2.46 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 193.57, 153.76, 148.71, 145.91, 143.80, 138.64, 137.94, 133.73, 132.08, 131.59, 130.63, 129.54, 128.85, 128.60, 128.48, 127.38, 124.49, 121.09, 21.99, 21.74; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{20}\text{NO}$ [$\text{M}+\text{H}$] $^+$ 338.1539, found 338.1534.

(6-Methyl-4-phenylquinolin-2-yl)(*m*-tolyl)methanone (**3e**): 24.6 mg, 73% yield. Yellow solid, m.p. 114~116 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.16 (d, $J=8.4$ Hz, 1H), 8.06~8.03 (m, 2H), 8.00 (s, 1H), 7.75 (s, 1H), 7.62 (dd, $J=8.4$, 1.8 Hz, 1H), 7.56 (s, 2H), 7.56 (d, $J=1.2$ Hz, 2H), 7.53~7.50 (m, 1H), 7.45~7.42 (m, 1H), 7.40 (t, $J=7.8$ Hz, 1H), 2.51 (s, 3H), 2.45 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.22, 153.63, 148.73, 145.95, 138.71, 137.93, 137.86, 136.37, 133.73, 132.10, 131.67, 130.68, 129.54, 128.82, 128.61, 128.49, 127.94, 127.42, 124.50, 121.05, 21.99, 21.39; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{20}\text{NO}$ [$\text{M}+\text{H}$] $^+$ 338.1539, found 338.1534.

(4-Methoxyphenyl)(6-methyl-4-phenylquinolin-2-yl)-methanone (**3f**): 27.9 mg, 79% yield. Yellow solid, m.p. 131~133 °C (lit.^[11d] 127~129 °C); ^1H NMR (400 MHz, CDCl_3) δ : 8.33 (d, $J=8.4$ Hz, 2H), 8.17 (d, $J=8.4$ Hz, 1H), 7.99 (s, 1H), 7.76 (s, 1H), 7.62 (dd, $J=8.4$, 2.0 Hz, 1H), 7.57~7.50 (m, 5H), 7.01 (d, $J=8.4$ Hz, 2H), 3.90 (s, 3H), 2.51 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 192.31, 163.65, 154.08, 148.71, 145.84, 138.57, 137.94, 133.92, 132.11, 130.55, 129.56, 129.12, 128.62, 128.50, 127.30, 124.51, 121.18, 113.50, 55.48, 22.00; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{20}\text{NO}_2$ [$\text{M}+\text{H}$] $^+$ 354.1489, found 354.1490.

(4-Bromophenyl)(6-methyl-4-phenylquinolin-2-yl)-methanone (**3g**): 28.5 mg, 71% yield. Yellow solid, m.p.

179~180 °C (lit.^[11d] 169~72 °C); ¹H NMR (600 MHz, CDCl₃) δ: 8.18 (d, *J*=8.4 Hz, 2H), 8.13 (d, *J*=8.4 Hz, 1H), 8.03 (s, 1H), 7.74 (s, 1H), 7.66 (d, *J*=8.4 Hz, 2H), 7.62 (dd, *J*=8.4, 1.8 Hz, 1H), 7.58~7.50 (m, 5H), 2.51 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 192.70, 152.82, 148.96, 145.84, 139.10, 137.76, 135.09, 133.02, 132.29, 131.38, 130.60, 129.51, 128.65, 128.58, 128.21, 127.53, 124.54, 120.98, 22.05; HRMS (ESI) calcd for C₂₃H₁₇BrNO [M+H]⁺ 402.0488, found 402.0485.

(3-Bromophenyl)(6-methyl-4-phenylquinolin-2-yl)-methanone (**3h**): 28.1 mg, 70% yield. Yellow solid, m.p. 153~154 °C; ¹H NMR (600 MHz, CDCl₃) δ: 8.45 (t, *J*=1.8 Hz, 1H), 8.23 (dt, *J*=7.8, 1.2 Hz, 1H), 8.15 (d, *J*=8.4 Hz, 1H), 8.05 (s, 1H), 7.75~7.73 (m, 2H), 7.63 (dd, *J*=9.0, 2.4 Hz, 1H), 7.56~7.52 (m, 5H), 7.39 (t, *J*=7.8 Hz, 1H), 2.51 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 192.17, 152.64, 148.99, 145.91, 139.16, 138.23, 137.78, 135.58, 134.30, 132.29, 130.72, 130.02, 129.58, 129.51, 128.65, 128.58, 127.59, 124.54, 122.25, 120.92, 22.03; HRMS (ESI) calcd for C₂₃H₁₇BrNO [M+H]⁺ 402.0488, found 402.0485.

(2-Bromophenyl)(6-methyl-4-phenylquinolin-2-yl)-methanone (**3i**): 26.9 mg, 67% yield. Yellow solid, m.p. 145~147 °C; ¹H NMR (600 MHz, CDCl₃) δ: 8.13 (s, 1H), 8.03 (d, *J*=9.0 Hz, 1H), 7.73 (s, 1H), 7.66 (dd, *J*=8.4, 1.2 Hz, 1H), 7.59 (dd, *J*=7.2, 1.2 Hz, 1H), 7.57 (d, *J*=1.8 Hz, 1H), 7.56 (d, *J*=4.2 Hz, 4H), 7.54~7.50 (m, 1H), 7.47~7.43 (m, 1H), 7.40~7.36 (m, 1H), 2.49 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 196.09, 151.73, 148.76, 146.48, 140.43, 139.33, 137.88, 133.00, 132.07, 131.30, 131.04, 130.44, 129.56, 128.62, 128.53, 127.85, 126.81, 124.49, 120.59, 120.11, 22.03; HRMS (ESI) calcd for C₂₃H₁₇BrNO [M+H]⁺, 402.0488, found 402.0484.

(4-Iodophenyl)(6-methyl-4-phenylquinolin-2-yl)-methanone (**3j**): 26.5 mg, 66% yield. Yellow solid, m.p. 174~176 °C; ¹H NMR (600 MHz, CDCl₃) δ: 8.13 (d, *J*=8.4 Hz, 1H), 8.03 (s, 1H), 8.01 (d, *J*=8.4 Hz, 2H), 7.89 (d, *J*=8.4 Hz, 2H), 7.74 (s, 1H), 7.62 (dd, *J*=8.4, 1.8 Hz, 1H), 7.58~7.50 (m, 5H), 2.51 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 192.93, 152.84, 148.94, 145.88, 139.05, 137.81, 137.36, 135.72, 132.86, 132.25, 130.63, 129.50, 128.63, 128.55, 127.55, 124.54, 120.94, 101.01, 22.01; HRMS (ESI) calcd for C₂₃H₁₇I₂NO [M+H]⁺ 450.0349, found 450.0353.

(6-Methyl-4-phenylquinolin-2-yl)(4-nitrophenyl)-methanone (**3k**): 22.6 mg, 61% yield. Yellow solid, m.p. 174~176 °C (lit.^[11d] 157~160 °C); ¹H NMR (600 MHz, CDCl₃) δ: 8.44 (d, *J*=9.0 Hz, 2H), 8.36 (d, *J*=9.0 Hz, 2H), 8.13 (s, 1H), 8.11 (d, *J*=8.4 Hz, 1H), 7.76 (s, 1H), 7.64 (dd, *J*=9.0, 1.8 Hz, 1H), 7.59~7.51 (m, 5H), 2.52 (s, 4H); ¹³C NMR (150 MHz, CDCl₃) δ: 192.02, 151.90, 149.96, 149.22, 145.95, 141.71, 139.64, 137.64, 132.48, 132.30, 130.67, 129.48, 128.70, 128.69, 127.78, 124.62, 123.02, 120.73, 22.06; HRMS (ESI) calcd for C₂₃H₁₇N₂O₃ [M+H]⁺ 369.1234, found 369.1239.

(4-(Dimethylamino)phenyl)(6-methyl-4-phenylquinolin-2-yl)methanone (**3l**): 25.6 mg, 70% yield. Yellow solid,

m.p. 145~147 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.25 (d, *J*=8.8 Hz, 2H), 8.18 (d, *J*=8.8 Hz, 1H), 7.92 (s, 1H), 7.74 (s, 1H), 7.61 (dd, *J*=8.8, 2.0 Hz, 1H), 7.57~7.52 (m, 5H), 6.72 (d, *J*=9.2 Hz, 2H), 3.09 (s, 6H), 2.51 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 191.79, 155.26, 153.63, 148.43, 145.84, 138.10, 138.01, 133.81, 131.87, 130.46, 129.57, 128.55, 128.36, 127.07, 124.45, 123.85, 121.30, 110.60, 40.03, 21.96; HRMS (ESI) calcd for C₂₅H₂₂N₂O [M+H]⁺ 367.1805, found 367.1803.

(6-Methyl-4-phenylquinolin-2-yl)(naphthalen-1-yl)-methanone (**3m**): 20.9 mg, 56% yield. Yellow solid, m.p. 154~156 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.50~8.46 (m, 1H), 8.18 (s, 1H), 8.07 (d, *J*=8.8 Hz, 2H), 7.96~7.93 (m, 2H), 7.78 (s, 1H), 7.60~7.52 (m, 9H), 2.51 (s, 3H). ¹³C NMR (150 MHz, CDCl₃) δ: 196.74, 153.95, 148.73, 146.31, 139.03, 137.93, 134.47, 133.93, 132.32, 132.15, 131.62, 131.14, 130.91, 129.59, 128.66, 128.56, 128.43, 127.61, 127.49, 126.24, 125.97, 124.48, 124.13, 121.00, 22.05; HRMS (ESI) calcd for C₂₇H₂₀NO [M+H]⁺ 374.1539, found 374.1542.

(Furan-2-yl)(6-methyl-4-phenylquinolin-2-yl)methanone (**3n**): 21.0 mg, 67% yield. Yellow solid, m.p. 137~139 °C (lit.^[11d] 135~138 °C); ¹H NMR (400 MHz, CDCl₃) δ: 8.32 (dd, *J*=3.6, 1.2 Hz, 1H), 8.19 (d, *J*=8.8 Hz, 1H), 8.16 (s, 1H), 7.79 (dd, *J*=1.6, 0.8 Hz, 1H), 7.74 (s, 1H), 7.64 (dd, *J*=8.8, 2.0 Hz, 1H), 7.57~7.48 (m, 5H), 6.68 (dd, *J*=3.6, 1.6 Hz, 1H), 2.51 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 179.49, 152.22, 151.26, 148.77, 147.70, 146.15, 139.02, 137.80, 132.22, 130.55, 129.54, 128.61, 128.53, 127.80, 124.62, 120.23, 112.47, 22.04; HRMS (ESI) calcd for C₂₁H₁₆NO₂ [M+H]⁺ 314.1176, found 314.1180.

(8-Methyl-4-phenylquinolin-2-yl)(*p*-tolyl)methanone (**3o**): 24.6 mg, 73% yield. Yellow solid, m.p. 114~116 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.36 (d, *J*=8.0 Hz, 2H), 8.14 (s, 1H), 7.86 (d, *J*=8.4 Hz, 1H), 7.65 (d, *J*=7.2 Hz, 1H), 7.58~7.48 (m, 6H), 7.36 (d, *J*=8.0 Hz, 2H), 2.87 (s, 3H), 2.50 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 193.10, 152.85, 149.66, 146.38, 143.55, 139.04, 138.21, 133.94, 131.90, 129.87, 129.63, 128.65, 128.54, 128.44, 128.08, 127.44, 123.67, 120.68, 21.76, 18.47; HRMS (ESI) calcd for C₂₄H₂₀NO [M+H]⁺ 338.1539, found 338.1543.

(6-Ethyl-4-phenylquinolin-2-yl)(*p*-tolyl)methanone (**3p**): 26.7 mg, 76% yield. Yellow solid, m.p. 125~127 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.19 (dd, *J*=8.8, 5.6 Hz, 3H), 8.01 (s, 1H), 7.78 (s, 1H), 7.66 (dd, *J*=8.8, 2.0 Hz, 1H), 7.60~7.51 (m, 5H), 7.33 (d, *J*=7.9 Hz, 2H), 2.81 (q, *J*=7.8 Hz, 2H), 2.46 (s, 3H), 1.29 (t, *J*=7.8 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 193.90, 154.03, 149.11, 146.36, 145.11, 144.11, 138.19, 133.93, 131.87, 131.23, 131.05, 129.82, 129.13, 128.88, 128.76, 127.65, 123.57, 121.37, 29.54, 22.03, 15.74; HRMS (ESI) calcd for C₂₅H₂₂NO [M+H]⁺ 352.1696, found 352.1698.

(6-Methoxy-4-phenylquinolin-2-yl)(phenyl)methanone (**3q**): 26.5 mg, 78% yield. Yellow solid, m.p. 146~148 °C (lit.^[11d] 144~146 °C); ¹H NMR (400 MHz, CDCl₃) δ: 8.28~8.25 (m, 2H), 8.16 (d, *J*=9.2 Hz, 1H), 8.05 (s, 1H), 7.63~7.50 (m, 8H), 7.44 (dd, *J*=9.2, 2.8 Hz, 1H), 7.27 (d,

$J=2.8$ Hz, 1H), 3.83 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 193.74, 159.51, 151.97, 147.92, 143.38, 138.03, 136.51, 132.77, 132.51, 131.42, 129.31, 128.80, 128.73, 128.54, 128.03, 122.48, 121.47, 103.56, 55.53; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{18}\text{NO}_2$ $[\text{M}+\text{H}]^+$ 340.1332, found 340.1336.

(6-Methoxy-4-phenylquinolin-2-yl)(*p*-tolyl)methanone (**3r**): 28.3 mg, 80% yield. Yellow solid, m.p. 143 ~ 145 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.18 (d, $J=7.8$ Hz, 2H), 8.15 (d, $J=9.0$ Hz, 1H), 8.02 (s, 1H), 7.59 ~ 7.48 (m, 5H), 7.42 (dd, $J=9.0, 3.0$ Hz, 1H), 7.32 (d, $J=7.8$ Hz, 2H), 7.26 (d, $J=3.0$ Hz, 1H), 3.82 (s, 3H), 2.45 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 193.40, 159.42, 152.31, 147.84, 143.65, 143.37, 138.07, 133.87, 132.48, 131.58, 129.32, 128.81, 128.72, 128.52, 122.41, 121.50, 103.56, 55.51, 21.73; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{20}\text{NO}_2$ $[\text{M}+\text{H}]^+$ 354.1489, found 354.1485.

(6-Methoxy-4-phenylquinolin-2-yl)(*m*-tolyl)methanone (**3s**): 26.9 mg, 76% yield. Yellow solid, m.p. 116 ~ 118 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.16 (d, $J=9.6$ Hz, 1H), 8.05 ~ 8.03 (m, 2H), 8.02 (s, 1H), 7.59 ~ 7.53 (m, 4H), 7.52 ~ 7.48 (m, 1H), 7.43 (dd, $J=9.0, 3.0$ Hz, 2H), 7.39 (t, $J=7.8$ Hz, 1H), 7.26 (d, $J=3.0$ Hz, 1H), 3.82 (s, 3H), 2.44 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.13, 159.47, 152.13, 147.87, 143.39, 138.03, 137.85, 136.48, 133.65, 132.52, 131.65, 129.33, 128.81, 128.76, 128.74, 128.55, 127.91, 122.48, 121.49, 103.52, 55.53, 21.42. HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{20}\text{NO}_2$ $[\text{M}+\text{H}]^+$ 354.1489, found 354.1491.

(4-Bromophenyl)(6-methoxy-4-phenylquinolin-2-yl)-methanone (**3t**): 33.0 mg, 79% yield. Yellow solid, m.p. 184 ~ 186 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.18 (d, $J=8.4$ Hz, 2H), 8.13 (d, $J=9.0$ Hz, 1H), 8.06 (s, 1H), 7.66 (d, $J=8.4$ Hz, 2H), 7.58 ~ 7.50 (m, 5H), 7.43 (dd, $J=9.0, 3.0$ Hz, 1H), 7.26 (d, $J=3.0$ Hz, 1H), 3.82 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 192.51, 159.67, 151.41, 148.06, 143.31, 137.91, 135.27, 133.00, 132.47, 131.32, 129.29, 128.93, 128.76, 128.61, 128.05, 122.65, 121.39, 103.54, 55.55; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{17}\text{BrNO}_2$ $[\text{M}+\text{H}]^+$ 418.0437, found 418.0436.

(3-Bromophenyl)(6-methoxy-4-phenylquinolin-2-yl)-methanone (**3u**): 30.9 mg, 74% yield. Yellow solid, m.p. 125 ~ 127 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.44 (t, $J=1.8$ Hz, 1H), 8.21 (dt, $J=7.8, 1.8$ Hz, 1H), 8.14 (d, $J=9.6$ Hz, 1H), 8.06 (s, 1H), 7.74 ~ 7.71 (m, 1H), 7.58 ~ 7.49 (m, 5H), 7.43 (dd, $J=9.0, 3.0$ Hz, 1H), 7.39 (t, $J=7.8$ Hz, 1H), 7.26 (d, $J=3.0$ Hz, 1H), 3.82 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 192.05, 159.75, 151.16, 148.08, 143.35, 138.35, 137.89, 135.48, 134.28, 132.56, 130.02, 129.57, 129.30, 128.99, 128.78, 128.64, 122.69, 122.21, 121.35, 103.53, 55.56; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{17}\text{BrNO}_2$ $[\text{M}+\text{H}]^+$ 418.0437, found 418.0436.

(2-Bromophenyl)(6-methoxy-4-phenylquinolin-2-yl)-methanone (**3v**): 27.5 mg, 66% yield. Yellow solid, m.p. 124 ~ 126 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.13 (s, 1H), 8.03 (d, $J=9.0$ Hz, 1H), 7.65 (dd, $J=7.8, 1.2$ Hz, 1H), 7.59 ~ 7.49 (m, 6H), 7.46 ~ 7.42 (m, 1H), 7.39 ~ 7.35 (m, 2H), 7.24 (d, $J=2.4$ Hz, 1H), 3.80 (s, 3H); ^{13}C NMR (150

MHz, CDCl_3) δ : 195.96, 159.89, 150.30, 147.85, 143.95, 140.51, 137.97, 132.98, 132.91, 131.23, 130.35, 129.34, 129.28, 128.76, 128.60, 126.81, 122.51, 120.55, 120.53, 103.49, 55.53; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{17}\text{BrNO}_2$ $[\text{M}+\text{H}]^+$ 418.0437, found 418.0439.

(4-Iodophenyl)(6-methoxy-4-phenylquinolin-2-yl)-methanone (**3w**): 29.3 mg, 63% yield. Yellow solid, m.p. 182 ~ 184 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.12 (d, $J=9.0$ Hz, 1H), 8.05 (s, 1H), 8.01 (d, $J=8.4$ Hz, 2H), 7.88 (d, $J=8.4$ Hz, 2H), 7.56 (s, 2H), 7.55 (d, $J=4.2$ Hz, 2H), 7.54 ~ 7.49 (m, 2H), 7.43 (dd, $J=9.0, 3.0$ Hz, 1H), 3.82 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 192.83, 159.67, 151.37, 148.05, 143.32, 137.91, 137.33, 135.83, 132.87, 132.48, 129.29, 128.94, 128.77, 128.62, 122.66, 121.38, 103.54, 100.92, 55.56; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{17}\text{INO}_2$ $[\text{M}+\text{H}]^+$ 466.0298, found 466.0296.

(4-Hydroxyphenyl)(6-methoxy-4-phenylquinolin-2-yl)-methanone (**3x**): 18.8 mg, 53% yield. Yellow solid, m.p. 99 ~ 101 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.17 (d, $J=9.0$ Hz, 1H), 8.09 (d, $J=9.0$ Hz, 2H), 7.94 (s, 1H), 7.58 ~ 7.47 (m, 5H), 7.43 (dd, $J=9.0, 3.0$ Hz, 1H), 7.26 (d, $J=3.0$ Hz, 1H), 6.86 (d, $J=8.4$ Hz, 2H), 3.81 (s, 3H), 2.05 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 193.14, 161.38, 159.58, 153.01, 148.62, 143.25, 138.07, 134.16, 131.93, 129.55, 129.02, 128.99, 128.90, 128.85, 123.08, 121.90, 115.81, 103.88, 55.80; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{18}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 356.1281, found 356.1286.

(6-Chloro-4-phenylquinolin-2-yl)(*p*-tolyl)methanone (**3y**): 24.7 mg, 69% yield. Yellow solid, m.p. 120 ~ 122 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.19 (d, $J=9.0$ Hz, 1H), 8.16 (d, $J=8.4$ Hz, 2H), 8.04 (s, 1H), 7.96 (d, $J=2.4$ Hz, 1H), 7.71 (dd, $J=9.0, 1.8$ Hz, 1H), 7.59 ~ 7.51 (m, 5H), 7.32 (d, $J=7.8$ Hz, 2H), 2.46 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 193.13, 154.74, 148.84, 145.69, 144.13, 137.03, 134.49, 133.38, 132.44, 131.54, 130.85, 129.44, 128.95, 128.93, 128.86, 128.07, 124.64, 121.75, 21.78. HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{17}\text{ClNO}$ $[\text{M}+\text{H}]^+$ 358.0993, found 358.0990.

(6-Chloro-4-phenylquinolin-2-yl)(*m*-tolyl)methanone (**3z**): 22.9 mg, 64% yield. Yellow solid, m.p. 108 ~ 110 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.20 (d, $J=8.8$ Hz, 1H), 8.05 (s, 1H), 8.03 ~ 8.00 (m, 2H), 7.97 (d, $J=2.4$ Hz, 1H), 7.72 (dd, $J=9.2, 2.4$ Hz, 1H), 7.60 ~ 7.51 (m, 5H), 7.47 ~ 7.44 (m, 1H), 7.41 (t, $J=7.8$ Hz, 1H), 2.45 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 193.82, 154.59, 148.87, 145.72, 138.02, 137.01, 135.99, 134.56, 134.00, 132.48, 131.59, 130.89, 129.44, 128.95, 128.87, 128.78, 128.12, 128.03, 124.64, 121.71, 21.41; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{17}\text{ClNO}$ $[\text{M}+\text{H}]^+$ 358.0993, found 358.0995.

(3-Bromophenyl)(6-chloro-4-phenylquinolin-2-yl)-methanone (**3aa**): 25.2 mg, 60% yield. Yellow solid, m.p. 116 ~ 117 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.43 (t, $J=2.0$ Hz, 1H), 8.20 (d, $J=8.8$ Hz, 2H), 8.10 (s, 1H), 7.98 (d, $J=2.0$ Hz, 1H), 7.78 ~ 7.72 (m, 2H), 7.61 ~ 7.52 (m, 5H), 7.41 (t, $J=8.0$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 191.82, 153.57, 149.15, 145.67, 137.84, 136.87, 135.84, 134.97, 134.23, 132.51, 131.09, 129.96, 129.69, 129.41,

129.03, 128.91, 128.28, 124.69, 122.33, 121.59; HRMS (ESI) calcd for $C_{22}H_{14}BrClNO$ $[M+H]^+$ 421.9942, found 421.9945.

(8-Chloro-4-phenylquinolin-2-yl)(*p*-tolyl)methanone (**3ab**): 22.5 mg, 63% yield. Yellow solid, m.p. 134~136 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.50 (d, $J=8.4$ Hz, 2H), 8.20 (s, 1H), 7.92 (dd, $J=8.0, 2.4$ Hz, 2H), 7.57~7.49 (m, 6H), 7.37 (d, $J=8.0$ Hz, 2H), 2.48 (s, 3H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 191.80, 154.32, 150.20, 143.99, 143.67, 137.42, 135.45, 133.42, 132.20, 129.91, 129.57, 128.86, 128.84, 128.82, 128.69, 128.06, 124.86, 121.77, 21.78; HRMS (ESI) calcd for $C_{23}H_{17}ClNO$ $[M+H]^+$ 358.0993, found 358.0996.

(4-Phenylbenzo[*g*]quinolin-2-yl)(*p*-tolyl)methanone (**3ac**): 19.4 mg, 52% yield. Yellow solid, m.p. 165~167 °C (lit.^[11d] 165~168 °C); 1H NMR (400 MHz, $CDCl_3$) δ : 8.21 (d, $J=8.0$ Hz, 2H), 8.10 (d, $J=9.2$ Hz, 1H), 8.05 (s, 1H), 8.03 (d, $J=8.8$ Hz, 1H), 7.90 (d, $J=7.2$ Hz, 1H), 7.78 (d, $J=8.4$ Hz, 1H), 7.56~7.51 (m, 4H), 7.49~7.45 (m, 2H), 7.34 (d, $J=7.6$ Hz, 2H), 7.24~7.18 (m, 1H), 2.47 (s, 3H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 193.24, 153.12, 149.44, 148.48, 143.92, 142.28, 133.70, 133.64, 131.93, 131.59, 129.38, 129.30, 129.00, 128.90, 128.66, 128.60, 128.35, 128.30, 127.45, 125.77, 125.23, 124.34, 21.78; HRMS (ESI) calcd for $C_{27}H_{20}NO$ $[M+H]^+$ 374.1539, found 374.1536.

(6-Methyl-4-(*p*-tolyl)quinolin-2-yl)(*p*-tolyl)methanone (**3ad**): 23.2 mg, 66% yield. Yellow solid, m.p. 119~121 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.17 (dd, $J=12.0, 8.0$ Hz, 3H), 8.00 (s, 1H), 7.79 (s, 1H), 7.61 (d, $J=8.4$ Hz, 1H), 7.47 (d, $J=8.0$ Hz, 2H), 7.35 (dd, $J=16.8, 7.6$ Hz, 4H), 2.51 (s, 3H), 2.49 (s, 3H), 2.46 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 193.94, 153.98, 149.02, 146.17, 144.07, 138.79, 138.69, 135.23, 133.95, 132.30, 131.87, 130.86, 129.73, 129.60, 129.12, 127.72, 124.83, 121.33, 22.27, 22.04, 21.59; HRMS (ESI) calcd for $C_{25}H_{22}NO$ $[M+H]^+$ 352.1696, found 352.1699.

(4-(4-Fluorophenyl)-6-methylquinolin-2-yl)(*p*-tolyl)methanone (**3ae**): 21.0 mg, 59% yield. Yellow solid, m.p. 176~178 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.17 (t, $J=8.0$ Hz, 3H), 7.97 (s, 1H), 7.70 (s, 1H), 7.62 (dd, $J=8.8, 2.0$ Hz, 1H), 7.57~7.51 (m, 2H), 7.33 (d, $J=8.0$ Hz, 2H), 7.29~7.22 (m, 3H), 2.52 (s, 3H), 2.46 (s, 3H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 193.48, 163.77, 162.13, 153.70, 147.62, 143.89, 138.86, 133.86 (d, $J=3.4$ Hz), 133.63, 132.19, 131.60, 131.26 (d, $J=8.1$ Hz), 130.71, 128.87, 127.35, 124.23, 121.14, 115.72 (d, $J=21.5$ Hz), 22.03, 21.76; HRMS (ESI) calcd for $C_{24}H_{19}FNO$ $[M+H]^+$ 356.1445, found 356.1437.

(4-(2-Fluorophenyl)-6-methylquinolin-2-yl)(*p*-tolyl)methanone (**3af**): 18.8 mg, 53% yield. Yellow solid, m.p. 161~163 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.23~8.17 (m, 3H), 8.06 (s, 1H), 7.64 (d, $J=8.4$ Hz, 1H), 7.56~7.43 (m, 3H), 7.37~7.26 (m, 4H), 2.52 (s, 3H), 2.48 (s, 3H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 193.35, 160.43, 158.79, 153.62, 145.63, 143.86, 142.93, 138.94, 133.66, 132.31, 131.68 (d, $J=3.3$ Hz), 131.63, 130.66 (d, $J=8.3$ Hz),

130.62, 128.88, 127.54, 125.38 (d, $J=15.9$ Hz), 124.46 (d, $J=3.6$ Hz), 124.41 (d, $J=1.1$ Hz), 122.01, 116.02 (d, $J=21.8$ Hz), 22.01, 21.77; HRMS (ESI) calcd for $C_{24}H_{19}FNO$ $[M+H]^+$ 356.1445, found 356.1438.

(4-(4-Chlorophenyl)-6-methylquinolin-2-yl)(*p*-tolyl)methanone (**3ag**): 23.8 mg, 64% yield. Yellow solid, m.p. 171~173 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.17 (t, $J=8.0$ Hz, 3H), 7.97 (s, 1H), 7.68 (s, 1H), 7.62 (dd, $J=8.4, 2.0$ Hz, 1H), 7.51 (q, $J=8.4$ Hz, 4H), 7.32 (d, $J=8.0$ Hz, 2H), 2.52 (s, 3H), 2.46 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 193.67, 153.93, 147.64, 146.10, 144.18, 139.21, 136.54, 134.98, 133.81, 132.50, 131.85, 131.10, 130.99, 129.16, 129.14, 127.38, 124.39, 121.28, 22.30, 22.04; HRMS (ESI) calcd for $C_{24}H_{19}ClNO$ $[M+H]^+$ 372.1150, found 372.1152.

Supporting Information The spectroscopic data of 1H NMR, ^{13}C NMR for **3a~3z** and **3aa~3ag**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Solomon, V. R.; Haq, W.; Srivastava, K.; Puri, S. K.; Katti, S. B. *J. Med. Chem.* **2007**, *50*, 394.
(b) Michael, J. P. *Nat. Prod. Rep.* **2007**, *24*, 223.
(c) Bharate, J. B.; Vishwakarma, R. A.; Bharate, S. B. *RSC Adv.* **2015**, *5*, 42020.
(d) Solomon, V. R.; Lee, H. *Curr. Med. Chem.* **2011**, *18*, 1488.
(e) Singh, S.; Kaur, G.; Mangla, V.; Gupta, M. K. *J. Enzyme Inhib. Med. Chem.* **2015**, *30*, 492.
(f) Afzal, O.; Kumar, S.; Haider, M. R.; Ali, M. R.; Kumar, R.; Jaggi, M.; Bawa, S. *Eur. J. Med. Chem.* **2015**, *97*, 871.
(g) Andries, K.; Verhasselt, P.; Guillemont, J.; Cambau, E.; Truffot-Pernot, C.; Lounis, N.; Jarlier, V. *Science* **2005**, *307*, 223.
(h) Behenna, D. C.; Stockdill, J. L.; Stoltz, B. M. *Angew. Chem., Int. Ed.* **2008**, *47*, 2365.
- [2] (a) Rouffet, M.; F.de Oliveira, C. A.; Udi, Y.; Agrawal, A.; Sagi, I.; McCammon, J. A.; Cohen, S. M. *J. Am. Chem. Soc.* **2010**, *132*, 8232.
(b) Kim, J. I.; Shin, I. S.; Kim, H.; Lee, J. K. *J. Am. Chem. Soc.* **2005**, *127*, 1614.
(c) Bhalla, V.; Vij, V.; Kumar, M.; Sharma, P. R.; Kaur, T. *Org. Lett.* **2012**, *14*, 1012.
(d) Tao, S.; Li, L.; Yu, J.; Jiang, Y.; Zhou, Y.; Lee, C. S.; Lee, S. T.; Zhang, X.; Kwon, O. *Chem. Mater.* **2009**, *21*, 1284.
(e) Wang, M.; Shi, Y. H.; Luo, J. F.; Du, W.; Shi, X. X.; Fossey, J. S.; Deng, W.-P. *Catal. Sci. Technol.* **2011**, *1*, 100.
(f) Wang, M.; Wang, Z.; Shi, Y. H.; Shi, X. X.; Fossey, J. S.; Deng, W. P. *Angew. Chem., Int. Ed.* **2011**, *50*, 4897.
(g) Wang, Z.; Yu, X.; Tian, B. X.; Payne, D. T.; Yang, W. L.; Liu, Y. Z.; Fossey, J. S.; Deng, W. P. *Chem.-Eur. J.* **2015**, *21*, 10457.
(h) Ma, C.; Fang, P.; Mei, T. S. *ACS Catal.* **2018**, *8*, 7179.
(i) Nimmagadda, S. K.; Liu, M. Y.; Karunananda, M. K.; Gao, D. W.; Apolinar, O.; Chen, J. S.; Liu, P.; Engle, K. M. *Angew. Chem., Int. Ed.* **2019**, *58*, 3923.
(j) Takamatsu, K.; Hirano, K.; Miura, M.; *Angew. Chem., Int. Ed.* **2017**, *56*, 5353.
(k) Barday, M.; Janot, C.; Halcovitch, N. R.; Muir, J.; Aissa, C. *Angew. Chem., Int. Ed.* **2017**, *56*, 13117.
- [3] (a) Yao, R.; Rong, G.; Yan, B.; Qiu, L.; Xu, X. *ACS Catal.* **2016**, *6*, 1024.
(b) Wang, Q.; Huang, J.; Zhou, L. *Adv. Synth. Catal.* **2015**, *357*, 2479.

- (c) Zhang, L.; Chen, Su.; Gao, Y.; Zhang, P.; Wu, Y.; Tang, G.; Zhao, Y. *Org. Lett.* **2016**, *18*, 1286.
- (d) Jiang, H.; An, X.; Tong, K.; Zheng, T.; Zhang, Y.; Yu, S. *Angew. Chem., Int. Ed.* **2015**, *54*, 4055.
- (e) Zhou, W.; Lei, J. *Chem. Commun.* **2014**, *50*, 5583.
- (f) Li, H.; Wang, C.; Huang, H.; Xu, X.; Li, Y. *Tetrahedron Lett.* **2011**, *52*, 1108.
- (g) Matsubara, Y.; Hirakawa, S.; Yamaguchi, Y.; Yoshida, Z. I. *Angew. Chem., Int. Ed.* **2011**, *50*, 7670.
- (h) Xu, P.; Zhu, T.; Wei, T.; Wang, S.; Ji, S. *RSC Adv.* **2016**, *6*, 32467.
- (i) Zheng, W. P.; Yang, W. G.; Luo, D. P.; Min, L.; Wang, X. Y.; Hu, Y. F. *Adv. Synth. Catal.* **2019**, *361*, 1995.
- (j) Ye, J. L.; Zhu, Y. N.; Geng, H.; Huang, P. Q. *Sci. China Chem.* **2018**, *61*, 687.
- [4] (a) Corey, E. J.; Tramontano, A. *J. Am. Chem. Soc.* **1981**, *103*, 5599.
- (b) Laras, Y.; Hugues, V.; Chandrasekaran, Y.; Blanchard-Desce, M.; Acher, F. C.; Pietrancosta, N. *J. Org. Chem.* **2012**, *77*, 8294.
- (c) Wu, Y. C.; Liu, L.; Li, H. J.; Wang, D.; Chen, Y. J. *J. Org. Chem.* **2006**, *71*, 6592.
- [5] (a) Bergstroem, F. W. *Chem. Rev.* **1944**, *35*, 77.
- (b) Coscia, A. T.; Dickerman, S. C. *J. Am. Chem. Soc.* **1959**, *81*, 3098.
- [6] (a) Bello, D.; Ramon, R.; Lavilla, R. *Curr. Org. Chem.* **2010**, *14*, 332.
- (b) Kouznetsov, V. V. *Tetrahedron* **2009**, *65*, 2721.
- [7] Wei, W. T.; Teng, F.; Li, Y.; Song, R. J.; Li, J. H. *Org. Lett.* **2019**, *21*, 6285.
- [8] Zheng, J.; Li, Z.; Huang, L. B.; Wu, W. Q.; Li, J. X.; Jiang, H. F. *Org. Lett.* **2016**, *18*, 3514.
- [9] Chen, P.; Nan, J.; Hu, Y.; Ma, Q.; Ma, Y. M. *Org. Lett.* **2019**, *21*, 4812.
- [10] (a) Hirupathi, N.; Puri, S.; Reddy, T. J.; Sridhar, B.; Reddy, M. S. *Adv. Synth. Catal.* **2016**, *358*, 303.
- (b) Gadakh, S. K.; Dey, S.; Sudalai, A. *Org. Biomol. Chem.* **2016**, *14*, 2969.
- (c) Zheng, L.; Guo, B.; Hua, R. *J. Org. Chem.* **2014**, *79*, 11541.
- (d) Gu, Z. Y.; Liu, C. G.; Wang, S. Y.; Ji, S. J. *Org. Lett.* **2016**, *18*, 2379.
- (e) Wei, W. T.; Cheng, Y. J.; Hu, Y.; Chen, Y. Y.; Zhang, X. J.; Zou, Y.; Yan, M. *Adv. Synth. Catal.* **2015**, *357*, 3474.
- (f) Bao, L.; Liu, J. L.; Xu, L.; Hu, Z. Y.; Xu, X. X. *Adv. Synth. Catal.* **2018**, *360*, 1870.
- [11] (a) Wu, X.; Geng, X.; Zhao, P.; Zhang, J. J.; Gong, X. X.; Wu, Y. D.; Wu, A. X. *Org. Lett.* **2017**, *19*, 1550.
- (b) Geng, X.; Wu, X.; Zhao, P.; Zhang, J. J.; Wu, Y. D.; Wu, A. X. *Org. Lett.* **2017**, *19*, 4179.
- (c) Geng, X.; Wang, C.; Zhao, P.; Zhou, Y.; Wu, Y. D.; Wu, A. X. *Org. Lett.* **2019**, *21*, 4939.
- (d) Gao, Q. H.; Liu, S.; Wu, X.; Wu, A. X. *Org. Lett.* **2014**, *16*, 4582.
- [12] (a) Zhang, X.; Wang, T. L.; Huo, C. D.; Wang, X. C.; Quan, Z. J. *Chem. Commun.* **2018**, *54*, 3114.
- (b) Wang, T. L.; Liu, X. J.; Huo, C. D.; Wang, X. C.; Quan, Z. J. *Chem. Commun.* **2018**, *54*, 499.
- (c) Zhang, X.; Wang, T. L.; Liu, X. J.; Wang, X. C.; Quan, Z. J. *Org. Biomol. Chem.* **2019**, *17*, 2379.
- (d) Wang, T. L.; Liu, X. J.; Zhang, X.; Cao, B. Q.; Quan, Z. J.; Wang, X. C. *Tetrahedron Lett.* **2018**, *59*, 4426.
- (e) Cao, B. Q.; Qiu, Y. F.; Zhang, X.; Zhu, Z. H.; Quan, Z. J.; Wang, X. C. *Eur. J. Org. Chem.* **2019**, *2019*, 1208.
- [13] (a) Li, D. J.; Zeng, F. L. *Org. Lett.* **2017**, *19*, 6498.
- (b) Zhu, J. W.; Hu, W. M.; Sun, S.; Yu, J. T.; Cheng, J. *Adv. Synth. Catal.* **2017**, *359*, 3725.
- (c) Huang, Y. N.; Li, Y. L.; Li, J.; Deng, J. J. *J. Org. Chem.* **2016**, *81*, 4645.
- (d) Ferguson, J.; Zeng, F. L.; Alwis, N.; Alper, H. *Org. Lett.* **2013**, *15*, 1998.
- (e) Pang, X. B.; Wu, M. Z.; Ni, J. X.; Zhang, F. M.; Lan, J. F.; Chen, B. H.; Yan, R. L. *J. Org. Chem.* **2017**, *82*, 10110.
- (f) Zhang, Z.; Liao, L. L.; Yan, S. S.; Wang, L.; He, Y. Q.; Ye, J. H.; Li, J.; Zhi, Y. G.; Yu, D. G. *Angew. Chem., Int. Ed.* **2016**, *55*, 7068.
- (h) Zhang, X.; Xu, X. F.; Yu, L. T.; Zhao, Q. *Tetrahedron Lett.* **2014**, *55*, 2280.
- (g) Sun, S.; Zhou, C.; Yu, J. T.; Cheng, J. *Org. Lett.* **2019**, *21*, 6579.
- (h) Ni, J. X.; Jiang, Y.; An, Z. Y.; Yan, R. L. *Org. Lett.* **2018**, *20*, 1534.
- [14] (a) Čubiňák, M.; Eigner, V.; Tobrman, T. *Adv. Synth. Catal.* **2018**, *360*, 4604.
- (b) Isley, N. A.; Wang, Y.; Gallou, F.; Handa, S.; Aue, D. H.; Lipshutz, B. H. *ACS Catal.* **2017**, *7*, 8331.
- (c) Walker, B. R.; Sevov, C. S. *ACS Catal.* **2019**, *9*, 7197.
- (d) Grub, H.; Belu, C.; Bernhard, L. M.; Merschel, A.; Sewald, N. *Chem.-Eur. J.* **2019**, *25*, 5880.
- (e) Yan, B.; Zhou, Y. B.; Zhang, H.; Chen, J. J.; Liu, Y. H. *J. Org. Chem.* **2007**, *72*, 7783.
- (f) Smith, C. R.; Bunnelle, E. M.; Rhodes, A. J.; Sarpong, R. *Org. Lett.* **2007**, *9*, 1169.
- [15] (a) Deshidi, R.; Devari, S.; Shah, B. A. *Org. Chem. Front.* **2015**, *2*, 515.
- (b) Zhu, Y. P.; Liu, M. C.; Jia, F. C.; Yuan, J. J.; Gao, Q. H.; Lian, M.; Wu, A. X. *Org. Lett.* **2012**, *14*, 3392.
- (c) Kornblum, N.; Jones, W. J.; Anderson, G. J. *J. Am. Chem. Soc.* **1959**, *81*, 4113.
- (d) Kornblum, N.; Powers, J. W.; Anderson, G. J.; Jones, W. J.; Larson, H. O.; Levand, O.; Weaver, W. M. *J. Am. Chem. Soc.* **1957**, *79*, 6562.
- (e) Monga, A.; Bagchi, S.; Sharma, A. *New J. Chem.* **2018**, *42*, 1551.
- (f) Mohammed, S.; Vishwakarma, R. A.; Bharate, S. B. *J. Org. Chem.* **2015**, *80*, 6915.
- (g) Yan, R. L.; Yan, H.; Ma, C.; Ren, Z. Y.; Gao, X. A.; Huang, G. S.; Liang, Y. M. *J. Org. Chem.* **2012**, *77*, 2024.
- (h) An, Z. Y.; Jiang, Y.; Guan, X.; Yan, R. L. *Chem. Commun.* **2018**, *54*, 10738.

(Cheng, F.)