“水上”吡唑啉酮与三氟甲基酮的亲核加成

罗 亮 曹晓梅 赖国伟 刘金香 罗海清
路东亮 张 勇*

(赣南师范大学有机药物化学重点实验室 江西赣州 341000)

摘要 基于水相条件下发展了一种绿色高效的吡唑啉酮与三氟甲基酮的亲核加成的方法, 且以高收率获得吡唑啉酮取代的叔三氟甲基醇类化合物. 该方法具有的无催化剂、无色谱柱分离、环境友好以及易于后处理的优点使其在制备各种吡唑啉酮取代的叔三氟甲基醇类化合物中很有应用前景.

关键词 “水上”反应; 亲核加成; 吡唑啉酮; 三氟甲基酮

“On Water” Nucleophilic Addition of Pyrazolones to Trifluoromethyl Ketones

Luo, Liang Cao, Xiaomei Lai, Guowei Liu, Jinxiang Luo, Haiqing
Lu, Dongliang Zhang, Yong*

(Key Laboratory of Organo-pharmaceutical Chemistry, Gannan Normal University, Ganzhou, Jiangxi 341000)

Abstract A green and efficient nucleophilic addition reaction of trifluoromethyl ketone with pyrazolone was developed under “on water” conditions, affording pyrazolone substituted tertiary trifluoromethyl alcohols in high yields. The advantages of being catalyst-free, column chromatography-free, environmentally benign and easy workup make it a promising method for preparation of a variety of pyrazolone substituted tertiary trifluoromethyl alcohols.

Keywords “on water” reaction; nucleophilic addition; pyrazolone; trifluoromethyl ketone

1 Introduction

The development of practical, atom-economical, and environmentally benign processes for the synthesis of bio-active heterocyclic compounds represents an important goal of current chemical and pharmaceutical research.^[1] In this context, the use of water as a reaction medium^[2] has drawn significant attention, because water is cheap, non-flammable, nontoxic and the most abundant liquid on the planet. In addition to being a green solvent, water has unique physical and chemical properties, which shows incomparable effect on the rate and selectivity of organic reactions.^[3] Since the landmark work of Breslow and Sharpless,^[4] “on water” reaction has been established as a powerful strategy in organic synthesis. While many reactions have been conducted successfully under “on water” conditions, it is still highly desirable to explore practical “on water” reactions for the goal of green and sustainable

chemistry.

Fluorine-containing molecules play a significant role in agrochemicals, medicinal chemistry, and material science, owing to their unique properties.^[5] It is well established that the incorporation of trifluoromethyl (CF₃) group into drug candidates can significantly alter their lipophilicity, metabolic stability, and bioavailability.^[6] Thus, the development of efficient and practical synthetic strategies to generate a novel trifluoromethylated heterocyclic system has received much attention.^[7] Pyrazolone, a special five-membered heterocycle containing two adjacent nitrogen atoms, is a significant pharmacophore with a broad spectrum of biological activities, such as antipyretic, analgesic, antimicrobial, antiviral, antioxidant, anticancer, and anti-inflammatory effects.^[8] In view of the great medicinal significance of pyrazolones and their derivatives, we became interested in the synthesis of trifluoromethylated py-

* Corresponding author. E-mail: yong_zhangnnu@126.com

Received October 29, 2019; revised January 14, 2020; published online February 14, 2020.

Project supported by the National Natural Science Foundation of China (No. 21562003), the Jiangxi Provincial Department of Science and Technology Fund (20192BAB213007 and 20144BAB2130003), and the Education Department of Jiangxi Province (No. GJJ170835).

国家自然科学基金(No. 21562003)、江西省科学技术厅基金(Nos. 20192BAB213007 和 20144BAB2130003)和江西省教育厅(No. GJJ170835)资助项目.

razolone derivatives. Herein, we wish to report an “on water” novel, catalyst-free and column chromatography-free protocol for the synthesis of pyrazolone substituted tertiary trifluoromethyl alcohols (Scheme 1).

Scheme 1 Synthesis of pyrazolone substituted tertiary trifluoromethyl alcohols under “on water” condition

2 Results and discussion

Initially, trifluoroacetophenone (**1a**) and 1-phenyl-3-methyl-5-pyrazolone (**2a**) were selected as model substrates to investigate the “on water” reaction (Table 1). To our delight, the nucleophilic addition reaction of **2a** to **1a** proceeded smoothly to afford the desired product **3a** in 82% yield without any catalyst under ambient temperature (Entry 1). Increasing the reaction temperature in water from room temperature to 50 °C and further to 80 °C led to the formation of product **3a** in excellent yields (Entries 2, 3). The use of organic solvents such as CH₃CN, tetrahydrofuran (THF), *N,N*-dimethylformamide (DMF), 1,4-dioxane, dimethyl sulfoxide (DMSO), CHCl₃, toluene and CH₃OH did not improved the reaction efficiency as compared to pure water (Entries 4~11). It is noteworthy that the yield of **3a** rose dramatically with the increase of proportion of water in the co-solvent of CH₃OH-H₂O (Entries 12, 13). Notably, this “on water” reaction could be used for preparative synthesis, as demonstrated by a scale up reaction using 5.0 mmol of **2a** under the similar reaction conditions described above (Entry 14). After completion of the reaction, the desired product **3a** (>99% purity) was obtained in 1.585 g with 91% yield by simple filtration and washing with water. This procedure effectively avoids the use of volatile organic solvents for post-process.

Having optimized the reaction conditions, the substrate scope of trifluoromethyl ketones **1** was then explored, and the results are presented in Table 2. Generally, the reaction proceeded well for a wide range of trifluoromethyl ketones to afford the desired products **3** in good to excellent yields. It was found that the electronic nature of the substituents and their position on the phenyl ring have no obvious impact on the yields of the processes. The reactions of poly-substituted aryl substrates (2,4-Cl₂C₆H₄ and C₆F₅) worked as well and provided products **3k** and **3l** in 81% and 88% yields, respectively. The 2-naphthyl substrate also led to the corresponding product (**3m**) in high yield. Notably, trifluoromethyl ketones bearing heterocyclic groups (*N*-Ts protected indole and thiophene) were also tolerable to give the corresponding products (**3n** and **3o**) in excellent yields. Moreover, the alkyl trifluoromethyl ketone (1,1,1-trifluorobutan-2-one as an example) was also applicable, affording the trifluoromethyl alcohols adduct **3p** in 88% yield.

Table 1 Optimization of reaction conditions^a

Entry	Temp./°C	Solvent	T/h	Yield/%
1	r.t.	H ₂ O	24	82
2	50	H ₂ O	18	99 (95 ^b)
3	80	H ₂ O	15	95 ^b
4	50	CH ₃ CN	48	90
5	50	THF	48	85
6	50	DMF	48	15
7	50	1,4-Dioxane	48	56
8	50	DMSO	48	44
9	50	CHCl ₃	48	62
10	50	Toluene	48	64
11	50	CH ₃ OH	48	52
12	50	H ₂ O-CH ₃ OH (1:3)	18	75
13	50	H ₂ O-CH ₃ OH (3:1)	18	90
14 ^c	50	H ₂ O	24	91 ^b

^a Unless otherwise noted, the reaction was performed with 0.22 mmol of **1a** and 0.20 mmol of **2a** in 1 mL of solvent. The yield was determined by ¹H NMR analysis of the crude reaction mixture. ^b Isolated yield by washing process. ^c The reaction was performed at a gram scale.

Next, the substrate scope of 2-pyrazolin-5-one nucleophiles was briefly investigated (Table 3). Both electron-donating (4-Me and 4-OMe) and electron-withdrawing (4-F, 4-Br and 4-CF₃) groups on the phenyl ring were applicable in this “on water” process to afford the desired nucleophilic addition products **4a**~**4e** in good to excellent yields. In addition, pyrazolone with a hydrogen on nitrogen (*N*₁) was also suitable for this transformation and gave the product **4f** in high yield.

On the basis of previous reports,^[9] a plausible transition state involving an oil-water interface was assumed to explain the high efficiency of this process. As shown in Figure 1, the formation of tautomer of **2a** was promoted by water through hydrogen bonding. The nucleophilic addition reaction between tautomer of **2a** and trifluoromethyl ketones **1a** may take place at the oil-water interface. Through the interactions with hydrogen-bond networks at this interface, the carbonyl group of the electrophile and the hydroxyl group of nucleophile might be organized at the periphery of the oil droplet, while the hydrophobic part is positioned within the hydrophobic interior. As a consequence, the free hydroxyl groups at the phase boundary can effectively activate the reactants and stabilize the transition state.

3 Conclusions

In summary, an “on water” nucleophilic addition of pyrazolones to trifluoromethyl ketones has been developed using a catalyst-free and column chromatography-free

Table 2 Scope of trifluoromethyl ketones^a

$\text{R}^1\text{C(=O)CF}_3 + \text{2a} \xrightarrow[50 \sim 80\text{ }^\circ\text{C}]{1\text{ mL H}_2\text{O}} \text{3}$
3a 50 °C, 18 h, 95% yield	3b 50 °C, 26 h, 80% yield	3c 50 °C, 32 h, 94% yield	3d 80 °C, 32 h, 94% yield
3e 80 °C, 30 h, 88% yield	3f 80 °C, 26 h, 85% yield	3g^b 80 °C, 48 h, 81% yield	3h 80 °C, 25 h, 86% yield
3i 50 °C, 24 h, 91% yield	3j 50 °C, 24 h, 91% yield	3k 50 °C, 36 h, 81% yield	3l 50 °C, 19 h, 88% yield
3m^b 50 °C, 24 h, 85% yield	3n^c 80 °C, 40 h, 94% yield	3o 50 °C, 24 h, 96% yield	3p 50 °C, 20 h, 88% yield

^a Unless otherwise noted, the reaction was performed with 0.22 mmol of **1** and 0.20 mmol of **2a** in 1 mL of water. The yield refers to the isolated product. ^b 0.20 mmol of **1g** or **1m** and 0.22 mmol of **2a** were used. ^c 0.20 mmol of **1n** was used.

Figure 1 Proposed transition state

protocol. The reaction rate was greatly enhanced under “on water” conditions. Moreover, this process provides a highly efficient and environmentally benign approach for the synthesis of a variety of pyrazolone substituted tertiary trifluoromethyl alcohols.

4 Experimental section

4.1 General information

NMR spectra were recorded on a liquid NMR spectrometer (400 MHz for ¹H, 100 MHz for ¹³C and 376 MHz for ¹⁹F) using DMSO-*d*₆ as the solvent. The residual proton in DMSO-*d*₆ (δ 2.50) served as an internal standard for ¹H NMR, the ¹³C-atom of DMSO-*d*₆ was used as an internal

standard (δ 39.50) for ¹³C NMR. Infrared spectra were recorded on an FT-IR spectrometer, and only major peaks were reported in cm⁻¹. The high-resolution mass spectra (HRMS) were recorded on an electrospray ionization (ESI) apparatus using Orbitrap mass spectrometry. Unless otherwise noted, analytical grade solvents and commercially available reagents were used without further purification. Trifluoromethyl ketone **1** was obtained from commercial sources whereas **1j** was prepared as reported literature.^[10] 2,2,2-Trifluoro-1-(1-tosyl-1*H*-indol-3-yl)ethanone (**1n**) was prepared following literature procedure.^[10b] 2-Aryl-1,5-dimethyl-1*H*-pyrazol-3(2*H*)-ones **2b**, **2c**, **2d** and **2e** were readily prepared according to the reported procedures with slight modifications.

4.2 General procedure for the synthesis of 2b~2e

A mixture containing corresponding substituted phenylhydrazine hydrochloride (4.47 mmol), ethyl acetoacetate (565 μ L, 4.47 mmol) and 2.0 mL of acetic acid was refluxed for 8 h. The solution was then cooled to ambient temperature and concentrated *in vacuo*. The crude product was purified by silica gel flash chromatography (*V*(petroleum ether)/*V*(ethyl acetate)=3/1) to afford **2b**~**2e**.^[11a]

2-(4-Methoxyphenyl)-5-methyl-2,4-dihydro-3*H*-pyrazol-3-one (**2b**): Yellow solid, 75% yield. m.p. 126~127 °C

Table 3 Scope of pyrazolones^a

^a Unless otherwise noted, the reaction was performed with 0.22 mmol of **1a** and 0.20 mmol of **2** in 1 mL water. The yield refers to the isolated product.

^b 0.40 mmol of **1a** was used.

(lit.^[11a] 129–130 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.71 (d, *J*=9.2 Hz, 2H), 6.91 (d, *J*=9.2 Hz, 2H), 3.80 (s, 3H), 3.39 (s, 2H), 2.17 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 170.4, 157.0, 156.3, 131.4, 120.9, 114.0, 55.5, 43.0, 17.1.

2-(4-Fluorophenyl)-5-methyl-2,4-dihydro-3H-pyrazol-3-one (**2c**):^[11b] Yellow solid, 70% yield. m.p. 148–149 °C (lit.^[11c] 148–149 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.84–7.80 (m, 2H), 7.07 (t, *J*=8.8 Hz, 2H), 3.43 (s, 2H), 2.19 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 170.5, 159.9 (d, ¹*J*_{CF}=244.3 Hz), 156.6, 134.3 (d, ¹*J*_{CF}=2.7 Hz), 120.7 (d, ¹*J*_{CF}=7.9 Hz), 115.6 (d, ¹*J*_{CF}=22.6 Hz), 43.1, 17.1.

2-(4-Bromophenyl)-5-methyl-2,4-dihydro-3H-pyrazol-3-one (**2d**):^[11c] Yellow solid, 70% yield. m.p. 168–169 °C (lit.^[11a] 160–162 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.80–7.78 (m, 2H), 7.50–7.48 (m, 2H), 3.43 (s, 2H), 2.20 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 170.6, 156.7, 137.2, 131.9, 120.2, 117.8, 43.2, 17.2.

5-Methyl-2-(4-(trifluoromethyl)phenyl)-2,4-dihydro-3H-pyrazol-3-one (**2e**):^[11d] Reddish-brown solid, 68% yield. m.p. 171–172 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.05 (d, *J*=8.0 Hz, 2H), 7.65 (d, *J*=8.0 Hz, 2H), 3.47 (s, 2H), 2.23 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 170.9, 157.0, 140.9, 126.7, 126.4, 126.2, 126.2, 126.1, 125.5, 122.8, 118.2, 43.8, 17.4.

4.3 Representative procedure for nucleophilic addition reaction

The mixture of trifluoroacetophenone (**1a**) (38.3 mg, 0.22 mmol) and 1-phenyl-3-methyl-5-pyrazolone (**2a**) (34.8 mg, 0.20 mmol) in 1.0 mL of H₂O were stirred at 50 °C.

Once the reaction completed, the solid mixture was filtered, washed by water and dried under vacuum to afford the analytically pure product **3a**.

3-Methyl-1-phenyl-4-(2,2,2-trifluoro-1-hydroxy-1-phenylethyl)-1H-pyrazol-5-ol (**3a**): 66.2 mg, 95% yield. White solid, m.p. 124–125 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.71–7.68 (m, 2H), 7.56–7.54 (m, 2H), 7.51–7.47 (m, 2H), 7.45–7.36 (m, 3H), 7.31–7.27 (m, 1H), 1.64 (s, 3H); ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ: –74.8 (s, 3F); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 147.0, 139.1, 136.3, 129.1, 128.4, 128.1, 127.3, 125.9, 123.1 (q, *J*=287.0 Hz), 120.4, 100.9, 76.1 (q, *J*=29.9 Hz), 12.4; IR (KBr) ν: 3423, 3071, 2977, 1600, 1569, 1499, 1171, 1156, 940, 835 cm^{–1}. HRMS (ESI/Orbitrap) calcd for C₁₈H₁₆F₃N₂O₂ [M+H]⁺ 349.1158; found 349.1153.

3-Methyl-1-phenyl-4-(2,2,2-trifluoro-1-(4-fluorophenyl)-1-hydroxyethyl)-1H-pyrazol-5-ol (**3b**): 58.6 mg, 80% yield. White solid, m.p. 125–126 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.69 (d, *J*=8.0 Hz, 2H), 7.60–7.57 (m, 2H), 7.50–7.46 (m, 2H), 7.30–7.22 (m, 3H), 1.71 (s, 3H); ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ: –75.0 (s, 3F), –113.9 (s, 1F); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 161.7 (d, *J*=245.3 Hz), 147.0, 136.2, 135.4 (d, *J*=3.1 Hz), 129.5 (d, *J*=8.5 Hz), 129.0, 125.8 (q, *J*=283.5 Hz), 125.8, 120.4, 114.8 (d, *J*=21.5 Hz), 100.9, 75.6 (q, *J*=28.4 Hz), 12.5; IR (KBr) ν: 3089, 2928, 1600, 1562, 1498, 1181, 1166, 1145, 824, 717 cm^{–1}. HRMS (ESI/Orbitrap) calcd for C₁₈H₁₅F₄N₂O₂ [M+H]⁺ 367.1064; found 367.1057.

4-(1-(4-Chlorophenyl)-2,2,2-trifluoro-1-hydroxyethyl)-3-methyl-1-phenyl-1H-pyrazol-5-ol (**3c**): 72.0 mg, 94% yield. White solid, m.p. 135–136 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.69 (d, *J*=7.6 Hz, 2H), 7.58–7.55 (m, 2H), 7.49–7.46 (m, 4H), 7.22 (t, *J*=7.2 Hz, 1H), 1.75 (s, 3H); ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ: –74.9 (s, 3F); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 147.1, 138.1, 136.5, 133.0, 129.2, 129.0, 128.0, 125.8, 125.6 (q, *J*=288.1 Hz), 120.4, 100.7, 75.6 (q, *J*=29.7 Hz), 12.6; IR (KBr) ν: 3317, 2999, 1597, 1580, 1544, 1497, 1173, 1153, 941, 812 cm^{–1}. HRMS (ESI/Orbitrap) calcd for C₁₈H₁₅ClF₃N₂O₂ [M+H]⁺ 383.0769; found 383.0764.

4-(1-(4-Bromophenyl)-2,2,2-trifluoro-1-hydroxyethyl)-3-methyl-1-phenyl-1H-pyrazol-5-ol (**3d**): 80.3 mg, 94% yield. White solid, m.p. 135–136 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.69 (d, *J*=8.0 Hz, 2H), 7.62 (d, *J*=8.4 Hz, 2H), 7.51–7.46 (m, 4H), 7.30–7.26 (m, 1H), 1.75 (s, 3H); ¹⁹F NMR (376 MHz, DMSO-*d*₆) δ: –74.9 (s, 3F); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 147.1, 138.6, 136.4, 130.9, 129.5, 129.0, 125.6 (q, *J*=283.9 Hz), 125.8, 121.6, 120.4, 100.7, 75.6 (q, *J*=30.5 Hz), 12.6; IR (KBr) ν: 3282, 3065, 2927, 1600, 1567, 1503, 1166, 1152, 841, 724 cm^{–1}. HRMS (ESI/Orbitrap) calcd for C₁₈H₁₅BrF₃N₂O₂ [M+H]⁺ 427.0264; found 427.0261.

3-Methyl-1-phenyl-4-(2,2,2-trifluoro-1-hydroxy-1-(*p*-tolylethyl)-1H-pyrazol-5-ol (**3e**): 63.8 mg, 88% yield. White solid, m.p. 137–138 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.69 (d, *J*=7.6 Hz, 2H), 7.51–7.47 (m, 2H), 7.42 (d, *J*=8.0 Hz, 2H), 7.31–7.27 (m, 1H), 7.22 (d, *J*=8.0 Hz,

2H); 2.32 (s, 3H), 1.63 (s, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -74.9 (s, 3F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 147.0, 137.6, 136.0, 129.0, 128.5, 127.1, 125.9 (q, $J=285.2$ Hz), 125.8, 120.2, 100.9, 75.9 (q, $J=29.1$ Hz), 20.6, 12.3; IR (KBr) ν : 3314, 3036, 2928, 1595, 1580, 1544, 1500, 1395, 1166, 1151, 944, 928, 812 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{19}\text{H}_{18}\text{F}_3\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 363.1315; found 363.1308.

3-Methyl-1-phenyl-4-(2,2,2-trifluoro-1-hydroxy-1-(4-methoxyphenyl)ethyl)-1H-pyrazol-5-ol (**3f**): 64.3 mg, 85% yield. White solid, m.p. 112~113 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 7.70 (d, $J=7.2$ Hz, 2H), 7.49~7.45 (m, 4H), 7.30~7.27 (m, 1H), 6.97 (d, $J=8.0$ Hz, 2H), 3.77 (s, 3H), 1.65 (s, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -75.0 (s, 3F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 158.9, 147.0, 130.9, 129.0, 128.6, 126.0 (q, $J=283.7$ Hz), 125.8, 120.4, 113.3, 100.9, 75.8 (q, $J=28.8$ Hz), 55.1, 12.2; IR (KBr) ν : 3304, 3004, 2845, 1611, 1581, 1550, 1516, 1260, 1168, 1150, 941, 925, 827 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{19}\text{H}_{18}\text{F}_3\text{N}_2\text{O}_3$ $[\text{M} + \text{H}]^+$ 379.1264; found 379.1260.

4-(1-([1,1'-Biphenyl]-4-yl)-2,2,2-trifluoro-1-hydroxyethyl)-3-methyl-1-phenyl-1H-pyrazol-5-ol (**3g**): 68.7 mg, 81% yield. White solid, m.p. 119~120 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 7.73~7.64 (m, 7H), 7.49~7.27 (m, 7H), 1.74 (s, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -74.8 (s, 3F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 147.1, 140.0, 139.4, 138.2, 129.1, 129.0, 127.9, 127.7, 126.7, 126.3, 126.0 (q, $J=284.4$ Hz), 125.9, 120.3, 100.9, 75.9 (q, $J=34.0$ Hz), 12.5; IR (KBr) ν : 3420, 3285, 1602, 1572, 1497, 1168, 1156, 843, 701 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{24}\text{H}_{20}\text{F}_3\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 425.1471; found 425.1473.

3-Methyl-1-phenyl-4-(2,2,2-trifluoro-1-(3-fluorophenyl)-1-hydroxyethyl)-1H-pyrazol-5-ol (**3h**): 63.0 mg, 86% yield. White solid, m.p. 142~143 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 7.69 (d, $J=7.6$ Hz, 2H), 7.50~7.44 (m, 3H), 7.38 (d, $J=7.6$ Hz, 1H), 7.33~7.22 (m, 3H), 1.76 (s, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -74.9 (s, 3F), -113.0 (s, 1F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 161.7 (d, $J=241.6$ Hz), 147.1, 141.9 (d, $J=6.5$ Hz), 136.4, 130.0 (d, $J=8.0$ Hz), 129.0, 125.8, 125.6 (q, $J=283.0$ Hz), 123.3, 120.3, 115.2 (d, $J=20.8$ Hz), 114.2 (d, $J=23.4$ Hz), 100.7, 75.6 (q, $J=31.9$ Hz), 12.5; IR (KBr) ν : 3282, 3089, 2925, 1600, 1562, 1500, 1181, 1166, 1143, 827, 717 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{18}\text{H}_{15}\text{F}_4\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 367.1064; found 367.1064.

4-(1-(3-Bromophenyl)-2,2,2-trifluoro-1-hydroxyethyl)-3-methyl-1-phenyl-1H-pyrazol-5-ol (**3i**): 77.8 mg, 91% yield. White solid, m.p. 157~158 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 7.70~7.68 (m, 3H), 7.60 (d, $J=8.0$ Hz, 1H), 7.55 (d, $J=7.6$ Hz, 1H), 7.50~7.46 (m, 2H), 7.41~7.37 (m, 1H), 7.30~7.26 (m, 1H), 1.79 (s, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -74.8 (s, 3F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 147.1, 141.8, 131.2, 130.2, 129.9, 129.0, 126.2, 125.8, 125.5 (q, $J=285.9$ Hz), 121.2, 120.4, 100.7, 75.4 (q, $J=30.2$ Hz), 12.7; IR (KBr) ν : 3306, 3079,

2925, 1602, 1564, 1499, 1172, 1160, 835, 715 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{18}\text{H}_{15}\text{BrF}_3\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 427.0264; found 427.0265.

3-Methyl-1-phenyl-4-(2,2,2-trifluoro-1-hydroxy-1-(*m*-tolyl)ethyl)-1H-pyrazol-5-ol (**3j**): 65.9 mg, 91% yield. White solid, m.p. 121~122 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 7.70~7.68 (m, 2H), 7.51~7.47 (m, 2H), 7.36~7.27 (m, 4H), 7.20 (d, $J=6.8$ Hz, 1H), 2.33 (s, 3H), 1.64 (s, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -74.7 (s, 3F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 146.9, 138.9, 137.1, 129.0, 128.9, 127.8, 125.8, 124.2, 120.1, 100.8, 76.0 (q, $J=28.8$ Hz), 21.1, 11.9; IR (KBr) ν : 3314, 3065, 2928, 1602, 1565, 1498, 1410, 1175, 1149, 816, 716 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{19}\text{H}_{18}\text{F}_3\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 363.1315; found 363.1310.

4-(1-(2,5-Dichlorophenyl)-2,2,2-trifluoro-1-hydroxyethyl)-3-methyl-1-phenyl-1H-pyrazol-5-ol (**3k**): 67.6 mg, 81% yield. White solid, m.p. 158~159 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 8.75 (br, 1H), 7.70~7.66 (m, 4H), 7.55~7.46 (m, 3H), 7.30~7.26 (m, 1H), 1.70 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 146.3, 136.0, 135.2, 134.1, 134.0, 131.5, 129.3, 129.2, 126.7, 125.9, 125.7 (q, $J=285.0$ Hz), 120.0, 98.0, 76.0 (q, $J=29.2$ Hz), 11.7; IR (KBr) ν : 3325, 3076, 2938, 1596, 1580, 1546, 1450, 1174, 1151, 939, 927 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{18}\text{H}_{14}\text{Cl}_2\text{F}_3\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 417.0379; found 417.0380.

3-Methyl-1-phenyl-4-(2,2,2-trifluoro-1-hydroxy-1-(perfluorophenyl)ethyl)-1H-pyrazol-5-ol (**3l**): 77.1 mg, 88% yield. White solid, m.p. 148~149 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 7.65 (d, $J=8.0$ Hz, 2H), 7.48~7.44 (m, 2H), 7.28~7.24 (m, 1H), 2.12 (s, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -77.7 (s, 3F), -136.9 (s, 2F), -153.8~-153.9 (m, 1F), -162.7~-162.8 (m, 2F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 147.1, 145.9, 143.4, 141.6, 139.1, 138.4, 136.1, 129.0, 128.7, 126.1, 125.6, 123.2, 119.8, 114.5, 98.8, 73.9 (q, $J=34.0$ Hz), 12.2; IR (KBr) ν : 3425, 3177, 2922, 1604, 1571, 1525, 1492, 1278, 1193, 1183, 1125, 992, 926, 795 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{18}\text{H}_{11}\text{F}_8\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 439.0687; found 439.0678.

3-Methyl-1-phenyl-4-(2,2,2-trifluoro-1-hydroxy-1-(naphthalen-2-yl)ethyl)-1H-pyrazol-5-ol (**3m**): 67.7 mg, 85% yield. White solid, m.p. 130~131 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 8.10~7.94 (m, 4H), 7.72~7.29 (m, 8H), 1.63 (s, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -75.6 (s, 3F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 147.0, 136.2, 132.4, 132.0, 129.0, 128.5, 127.8, 127.3, 126.7, 126.4, 126.0 (q, $J=285.2$ Hz), 125.8, 125.6, 120.3, 100.6, 76.2 (q, $J=31.1$ Hz), 12.3; IR (KBr) ν : 3264, 3063, 2920, 1560, 1568, 1500, 1154, 827, 751, 711 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{22}\text{H}_{18}\text{F}_3\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 399.1315; found 399.1314.

3-Methyl-1-phenyl-4-(2,2,2-trifluoro-1-hydroxy-1-(1-tosyl-1H-indol-3-yl)ethyl)-1H-pyrazol-5-ol (**3n**): 101.8 mg, 94% yield. White solid, m.p. 148~149 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 8.00 (d, $J=8.4$ Hz, 1H), 7.92 (d, $J=8.4$ Hz, 2H), 7.69~7.65 (m, 3H), 7.49~7.45 (m, 3H),

7.41~7.35 (m, 3H), 7.30~7.26 (m, 1H), 7.22~7.19 (m, 1H), 2.32 (s, 3H), 1.43 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 145.8, 134.3, 133.9, 130.3, 129.1, 128.5, 126.8, 125.9, 125.6 (q, $J=284.7$ Hz), 125.3, 123.6, 123.0, 121.4, 120.1, 119.1, 113.4, 96.9, 72.9 (q, $J=34.0$ Hz), 21.0, 11.7; IR (KBr) ν : 3301, 3076, 2920, 1598, 1560, 1496, 1380, 1177, 1162, 1127, 833, 715, 662 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{27}\text{H}_{23}\text{F}_3\text{N}_3\text{O}_4\text{S}$ $[\text{M} + \text{H}]^+$ 542.1356; found 542.1355.

3-Methyl-1-phenyl-4-(2,2,2-trifluoro-1-hydroxy-1-(thiophen-2-yl)ethyl)-1H-pyrazol-5-ol (**3o**): 68.0 mg, 96% yield. White solid, m.p. 111~112 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 7.70 (d, $J=7.6$ Hz, 2H), 7.65 (d, $J=5.2$ Hz, 1H), 7.50 (t, $J=8.0$ Hz, 2H), 7.33~7.29 (m, 1H), 7.22 (s, 1H), 7.05~7.03 (m, 1H), 1.79 (s, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -77.2 (s, 3F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 146.9, 142.8, 135.8, 129.1, 127.3, 126.2, 126.0, 125.5 (q, $J=285.5$ Hz), 120.3, 99.4, 74.4 (q, $J=31.1$ Hz), 11.9; IR (KBr) ν : 3327, 2933, 1598, 1582, 1544, 1500, 1178, 1161, 1153, 928, 707 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{16}\text{H}_{14}\text{F}_3\text{N}_2\text{O}_2\text{S}$ $[\text{M} + \text{H}]^+$ 355.0723; found 355.0723.

3-Methyl-1-phenyl-4-(1,1,1-trifluoro-2-hydroxybutan-2-yl)-1H-pyrazol-5-ol (**3p**): 53.0 mg, 88% yield. White solid, m.p. 114~115 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 7.69 (d, $J=7.6$ Hz, 2H), 7.69~7.45 (m, 2H), 7.27~7.24 (m, 1H), 2.41~2.32 (m, 1H), 2.26 (s, 3H), 1.77~1.68 (m, 1H), 0.87 (t, $J=7.2$ Hz, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -81.0 (s, 3F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 148.0, 136.3, 129.8 (q, $J=285.0$ Hz), 129.1, 125.5, 119.8, 97.6, 75.2 (q, $J=28.5$ Hz), 24.4, 13.0, 7.2; IR (KBr) ν : 3268, 2981, 1605, 1562, 1500, 1184, 1148, 731, 704 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{14}\text{H}_{15}\text{F}_3\text{N}_2\text{O}_2\text{Na}$ $[\text{M} + \text{Na}]^+$ 323.0978; found 323.0975.

3-Methyl-1-(*p*-tolyl)-4-(2,2,2-trifluoro-1-hydroxy-1-phenylethyl)-1H-pyrazol-5-ol (**4a**): 65.2 mg, 90% yield. White solid, m.p. 143~144 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 7.57~7.53 (m, 4H), 7.44~7.36 (m, 3H), 7.29 (d, $J=8.4$ Hz, 2H), 2.33 (s, 3H), 1.61 (s, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -74.8 (s, 3F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 146.4, 139.1, 135.3, 129.5, 128.4, 128.1, 127.3, 126.0 (q, $J=283.0$ Hz), 120.5, 100.7, 76.1 (q, $J=31.7$ Hz), 20.5, 12.1; IR (KBr) ν : 3261, 3036, 2928, 1593, 1580, 1544, 1516, 1158, 937, 695 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{19}\text{H}_{18}\text{F}_3\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 363.1315; found 363.1310.

1-(4-Methoxyphenyl)-3-methyl-4-(2,2,2-trifluoro-1-hydroxy-1-phenylethyl)-1H-pyrazol-5-ol (**4b**): 63.6 mg, 84% yield. White solid, m.p. 141~142 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 7.58~7.56 (m, 4H), 7.42~7.40 (m, 3H), 7.05 (d, $J=8.0$ Hz, 2H), 3.79 (s, 3H), 1.60 (s, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -74.9 (s, 3F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 157.5, 145.8, 139.1, 128.3, 128.0, 127.3, 122.9, 114.2, 100.4, 76.2, 55.4, 12.2. IR (KBr) ν : 3312, 2973, 1595, 1566, 1510, 1252, 1150, 831, 701 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{19}\text{H}_{18}\text{F}_3\text{N}_2\text{O}_3$ $[\text{M} + \text{H}]^+$ 379.1264; found 379.1258.

1-(4-Fluorophenyl)-3-methyl-4-(2,2,2-trifluoro-1-hydroxy-1-phenylethyl)-1H-pyrazol-5-ol (**4c**): 62.3 mg, 85% yield. White solid, m.p. 131~132 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 7.73~7.35 (m, 9H), 1.64 (s, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -74.8 (s, 3F), -116.2 (s, 1F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 159.9 (d, $J=241.6$ Hz), 147.1, 139.1, 128.4, 128.1, 127.3, 122.8 (d, $J=4.7$ Hz), 115.9 (d, $J=22.8$ Hz), 100.9, 76.1 (q, $J=32.1$ Hz), 12.5; IR (KBr) ν : 3423, 2925, 1563, 1507, 1174, 1153, 846, 711 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{18}\text{H}_{15}\text{F}_4\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 367.1064; found 367.1057.

1-(4-Bromophenyl)-3-methyl-4-(2,2,2-trifluoro-1-hydroxy-1-phenylethyl)-1H-pyrazol-5-ol (**4d**): 80.3 mg, 94% yield. White solid, m.p. 140~141 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 7.68~7.40 (m, 9H), 1.65 (s, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -74.7 (s, 3F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 147.9, 139.0, 131.9, 128.3, 128.0, 127.2, 121.9, 117.9, 101.4, 76.0, 12.5; IR (KBr) ν : 3317, 2928, 1600, 1555, 1490, 1172, 1154, 830, 718 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{18}\text{H}_{15}\text{BrF}_3\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 427.0264; found 427.0262.

3-Methyl-4-(2,2,2-trifluoro-1-hydroxy-1-phenylethyl)-1-(4-(trifluoromethyl)phenyl)-1H-pyrazol-5-ol (**4e**): 71.6 mg, 86% yield. Pale yellow solid, m.p. 186~187 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 7.98~7.96 (m, 2H), 7.86~7.84 (m, 2H), 7.56~7.54 (m, 2H), 7.43~7.36 (m, 3H), 1.70 (s, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -60.7 (s, 3F), -74.7 (s, 3F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 149.0, 139.8, 138.9, 128.3, 127.9, 127.2, 126.3, 126.2, 126.1, 125.5, 125.4, 125.2, 124.3, 122.7, 119.6, 117.4, 101.8, 75.8 (q, $J=30.3$ Hz), 12.6; IR (KBr) ν : 3139, 3089, 1628, 1325, 1159, 1127, 707 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{19}\text{H}_{15}\text{F}_6\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 417.1032; found 417.1024.

3-Methyl-4-(2,2,2-trifluoro-1-hydroxy-1-phenylethyl)-1H-pyrazol-5-ol (**4f**): 49.5 mg, 91% yield. White solid, m.p. 204.5 $^{\circ}\text{C}$ (dec.); ^1H NMR (400 MHz, DMSO- d_6) δ : 10.79 (br, 2H), 7.47~7.45 (m, 2H), 7.37~7.29 (m, 3H), 7.07 (s, 1H), 1.70 (s, 3H); ^{19}F NMR (376 MHz, DMSO- d_6) δ : -74.9 (s, 3F); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 159.0, 140.2, 139.6, 127.9, 127.7, 127.3, 125.7 (q, $J=285.2$ Hz), 100.2, 75.3 (q, $J=30.1$ Hz), 11.5; IR (KBr) ν : 3292, 3184, 1607, 1507, 1179, 1158, 698 cm^{-1} . HRMS (ESI/Orbitrap) calcd for $\text{C}_{12}\text{H}_{12}\text{F}_3\text{N}_2\text{O}_2$ $[\text{M} + \text{H}]^+$ 273.0845; found 273.0840.

4.4 Procedure for the gram-scale synthesis of 3a

The mixture of trifluoroacetophenone (**1a**) (957.7 mg, 5.5 mmol), and 1-phenyl-3-methyl-5-pyrazolone (**2a**) (871.0 mg, 5.0 mmol) in 9.0 mL H_2O were stirred at 50 $^{\circ}\text{C}$. Once the reaction completed, the solid mixture was filtered, washed by water and dried under vacuum to afford 1.585 g (91%) of the desired product **3a**.

Supporting Information ^1H NMR, ^{13}C NMR and ^{19}F NMR spectra of compounds, details for the gram-scale synthesis of **3a**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>. (Please delete this text if not appropriate).

References

- [1] (a) Trost, B. M. *Angew. Chem., Int. Ed.* **1995**, *34*, 259.
 (b) Anastas, P. T.; Williamson, T. C. *Green Chemistry: Frontiers in Chemical Synthesis and Processes*, Oxford University Press, Oxford, **1998**.
 (c) Li, C. J. *Chem. Rev.* **2005**, *105*, 3095.
 (d) Sheldon, R. A. *Green Chem.* **2005**, *7*, 267.
 (e) Gawande, M. B.; Bonifacio, V. D.; Luque, R.; Branco, P. S.; Varma, R. S. *Chem. Soc. Rev.* **2013**, *42*, 5522.
- [2] (a) Lindstrom, U. M. *Chem. Rev.* **2002**, *102*, 2751.
 (b) Kobayashi, S.; Manabe, K. *Acc. Chem. Res.* **2002**, *35*, 209.
 (c) Horvath, I. T.; Anastas, P. T. *Chem. Rev.* **2007**, *107*, 2167.
 (d) Niu, R.; Xiao, J.; Liang, T.; Li, X. *Org. Lett.* **2012**, *14*, 676.
 (e) Shen, Z. L.; Wang, S. Y.; Chok, Y. K.; Xu, Y. H.; Loh, T. P. *Chem. Rev.* **2013**, *113*, 271.
 (f) Butler, R. N.; Coyne, A. G. *Org. Biomol. Chem.* **2016**, *14*, 9945.
 (g) Xiao, J.; Wen, H.; Wang, L.; Xu, L.; Hao, Z.; Shao, C.-L.; Wang, C.-Y. *Green Chem.* **2016**, *18*, 1032.
 (h) Guo, W.; Liu, X.; Liu, Y.; Li, C. *ACS Catal.* **2017**, *8*, 328.
 (i) Kitanosono, T.; Masuda, K.; Xu, P.; Kobayashi, S. *Chem. Rev.* **2018**, *118*, 679.
 (j) Xie, P.; Wang, J.; Liu, Y.; Fan, J.; Wo, X.; Fu, W.; Sun, Z.; Loh, T. P. *Nat. Commun.* **2018**, *9*, 1321.
 (k) Han, J.; Zhang, J.-L.; Zhang, W.-Q.; Gao, Z.; Xu, L.-W.; Jian, Y. *J. Org. Chem.* **2019**, *84*, 7642.
 (l) Wang, W.; Wang, S. X.; Qin, X. Y.; Li, J. T. *Synth. Commun.* **2005**, *35*, 1263.
 (m) Wang, A.-Q.; Jin, T.-S.; Cheng, Z.-L.; Li, T.-S. *Asian J. Chem.* **2010**, *22*, 1973.
 (n) Thakur, P. B.; Meshram, H. M. *RSC Adv.* **2014**, *4*, 6019.
 (o) Zhou, Z.; Duan, J. F.; Mu, X. J.; Xiao, X. Y. *Chin. J. Org. Chem.* **2018**, *38*, 585 (in Chinese).
 (周昱, 段建凤, 穆小静, 肖尚友, 有机化学, **2018**, *38*, 585.)
 (p) Hu, X. Y.; Yu, H. F.; Wang, W. J.; Jiang, S. Q.; Liu, Q.; He, J. *Chin. J. Org. Chem.* **2019**, *39*, 3183 (in Chinese).
 (胡小宇, 于海丰, 王文举, 姜思傲, 刘奇, 何洁, 有机化学, **2019**, *39*, 3183.)
- [3] (a) Narayan, S.; Muldoon, J.; Finn, M. G.; Fokin, V. V.; Kolb, H. C.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2005**, *44*, 3275.
 (b) Paladhi, S.; Chauhan, A.; Dhara, K.; Tiwari, A. K.; Dash, J. *Green Chem.* **2012**, *14*, 2990.
 (c) Paladhi, S.; Bhati, M.; Panda, D.; Dash, J. *J. Org. Chem.* **2014**, *79*, 1473.
 (d) Yu, J. S.; Liu, Y. L.; Tang, J.; Wang, X.; Zhou, J. *Angew. Chem., Int. Ed.* **2014**, *53*, 9512.
 (e) Zhang, Y.; Wei, B. W.; Lin, H.; Zhang, L.; Liu, J. X.; Luo, H. Q.; Fan, X. L. *Green Chem.* **2015**, *17*, 3266.
 (f) SaiPrathima, P.; Srinivas, K.; Rao, M. M. *Green Chem.* **2015**, *17*, 2339.
 (g) Luque-Agudo, V.; Gil, M. V.; Roman, E.; Serrano, J. A. *Green Chem.* **2016**, *18*, 3844.
 (h) Balaraman, K.; Ding, R.; Wolf, C. *Adv. Synth. Catal.* **2017**, *359*, 4165.
 (i) Dilauro, G.; Dell'Aera, M.; Vitale, P.; Capriati, V.; Perna, F. M. *Angew. Chem., Int. Ed.* **2017**, *56*, 10200.
 (j) Zhu, S.; Chen, C. Q.; Xiao, M. Y.; Yu, L. P.; Wang, L.; Xiao, J. *Green Chem.* **2017**, *19*, 5653.
 (k) Hajra, S.; Singha Roy, S.; Aziz, S. M.; Das, D. *Org. Lett.* **2017**, *19*, 4082.
 (l) Hikawa, H.; Ichinose, R.; Kikkawa, S.; Azumaya, I. *Green Chem.* **2018**, *20*, 1297.
 (m) Hikawa, H.; Ichinose, R.; Kikkawa, S.; Azumaya, I. *A. J. Org. Chem.* **2018**, *7*, 416.
 (n) Wang, Z.; Cui, Y.-T.; Xu, Z.-B.; Qu, J. *J. Org. Chem.* **2008**, *73*, 2270.
- [4] (a) Rideout, D. C.; Breslow, R. *J. Am. Chem. Soc.* **1980**, *102*, 7816.
 (b) Narayan, S.; Muldoon, J.; Finn, M. G.; Fokin, V. V.; Kolb, H. C.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2005**, *44*, 3275.
 [5] (a) Gouverneur, V.; Müller, K. In *Fluorine in Pharmaceutical and Medicinal Chemistry: From Biophysical Aspects to Clinical Applications*, Imperial College Press, London, **2012**.
 (b) Jeschke, P. *Pest Manage. Sci.* **2010**, *66*, 10.
 (c) Lectard, S.; Hamashima, Y.; Sodeoka, M. *Adv. Synth. Catal.* **2010**, *352*, 2708.
 (d) Ni, C.; Hu, J. *Chem. Soc. Rev.* **2016**, *45*, 5441.
 (e) Politanskaya, L. V.; Selivanova, G. A.; Panteleeva, E. V.; Tretyakov, E. V.; Platonov, V. E.; Nikul'shin, P. V.; Vinogradov, A. S.; Zonov, Y. V.; Karpov, V. M.; Mezhenkova, T. V.; Vasilyev, A. V.; Koldobskii, A. B.; Shilova, O. S.; Morozova, S. M.; Burgart, Y. V.; Shchegolkov, E. V.; Saloutin, V. I.; Sokolov, V. B.; Aksinenko, A. Y.; Nenajdenko, V. G.; Moskalik, M. Y.; Astakhova, V. V.; Shainyan, B. A.; Tabolin, A. A.; Ioffe, S. L.; Muzalevskiy, V. M.; Balenkova, E. S.; Shastin, A. V.; Tyutyunov, A. A.; Boiko, V. E.; Igumnov, S. M.; Dilman, A. D.; Adonin, N. Y.; Bardin, V. V.; Masoud, S. M.; Vorobyeva, D. V.; Osipov, S. N.; Nosova, E. V.; Lipunova, G. N.; Charushin, V. N.; Prima, D. O.; Makarov, A. G.; Zibarev, A. V.; Trofimov, B. A.; Sobenina, L. N.; Belyaeva, K. V.; Sosnovskikh, V. Y.; Obyednenov, D. L.; Usachev, S. A. *Russ. Chem. Rev.* **2019**, *88*, 425.
 (f) Ren, Z. W.; Ren, N.; Zhang, F. G.; Ma, J. A. *Acta Chim. Sin.* **2018**, *76*, 940 (in Chinese).
 (任智雯, 任楠, 张发光, 马军安, 化学学报, **2018**, *76*, 940.)
 (g) Hu, J. J.; Huang, Y. G.; Xu, X. H.; Qin, F. L. *Chin. J. Org. Chem.* **2019**, *39*, 177 (in Chinese).
 (胡娟娟, 黄焰根, 徐修华, 卿凤翎, 有机化学, **2019**, *39*, 177.)
- [6] (a) Müller, K.; Faeh, C.; Diederich, F. *Science* **2007**, *317*, 1881.
 (b) Purser, S.; Moore, P. R.; Swallow, S.; Gouverneur, V. *Chem. Soc. Rev.* **2008**, *37*, 320.
 (c) Zhou, Y.; Wang, J.; Gu, Z.; Wang, S.; Zhu, W.; Aceña, J. L.; Soloshonok, V. A.; Izawa, K.; Liu, H. *Chem. Rev.* **2016**, *116*, 422.
- [7] (a) Kawai, H.; Okusu, S.; Tokunaga, E.; Sato, H.; Shiro, M.; Shibata, N. *Angew. Chem., Int. Ed.* **2012**, *51*, 4959.
 (b) Merino, E.; Nevado, C. *Chem. Soc. Rev.* **2014**, *43*, 6598.
 (c) Feng, Z.; Min, Q.-Q.; Xiao, Y.-L.; Zhang, B.; Zhang, X. *Angew. Chem., Int. Ed.* **2014**, *53*, 1669.
 (d) Zhou, D.; Huang, Z.; Yu, X.; Wang, Y.; Li, J.; Wang, W.; Xie, H. *Org. Lett.* **2015**, *17*, 5554.
 (e) Xu, X.-H.; Matsuzaki, K.; Shibata, N. *Chem. Rev.* **2015**, *115*, 731.
 (f) Wang, L.; Chen, J.; Huang, Y. *Angew. Chem., Int. Ed.* **2015**, *54*, 15414.
 (g) Huang, Y.-Y.; Yang, X.; Chen, Z.; Verpoort, F.; Shibata, N. *Chem.-Eur. J.* **2015**, *21*, 8664.
 (h) Montesinos-Magraner, M.; Vila, C.; Blay, G.; Fernández, I.; Muñoz, M. C.; Pedro, J. R. *Adv. Synth. Catal.* **2015**, *357*, 3047.
 (i) Feng, Z.; Min, Q.-Q.; Zhao, H.-Y.; Gu, J.-W.; Zhang, X. *Angew. Chem., Int. Ed.* **2015**, *54*, 1270.
 (j) Lou, H. Q.; Wang, Y.; Jin, E.; Lin, X. *J. Org. Chem.* **2016**, *81*, 2019.
 (k) Cook, A. M.; Wolf, C. *Angew. Chem., Int. Ed.* **2016**, *55*, 2929.
 (l) Zhang, Z. M.; Xu, B.; Xu, S.; Wu, H. H.; Zhang, J. *Angew. Chem., Int. Ed.* **2016**, *55*, 6324.
 (m) Feng, Z.; Min, Q.-Q.; Zhang, X. *Org. Lett.* **2016**, *18*, 44.
 (n) Duan, J.; Cheng, Y.; Cheng, J.; Li, R.; Li, P. *Chem.-Eur. J.* **2017**, *23*, 519.
 (o) Holmes, M.; Nguyen, K. D.; Schwartz, L. A.; Luong, T.; Krusche, M. *J. Am. Chem. Soc.* **2017**, *139*, 8114.
 (p) Xie, E.; Rahman, A.; Lin, X. *Org. Chem. Front.* **2017**, *4*, 1407.
 (q) Xu, B.; Zhang, Z.-M.; Xu, S.; Liu, B.; Xiao, Y.; Zhang, J. *ACS Catal.* **2017**, *7*, 210.
 (r) Feng, Z.; Min, Q.-Q.; Fu, X.-P.; An, L.; Zhang, X. *Nat. Chem.* **2017**, *9*, 918.
 (s) Feng, Z.; Xiao, Y.-L.; Zhang, X. *Acc. Chem. Res.* **2018**, *51*, 2264.
 (t) Qing, F.-L. *Chin. J. Org. Chem.* **2012**, *32*, 815 (in Chinese).

- (卿凤翎, 有机化学, **2012**, 32, 815.)
(u) Zhang, F. G.; Peng, X.; Ma, J. A. *Chin. J. Org. Chem.* **2019**, 39, 109 (in Chinese).
(张发光, 彭星, 马军安, 有机化学, **2019**, 39, 109.)
- [8] (a) Gursoy, A.; Demirayak, S.; Capan, G.; Erol, K.; Vural, K. *Eur. J. Med. Chem.* **2000**, 35, 359.
(b) Rubtsov, A. E.; Makhmudov, R. R.; Kovylyayeva, N. V.; Prosyaniuk, N. I.; Bobrov, A. V.; Zalesov, V. V. *Pharm. Chem. J.* **2002**, 36, 608.
(c) Nishio, M.; Matsuda, M.; Ohyanagi, F.; Sato, Y.; Okumura, S.; Tabata, D.; Morikawa, A.; Nakagawa, K.; Horai, T. *Lung Cancer* **2005**, 49, 245.
(d) Bondock, S.; Rabie, R.; Etman, H. A.; Fadda, A. A. *Eur. J. Med. Chem.* **2008**, 43, 2122.
(e) Rostom, S. A.; el-Ashmawy, I. M.; Abd el Razik, H. A.; Badr, M. H.; Ashour, H. M. *Bioorg. Med. Chem.* **2009**, 17, 882.
(f) Bayrak, H.; Demirbas, A.; Demirbas, N.; Karaoglu, S. A. *Eur. J. Med. Chem.* **2010**, 45, 4726.
(g) Fustero, S.; Sanchez-Rosello, M.; Barrio, P.; Simon-Fuentes, A. *Chem Rev.* **2011**, 111, 6984.
(h) Li, J.-H.; Du, D.-M. *Org. Biomol. Chem.* **2013**, 11, 6215.
(i) Ansari, A.; Ali, A.; Asif, M.; Shamsuzzaman. *New J. Chem.* **2017**, 41, 16.
(j) Liu, S.; Bao, X.; Wang, B. *Chem. Commun.* **2018**, 54, 11515.
- [9] (a) Jung, Y. S.; Marcus, R. A. *J. Am. Chem. Soc.* **2007**, 129, 5492.
(b) Thomas, L. L.; Tirado-Rives, J.; Jorgensen, W. L. *J. Am. Chem. Soc.* **2010**, 132, 3097.
(c) Manna, A.; Kumar, A. *J. Phys. Chem. A* **2013**, 117, 2446.
(d) Sela, T.; Vigalok, A. *Org. Lett.* **2014**, 16, 1964.
- [10] (a) Kelly, C. B.; Mercadante, M. A.; Hamlin, T. A.; Fletcher, M. H.; Leadbeater, N. E. *J. Org. Chem.* **2012**, 77, 8131.
(b) Martinelli, E.; Vicini, A. C.; Mancinelli, M.; Mazzanti, A.; Zani, P.; Bernardi, L.; Fochi, M. *Chem. Commun.* **2015**, 51, 658.
- [11] (a) Rassu, G.; Zambrano, V.; Pinna, L.; Curti, C.; Battistini, L.; Sartori, A.; Pelosi, G.; Casiraghi, G.; Zanardi, F. *Adv. Synth. Catal.* **2014**, 356, 2330.
(b) Zhu, Y.-F.; Wei, B.-L.; Wei, J.-J.; Wang, W.-Q.; Song, W.-B.; Xuan, L.-J. *Tetrahedron Lett.* **2019**, 60, 1202.
(c) Stefane B.; Polanc S. *Tetrahedron* **2007**, 63, 10902.
(d) MacLean, M. A.; Diez-Cecilia, E.; Lavery, C. B.; Reed, M. A.; Wang, Y. F.; Weaver, D. F.; Stradiotto, M. *Bioorg. Med. Chem. Lett.* **2016**, 26, 100.
(e) Nishi, H.; Watanabe, T.; Yuki, S.; Morinaka, Y.; Iseki, K.; Sakurai, H. *EP 208874*, **1987**.

(Li, L.; Fan, Y.)