钌(II)-催化偶氮苯与乙醛酸乙酯的环化反应构建 3-羧酸酯吲唑

陈 训^{*,a,b} 王 颖^b 王烁今^{a,b} 孔杜林^b 文丽君^b
翟锐锐^b 赵 珂^b 白丽丽^{*,b} 李友宾^a

(^a 海南医学院 热带转化医学教育部重点实验室 海南省热带药用植物研究开发重点实验室 海口 571199)

(^b 海南医学院药学院 海口 571199)

摘要 报道了一种简便有效的 Ru(II)催化偶氮苯与乙醛酸乙酯的高选择性[4+1]环化反应, 该反应历程涉及 C—H 键的活化. 这种策略能够以中等至良好的产率去高效构建多样性的吲唑骨架结构. 同时研究了该反应的动力学同位素效应, 实验结果表明 C—H 键活化可能不属于整个反应历程的决速步骤.

关键词 钌(II); C—H 键活化; 环化反应; 吲唑

Synthesis of 3-Carboxylate Indazoles via Ru(II)-Catalyzed Annulation of Azobenzenes with Ethyl Glyoxalate

Chen, Xun^{*,a,b} Wang, Ying^b Wang, Shuojin^{a,b} Kong, Dulin^b Wen, Lijun^b
Zhai, Ruirui^b Zhao, Ke^b Bai, Lili^{*,b} Li, Youbin^a

(^a Key Laboratory of Tropical Translational Medicine of Ministry of Education, Hainan Key Laboratory for Research and Development of Tropical Herbs, Hainan Medical University, Haikou 571199)

(^b School of Pharmacy, Hainan Medical University, Haikou 571199)

Abstract The Ru(II)-catalyzed regioselective [4+1] cycloaddition of azobenzenes with ethyl glyoxalate through C—H bond activation has been developed. This method provides a facile approach to various 3-carboxylate indazoles in moderate to good yields. Meantime, the kinetic isotope effect was further investigated and the results indicated that the C—H bond-breaking was possibly not involved in the rate-limiting step of this transformation.

Keywords Ru(II), C—H bond activation, cycloaddition, Indazoles

1 Introduction

Indazole heterocycles have attracted significant interest among the medicinal chemistry community because of their promising pharmaceutical applications and biological activities.^[1] In particular, the 3-carboxylate indazole scaffolds are constituents of key privileged molecules, acting as anti-cancer drug, viral polymerase inhibition and anti-inflammatory agent, *etc.* (Figure 1).^[2] Thus, the development of robust synthetic approaches for affording substituted 3-carboxylate indazoles has been the focus of intensive research. The classical routes to 3-carboxylate indazoles involve direct lithiation at the C3-position of indazoles followed by the addition of electrophiles, but these

strategies are limited by their low yields and harsh reaction conditions.^[3] Therefore, there is a continued strong demand for constructing 3-carboxylate indazoles in simple operation from cheap starting materials.

Transition-metal-catalyzed C—H bond functionalization has emerged a powerful strategy to the rapid assembly of various heterocycles.^[4] In this regard, the recent progress has been focused on the synthesis of indazoles via C—H bond activation process under Rh(III), Co(III) and Re(I) catalysts. For example, the pioneering work by Ellman took advantage of the directing character of azo group to enable a Rh(III)-catalyzed C—H activation/cyclization of azobenzenes with aryl aldehydes for the synthesis of 2,3-diaryl-2*H*-indazoles (Scheme 1).^[5] Moreover,

* Corresponding authors. E-mail: chenxunchenpei@163.com; lisabll@sina.com

Received September 15, 2019; revised November 4, 2019; published online November 21, 2019.

Project supported by the Program of Hainan Association for Science and Technology Plans to Youth R&D Innovation (No. QCXM201808) and the Natural Science Foundation of Hainan Province (No. 819QN227).

海南省科协青年科技英才学术创新计划(No. QCXM201808)和海南省自然科学基金(No. 819QN227)资助项目.

Figure 1 Selected bioactive indazole compounds

Ellman and Wang^[6] also reported representative works on the synthesis of 2,3-diaryl-2H-indazoles via the Co(III)- or Re(I)-catalyzed redox-neutral coupling of azobenzenes with aryl aldehydes, respectively. However, these methods are only limited to furnishing 3-aryl indazole under expensive catalysts. On the other hand, Ru(II) complexes have been shown to efficiently catalyze arene C—H functionalization with outstanding catalytic efficiency.^[7] In continuation of our effort in developing efficient Ru(II)-catalyzed C—H functionalization for the assembly of various heterocycles,^[8] we herein report an efficient Ru(II)-catalyzed C—H bond [4 + 1] cycloaddition of azobenzenes with ethyl glyoxalate for the production of the desired 3-carboxylate indazoles (Scheme 1).

Scheme 1 Indazole synthesis via C—H functionalization

2 Results and discussion

Initially, we commenced our studies by screening reaction conditions for the envisioned C—H bond [4 + 1] cycloaddition of azobenzenes **1a** with ethyl glyoxalate **2a** (Table 1). To our delight, the desired 3-carboxylate indazole **3a** was obtained in 24% yield by using [Ru(*p*-cymene)Cl₂]₂/AgSbF₆ as the co-catalyst (Entry 1). It is well-

known that the OAc source is important for the current C—H functionalization reactions. We next investigated the effects of additives for this reaction optimization. Further investigation revealed that additives such as NaOAc, KOAc, LiOAc, CsOAc and AgOAc led to a significantly increase in the product yield (38%~51%, Entries 2~6), whereas the reaction of **1a** with **2a** in the presence of acid additives such as AcOH and PivOH furnished the product **3a** in slightly decreased yields (41% and 26%, Entries 6, 7). Subsequently, a variety of solvents were screened, and these results showed that 1,2-dichloroethane (DCE) was the optimal solvent (Entries 8~12 vs 13). In addition, **3a** was formed in 76% yield when the reaction temperature increased to 100 °C (Table 1, Entry 14). To our delight, we soon found that extending the reaction time could afford the improved reaction yield of 82% (Entries 15). Finally, various Ru complexes were tested, but only [Ru(*p*-cymene)Cl₂]₂ was active. The other complexes, including RuCl₂(PPh₃)₄, Ru₃(CO)₁₂ and RuHCl(CO)(PPh₃)₃ were totally ineffective (Entries 16~18).

Table 1 Optimization of the reaction parameters^a

Entry	Catalyst	Additive	Solvent	Yield ^b /%
1	[Ru(<i>p</i> -cymene)Cl ₂] ₂	None	Dioxane	24
2	[Ru(<i>p</i> -cymene)Cl ₂] ₂	NaOAc	Dioxane	51
3	[Ru(<i>p</i> -cymene)Cl ₂] ₂	KOAc	Dioxane	46
4	[Ru(<i>p</i> -cymene)Cl ₂] ₂	LiOAc	Dioxane	47
5	[Ru(<i>p</i> -cymene)Cl ₂] ₂	CsOAc	Dioxane	40
6	[Ru(<i>p</i> -cymene)Cl ₂] ₂	AgOAc	Dioxane	38
7	[Ru(<i>p</i> -cymene)Cl ₂] ₂	PivOH	Dioxane	41
8	[Ru(<i>p</i> -cymene)Cl ₂] ₂	AcOH	CH ₃ CN	26
9	[Ru(<i>p</i> -cymene)Cl ₂] ₂	NaOAc	Toluene	Trace
10	[Ru(<i>p</i> -cymene)Cl ₂] ₂	NaOAc	THF	48
11	[Ru(<i>p</i> -cymene)Cl ₂] ₂	NaOAc	TFE	21
12	[Ru(<i>p</i> -cymene)Cl ₂] ₂	NaOAc	MeOH	Trace
13	[Ru(<i>p</i> -cymene)Cl ₂] ₂	NaOAc	DCE	64
14	[Ru(<i>p</i> -cymene)Cl ₂] ₂	NaOAc	DCE	76 ^c
15	[Ru(<i>p</i> -cymene)Cl ₂] ₂	NaOAc	DCE	82 ^d
16	RuCl ₂ (PPh ₃) ₄	NaOAc	DCE	0
17	Ru ₃ (CO) ₁₂	NaOAc	DCE	0
18	RuHCl(CO)(PPh ₃) ₃	NaOAc	DCE	0

^a Unless otherwise noted, all the reactions were carried out using azobenzenes (**1a**, 0.20 mmol) and ethyl glyoxalate (**2a**, 0.40 mmol) with catalysts (5 mol%) in the presence of AgSbF₆ (20 mol%) and additive (50 mol%) in solvent (2.0 mL) at 80 °C for 12 h under Ar in a sealed reaction tube, followed by flash chromatography on SiO₂. ^b Isolated yield. ^c The reaction temperature is 100 °C. ^d The reaction time is 24 h. TFE = 2,2,2-trifluoroethanol.

With an optimized catalytic system in hand, we then

turned our attention toward exploring scope and limitation of Ru(II)-catalyzed C—H bond [4+1] cycloaddition of azobenzenes (**1**) with ethyl glyoxalate (**2a**). As shown in Table 2, the reactivity of substituted azobenzenes **1** was obviously dependent on the electronic properties of substituents. Azobenzenes bearing electron-donating groups (Me, MeO) on the *para*-position afforded excellent yields of **3b** (84% yield) and **3c** (88% yield), respectively. In contrast, electron-withdrawing halogen group (Cl), ester group (COOEt) and cyano group (CN) slightly decreased the reaction conversion and produced **3d**~**3f** in moderate to good yields (59%~77% yields), which indicated that C—H activation was involved in an electrophilic aromatic substitution process. Notably, this reaction protocol could also smoothly convert *ortho*- or *meta*-substituted azobenzenes to the desired **3g** or **3h** with excellent yields (80% and 83%) in spite of steric congestion. Moreover, 3,5-dimethyl-substituted azobenzene **1i** could afford the desired product **3i** with 78% yield, in which no significant effect of steric hindrance from methyl moiety was observed in this transformation. In addition, the unsymmetrical azobenzene **1j** having both methyl group and methoxy group on one aryl ring is also allowed for this transformation to assemble the corresponding indazole **3j** in 76% yield. Beside, methyl 2-oxoacetate and 2-oxo-2-phenylacetaldehyde could also react with **1a** to afford the corresponding products **3k** and **3l**, respectively. Unfortunately, other aldehydes such as acetaldehyde were not allowed for this transformation, and the starting materials were recovered completely. When we moved to employ 2-arylpyridine as an arene substrate (Eq. 1), indolizine products **5a**~**5c** were also isolated in accepted yield under slightly modified reaction condition. Finally, we converted **3k** to the corresponding product **6a** in 78% yield via a hydrolysis and subsequent decarboxylation process (Eq. 2).

To gain more insights into the mechanism, some experiments were conducted (Scheme 2). A competition reaction was performed to determine the electronic preference of this transformation. When **1c** and **1f** were treated at the same time, the reaction favored the electron-rich azobenzenes in 3.2 : 1 ratio (Scheme 2a), thus implying

 Table 2 Substrate scope^{a,b}

^a All the reactions were carried out using azobenzenes (**1**) (0.20 mmol) and aldehyde (**2**) (0.40 mmol) with [Ru(*p*-cymene)Cl₂] (5 mol%) in the presence of AgSbF₆ (20 mol%) and NaOAc (50 mol %) in DCE (2.0 mL) at 100 °C for 24 h under Ar in a sealed reaction tube. Followed by flash chromatography on SiO₂. ^b Isolated yield.

that the C—H activation step might be an electrophilic aromatic substitution process.^[9] Subsequently, the intermolecular isotope effect ($K_H/K_D=1.1$) further indicated that C—H bond-breaking was not involved in the rate-limiting step of this transformation (Scheme 2b).^[10] On the basis of the above-mentioned results, a plausible mechanism is proposed in Scheme 3. First, coordination of an azo group in azobenzene **1a** to a cationic Ru species to form a five-membered cobaltacycle intermediate **I**. Subsequently, an insertion of C=O to Ru—C bond of interme-

(a) Azobenzenes substrate competition

(b) Kinetic isotope study

Scheme 2 Preliminary mechanistic studies

diolate **A** forms the seven-membered intermediate **II**, which undergoes protonation to give the alcohol **III** with release of Ru catalyst. Finally, indazole **3a** is obtained through an intramolecular nucleophilic substitution/cyclization process.

Scheme 3 Proposed mechanism

3 Conclusions

In conclusion, we have developed a novel Ru(II)-catalyzed regioselective C—H bond cycloaddition of azobenzenes with ethyl glyoxalate for the rapid assembly

of 3-carboxylate indazole. This catalytic system tolerates a wide range of azobenzene substrates with high reaction efficiency. Further synthetic applications to bioactive molecules are underway in our laboratory.

4 Experimental section

4.1 General information

The azobenzene substrates were prepared according to the previously reported procedure.^[5] Solvents were treated with 4 Å molecular sieves or sodium and distilled prior to use. Purifications of reaction products were carried out by flash chromatography using silica gel (200~300 mesh). ¹H NMR and ¹³C NMR spectra were recorded with tetramethylsilane (TMS) as internal standard at ambient temperature unless otherwise indicated on a Bruker Avance DPX 600 fourier Transform spectrometer operating at 400 MHz for ¹H NMR and 100 MHz for ¹³C NMR. High resolution mass spectra (HRMS) were recorded on an IF-TOF spectrometer (Micromass).

4.2 General procedure for the synthesis of 3-carboxylate indazole derivatives 3

A 10 mL of reaction tube was charged with azobenzenes **1** (0.2 mmol), [Ru(*p*-cymene)Cl₂] (5 mol%), AgSbF₆ (20 mol%), NaOAc (50 mol%) and DCE (1.5 mL) under Ar atmosphere. Then ethyl glyoxalate **2a** (0.4 mmol) in DCE (0.5 mL) was added in one-pot under Ar and the mixture was stirred at 100 °C for 24 h. The corresponding reaction mixture was filtered through a pad of Celite, washed with DCE and concentrated under reduced pressure. The residue was purified by flash chromatography on silica gel using ethyl acetate/petroleum ether (*V*:*V*=1:8) as eluent to afford the products **3**.

Ethyl 2-phenyl-2*H*-indazole-3-carboxylate (**3a**): Brown solid, 82% yield. m.p. 80~82 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.12 (d, *J*=8.3 Hz, 1H), 7.85 (d, *J*=8.5 Hz, 1H), 7.52 (s, 5H), 7.41 (t, *J*=7.1 Hz, 1H), 7.34 (t, *J*=7.4 Hz, 1H), 4.36 (q, *J*=13.1, 2H), 1.33 (t, *J*=6.2 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ: 159.5, 148.4, 141.0, 129.3, 128.6, 127.1, 126.4, 125.5, 125.1, 124.01, 121.6, 118.6, 61.1, 14.1; HRMS (ESI) calcd for C₁₆H₁₅N₂O₂ [M+H]⁺ 267.1132, found 267.1135.

Ethyl 5-methyl-2-(*p*-tolyl)-2*H*-indazole-3-carboxylate (**3b**): Brown solid, 84% yield. m.p. 68~70 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.85 (s, 1H), 7.73 (d, *J*=8.6 Hz, 1H), 7.39 (d, *J*=7.3 Hz, 2H), 7.30 (d, *J*=7.5 Hz, 2H), 7.24 (d, *J*=9.1 Hz, 1H), 4.36 (q, *J*=6.6 Hz, 2H), 2.50 (s, 3H), 2.44 (s, 3H), 1.34 (t, *J*=6.8 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ: 159.7, 147.4, 139.2, 138.7, 135.3, 129.8, 129.2, 126.1, 124.4, 119.7, 118.2, 60.9, 22.2, 21.3, 14.2; HRMS (ESI) calcd for C₁₈H₁₉N₂O₂ [M+H]⁺ 295.1631, found 295.1636.

Ethyl 5-methoxy-2-(4-methoxyphenyl)-2*H*-indazole-3-carboxylate (**3c**): Brown solid, 88% yield. m.p. 88~90 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.64 (d, *J*=9.1 Hz, 1H), 7.35 (d, *J*=7.6 Hz, 2H), 7.26 (s, 1H), 7.01 (d, *J*=9.2

H₂, 1H), 6.93 (d, $J=7.6$ Hz, 2H), 4.30~4.21 (q, 2H), 3.83 (d, $J=8.6$ Hz, 3H), 3.80 (s, 3H), 1.25 (t, $J=6.5$ Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 160.0, 159.8, 157.9, 145.0, 134.2, 127.4, 125.1, 124.1, 121.8, 119.9, 113.7, 97.9, 60.8, 55.6, 55.4, 14.2; HRMS (ESI) calcd for C₁₈H₁₉N₂O₄ [M+H]⁺ 327.1328, found 327.1332.

Ethyl 5-chloro-2-(4-chlorophenyl)-2H-indazole-3-carboxylate (**3d**): Brown solid, 77% yield. m.p. 186~188 °C; ¹H NMR (400 MHz, CDCl₃) δ : 8.08 (s, 1H), 7.77 (d, $J=9.1$ Hz, 1H), 7.56~7.44 (m, 5H), 7.36 (d, $J=9.0$ Hz, 1H), 4.39 (q, $J=7.0$ Hz, 2H), 1.38 (t, $J=7.0$ Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 159.0, 146.9, 139.1, 135.6, 131.7, 129.4, 128.9, 127.6, 124.4, 124.2, 120.5, 120.1, 61.5, 14.2; HRMS (ESI) calcd for C₁₆H₁₃Cl₂N₂O₂ [M+H]⁺ 335.0355, found 335.0359.

Diethyl 2-(4-(ethoxycarbonyl)phenyl)-2H-indazole-3,5-dicarboxylate (**3e**): Brown solid, 68% yield. m.p. 139~141 °C; ¹H NMR (400 MHz, CDCl₃) δ : 8.91 (s, 1H), 8.23 (d, $J=7.3$ Hz, 2H), 8.05 (d, $J=8.9$ Hz, 1H), 7.86 (d, $J=8.8$ Hz, 1H), 7.63 (d, $J=7.8$ Hz, 2H), 4.51~4.36 (m, 6H), 1.42 (m, 9H); ¹³C NMR (101 MHz, CDCl₃) δ : 166.5, 165.6, 158.9, 149.8, 143.9, 131.6, 130.6, 130.1, 127.8, 127.3, 127.2, 126.3, 125.7, 123.4, 122.9, 118.5, 61.7, 61.4, 61.3, 14.3, 14.1; HRMS (ESI) calcd for C₂₂H₂₃N₂O₆ [M+H]⁺ 411.1501, found 411.1508.

Ethyl 5-cyano-2-(4-cyanophenyl)-2H-indazole-3-carboxylate (**3f**): Brown solid, 59% yield. m.p. 192~194 °C; ¹H NMR (400 MHz, CDCl₃) δ : 9.81 (s, 1H), 8.07 (s, 1H), 7.93 (d, $J=8.5$ Hz, 1H), 7.84 (d, $J=8.6$ Hz, 1H), 7.24~7.03 (m, 3H), 4.40 (q, 2H), 1.42 (t, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 162.5, 142.2, 141.1, 132.7, 131.4, 128.8, 127.7, 127.3, 121.9, 120.9, 119.9, 118.0, 117.9, 117.5, 117.3, 104.4, 59.6, 14.9; HRMS (ESI) calcd for C₁₈H₁₃N₄O₂ [M+H]⁺ 317.1064, found 317.1070.

Ethyl 7-methyl-2-(o-tolyl)-2H-indazole-3-carboxylate (**3g**): Brown solid, 80% yield. m.p. 74~76 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.97 (d, $J=8.4$ Hz, 1H), 7.42 (dd, $J=10.5, 4.2$ Hz, 1H), 7.38~7.26 (m, 4H), 7.18 (d, $J=6.8$ Hz, 1H), 4.30 (q, 2H), 2.69 (s, 3H), 2.01 (s, 3H), 1.27 (t, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 159.4, 148.8, 140.8, 135.0, 130.5, 129.6, 128.9, 127.0, 126.3, 126.1, 125.7, 123.3, 119.0, 60.9, 17.1, 17.0, 14.0; HRMS (ESI) calcd for C₁₈H₁₉N₂O₂ [M+H]⁺ 295.1446, found 295.1453.

Ethyl 6-methyl-2-(m-tolyl)-2H-indazole-3-carboxylate (**3h**): Brown solid, 83% yield. m.p. 85~87 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.99 (d, $J=8.6$ Hz, 1H), 7.58 (s, 1H), 7.41~7.26 (m, 4H), 7.17 (d, $J=8.6$ Hz, 1H), 4.35 (q, $J=7.0$ Hz, 2H), 2.49 (s, 3H), 2.43 (s, 3H), 1.33 (t, $J=7.0$ Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 159.5, 149.0, 141.0, 138.7, 137.0, 130.0, 128.4, 128.3, 126.8, 124.9, 123.5, 122.4, 121.0, 116.8, 61.0, 22.1, 21.3, 14.2; HRMS (ESI) calcd for C₁₈H₁₉N₂O₂ [M+H]⁺ 295.1508, found 295.1514.

Ethyl 2-(3,5-dimethylphenyl)-4,6-dimethyl-2H-indazole-3-carboxylate (**3i**): Brown solid, 78% yield. m.p. 67~69 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.38 (s, 1H), 7.26 (s, 1H), 7.12 (s, 2H), 6.85 (s, 1H), 4.25 (q, $J=7.1$ Hz, 2H),

2.61 (s, 3H), 2.42 (s, 3H), 2.37 (s, 6H), 1.13 (t, $J=7.1$ Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 165.8, 142.0, 132.4, 131.2, 130.0, 123.3, 121.9, 119.1, 115.8, 104.3, 98.2, 60.8, 17.9, 17.5, 13.7, 11.0; HRMS (ESI) calcd for C₂₀H₂₃N₂O₂ [M+H]⁺ 323.1791, found 323.1795.

Ethyl 2-(3-methoxy-5-methylphenyl)-2H-indazole-3-carboxylate (**3j**): Brown solid, 76% yield. m.p. 95~97 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.73 (d, $J=9.2$ Hz, 1H), 7.33 (dt, $J=30.7, 9.0$ Hz, 5H), 7.09 (d, $J=9.0$ Hz, 1H), 4.38 (q, 2H), 3.92 (s, 3H), 2.45 (s, 3H), 1.32 (t, $J=6.7$ Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 159.8, 157.9, 145.1, 139.1, 138.7, 129.1, 126.0, 125.1, 121.9, 120.0, 97.9, 60.8, 55.4, 21.3, 14.2; HRMS (ESI) calcd for C₁₈H₁₉N₂O₃ [M+H]⁺ 311.1338, found 311.1342.

Methyl 2-phenyl-2H-indazole-3-carboxylate (**3k**): Brown solid, 74% yield. m.p. 79~81 °C; ¹H NMR (500 MHz, CDCl₃) δ : 8.11 (d, $J=8.2$ Hz, 1H), 7.86 (d, $J=8.3$ Hz, 1H), 7.53 (s, 5H), 7.42 (t, $J=7.2$ Hz, 1H), 7.35 (t, $J=7.2$ Hz, 1H), 3.92 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ : 160.0, 148.4, 140.9, 129.4, 128.7, 127.1, 126.33, 125.6, 124.83, 123.9, 121.5, 118.6, 52.0; HRMS (ESI) calcd for C₁₅H₁₃N₂O₂ [M+H]⁺ 253.1901, found 253.1908.

Phenyl(2-phenyl-2H-indazol-3-yl)methanone (**3l**): Brown solid, 32% yield. m.p. 153~155 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.89~7.85 (m, 3H), 7.60 (t, $J=7.5$ Hz, 1H), 7.55~7.52 (m, 2H), 7.47~7.35 (m, 7H), 7.19~7.16 (m, 1H); ¹³C NMR (101 MHz, CDCl₃) δ : 168.9, 138.9, 132.5, 130.3, 126.9, 125.9, 124.0, 123.3, 123.2, 122.9, 121.7, 120.5, 120.1, 119.3, 116.5, 114.9; HRMS (ESI) calcd for C₂₀H₁₅N₂O [M+H]⁺ 299.3208, found 299.3213.

Ethyl pyrido[2,1-*a*]isoindole-6-carboxylate (**5a**): Yellow solid, 36% yield. m.p. 65~67 °C; ¹H NMR (400 MHz, CDCl₃) δ : 9.98 (d, $J=6.6$ Hz, 1H), 8.27 (d, $J=8.5$ Hz, 1H), 8.10 (ddt, $J=12.0, 8.2, 1.1$ Hz, 2H), 7.55 (m, 1H), 7.29~7.27 (m, 1H), 7.27~7.25 (m, 1H), 7.21 (td, $J=6.9, 1.5$ Hz, 1H), 4.51 (q, 2H), 1.52 (t, $J=7.2$ Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 150.1, 126.3, 124.8, 122.5, 122.1, 117.3, 116.6, 115.9, 115.6, 114.9, 114.0, 113.9, 103.3, 67.6, 31.9; HRMS (ESI) calcd for C₁₅H₁₄N₂O₂ [M+H]⁺ 240.3609, found 240.3612.

Ethyl 8-methoxypyrido[2,1-*a*]isoindole-6-carboxylate (**5b**): Yellow solid, 27% yield. m.p. 94~96 °C; ¹H NMR (400 MHz, CDCl₃) δ : 9.88 (s, 1H), 8.00~7.91 (m, 2H), 7.60 (s, 1H), 7.25 (m, 1H), 7.12 (td, $J=7.0, 1.4$ Hz, 1H), 6.90 (dd, $J=8.8, 2.3$ Hz, 1H), 4.49 (q, $J=7.1$ Hz, 3H), 3.95 (s, 3H), 1.51 (t, $J=7.1$ Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 150.1, 148.3, 126.5, 122.3, 117.8, 116.7, 113.4, 112.9, 111.1, 110.9, 103.1, 99.0, 67.5, 64.2, 31.9; HRMS (ESI) calcd for C₁₆H₁₆N₂O₃ [M+H]⁺ 270.1681, found 270.1685.

Ethyl 8-acetylpyrido[2,1-*a*]isoindole-6-carboxylate (**5c**): Yellow solid, 21% yield. m.p. 113~115 °C; ¹H NMR (400 MHz, CDCl₃) δ : 9.90 (d, $J=6.7$ Hz, 1H), 8.85 (s, 1H), 8.11~8.07 (m, 2H), 7.79 (dd, $J=8.6, 1.5$ Hz, 1H), 7.33~7.27 (m, 1H), 7.24 (td, $J=6.9, 1.5$ Hz, 1H), 4.51 (q, $J=7.1$ Hz, 2H), 2.72 (s, 3H), 1.54 (t, $J=7.2$ Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 179.0, 149.7, 129.2, 125.7,

123.9, 122.1, 118.1, 117.6, 116.2, 115.7, 115.2, 114.6, 114.5, 104.1, 67.9, 41.4, 31.8; HRMS (ESI) calcd for $C_{17}H_{15}NO_3 [M+H]^+$ 282.1131, found 282.1134.

2-Phenyl-2H-indazole (**6a**): Yellow solid, 36% yield. m.p. 83~85 °C; 1H NMR (500 MHz, $CDCl_3$) δ : 8.43 (s, 1H), 7.93 (d, $J=7.7$ Hz, 2H), 7.83 (d, $J=8.8$ Hz, 1H), 7.73 (d, $J=8.5$ Hz, 1H), 7.55 (t, $J=7.9$ Hz, 2H), 7.42 (t, $J=7.4$ Hz, 1H), 7.38~7.33 (m, 1H), 7.15 (dd, $J=8.0, 7.0$ Hz, 1H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 149.8, 140.5, 129.6, 127.9, 126.8, 122.8, 122.4, 121.0, 120.4, 120.4, 118.0; MS (ESI) m/z : 195.09.

Supporting Information The kinetic isotope effect (KIE) experiments and the 1H NMR and ^{13}C NMR spectra of all products. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- (a) De Angelis, M.; Stossi, F.; Carlson, K. A.; Katzenellenbogen, B. S.; Katzenellenbogen, K. A. *J. Med. Chem.* **2005**, *48*, 1132.
(b) A Haddadin, M. J.; Conrad, W. E.; Kurth, M. J. *Mini-Rev. Med. Chem.* **2012**, *12*, 1293.
(c) Hu, J.-T.; Cheng, Y.-F.; Yang, Y.-Q.; Rao, Y. *Chem. Commun.* **2011**, *47*, 10133.
(d) Magano, J.; Waldo, M.; Greene, D.; Nord, E. *Org. Process Res. Dev.* **2008**, *12*, 877.
- (a) De Lena, M.; Lorusso, V.; Latorre, A.; Fanizza, G.; Gargano, G.; Caporusso, L.; Guida, M.; Catino, A.; Crucitta, E.; Sambiasi, D.; Mazzei, A. *Eur. J. Cancer* **2001**, *37*, 364.
(b) Atta-ur-Rahman; Malik, S.; Cun-heng, H.; Clardy, J. *Tetrahedron Lett.* **1985**, *26*, 2759.
(c) Halim, R.; Harding, M.; Hufton, R.; Morton, C. J.; Jahangiri, S.; Pool, B. R.; Jaynes, T. P.; Draffan, A. G.; Lilly, M. J.; Frey, B. *WO 2012051659*, **2011**.
(d) Steffan, R. J.; Matelan, E. M. *WO 2006050006*, **2006**.
- (a) Luo, G.; Chen, L.; Dubowchik, G. J. *Org. Chem.* **2006**, *71*, 5392.
(b) Unsinn, A.; Knochel, P. *Chem. Commun.* **2012**, *48*, 2680.
(c) Bunnell, A.; Yang, C.; Petrica, A.; Soth, M. J. *Synth. Commun.* **2006**, *36*, 285.
- For selected reviews, see:
(a) Ackermann, C. *Acc. Chem. Res.* **2014**, *47*, 281.
(b) Song, G.; Wang, F.; Li, X. *Chem. Soc. Rev.* **2012**, *41*, 3651.
(c) Huang, Z.; Lim, H. N.; Mo, F.; Young, M. C.; Dong, G. *Chem. Soc. Rev.* **2015**, *44*, 7764.
(d) Shen, C.; Zhang, P.; Sun, Q.; Bai, S.; Hor, T. S. A.; Liu, X. *Chem. Soc. Rev.* **2015**, *44*, 291.
(e) Chen, Z.; Wang, B.; Zhang, J.; Yu, W.; Liu, Z.; Zhang, Y. *Org. Chem. Front.* **2015**, *2*, 1107.
- (f) Song, G.; Li, X. *Acc. Chem. Res.* **2015**, *48*, 1007.
(g) Zhang, F.; Spring, D. R. *Chem. Soc. Rev.* **2014**, *43*, 6906.
(h) Wencel-Delord, J.; Glorius, F. *Nat. Chem.* **2013**, *5*, 369.
(i) Chen, X.; Bai, L.; Zeng, W. *Chin. J. Org. Chem.* **2018**, *38*, 1859 (in Chinese).
(陈训, 白丽丽, 曾伟, 有机化学, **2018**, *38*, 1859.)
(j) Zhang, Z.; Butt, N. A.; Zhou, M.; Liu, D.; Zhang, W. *Chin. J. Chem.* **2018**, *36*, 443.
(k) Jeong, T.; Han, S. H.; Han, S.; Sharma, S.; Park, J.; Lee, J.; Kwak, J. H.; Jung, Y. H.; Kim, I. S. *Org. Lett.* **2016**, *18*, 232.
(l) Chen, W.; Yang, W.; Wu, R.; Yang, D. *Green Chem.* **2018**, *20*, 2512.
(m) Yang, W.; Chen, Y.; Yao, Y.; Yang, X.; Lin, Q.; Yang, D. *J. Org. Chem.* **2019**, *84*, 11080.
(n) Yao, Y.; Yang, W.; Lin, Q.; Yang, W.; Li, H.; Wang, L.; Gu, F.; Yang, D. *Org. Chem. Front.* **2019**, *6*, 3360.
- Lian, Y.; Bergman, R. G.; Lavis, L. D.; Ellman, J. A. *J. Am. Chem. Soc.* **2013**, *135*, 7122.
- (a) Hummel, J. R.; Ellman, J. A. *J. Am. Chem. Soc.* **2015**, *137*, 490.
(b) Geng, X.; Wang, C. *Org. Lett.* **2015**, *17*, 2434.
- (a) T. Gensch, M. N. Hopkinson, F. Glorius, J. Wencel-Delord, *Chem. Soc. Rev.* **2016**, *45*, 2900.
(b) Suzuki, C.; Morimoto, K.; Hirano, K.; Satoh, T.; Miura, M. *Adv. Synth. Catal.* **2014**, *356*, 1521.
(c) Itoh, M.; Hashimoto, Y.; Hirano, K.; Satoh, T.; Miura, M. *J. Org. Chem.* **2013**, *78*, 8098.
(d) Suzuki, C.; Hirano, K.; Satoh, T.; Miura, M. *Org. Lett.* **2013**, *15*, 3990.
(e) Kong, D.; Lu, G.; Wu, M.; Shi, Z.; Lin, Q. *ACS Sustainable Chem. Eng.* **2017**, *5*, 3465.
(f) De Sarkar, S.; Liu, W.; Kozhushkov, S. I.; Ackermann, L. *Adv. Synth. Catal.* **2014**, *356*, 1461.
(g) Zhang, Y.; Yang, Z.; Yu, X.; Cheng, X.; Li, W.; Guo, L.; Hai, L.; Guo, L.; Wu, Y. *Chin. J. Org. Chem.* **2018**, *38*, 3211 (in Chinese).
(张勇, 杨中振, 余昕玲, 程旭, 李伟剑, 郭令妹, 海俐, 郭丽, 吴勇, 有机化学, **2018**, *38*, 3211.)
(h) Xu, W.; Wang, N.; Zhang, M.; Shi, D. *Chin. J. Org. Chem.* **2019**, *39*, 1735 (in Chinese).
(徐文韬, 王宁, 张梦烨, 史达清, 有机化学, **2019**, *39*, 1735.)
- Bai, L.; Wang, Y.; Wang, S.; Kong, D.; Fu, Y.; Peng, D.; Wen, L.; Chen, X. *Chin. J. Org. Chem.* **2018**, *38*, 3242 (in Chinese).
(白丽丽, 王颖, 王烁今, 孔杜林, 付艳, 彭德乾, 文丽君, 陈训, 有机化学, **2018**, *38*, 3242.)
- (a) Li, B.; Xu, H.; Wang, H.; Wang, B. *ACS Catal.* **2016**, *6*, 3856.
(b) Huang, X.; Huang, J.; Du, C.; Zhang, X.; Song, F.; You, J. *Angew. Chem., Int. Ed.* **2013**, *52*, 12970.
(c) Thirunavukkarasu, V. S.; Raghuvanshi, K.; Ackermann, L. *Org. Lett.* **2013**, *15*, 3286.
- (a) Chen, X.; Hu, X.; Deng, Y.; Jiang, H.; Zeng, W. *Org. Lett.* **2016**, *18*, 4742.
(b) Jeon, B.; Yeon, U.; Son, J. Y.; Lee, P. H. *Org. Lett.* **2016**, *18*, 4610.

(Li, L.; Fan, Y.)