三嗪衍生物荧光探针对 Zr^{4+} , Fe^{3+} 和丙酮的识别

马学林^{a,b} 韩利民^{*,a} 张骁勇^{*,b} 张玉恒^b 王 丽^b
杨 坤^b 冀 婕^b

(^a 内蒙古工业大学化工学院 呼和浩特 010051)

(^b 包头师范学院化学学院 内蒙古包头 014030)

摘要 一锅法合成了 2,2',2''-(1,3,5-三嗪-2,4,6-三亚胺基)三苯甲酸荧光响应化学传感器(L, **H₃TATIB**), 系统地研究了化合物 L 对过渡金属离子的识别性能. 实验结果显示, 在 *N,N*-二甲基甲酰胺(DMF)水溶液中实现了对 Fe^{3+} 和 Zr^{4+} 的识别; 化合物 L 在有机溶剂识别中能够有效地识别丙酮. 化合物 L 对 Zr^{4+} 和 Fe^{3+} 的检测限分别为 3.60×10^{-6} 和 1.33×10^{-6} mol/L. 实验初步探究了化合物 L 作为荧光探针对人体尿液和水样中 Fe^{3+} 离子的识别.

关键词 荧光传感器; Zr^{4+} 离子; Fe^{3+} 离子; 丙酮; 1,3,5-三嗪衍生物

Triazine Derivative for Fluorescence Sensing of Zr^{4+} , Fe^{3+} Ions and Acetone

Ma, Xuelin^{a,b} Han, Limin^{*,a} Zhang, Xiaoyong^{*,b} Zhang, Yuheng^b
Wang, Li^b Yang, Kun^b Ji, Jie^b

(^a Chemical Engineering College, Inner Mongolia University of Technology, Hohhot 010051)

(^b Department of Chemistry, Baotou Teachers' College, Baotou, Inner Mongolia 014030)

Abstract 2,2',2''-(1,3,5-Triazine-2,4,6-triimino)tribenzoic acid (L, **H₃TATIB**) fluorescence response chemosensor has been synthesized by one pot reaction and its fluorescent behaviors to transition metal ions were systematically investigated. The experimental results show that the recognition of Zr^{4+} and Fe^{3+} is realized in *N,N*-dimethylformamide (DMF) aqueous solution; The compound L can effectively detect acetone in organic solvent. The detection limits of Zr^{4+} and Fe^{3+} are 3.60×10^{-6} and 1.33×10^{-6} mol/L, respectively. It was preliminarily explored that compound L was used as a fluorescent probe to identify Fe^{3+} ion in human urine and water samples.

Keywords fluorescent sensor; Zr^{4+} ion; Fe^{3+} ion; acetone; 1,3,5-triazine derivatives

1 Introduction

The sensing and detection of hazardous substances plays an important role in life and environmental sciences. The past decade has witnessed a fast growth for detection of Fe^{3+} ion because massive utilization of Fe^{3+} ion causes environmental pollution and health problems.^[1~7] It is well known that the Fe^{3+} ion plays an important role in biological metabolism and are recognized as an industrial pollutant. However, these widely used instrumental analytical techniques have some limitations, including expensive

equipment, complication in sample preparation and well-trained person. To defeat this problem, the large number of well-designed sensors based on this Fe^{3+} ion have been developed by fluorescent^[8~18] and colorimetric^[19] sensing methods, of which sensitive and selective detections of Fe^{3+} ion by fluorescent methods are most competitive and have attracted great attention in recent years. Although metal-organic frameworks (MOFs) as fluorescent probe have been widely investigated for their potential application in the sensitive and selective detection of Fe^{3+} ion, many MOFs are limited in wide application

* Corresponding authors. E-mail: 66123@bttc.edu.cn; hanlimin@imut.edu.cn

Received December 6, 2019; revised February 27, 2020; published online March 31, 2020.

Project supported by the Natural Science Foundation of Inner Mongolia (Nos. 2018BS02009, 2019MS02031), the Young Innovative Talents in Baotou City (No. 30324001), the National Natural Science Foundation of China (Nos. 21767021, 51563019) and the Education Department Project of Inner Mongolia (No. NJZY19187).

内蒙古自然科学基金(Nos. 2018BS02009, 2019MS02031)、包头市青年创新人才(No. 30324001)、国家自然科学基金(Nos. 21767021, 51563019)和内蒙古教育厅(No. NJZY19187)资助项目.

due to complicate operations, slow signal response, poor stability,^[20] expensive raw materials,^[21–24] toxic (including heavy metal)^[25,26] and not suitable for aqueous and biological environment. In addition, though Zr^{4+} is not classified as one of the essential elements for human and cannot also produce toxic effects for environment, it is a rare metal and very important atomic energy elements which is widely used in military industry, smelting, atomic energy industry and battery.^[27,28] However, up to now, there is only a small amount of detection of Zr^{4+} ion,^[29] mostly the recognition of Fe^{3+} ion by Zr(IV)-MOFs .^[30,31] Besides, MOFs are also used as fluorescent probes to detect small molecular compounds, such as H_2S ,^[32] biomolecules,^[33] acetone^[34,35] and nitroaromatic compounds.^[36–38] Therefore, there is an urgent need to develop a new chemosensor for the rapid selective and sensitive detection of Zr^{4+} , Fe^{3+} ion and acetone as the replacement.

In this work, it reports an organic fluorescent probe **L** (no heavy metal ions and low toxicity) which is rationally designed on the basis of 2,4,6-trichloro-1,3,5-triazine and contains nine coordination centers (Figure 1). It exhibits strong fluorescence and highly sensitive and selective toward Zr^{4+} , Fe^{3+} ions and acetone in room temperature. With the addition of Fe^{3+} ion, compound **L** could give rise to a visible colorless-to-yellow change and clear fluorescence quenching. In contrast to this, with the addition of Zr^{4+} ion, the compound shows fluorescence enhancement. This is the first example of fluorescent organic compound chemosensor for the highly selective detection of Zr^{4+} , Fe^{3+} ions and acetone simultaneously based on 1,3,5-triazine derivatives with the highest density of uncoordinated N and carboxylate O atoms. This study might provide a new method for the rapid detection of Zr^{4+} , Fe^{3+} ions and acetone with the advantages of simple sample pretreatment processes, highly sensitive and selective sensing.

Figure 1 Structure and fluorescent of **L**

2 Results and discussion

2.1 Metal ions detection

The fluorescent properties of the sensor **L** in *N,N*-dimethylformamide (DMF) are measured at room temperature. **L** exhibits strong fluorescent emission at 415 nm. Furthermore, **L** is explored for the application in the detection of metal ions in DMF and water system based on fluorescent sensing. In the measurement, 1.2161 mg of freshly prepared samples of **L** are dissolved in 250 mL DMF solution ($1 \times 10^{-5} \text{ mol} \cdot \text{L}^{-1}$), to which is dropwise added

different metal ions of aqueous solution. In this regard, the responses of the fluorescence of **L**/DMF solution toward 15 different metal cations (Zr^{4+} , Fe^{3+} , Co^{2+} , Cr^{3+} , Ni^{2+} , Cu^{2+} , Cd^{2+} , Pb^{2+} , Ag^{+} , Sr^{2+} , Ba^{2+} , Ca^{2+} , Mg^{2+} , Na^{+} , K^{+}) in aqueous solution were studied and obtained experimental data finally (Figure 2). Obviously, most cations display a negligible influence on fluorescence intensity of **L**. Specifically, the fluorescence intensity of **L** at 415 nm changes slightly while Co^{2+} , Cr^{3+} , Ni^{2+} , Cu^{2+} , Pb^{2+} , Ag^{+} , Sr^{2+} , Ba^{2+} , Ca^{2+} , Na^{+} and K^{+} ions were added respectively. It can see clearly that is that the fluorescence intensity of **L** decreases significantly, to which is added Fe^{3+} ion. Conversely, the fluorescence intensity of **L** increases significantly while Zr^{4+} ion is added. In addition, the compound **L** could rapidly give rise to a visible colorless-to-yellow change after adding Fe^{3+} ion and the experiments show that yellow is not caused by the color of Fe^{3+} ion itself.

Figure 2 Fluorescence spectra of **L** (2.9 mL , $1 \times 10^{-5} \text{ mol} \cdot \text{L}^{-1}$) with different metal cations (0.1 mL , $10^{-1} \text{ mol} \cdot \text{L}^{-1}$) ($\lambda_{\text{ex}} = 365 \text{ nm}$)

2.1.1 Fe^{3+} ion detection

To explore the complexation ratio of **L** (2.9 mL , $1 \times 10^{-5} \text{ mol} \cdot \text{L}^{-1}$) with Fe^{3+} ion in DMF solution, concentration-dependent experiments were further taken by UV method (Figure 3a). The absorption intensity of **L** gradually decreased with the increasing concentration of Fe^{3+} ion. The Job's curve was given according to the UV concentration curve of **L** and Fe^{3+} ions (Figure 3b). The results showed that the molar fraction of $[\text{Fe}^{3+}]/[\text{L} + \text{Fe}^{3+}]$ was 0.4 when the curve had inflection point, demonstrating 2 : 1 binding stoichiometry ration for **L** + Fe^{3+} . Moreover, to evaluate the detection effects of **L** to Fe^{3+} ion, the fluorescence spectra of **L** in DMF (2.9 mL , $1 \times 10^{-5} \text{ mol} \cdot \text{L}^{-1}$) with different concentrations of Fe^{3+} ion were carried out (Figure 4a). The experimental result showed that with the addition of different concentrations Fe^{3+} ion, the fluorescence intensity of **L** gradually decreases and shows a highly broad linear relationship with the concentration of Fe^{3+} ion from 1×10^{-6} to $1 \times 10^{-5} \text{ mol} \cdot \text{L}^{-1}$ expressed as an equation of $y = 41790.2576x + 0.0406$ and $R^2 = 0.99905$,

where y is the ratio of $(F_0 - F)/F_0$ and x is the concentration of Fe^{3+} ion. The other reported sensors for Fe^{3+} ion detection are generally within $1.25 \times 10^{-3} \text{ mol} \cdot \text{L}^{-1}$ of Fe^{3+} .^[39, 40] According to the fluorescence intensity of **L**, different concentration of Fe^{3+} and Benesi-Hildebrand equation, the detection limit of Fe^{3+} in **L** solution is calculated to be $1.33 \times 10^{-6} \text{ mol/L}$ (Figure 4b).

Figure 3 (a) UV-vis spectra of **L** with different concentration Fe^{3+} and (b) Job's plot of **L** with Fe^{3+} (a) $[\text{L}] = 2.9 \text{ mL}, 1 \times 10^{-5} \text{ mol} \cdot \text{L}^{-1}$; $[\text{Fe}^{3+}] = 0.1 \text{ mL}, 1 \times 10^{-6} \sim 10 \times 10^{-5} \text{ mol} \cdot \text{L}^{-1}$. (b) complexation ratio of **L** and Fe^{3+} , $[\text{Fe}^{3+}] = 0.1 \text{ mL}, 1 \times 10^{-6} \sim 10 \times 10^{-5} \text{ mol} \cdot \text{L}^{-1}$

Obviously, the addition of different concentration Fe^{3+} ion to the DMF solution of **L** could give rise to a visible colorless-to-yellow change, and clear fluorescence quenching while the concentration of Fe^{3+} is up to $10 \times 10^{-3} \text{ mol} \cdot \text{L}^{-1}$ (Figure 5). The initial fluorescence intensity of **L** ($1 \times 10^{-5} \text{ mol} \cdot \text{L}^{-1}$) was decreased by 99.9% when only 0.1 mL ($1 \times 10^{-1} \text{ mol} \cdot \text{L}^{-1}$) Fe^{3+} was added. Further, the quenching efficiency can be rationalized by the Stern-Volmer (SV) equation, $F_0/F = 1 + K_{SV}[\text{M}]$, where F_0 and F are the relative fluorescence intensity before and after adding Fe^{3+} , $[\text{M}]$ is the molar concentration of the Fe^{3+} ion, and K_{SV} ($\text{L} \cdot \text{mol}^{-1}$) is the Stern-Volmer quenching constant. The K_{SV} (Fe^{3+}) value is calculated as $7.111 \times 10^4 \text{ L} \cdot \text{mol}^{-1}$, suggesting a strong quenching effect on fluorescence. Under the same conditions, the quenching constants of other metals are shown in Table 1. The reason of fluo-

Figure 4 Fluorescence spectra (a) of **L** with different concentration Fe^{3+} and detection limit (b) of **L** with Fe^{3+} ($\lambda_{\text{ex}} = 324 \text{ nm}$) (a) $[\text{L}] = 2.9 \text{ mL}, 1 \times 10^{-5} \text{ mol} \cdot \text{L}^{-1}$; $[\text{Fe}^{3+}] = 0.1 \text{ mL}, 1 \times 10^{-6} \sim 10 \times 10^{-6} \text{ mol} \cdot \text{L}^{-1}$; (b) Detection limit of Fe^{3+} , $[\text{Fe}^{3+}] = 0.1 \text{ mL}, 1 \times 10^{-6} \sim 10 \times 10^{-6} \text{ mol} \cdot \text{L}^{-1}$

Figure 5 Photograph of **L** containing different concentrations of Fe^{3+} ion

rescence quenching may be that when Fe^{3+} and **L** form 2 : 1 complex, the structure is isomerized, the imine (NH) bond is broken, and the bond length is shortened, which makes ligand **L** show color change when encountering Fe^{3+} ions. And there may be electron and energy transfer between the paramagnetism of Fe^{3+} and the unfilled d orbital in **L**, which makes the fluorescence quenching.

Meanwhile, the antijamming experiments of Fe^{3+} ion specifically demonstrate that **L** can detect Fe^{3+} ion in the presence of other metal ions (Co^{2+} , Ni^{2+} , Cu^{2+} , Pb^{2+} , Ag^+ , Sr^{2+} , Ba^{2+} , Ca^{2+} , Na^+ , K^+ , Cd^{2+} , Mg^{2+} , Cr^{2+}) (Figure 6).

Table 1 Quenching constants of different metals in DMF solution of **L**^a

Cation	Quenching constant/ ($\text{L}\cdot\text{mol}^{-1}$)	Cation	Quenching constant/ ($\text{L}\cdot\text{mol}^{-1}$)
K^+	0.424	Na^+	0.071
Ag^+	-0.323	Ca^{2+}	-0.471
Ba^{2+}	0.071	Sr^{2+}	-0.212
Co^{2+}	1.099	Cr^{2+}	2.000
Cu^{2+}	2.779	Ni^{2+}	2.036
Mg^{2+}	0.899	Cd^{2+}	-1.519
Pb^{2+}	-0.011	Zr^{4+}	-2.904
Fe^{3+}	7.111×10^4		

^a **L**: 2.9 mL, 1×10^{-3} mol/L; ^b Concentration of metal ions: 0.1 mL, 1×10^{-1} mol/L; $\lambda_{\text{ex}}=365$ nm.

Figure 6 Effect of the disturbing metal ions on Fluorescence recognition of **L** to Fe^{3+}

2.1.2 Zr^{4+} ion detection

Compared with Fe^{3+} ion, Zr^{4+} ion exhibits superior fluorescence enhancement effect on **L**. To explore the detection effects of **L** in DMF (2.9 mL , $1 \times 10^{-6} \text{ mol}\cdot\text{L}^{-1}$) as the fluorescence probe of Zr^{4+} , concentration-dependent experiments were further taken (Figure 7). The fluorescence intensity of **L** gradually enhanced with the increasing concentration of Zr^{4+} ion. The fluorescence enhancement may be attributed to the formation of $\text{Zr}-\text{O}$ and $\text{Zr}-\text{N}$ bonds in **L** and Zr^{4+} ions, which enhance the rigid structure of the complexes. Simultaneously, the transition from $n-\pi^*$ to $\pi-\pi^*$ in the complexes may also be the reason of fluorescence enhancement. The Job's curve was given according to the fluorescence concentration curve of **L** and Zr^{4+} ions (Figure 7b). The results showed that the molar fraction of $[\text{Zr}^{4+}]/[\text{L}+\text{Zr}^{4+}]$ was 0.5 when the curve has inflection point, demonstrating 1 : 1 binding stoichiometry ratio for **L**+ Zr^{4+} . Moreover, to evaluate the detection effects of **L** to Zr^{4+} ion, the fluorescence spectra of **L** in DMF (2.9 mL , $1 \times 10^{-5} \text{ mol}\cdot\text{L}^{-1}$) with different concentrations of Zr^{4+} ion were taken (Figure 8). The experimental results show that with the addition of different concentrations Zr^{4+} ion, the fluorescence intensity of **L** gradually enhanced and showed a highly broad linear relationship

with the concentration of Zr^{4+} ion from 1×10^{-5} to $1 \times 10^{-4} \text{ mol}\cdot\text{L}^{-1}$ expressed as an equation of $y=3.592 \times 10^6 x + 39.24337$ and $R^2=0.99437$, where y is fluorescence intensity of **L** with different concentration Zr^{4+} and x is the concentration of Zr^{4+} ion. According to the fluorescence intensity of **L**, different concentration of Zr^{4+} and benesi-Hildebrand equation, the detection limit of Zr^{4+} in **L** solution is calculated to be $3.60 \times 10^{-6} \text{ mol/L}$.

Figure 7 (a) Fluorescence spectra of **L** with different concentration Zr^{4+} and (b) Job's plot of **L** with Zr^{4+} ($\lambda_{\text{ex}}=300 \text{ nm}$)

(a) **L**: 2.9 mL , $1 \times 10^{-5} \text{ mol}\cdot\text{L}^{-1}$; $[\text{Zr}^{4+}] = 0.1 \text{ mL}$, $1 \times 10^{-6} \sim 10 \times 10^{-6} \text{ mol}\cdot\text{L}^{-1}$. (b) Complexation ratio of **L** and Zr^{4+} , $[\text{Zr}^{4+}] = 0.1 \text{ mL}$, $1 \times 10^{-6} \sim 10 \times 10^{-6} \text{ mol}\cdot\text{L}^{-1}$

Of course, the antiinterference ability is also very important for sensors, so a series of antijamming experiments were performed for **L** (Figure 9). Initially, the fluorescence intensity of **L** containing Zr^{4+} ion shows negligible change with the addition of other metal ions except Cr^{2+} ion. The negligible change of the fluorescence intensity indicates that **L** can detect Zr^{4+} ion in the presence of other metal ions (Co^{2+} , Ni^{2+} , Cu^{2+} , Pb^{2+} , Ag^+ , Sr^{2+} , Ba^{2+} , Ca^{2+} , Na^+ , K^+ , Cd^{2+} , Mg^{2+}).

2.2 Acetone detection

The solvents play an important role in the fluorescence of compound **L**. Different organic solvents have different effects on the fluorescence of compound **L**. The fourteen different organic solvents, DMF, acetone (CP), xylene

Figure 8 (a) Fluorescence spectra of **L** with different concentration Zr^{4+} and (b) detection limit of **L** with Zr^{4+} ($\lambda_{\text{ex}} = 348 \text{ nm}$) (a) $[\text{L}] = 2.9 \text{ mL}, 1 \times 10^{-5} \text{ mol} \cdot \text{L}^{-1}$; $[\text{Zr}^{4+}] = 0.1 \text{ mL}, 1 \times 10^{-5} \sim 10 \times 10^{-5} \text{ mol} \cdot \text{L}^{-1}$. (b) Detection limit of Zr^{4+} ; $[\text{Zr}^{4+}] = 0.1 \text{ mL}, 1 \times 10^{-5} \sim 10 \times 10^{-5} \text{ mol} \cdot \text{L}^{-1}$

Figure 9 Effect of the disturbing metal ions on Fluorescence recognition of **L** to Zr^{4+}

(XY), acetonitrile (ACN), ethyl acetate (EAC), isopropyl alcohol (IPA), dichloromethane (MC), benzene (Benzol), dimethyl sulfoxide (DMSO), tetrahydrofuran (THF), cyclohexane (CYC), 1,4-dioxane (Diox), methanol (MT) and ethanol (EA), in DMF solution were investigated and obtained experimental data finally (Figure 10). The solvents Diox, MT and EA have enhanced fluorescence effect on compound **L**, and the intensity exceeds the upper limit of instrument detection. Compared with DMF, it may be due

to the strong polarity and weak p- π conjugation of 1,4-dioxane, which lead to the fluorescence enhancement and blue shift of **L**, while the fluorescence enhancement and red shift of **L** may be due to the strong polarity of methanol and ethanol, forming hydrogen bond with **L**. The results show that the initial fluorescence intensity of **L** ($1 \times 10^{-5} \text{ mol} \cdot \text{L}^{-1}$) was decreased by 94.48% when only 0.1 mL of acetone was added. The acetone showed a strong quenching effect on fluorescence of compound **L**. It may be that there is energy transfer between compound **L** and acetone, which makes the excitation of compound **L** and the absorption of acetone has some competitive effect, which leads to the rapid decrease of fluorescence intensity of **L**. It was subsequently studied a possible interference of various organic solvents on selective quenching of the fluorescence of **L** caused by acetone (Figure 11). It is found that there is no significant impact of other solvent on the quenching of the fluorescence caused by the presence of acetone. Interestingly, when the λ_{ex} is larger than 300 nm, the fluorescence quenching of compound **L** by acetone is not obvious. To some extent, the fluorescence effect of solvent on compound **L** depends on the specific λ_{ex} .

Figure 10 Fluorescence effect of organic solvent on **L** ($\lambda_{\text{ex}} = 300 \text{ nm}$)

2.3 Application to the determination of Fe^{3+} in urine and water samples

The high selectivity and sensitivity make **L** promising as a fluorescent probe for detecting Fe^{3+} in aqueous solution. To show its potential for application in biological samples, the fluorescent **L** probe was applied to the detection of Fe^{3+} in human urine samples. Six human urine samples were donated by healthy volunteers. The urine samples were acidified with nitric acid and centrifuged at 5000 r/min for 10 min. Add 0.1 mL of urine directly to DMF solution of compound **L** without any other treatment before test analysis. The concentrations of Fe^{3+} ions in urine were calculated by Stern-Volmer (SV) equation and the values of $y = 41790.2576x + 0.0406$ were shown in Table 2. The range of Fe^{3+} ions in six urine samples was about $1.68 \sim 5.78 \mu\text{mol} \cdot \text{L}^{-1}$ by calculation. In a sense, compound **L** can be used as a fluorescence probe to measure

the molarity of Fe^{3+} ion conveniently. Of course, the flow injection ICPMS (inductively coupled plasma mass spectrometry) method can also be used to accurately analyze Fe^{3+} in human urine.

Figure 11 Interference experiment of organic solvent to **L** fluorescence ($\lambda_{\text{ex}} = 270 \text{ nm}$)

Table 2 Detection of Fe^{3+} ion concentration in human urine by fluorescence of compound **L**^a

Urine	F_0	F	$[\text{Fe}^{3+}]/(\mu\text{mol}\cdot\text{L}^{-1})$		Error
			F_0/F^b	Y^c	
1	573.90	411.89	5.53	5.78	± 0.25
2	573.90	449.90	3.86	4.19	± 0.33
3	573.90	505.17	1.91	1.89	± 0.02
4	573.90	510.28	1.75	1.68	± 0.07
5	573.90	435.67	4.46	4.79	± 0.33
6	573.90	475.64	2.91	3.12	± 0.21

^a **L**: 2.9 mL, $1 \times 10^{-5} \text{ mol/L}$; sample: 0.1 mL; $\lambda_{\text{ex}} = 300 \text{ nm}$. ^b $F_0/F = 1 + K_{\text{SV}}[\text{Fe}^{3+}]$; ^c $y = 41790.2576x + 0.0406$.

Besides, the tap-water and drinking-water were also tested by the above detection methods, and the concentrations of Fe^{3+} ions were 10.7 (9.38) and 2.28 (2.36) $\mu\text{mol}\cdot\text{L}^{-1}$, respectively. It should be mentioned that the Environmental Protection Ministry (P. R. China) established the standard (5.4 $\mu\text{mol}\cdot\text{L}^{-1}$) for the presence of Fe^{3+} ions in drinking H_2O . In this way, the tap-water cannot meet the drinking requirements, while the direct-drinking water is within the scope of national standards.

3 Conclusions

In summary, as a fluorescence probe, **L** shows high sensitivity and selectivity for Zr^{4+} , Fe^{3+} ions and acetone, and the detection of Fe^{3+} ion can be observed by naked eye. Meanwhile, the experimental results show that the trace of Fe^{3+} ion could remarkably decrease the fluorescence of **L**. In contrast to this, the trace of Zr^{4+} ion could remarkably increase the fluorescence of **L**. The detection of Fe^{3+} ion in human urine and water was realized above two detection methods. Therefore, It is expected that the compound **L** can be used as a tunable organic fluorescence material by Fe^{3+} and Zr^{4+} ions, and also as a good probe to identify acetone in organic solvent. This study might also provide a new idea of designing fluorescent organic compound

chemosensors for the sensitive and selective detection of metal ions and acetone.

4 Experimental section

4.1 General methods

All reagents were purchased commercially and used without further purification unless otherwise noted. ^1H NMR spectra were recorded using 500 MHz, and ^{13}C NMR spectra were recorded using 125 MHz. Elemental analyses of C, H, N in the solid samples were tested on a Vario EL analyzer (Elementar Analysensysteme GmbH). FT-IR spectra were recorded in the range of 500~4000 cm^{-1} on a Perkin-Elmer FTIR spectrometer using KBr pellets. The fluorescence spectra were acquired on an F4500 spectrometer with a 850W Xenon lamp as the excitation source at room temperature.

4.2 Synthesis of 2,2',2''-(1,3,5-triazine-2,4,6-triimino)tribenzoic acid (**L**)

The **L** was synthesized according to the literature method^[41~43] with some modification (Scheme 1). Generally speaking, the reaction of cyanuric chloride is step-by-step. Low temperature is conducive to the first substitution, the second substitution at 50~60 $^{\circ}\text{C}$, and the third substitution at 110~120 $^{\circ}\text{C}$. It is verified by the following two methods. The results show that they are both suitable for the synthesis of **L** by step-by-step and one pot reaction.

Scheme 1 Synthetic route of **L**

Method 1: 2.263 g (16.5 mmol) of 2-aminobenzoic acid and 25 mL of AcOH were added to a 100 mL round bottom flask. 0.922 g (5.0 mmol) of cyanuric chloride was dissolved in 20 mL of AcOH and slowly dripped into the round bottom flask under the ice bath. The mixture reacts at 0 $^{\circ}\text{C}$ for 5 h. Then, the reaction continued for 5 h at 50 $^{\circ}\text{C}$. Finally, the reaction refluxed at 120 $^{\circ}\text{C}$ for 20 min. The suspension was filtered and the product was washed several times with boiling water and ethanol, and then the product was dried in vacuum dryer at 80 $^{\circ}\text{C}$ for one day. 2.345 g of white powder was collected in 96.4% yield.

Method 2: 0.922 g (5.0 mmol) of cyanuric chloride and 2.263 g (16.5 mmol) of 2-aminobenzoic acid were dissolved in 50 mL of AcOH in a 100 mL round-bottom flask. The mixture was stirred and refluxed immediately at 120 °C for 15 min, generating a pink aqueous suspension. Then, the suspension was filtered and the product was washed several times with boiling water and ethanol, and then the mixture was dried in vacuum dryer at 80 °C for one day. 2.352 g of white powder was collected in 96.7% yield.

2,2',2''-(1,3,5-Triazine-2,4,6-triimino)tribenzoic acid (**L**): m.p. ≥ 330 °C (dec.); ^1H NMR (DMF- d_7 , 500 MHz) δ : 11.19 (s, 3H, COOH), 8.86 (d, $J=10.0$ Hz, 3H, ArH), 8.13~8.15 (m, 3H, ArH), 7.68~7.72 (m, 3H, ArH), 7.16~7.20 (m, 3H, ArH); IR (KBr) ν : 3440, 3312, 1685, 1610, 1567, 1523, 1413, 1161; ^{13}C NMR (DMF- d_7 , 125 MHz) δ : 170.1, 164.5, 142.2, 133.8, 131.5, 121.6, 120.9, 123.3, 116.3 cm^{-1} . Anal. calcd for $\text{C}_{24}\text{H}_{18}\text{N}_6\text{O}_6$: C 59.26, H 3.73, N 17.28; found C 58.31, H 3.58, N 16.57.

References

- [1] Wang, P.; Okamura, T.; Zhou, H. P.; Sun, W. Y.; Tian, Y. P. *Chin. Chem. Lett.* **2013**, 24(1), 20.
- [2] Xu, H.; Hu, H. C.; Cao, C. S.; Zhao, B. *Inorg. Chem.* **2015**, 54(10), 4585.
- [3] Wang, J.; Jiang, M.; Yan, L.; Peng, R.; Huangfu, M. J.; Guo, X. X.; Li, Y.; Wu, P. Y. *Inorg. Chem.* **2016**, 55(24), 12660.
- [4] Yan, W.; Zhang, C.; Chen, S. G.; Han, L. J.; Zheng, H. G. *ACS Appl. Mater. Interfaces* **2017**, 9(19), 1629.
- [5] Arici, M. *Cryst. Growth Des.* **2017**, 17(10), 5499.
- [6] Li, P.; Zhou, L. J.; Yang, N. N.; Sui, Q.; Gong, T.; Gao, E. Q. *Cryst. Growth Des.* **2018**, 18(11), 7191.
- [7] Tang, S. F.; Hou, X. M. *Cryst. Growth Des.* **2019**, 19(1), 45.
- [8] Zheng, M.; Tan, H. Q.; Xie, Z. G.; Zhang, L. G.; Jing, X. B.; Sun, Z. C. *ACS Appl. Mater. Interfaces* **2013**, 5(3), 1078.
- [9] Xu, X. Y.; Yan, B. *ACS Appl. Mater. Interfaces* **2015**, 7(1), 721.
- [10] Wen, G. X.; Wu, Y. P.; Dong, W. W.; Zhao, J.; Li, D. S.; Zhang, J. *Inorg. Chem.* **2016**, 55(20), 10114.
- [11] Rao, P. C.; Mandal, S. *Inorg. Chem.* **2018**, 57(19), 11855.
- [12] Gogoi, C.; Biswas, S. *Dalton Trans.* **2018**, 47(41), 14696.
- [13] Yu, C. Y.; Sun, X. D.; Zou, L. F.; Li, G. H.; Zhang, L. R.; Liu, Y. L. *Inorg. Chem.* **2019**, 58(6), 4026.
- [14] He, H. M.; Zhu, Q. Q.; Li, C. P.; Du, M. *Cryst. Growth Des.* **2019**, 19(2), 694.
- [15] Jia, H. J.; Han, L. M.; Zhu, N.; Gao, Y. Y.; Wang, Y. Q.; Suo, Q. L. *Chin. J. Org. Chem.* **2019**, 39(6), 1753 (in Chinese). (贾慧劫, 韩利民, 竺宁, 高媛媛, 王亚琦, 索全伶, 有机化学, **2019**, 39(6), 1753.)
- [16] Liu, M. L.; Chen, B. B.; Li, C. M.; Huang, C. Z. *Sci. China: Chem.* **2019**, 62(8), 968.
- [17] Yu, L.; Qiao, Y. M.; Miao, L. X.; He, Y. Q.; Zhou, Y. *Chin. Chem. Lett.* **2018**, 29(11), 1545.
- [18] Pang, C. M.; Luo, S. H.; Hao, Z. F.; Gao, J.; Huang, Z. H.; Yu, J. H.; Yu, S. M.; Wang, Z. Y. *Chin. J. Org. Chem.* **2018**, 38(10), 2606 (in Chinese). (庞楚明, 罗时荷, 郝志峰, 高健, 黄召昊, 余家海, 余思敏, 汪朝阳, 有机化学, **2018**, 38(10), 2606.)
- [19] Sheng, K.; Lu, H. F.; Sun, A. B.; Wan, Y. M.; Liu, Y. T.; Chen, F.; Bian, W. C.; Li, Y.; Kuang, R.; Sun, D. *Chin. Chem. Lett.* **2019**, 30(4), 895.
- [20] Wang, J. H.; Fan, Y. d.; Lee, H. W.; Yi, C. Q.; Cheng, C. M.; Zhao, X.; Yang, M. *ACS Appl. Nano Mater.* **2018**, 1(7), 3747.
- [21] Huang, J. J.; Yu, J. H.; Bai, F. Q.; Xu, J. Q. *Cryst. Growth Des.* **2018**, 18(9), 5353.
- [22] Lu, S. Q.; Liu, Y. Y.; Duan, Z. M.; Wang, Z. X.; Li, M. X.; He, X. *Cryst. Growth Des.* **2018**, 18(8), 4602.
- [23] Smith, J. A.; Singh-Wilmot, M. A.; Carter, K. P.; Cahill, C. L.; August Ridenour, J. *Cryst. Growth Des.* **2019**, 19(1), 305.
- [24] Fan, K.; Bao, S. S.; Nie, W. X.; Liao, C. H.; Zheng, L. M. *Inorg. Chem.* **2018**, 57(3), 1079.
- [25] Zhang, H. J.; Fan, R. Q.; Chen, W.; Fan, J. Z.; Dong, Y. W.; Song, Y.; Du, X.; Wang, P.; Yang, Y. L. *Cryst. Growth Des.* **2016**, 16(9), 5429.
- [26] Zhang, C. H.; Shi, H. Z.; Sun, L. B.; Yan, Y.; Wang, B. L.; Liang, Z. Q.; Wang, L.; Li, J. Y. *Cryst. Growth Des.* **2018**, 18(12), 7683.
- [27] Yoon, C. S.; Choi, M. J.; Jun, D. W.; Zhang, Q.; Kaghazchi, P.; Kim, K. H.; Sun, Y. K. *Chem. Mater.* **2018**, 30(5), 1808.
- [28] Yoon, C. S.; Kim, U. H.; Park, G. T.; Kim, S. J.; Kim, K. H.; Kim, J.; Sun, Y. K. *Energy Lett.* **2018**, 3(7), 1634.
- [29] Meng, H. M.; Fu, T.; Zhang, X. B.; Wang, N. N.; Tan, W. H.; Shen, G. L.; Yu, R. Q. *Anal. Chem.* **2012**, 84(5), 2124.
- [30] Wang, B.; Yang, Q.; Guo, C.; Sun, Y. X.; Xie, L. H.; Li, J. R. *ACS Appl. Mater. Interfaces* **2017**, 9(11), 10286.
- [31] Xu, M. M.; Kong, X. J.; He, T.; Wu, X. Q.; Xie, L. H.; Li, J. R. *Inorg. Chem.* **2018**, 57(22), 14260.
- [32] He, P.; Tang, L. J.; Zhong, K. L.; Hou, S. H.; Yan, X. M. *Chin. J. Org. Chem.*, **2017**, 37(2), 423 (in Chinese). (何平, 汤立军, 钟克利, 侯淑华, 燕小梅, 有机化学, **2017**, 37(2), 423.)
- [33] Yu, L.; Qiao, Y. M.; Miao, L. X.; He, Y. Q.; Zhou, Y. *Chin. Chem. Lett.* **2018**, 29(11), 1545.
- [34] Wang, T.; Liu, Q. H.; Gao, Y.; Yang, X. Y.; Yang, W. T.; Dang, S. *Chin. Chem. Lett.* **2016**, 27(4), 497.
- [35] Zhang, Q. S.; Wang, J.; Kirillov, A. M.; Dou, W.; Xu, C.; Xu, C. L.; Yang, L. Z.; Fang, R.; Liu, W. S. *ACS Appl. Mater. Interfaces* **2018**, 10(28), 23976.
- [36] Zhao, X. X.; Wang, S. L.; Zhang, L. Y.; Liu, S. Y.; Yuan, G. Z. *Inorg. Chem.* **2019**, 58(4), 2444.
- [37] Wang, X.; Fan, W. D.; Zhang, M.; Shang, Y. Z.; Wang, Y. T.; Liu, D.; Guo, H. L.; Dai, F. N.; Sun, D. F. *Chin. Chem. Lett.* **2019**, 30(3), 801.
- [38] Huang, W. H.; Ren, J.; Yang, Y. H.; Li, X. M.; Wang, Q.; Jiang, N.; Yu, J. Q.; Wang, F.; Zhang, J. *Inorg. Chem.* **2019**, 58(2), 1481.
- [39] Chen, Z.; Sun, Y. W.; Zhang, L. L.; Sun, D.; Liu, F. L.; Meng, Q. G.; Wang, R. M.; Sun, D. F. *Chem. Commun.* **2013**, 49(98), 11557.
- [40] Wang, R.; Dong, X. Y.; Xu, H.; Pei, R. B.; Ma, M. L.; Zang, S. Q.; Hou, H. W.; Mak, T. C. W. *Chem. Commun.* **2014**, 50(65), 9153.
- [41] Karuehanon, W.; Fanfuenha, W.; Rujiwatra, A.; Pattarawarapan, M. *Tetrahedron Lett.* **2012**, 53(27), 3486.
- [42] Wang, X. S.; Liang, J.; Li, L.; Lin, Z. J.; Bag, P. P.; Gao, S. Y.; Huang, Y. B.; Cao, R. *Inorg. Chem.* **2016**, 55(5), 2641.
- [43] Kolmakov, K. A. *J. Heterocycl. Chem.* **2008**, 45(2), 533.

(Li, L.; Fan, Y.)