钯催化氧杂降冰片烯与三氟硼酸钾盐的不对称开环反应

陶萍芳^{†,a} 黄俊^{†,a,b} 刘玉钊^a 韦光明^a 王艺菲^a
 韦贤生^a 黄国保^{*,a} 李秀英^{*,a}

(^a 玉林师范学院化学与食品科学学院 广西农产资源化学与生物技术重点实验室 广西玉林 537000)

(^b 南宁师范大学化学与材料学院 南宁 530001)

摘要 开发了一种新型高效的钯催化的三氟硼酸钾盐与氧杂降冰片烯的不对称开环反应的方法。在温和的条件下, 可得到相应的顺-2-芳基-1,2-二氢萘-1-醇产物, 其产率良好, 目标产物对映体选择性较高。

关键词 氧杂降冰片烯; 三氟钾硼酸盐; 不对称开环反应

Palladium-Catalyzed Asymmetric Ring-Opening Reactions of Oxabenzonorbornadienes with Potassium Trifluoroborate Salts

Tao, Pingfang^{†,a} Huang, Jun^{†,a,b} Liu, Yuzhao^a Wei, Guangming^a Wang, Yifei^a
 Wei, Xiansheng^a Huang, Guobao^{*,a} Li, Xiuying^{*,a}

(^a Guangxi Key Laboratory of Agricultural Resources Chemistry and Biotechnology, College of Chemistry and Food Science, Yulin Normal University, Yulin, Guangxi 537000)

(^b College of Chemistry and Materials, Nanning Normal University, Nanning 530001)

Abstract A new, versatile and highly efficient palladium-catalyzed asymmetric ring-opening reaction of oxabenzonorbornadienes with a wide range of potassium trifluoroborate salts was developed. The corresponding *cis*-2-aryl-1,2-dihydronaphthalen-1-ol products were obtained in good yields with moderate to good enantioselectivities under mild conditions.

Keywords oxabenzonorbornadiene; trifluoroborate salt; asymmetric ring-opening reaction

1 Introduction

The synthesis of substituted 1,2-dihydronaphthalenes is a theme of ongoing interest because these compounds possess a wide range of biological activities,^[1] and they are also important intermediates in organic synthesis.^[2] Among the available methods for hydronaphthalenes synthesis, the transition metal-catalyzed asymmetric ring-opening (ARO) reactions of hetero-bicyclic alkenes with various carbanion and heteroatom nucleophiles are valuable processes, which have been intensively investigated. Many metal catalysts such as Fe,^[3] Ni,^[4] Cu,^[5] Ru,^[6] Rh,^[7] Pd,^[8] Pt^[9] and Ir^[10] were explored for the ring-opening of oxa- and aza-bicyclic alkenes.

How to find a group of novel nucleophiles as the aryl source to react with oxabicyclic alkenes is a long-term challenge to the chemists. Various aryl donors such as aryl halides,^[11] organometallic reagents,^[12] arylboronic acids^[13] and arylsulfinate salts^[14] had been used in the ARO reactions. To the best of our knowledge, the study referring to the reaction of aryl halides with oxabenzonorbornadiene is relatively rare because of the low activity of these carbanion nucleophiles. On the contrary, high active organometallic reagents may incur some side effects to give the ring-opening product in low yields.^[12a] In recent years, arylsulfinate salts due to the stability and low toxicity, have been employed as effective carbanion nucleophiles in the ring-opening reactions of oxabicyclic alkenes

* Corresponding authors. E-mail: lzjx0915@163.com; lxys8401@163.com

Received December 22, 2019; revised March 11, 2020; published online March 31, 2020.

Project supported by the National Natural Science Foundation of China (No. 21961042), the Natural Science Foundation of Guangxi Province (No. 2018GXNSFAA294064) and the Open Fund of Guangxi Key Laboratory of Agricultural Resources Chemistry and Biotechnology Research Grant (No. KF04). 国家自然科学基金(No.21961042)、广西自然科学基金(No. 2018GXNSFAA294064)及广西农产资源化学与生物技术重点实验室开放基金(No. KF04)资助项目。

[†] 共同第一作者(These authors contributed equally to this work).

to afford the corresponding products, but with low enantioselectivities. Arylboronic acids are appropriate reagents in this type of ARO reactions due to their low cost, remarkable stability and ease of handling. However, arylboronic acids still have some drawbacks, such as having high nucleophile equivalence and low reactivity in the air atmosphere. Nevertheless, these results gave us good references.

Potassium trifluoroborate salts are a kind of ideal reagents in organic chemistry, for example, the construction of C—C bonds.^[15] Compared with boronic acids, potassium trifluoroborate salts not only own similar chemical properties, but also are more convenient and stable.^[16] To our surprise, so far there is no report on the ring opening reaction of oxybenzodiene with potassium trifluoroborate as the substrate. Herein a novel ARO addition of potassium trifluoroborate salts to oxabicyclic alkenes in the presence of palladium complex to afford the corresponding products of *cis*-2-aryl-1,2-dihydronaphthalene-1-ols in high yields (up to 98%) with moderate to good enantioselectivities (up to 98% *ee*) is reported. *cis*-Configuration, not *trans*-configuration, of the product was obtained through X-ray diffraction analysis.

2 Results and discussion

To explore the ring opening reaction, preliminary optimization of the reaction conditions was carried out using oxabenzonorborene (**1a**) and potassium phenyltrifluoroborate (**2a**) as model substrates in toluene/water (*V*: *V*=5:1) at room temperature. When **1a** reacted with **2a** in the presence of 5 mol% Pd(OAc)₂ and 10 mol% DPPP under nitrogen atmosphere, desired ring-opened product **3a** was obtained in 25% isolated yield after 24 h (Table 1, Entry 1). Encouraged by this result, several chiral ligands were screened for the ring-opening reaction of **1a** under similar conditions, including (*S,S*)-Me-DuPhos, (*S*)-DM-SegPhos, (*R*)-(*S*)-PPF-PtBu₂, (*R*)-BINAP and (*S*)-PipPhos. The results indicated that (*R*)-BINAP was the most effective ligand among the ligands screened for this reaction compared with the others (Table 1, Entries 2~6), and **3a** was afforded in 90% yield with 55% *ee* (Table 1, Entry 6). Meanwhile, the catalyst loading also had a significant impact. The results indicated that 5 mol% Pd(OAc)₂ with 10 mol% (*R*)-BINAP would be a more suitable catalyst loading in terms of the yield and enantioselectivity, whereas increasing or decreasing the amount of catalyst loading improved neither the yield nor *ee* value of **3a** (Table 1, Entries 7~9). Similar yield and enantioselectivity were obtained whether in a nitrogen or air atmosphere (Table 1, Entry 10). In addition, the effect of solvents was investigated. A mixture of water and CH₃CN was totally ineffective in this reaction (Table 1, Entry 11). When the reaction was carried out in a mixed solvent of water with tetrahydrofuran (THF), expected product **3a** was afforded in only 20% yield with 30% *ee* (Table 1, Entry 12). It was observed that the reaction in a mixed solvent of water with protic solvents MeOH gave satisfac-

Table 1 Optimization of reaction conditions for the ARO reaction^a

Entry	Solvent	Pd(OAc) ₂ /mol%	Ligand	Time/h	Yield/%	<i>ee</i> ^b /%
1	Toluene	5.0	L1	24	25	—
2	Toluene	5.0	L2	7	49	35
3	Toluene	5.0	L3	7	60	21
4	Toluene	5.0	L4	24	NR ^c	—
5	Toluene	5.0	L5	0.5	90	55
6	Toluene	5.0	L6	24	NR ^c	—
7	Toluene	2.5	L5	1	80	55
8	Toluene	1.0	L5	2	71	55
9	Toluene	10.0	L5	0.5	92	54
10 ^d	Toluene	5.0	L5	0.5	91	55
11 ^d	MeCN	5.0	L5	24	NR ^c	—
12 ^d	THF	5.0	L5	24	20	30
13 ^d	MeOH	5.0	L5	22	30	76
14 ^d	CH ₂ Cl ₂	5.0	L5	3	85	45
15 ^d	CHCl ₃	5.0	L5	3	80	50
16 ^d	DCE	5.0	L5	1	81	52
17 ^{d,e}	Toluene	5.0	L5	5	72	58
18 ^{d,f}	Toluene	5.0	L5	5	85	42

^a Reaction conditions: **1a** (0.2 mmol), **2a** (1.2 equiv.), Pd(OAc)₂ and ligand (10 mol%) in solvent and H₂O (*V*: *V*=5:1, 3 mL in total) under nitrogen at room temperature. ^b Determined by HPLC with a chiralcel OD-H column.

^c NR: no reaction. ^d Carried out in air. ^e The reaction at 0 °C. ^f The reaction at 50 °C.

tory enantioselectivity (up to 76% *ee*) but poor conversion (Table 1, Entry 13). Moderate to good yields and enantioselectivities were obtained when a mixed solvent of water with CH₂Cl₂, CHCl₃ and dichloroethane (DCE) were respectively employed as the solvents (Table 1, Entries 14~16). Importantly, the study on solvent effect indicated that the reaction worked very well in a mixed solvent of water with the aromatic solvent of toluene showed the best combination of reactivity and enantioselectivity among all solvents tested (91% yield, 55% *ee*). However, satisfactory results had not been achieved when the impact of reaction temperature was investigated. The reaction run at 0 °C gave a diminished yield and slightly improved *ee* (Table 1,

Entry 17). Raising the temperature to 50 °C somewhat decreased the yield and *ee* even after a prolonged reaction time (Table 1, Entry 18).

Next, with the optimized reaction conditions identified, the scope of potassium trifluoroborate salts was explored for this reaction to evaluate their effect on the reactivity and enantioselectivity (Table 2). From Table 2, it can be seen that the structures of potassium trifluoroborate salts have significant impact on the reactivity and enantioselectivities. In general, potassium trifluoroborate salts with electron-donating substituents on the phenyl rings gave higher yields but lower enantioselectivities than those with electron-withdrawing groups (Table 2, Entries 2 ~ 9). When it came to the position property of monosubstituted on the phenyl rings of potassium trifluoroborate salts, it had little effect on the reactivity, and the ring-opening of **1a** with parasubstituted potassium trifluoroborate salts offered better enantioselectivities than those with *meta*- and *ortho*-substituted potassium trifluoroborate salts (Table 2, Entries 2, 3, 5 ~ 7). It is noteworthy that the addition of alkyl-potassium trifluoroborate salt to **1a** was found to give the corresponding product **3j** successfully just prolonged the reaction time to 3 h (Table 2, Entry 10). Potassium trifluoroborate salts containing heterocycle were employed in the ring-opening of **1a**, the results was also satisfactory (Table 2, Entry 11). Thiophene-2-potassium trifluoroborate salt reacted with **1a** and **3k** was obtained in high yield (95% yield) with modest enantioselectivity (71% *ee*).

Table 2 Ring-opening of oxabenzonorbornadiene **1a** with various potassium trifluoroborate salts^a

Entry	R	Product	Time/h	Yield/%	<i>ee</i> ^b /%
1	C ₆ H ₅	3a	0.5	91	55
2	4-CH ₃ C ₆ H ₄	3b	0.5	94	62
3	2-CH ₃ C ₆ H ₄	3c	0.5	92	20
4	4- ⁱ BuC ₆ H ₄	3d	0.5	95	36
5	2-CH ₃ OC ₆ H ₄	3e	0.5	90	2
6	3-CH ₃ OC ₆ H ₄	3f	0.5	91	56
7	4-CH ₃ OC ₆ H ₄	3g	0.5	98	51
8	4-BrC ₆ H ₄	3h	0.5	84	64
9	4-FC ₆ H ₄	3i	0.5	84	60
10	CH ₃	3j	3	81	56
11	2-Thienyl	3k	0.5	95	71

^a Reaction conditions: **1a** (0.2 mmol), **2** (1.2 equiv.), Pd(OAc)₂ (5 mol%), and (R)-BINAP (10 mol%) in toluene and H₂O (V : V = 5 : 1, 3 mL in total) in air at room temperature. ^b Determined by HPLC with a chiralcel OD-H column.

The molecular configuration of **3d** was unambiguously confirmed by X-ray diffraction analysis (Figure 1). The single crystal was obtained by solvent evaporation from a mixture of CHCl₃ and hexane. The configuration of **3d** was assigned as (1*S*,2*R*) and confirmed as the 1,2-*cis*- configuration.

ration.^[13c]

Figure 1 X-ray structure of the product **3d**

The substrate scope of oxabenzonorbornadiene derivatives was further evaluated, and the results were compiled in Table 3. From Table 3, besides **1a**, it can be seen that the reaction of various oxabenzonorbornadienes with potassium trifluoroborate salts proceeded smoothly to generate the anticipated ring-opening products in high yields (up to 93%) with good to excellent enantioselectivity (up to 97% *ee*). Furthermore, the influence of different potassium trifluoroborate salts on the reaction of each substrate is similar to that of substrate **1a**, potassium trifluoroborate salts with electron-donating substituents on the phenyl rings gave higher yields but lower enantioselectivities than those with electron-withdrawing groups. The excellent performance of this catalyst system is highlighted by the reaction of **1b** with potassium cyclopropyl trifluoroborane, which furnished product **4a** in excellent yield (90%) with remarkable enantioselectivity (up to 97% *ee*) (Table 3, Entry 2).

Table 3 Ring-opening of oxabenzonorbornadiene **1b** and **1c** with various potassium trifluoroborate salts^a

1b : R ¹ = H, R ² = OMe 1c : R ¹ + R ² =					
Entry	Substrate	R	Product	Yield/%	<i>ee</i> ^b /%
1	1b	Cyclopropyl	4a	90	97
2	1b	4-FC ₆ H ₄	4b	84	52
3	1b	4-CH ₃ C ₆ H ₄	4c	91	51
4	1c	C ₆ H ₅	5a	85	55
5	1c	Cyclopropyl	5b	90	97
6	1c	4-FC ₆ H ₄	5c	84	68
7	1c	4-CH ₃ C ₆ H ₄	5d	91	65
8	1c	2-Thienyl	5e	93	87

^a Reaction conditions: **1b**, **1c** (0.2 mmol), **2** (1.2 equiv.), Pd(OAc)₂ (5 mol%), and (R)-BINAP (10 mol%) in toluene and H₂O (V : V = 5 : 1, 3 mL in total) in air at room temperature. ^b Determined by HPLC with a chiralcel OD-H column.

Based upon the experimental observations and literatures,^[17] a proposed mechanism for the palladium(II)-catalyzed asymmetric ring-opening of potassium phenyltrifluoroborate to oxabenzonorbornadiene **3a** is summarized in Scheme 1. The catalytic cycle would be initiated by the

coordination of $\text{Pd}(\text{OAc})_2$ and (*R*)-BINAP to form the active chiral palladium complex **A**, which leads to intermediate **B** after addition of potassium phenyltrifluoroborate (**2a**).^[18] **B** is followed by insertion of oxabenzonorbornadiene (**1a**) into the carbon-palladium bond from the *exo* side and produce intermediate **C**. β -Elimination of oxygen results in opening the furyl ring to give the ring-opened intermediate **D** which follows by the hydrolysis to liberate ring-opened product **3a**.^[19] Meanwhile, the palladium species **A** is regenerated to promote the next catalytic cycle.

Scheme 1 Proposed mechanism

3 Conclusions

In summary, a new palladium(II)-catalyzed asymmetric ring-opening reaction of oxabenzonorbornadienes with potassium trifluoroborate salts was successfully developed. $\text{Pd}(\text{OAc})_2$ and (*R*)-BINAP were used as the catalyst system in this reaction. A practical and efficient approach was provided to synthesize the optically active *cis*-2-aryl-1,2-dihydronaphthalen-1-ol derivatives in high yields with moderate to good enantioselectivities under mild conditions. Further investigations on the applications of palladium catalysts in the asymmetric ring-opening are in progress in our laboratory and will be reported.

4 Experimental section

4.1 General information

^1H NMR, ^{13}C NMR and ^{19}F NMR spectra were recorded on a Bruker Avance nuclear magnetic resonance spectrometer at 400/500 MHz and 100/125 MHz at room temperature in CDCl_3 , respectively. HRMS (ion trap) were recorded by Q EXACTIVE mass spectrometry using APCI or ESI. The HPLC analysis was performed on a Waters Alliance e2695 high performance liquid chromatography with a Chiralcel OD-H column or Chiralpak AD-H column. Unless otherwise indicated, all reagents were purchased from commercial suppliers and used without further purification. Toluene was used without any pretreatment.

All flasks were flame-dried under a stream of nitrogen and cooled to room temperature before use. Oxabenzonorbornadienes **1a**~**1c** were prepared according to the literature procedures.^[20]

4.2 General procedures for the products

All experiments were carried out under an air atmosphere. Palladium diacetate (2.2 mg, 5 mol%) and (*R*)-BINAP (12.4 mg, 10 mol%) were added to a 10 mL round-bottomed flask, followed by the addition of toluene/ H_2O (*V* : *V* = 5 : 1, 3 mL in total). After the mixture was stirred for about 5 min, oxabenzonorbornadienes **1a**~**1c** (0.2 mmol) and potassium trifluoroborate (1.2 equiv., 0.24 mmol) were put into the reaction system. The resulting mixture was stirred at room temperature until completion monitored by thin-layer chromatography (TLC). The solvent was removed *in vacuo*, and the crude mixture was then purified by column chromatography on silica gel (silica gel, 200~300 mesh) to afford the target product.

(1*S**,2*R**)-2-Phenyl-1,2-dihydronaphthalen-1-ol (**3a**): Colorless oil (40.0 mg, 90% yield). R_f =0.25 on silica gel (ethyl acetate/petroleum ether, *V* : *V* = 1 : 20). The *ee* was determined to be 55% using HPLC analysis on a Chiralcel OD-H column (hexane/2-propanol, *V* : *V* = 90 : 10, 1.0 mL/min, λ =254 nm). Retention times were 7.474 (major) and 11.708 (minor) min. ^1H NMR (500 MHz, CDCl_3) δ : 7.41~7.28 (m, 8H), 7.22 (d, *J*=7.2 Hz, 1H), 6.76 (dd, *J*=9.6, 1.7 Hz, 1H), 6.18 (dd, *J*=9.6, 4.0 Hz, 1H), 4.97 (d, *J*=5.6 Hz, 1H), 3.93~3.90 (m, 1H), 1.62 (s, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ : 137.7, 136.1, 132.6, 129.6, 129.2, 128.6, 128.3, 128.2, 128.0, 127.4, 126.7, 126.3, 71.3, 47.3; HRMS (APCI-ion trap) calcd for $\text{C}_{16}\text{H}_{12}\text{O}$ [*M*-2H]⁺ 220.0899, found 220.0889.

(1*S**,2*R**)-2-(4-Methylphenyl)-1,2-dihydronaphthalen-1-ol (**3b**): Colorless oil (44.4 mg, 94% yield). R_f =0.20 on silica gel (ethyl acetate/petroleum ether, *V* : *V* = 1 : 20). The *ee* was determined to be 62% using HPLC analysis on a Chiralcel OD-H column (hexane/2-propanol, *V* : *V* = 99 : 1, 1.0 mL/min, λ =254 nm). Retention times were 15.346 (major) and 31.153 (minor) min. ^1H NMR (500 MHz, CDCl_3) δ : 7.39~7.15 (m, 8H), 6.74 (dd, *J*=9.6, 1.8 Hz, 1H), 6.16 (dd, *J*=9.6, 4.1 Hz, 1H), 4.96 (t, *J*=6.9 Hz, 1H), 4.05~3.68 (m, 1H), 2.37 (s, 3H), 1.59 (s, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ : 137.0, 136.2, 134.4, 132.7, 129.9, 129.3, 129.1, 128.2, 128.0, 127.9, 126.6, 126.3, 71.2, 46.9, 21.0; HRMS (APCI-ion trap) calcd for $\text{C}_{17}\text{H}_{15}\text{O}$ [*M*-H]⁺ 235.1123, found 235.1123.

(1*S**,2*R**)-2-(2-Methylphenyl)-1,2-dihydronaphthalen-1-ol (**3c**): Colorless oil (43.7 mg, 92% yield). R_f =0.20 on silica gel (ethyl acetate/petroleum ether, *V* : *V* = 1 : 20). The *ee* was determined to be 20% using HPLC analysis on a Chiralcel OD-H column (hexane/2-propanol, *V* : *V* = 90 : 10, 1.0 mL/min, λ =254 nm). Retention times were 6.952 (minor) and 11.725 (major) min. ^1H NMR (500 MHz, CDCl_3) δ : 7.53~7.16 (m, 8H), 6.76 (dd, *J*=9.6, 2.2 Hz, 1H), 6.11 (dd, *J*=9.6, 3.1 Hz, 1H), 4.87 (s, 1H), 4.25~4.23 (m, 1H), 2.47 (s, 3H), 1.56 (s, 1H); ^{13}C NMR

(125 MHz, CDCl_3) δ : 136.7, 136.6, 135.4, 132.5, 130.7, 130.43, 129.3, 128.61, 127.9, 127.9, 127.6, 127.3, 126.5, 126.3, 69.6, 43.2, 19.8; HRMS (APCI-ion trap) calcd for $\text{C}_{17}\text{H}_{17}\text{O}$ $[\text{M}+\text{H}]^+$ 237.1279, found 237.1274.

(1*S**,2*R**)-2-(4-tertbutylphenyl)-1,2-dihydronaphthalen-1-ol (**3d**): Colorless oil (52.9 mg, 95% yield). R_f =0.20 on silica gel (ethyl acetate/petroleum ether, $V:V=1:20$). The *ee* was determined to be 36% using HPLC analysis on a Chiralcel OD-H column (hexane/2-propanol, $V:V=90:10$, 1.0 mL/min, $\lambda=254$ nm). Retention times were 5.040 (major) and 10.656 (minor) min. ^1H NMR (500 MHz, CDCl_3) δ : 7.42~7.12 (m, 8H), 6.70 (dd, $J=9.6$, 2.0 Hz, 1H), 6.14 (dd, $J=9.6$, 4.1 Hz, 1H), 5.01~4.82 (m, 1H), 3.87~3.85 (m, 1H), 1.57 (s, 1H), 1.31 (s, 9H); ^{13}C NMR (125 MHz, CDCl_3) δ : 150.3, 136.1, 134.4, 132.7, 129.9, 128.9, 128.2, 128.0, 127.9, 126.7, 126.3, 125.6, 71.3, 46.8, 34.4, 31.3; HRMS (APCI-ion trap) calcd for $\text{C}_{20}\text{H}_{22}\text{ONa}$ $[\text{M}+\text{Na}]^+$ 301.1569, found 301.1563.

(1*S**,2*R**)-2-(2-Methoxyphenyl)-1,2-dihydronaphthalen-1-ol (**3e**): Colorless oil (45.1 mg, 90 % yield). R_f =0.25 on silica gel (ethyl acetate/petroleum ether, $V:V=1:20$). The *ee* was determined to be 2% using HPLC analysis on a Chiralcel OD-H column (hexane/2-propanol, $V:V=90:10$, 1.0 mL/min, $\lambda=254$ nm). Retention times were 7.753 (major) and 12.228 (minor) min. ^1H NMR (400 MHz, CDCl_3) δ : 7.55~7.04 (m, 8H), 6.71 (dd, $J=9.6$, 2.5 Hz, 1H), 6.07 (dd, $J=9.6$, 3.2 Hz, 1H), 4.82 (t, $J=5.9$ Hz, 1H), 4.20 (dt, $J=5.6$, 2.9 Hz, 1H), 2.43 (s, 3H), 1.53 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 136.7, 136.6, 135.4, 132.5, 130.6, 130.4, 129.3, 128.6, 127.9, 127.9, 127.6, 127.2, 126.5, 126.3, 69.6, 43.2, 19.7; HRMS (APCI-ion trap) calcd for $\text{C}_{17}\text{H}_{15}\text{O}_2$ $[\text{M}-\text{H}]^-$ 251.1072, found 251.1071.

(1*S**,2*R**)-2-(3-Methoxyphenyl)-1,2-dihydronaphthalen-1-ol (**3f**): Colorless oil (45.9 mg, 91% yield). R_f =0.25 on silica gel (ethyl acetate/petroleum ether, $V:V=1:20$). The *ee* was determined to be 56% using HPLC analysis on a Chiralcel OD-H column (hexane/2-propanol, $V:V=90:10$, 1.0 mL/min, $\lambda=254$ nm). Retention times were 9.644 (major) and 14.246 (minor) min. ^1H NMR (400 MHz, CDCl_3) δ : 7.36 (d, $J=6.7$ Hz, 1H), 7.33~7.21 (m, 3H), 7.17 (dd, $J=7.2$, 1.4 Hz, 1H), 6.94~6.84 (m, 1H), 6.84~6.81 (m, 2H), 6.70 (dd, $J=9.6$, 2.0 Hz, 1H), 6.12 (dd, $J=9.6$, 4.0 Hz, 1H), 4.94 (d, $J=5.9$ Hz, 1H), 3.92~3.80 (m, 1H), 3.74 (s, 3H), 1.61 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 159.8, 139.3, 136.1, 132.6, 129.6, 129.5, 128.3, 128.2, 128.0, 126.7, 126.4, 121.5, 114.8, 112.9, 71.3, 55.1, 47.3; HRMS (APCI-ion trap) calcd for $\text{C}_{17}\text{H}_{15}\text{O}_2$ $[\text{M}-\text{H}]^-$ 251.1072, found 251.1077.

(1*S**,2*R**)-2-(4-Methoxyphenyl)-1,2-dihydronaphthalen-1-ol (**3g**): Colorless oil (49.4 mg, 98% yield). R_f =0.25 on silica gel (ethyl acetate/petroleum ether, $V:V=1:20$). The *ee* was determined to be 51% using HPLC analysis on a Chiralcel OD-H column (hexane/2-propanol, $V:V=90:10$, 1.0 mL/min, $\lambda=254$ nm). Retention times were 9.063 (major) and 14.563 (minor) min. ^1H NMR (400 MHz, CDCl_3) δ : 7.34 (dd, $J=4.3$, 3.6 Hz, 1H), 7.30~7.20

(m, 2H), 7.21~7.10 (m, 3H), 6.95~6.74 (m, 2H), 6.67 (dd, $J=9.6$, 1.9 Hz, 1H), 6.09 (dd, $J=9.6$, 4.3 Hz, 1H), 4.92 (t, $J=6.7$ Hz, 1H), 3.82~3.78 (m, 1H), 3.76 (s, 3H), 1.50 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 159.0, 136.3, 132.7, 130.3, 130.0, 129.1, 128.1, 128.0, 127.9, 126.5, 126.2, 114.1, 71.2, 55.2, 46.4; HRMS (APCI-ion trap) calcd for $\text{C}_{17}\text{H}_{15}\text{O}_2$ $[\text{M}-\text{H}]^-$ 251.1072, found 251.1074.

(1*S**,2*R**)-2-(4-Bromophenyl)-1,2-dihydronaphthalen-1-ol (**3h**): Colorless oil (50.4 mg, 84% yield). R_f =0.20 on silica gel (ethyl acetate/petroleum ether, $V:V=1:20$). The *ee* was determined to be 64% using HPLC analysis on a Chiralcel OD-H column (hexane/2-propanol, $V:V=90:10$, 1.0 mL/min, $\lambda=254$ nm). Retention times were 7.446 (major) and 11.562 (minor) min. ^1H NMR (500 MHz, CDCl_3) δ : 7.43 (d, $J=8.3$ Hz, 2H), 7.32 (dd, $J=13.6$, 7.2 Hz, 2H), 7.26 (t, $J=6.9$ Hz, 1H), 7.18 (d, $J=7.2$ Hz, 1H), 7.13 (d, $J=8.3$ Hz, 2H), 6.71 (dd, $J=9.6$, 1.5 Hz, 1H), 6.08 (dd, $J=9.6$, 4.0 Hz, 1H), 4.91 (s, 1H), 3.82~3.80 (m, 1H), 1.54 (s, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ : 136.9, 135.9, 132.4, 131.6, 131.0, 129.1, 128.5, 128.4, 128.2, 126.5, 126.4, 121.3, 71.1, 46.7; HRMS (APCI-ion trap) calcd for $\text{C}_{16}\text{H}_{12}\text{BrO}$ $[\text{M}-\text{H}]^-$ 299.0072, found 298.9905, 299.0066.

(1*S**,2*R**)-2-(4-Fluorophenyl)-1,2-dihydronaphthalen-1-ol (**3i**): White solid (40.4 mg, 84% yield). m.p. 59~60 °C. R_f =0.13 on silica gel (ethyl acetate/petroleum ether, $V:V=1:20$). The *ee* was determined to be 60% using HPLC analysis on a Chiralcel OD-H column (hexane/2-propanol, $V:V=90:10$, 1.0 mL/min, $\lambda=254$ nm). Retention times were 6.792 (major) and 11.332 (minor) min. ^1H NMR (400 MHz, CDCl_3) δ : 7.42~7.14 (m, 6H), 7.06~6.95 (m, 2H), 6.70 (dd, $J=9.6$, 2.0 Hz, 1H), 6.09 (dd, $J=9.6$, 4.1 Hz, 1H), 4.92 (d, $J=5.8$ Hz, 1H), 3.86~3.83 (m, 1H), 1.59 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.4, 161.0, 136.0, 133.32, 133.29, 132.5, 130.8, 130.7, 129.6, 128.4, 128.3, 128.1, 126.5, 126.4, 115.5, 115.3, 71.3, 46.5; ^{19}F NMR (376 MHz, CDCl_3) δ : -115.4; HRMS (APCI-ion trap) calcd for $\text{C}_{16}\text{H}_{12}\text{FO}$ $[\text{M}-\text{H}]^-$ 239.0872, found 239.0873.

(1*S**,2*R**)-2-Methyl-1,2-dihydronaphthalen-1-ol (**3j**): Colorless oil (25.9 mg, 81% yield). R_f =0.21 on silica gel (ethyl acetate/petroleum ether, $V:V=1:20$). The *ee* was determined to be 56% using HPLC analysis on a Chiralcel AD-H column (hexane/2-propanol, $V:V=90:10$, 1.0 mL/min, $\lambda=254$ nm). Retention times were 6.494 (minor) and 6.841 (major) min. ^1H NMR (400 MHz, CDCl_3) δ : 7.36 (dd, $J=7.2$, 1.4 Hz, 1H), 7.32~7.20 (m, 2H), 7.12 (dd, $J=7.1$, 1.5 Hz, 1H), 6.51 (dd, $J=9.5$, 2.4 Hz, 1H), 5.80 (dd, $J=9.5$, 3.1 Hz, 1H), 4.84~4.22 (m, 1H), 2.79~2.50 (m, 1H), 1.59 (s, 1H), 1.25 (d, $J=7.4$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 136.6, 132.4, 132.3, 128.4, 127.6, 127.3, 126.5, 126.4, 71.6, 35.2, 14.0; HRMS (APCI-ion trap) calcd for $\text{C}_{11}\text{H}_{13}\text{O}$ $[\text{M}+\text{H}]^+$ 161.0966, found 161.0961.

(1*S**,2*R**)-2-Thienyl-1,2-dihydronaphthalen-1-ol (**3k**): Colorless oil (43.3 mg, 95% yield). R_f =0.25 on silica gel (ethyl acetate/petroleum ether, $V:V=1:20$). The *ee* was determined to be 71% using HPLC analysis on a Chiralcel

OD-H column (hexane/2-propanol, $V:V=90:10$, 1.0 mL/min, $\lambda=254$ nm). Retention times were 8.842 (major) and 11.989 (minor) min. ^1H NMR (500 MHz, CDCl_3) δ : 7.40 (d, $J=7.0$ Hz, 1H), 7.34~7.23 (m, 2H), 7.17 (dd, $J=5.9$, 1.8 Hz, 2H), 6.97 (dd, $J=4.3$, 3.1 Hz, 2H), 6.65 (dd, $J=9.6$, 0.9 Hz, 1H), 6.16 (dd, $J=9.5$, 4.6 Hz, 1H), 5.01 (t, $J=6.5$ Hz, 1H), 4.34~3.95 (m, 1H), 1.86 (s, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ : 139.9, 136.2, 132.4, 129.5, 128.18, 128.15, 128.0, 126.9, 126.4, 126.2, 125.1, 71.0, 42.5; HRMS (APCI-ion trap) calcd for $\text{C}_{14}\text{H}_{11}\text{OS}$ $[\text{M}-\text{H}]^-$ 227.0531, found 227.0531.

(1*S**,2*R**)-5,8-Dimethoxy-2-cyclopropyl-1,2-dihydronaphthalen-1-ol (**4a**): Colorless oil (44.3 mg, 90% yield). $R_f=0.2$ on silica gel (ethyl acetate/petroleum ether, $V:V=1:10$). The *ee* was determined to be 97% using HPLC analysis on a Chiralcel OD-H column (hexane/2-propanol, $V:V=90:10$, 1.0 mL/min, $\lambda=254$ nm). Retention times were 11.343 (minor) and 18.273 (major) min. ^1H NMR (400 MHz, CDCl_3) δ : 6.91 (dd, $J=9.8$, 3.2 Hz, 1H), 6.77 (d, $J=3.4$ Hz, 2H), 6.01~5.94 (m, 1H), 5.04 (s, 1H), 3.84 (s, 3H), 3.79 (s, 3H), 1.77 (d, $J=6.3$ Hz, 1H), 1.57~1.52 (m, 1H), 1.32~1.14 (m, 2H), 0.47 (dd, $J=10.1$, 6.9 Hz, 1H), 0.39~0.17 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 150.6, 149.6, 130.3, 125.4, 122.8, 120.8, 111.3, 110.6, 63.4, 56.2, 56.1, 46.2, 11.1, 3.9, 3.7; HRMS (ESI-ion trap) calcd for $\text{C}_{15}\text{H}_{17}\text{O}_2$ $[\text{M}+\text{H}-\text{H}_2\text{O}]^+$ 229.1223, found 229.1216.

(1*S**,2*R**)-5,8-Dimethoxy-2-(4-Fluorophenyl)-1,2-dihydronaphthalen-1-ol (**4b**): Colorless oil (50.0 mg, 85% yield). $R_f=0.2$ on silica gel (ethyl acetate/petroleum ether, $V:V=1:10$). The *ee* was determined to be 52% using HPLC analysis on a Chiralcel OD-H column (hexane/2-propanol, $V:V=90:10$, 1.0 mL/min, $\lambda=254$ nm). Retention times were 17.871 (major) and 32.805 (minor) min. ^1H NMR (400 MHz, CDCl_3) δ : 7.39 (dd, $J=8.6$, 5.4 Hz, 1H), 7.19~6.98 (m, 1H), 6.83 (q, $J=9.0$ Hz, 1H), 6.16~5.95 (m, 1H), 5.06 (dd, $J=4.7$, 1.5 Hz, 1H), 3.83 (d, $J=5.1$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.2, 160.8, 150.7, 149.7, 136.3, 136.2, 130.7, 130.6, 128.8, 122.2, 115.4, 115.2, 111.5, 111.1, 64.4, 56.2, 46.5, 22.6; ^{19}F NMR (375 MHz, CDCl_3) δ : -116.10; HRMS (ESI-ion trap) calcd for $\text{C}_{18}\text{H}_{16}\text{O}_2\text{F}$ $[\text{M}+\text{H}-\text{H}_2\text{O}]^+$ 283.1129, found 283.1120.

(1*S**,2*R**)-5,8-Dimethoxy-2-(4-Methylphenyl)-1,2-dihydronaphthalen-1-ol (**4c**): Colorless oil (53.9 mg, 91% yield). $R_f=0.2$ on silica gel (ethyl acetate/petroleum ether, $V:V=1:10$). The *ee* was determined to be 51% using HPLC analysis on a Chiralcel OD-H column (hexane/2-propanol, $V:V=90:10$, 1.0 mL/min, $\lambda=254$ nm). Retention times were 15.041 (major) and 25.944 (minor) min. ^1H NMR (400 MHz, CDCl_3) δ : 7.34 (d, $J=8.0$ Hz, 2H), 7.22 (d, $J=7.8$ Hz, 2H), 7.09 (dd, $J=9.8$, 3.2 Hz, 1H), 6.82 (q, $J=9.0$ Hz, 2H), 6.16~6.12 (m, 1H), 5.08 (dd, $J=4.6$, 1.2 Hz, 1H), 3.84 (s, 1H), 3.83 (s, 1H), 3.80~3.75 (m, 1H), 2.38 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 150.8, 149.7, 137.2, 136.5, 129.2, 129.1, 128.9, 124.4, 122.6, 121.9, 111.4, 110.9, 64.3, 56.2, 56.2, 46.8, 21.1;

HRMS (ESI-ion trap) calcd for $\text{C}_{19}\text{H}_{19}\text{O}_2$ $[\text{M}+\text{H}-\text{H}_2\text{O}]^+$ 279.1380, found 279.1372.

(1*S**,2*R**)-2-Phenyl-1,2-dihydrotriphenylen-1-ol (**5a**): A white solid (54.8 mg, 85% yield). m.p. 157~158 °C. $R_f=0.23$ on silica gel (ethyl acetate/petroleum ether, $V:V=1:10$). The *ee* was determined to be 55% using HPLC analysis on a Chiralcel AD-H column (hexane/2-propanol, $V:V=90:10$, 1.0 mL/min, $\lambda=254$ nm). Retention times were 46.597 (major) and 49.684 (minor) min. ^1H NMR (500 MHz, CDCl_3) δ : 8.87~8.62 (m, 2H), 8.50~8.36 (m, 1H), 8.30~8.18 (m, 1H), 7.78~7.69 (m, 2H), 7.66 (d, $J=9.8$ Hz, 1H), 7.64~7.58 (m, 2H), 7.23~7.07 (m, 5H), 6.52~6.48 (m, 1H), 5.58 (d, $J=7.4$ Hz, 1H), 4.16 (d, $J=5.9$ Hz, 1H), 2.17 (d, $J=7.6$ Hz, 1H); ^{13}C NMR (126 MHz, CDCl_3) δ : 138.6, 130.8, 130.6, 130.1, 129.9, 128.7, 128.6, 127.8, 127.4, 127.3, 127.0, 126.97, 126.9, 126.5, 126.4, 123.8, 123.6, 123.1, 123.0, 122.3, 69.4, 48.8; HRMS (ESI-ion trap) calcd for $\text{C}_{24}\text{H}_{20}\text{O}$ $[\text{M}+\text{H}]^+$ 323.1431, found 323.1430.

(1*S**,2*R**)-2-Cyclopropyl-1,2-dihydrotriphenylen-1-ol (**5b**): Colorless oil (51.5 mg, 90% yield). $R_f=0.18$ on silica gel (ethyl acetate/petroleum ether, $V:V=1:10$). The *ee* was determined to be 97% using HPLC analysis on a Chiralcel AD-H column (hexane/2-propanol, $V:V=90:10$, 1.0 mL/min, $\lambda=254$ nm). Retention times were 25.025 (major) and 26.224 (minor) min. ^1H NMR (500 MHz, CDCl_3) δ : 8.82~8.69 (m, 2H), 8.40 (d, $J=7.8$ Hz, 1H), 8.32~8.23 (m, 1H), 7.78~7.58 (m, 4H), 7.43 (dd, $J=9.8$, 3.2 Hz, 1H), 6.43~6.27 (m, 1H), 5.45 (d, $J=4.5$ Hz, 1H), 3.69 (t, $J=6.8$ Hz, 1H), 1.91~1.76 (m, 1H), 0.87~0.71 (m, 2H), 0.56~0.31 (m, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ : 132.1, 130.5, 130.4, 129.8, 129.6, 128.7, 127.2, 126.8, 126.7, 126.3, 124.1, 123.7, 123.0, 122.9, 122.7, 66.5, 46.8, 22.6, 11.1, 3.8; HRMS (ESI-ion trap) calcd for $\text{C}_{21}\text{H}_{17}$ $[\text{M}+\text{H}]^+$ 269.1325, found 269.1320.

(1*S**,2*R**)-2-(4-Fluorophenyl)-1,2-dihydrotriphenylen-1-ol (**5c**): Colorless oil (57.2 mg, 84% yield). $R_f=0.15$ on silica gel (ethyl acetate/petroleum ether, $V:V=1:10$). The *ee* was determined to be 68% using HPLC analysis on a Chiralcel AD-H column (hexane/2-propanol, $V:V=90:10$, 1.0 mL/min, $\lambda=254$ nm). Retention times were 37.883 (major) and 67.269 (minor) min. ^1H NMR (500 MHz, CDCl_3) δ : 8.98~8.63 (m, 2H), 8.40~8.29 (m, 1H), 8.29~8.20 (m, 1H), 7.78~7.62 (m, 4H), 7.59 (dd, $J=9.9$, 3.3 Hz, 1H), 7.50 (dd, $J=8.5$, 5.4 Hz, 2H), 7.16 (t, $J=8.7$ Hz, 2H), 6.44~6.41 (m, 1H), 5.41 (d, $J=4.4$ Hz, 1H), 4.08~3.95 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ : 163.1, 161.1, 135.9, 130.7, 130.7, 129.7, 128.6, 128.58, 127.3, 127.0, 126.5, 126.4, 124.1, 123.7, 123.1, 123.09, 115.6, 115.5, 67.6, 47.3; ^{19}F NMR (375 MHz, CDCl_3) δ : -115.57; ^1H RMS (ESI-ion trap) calcd for $\text{C}_{24}\text{H}_{18}\text{OF}$ $[\text{M}+\text{H}]^+$ 341.1336, found 341.1331.

(1*S**,2*R**)-2-(4-methylphenyl)-1,2-dihydrotriphenylen-1-ol (**5d**): Colorless oil (61.2 mg, 91% yield). $R_f=0.20$ on silica gel (ethyl acetate/petroleum ether, $V:V=1:10$). The *ee* was determined to be 65% using HPLC analysis on a Chiralcel AD-H column (hexane/2-propanol, $V:V=$

90 : 10, 1.0 mL/min, λ =254 nm). Retention times were 38.762 (major) and 45.206 (minor) min. ^1H NMR (500 MHz, CDCl_3) δ : 8.77~8.70 (m, 2H), 8.54~8.36 (m, 1H), 8.33~8.15 (m, 1H), 7.77~7.70 (m, 2H), 7.66 (d, J =9.8 Hz, 1H), 7.63~7.58 (m, 2H), 7.01 (dd, J =13.5, 5.7 Hz, 2H), 6.97~6.90 (m, 2H), 6.65~6.33 (m, 1H), 5.58 (s, 1H), 4.13 (d, J =4.6 Hz, 1H), 2.20 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 138.7, 138.3, 130.8, 130.6, 130.1, 129.9, 128.7, 128.5, 127.8, 127.4, 127.3, 127.0, 126.9, 126.4, 126.4, 124.6, 123.9, 123.6, 123.1, 123.0, 122.2, 69.5, 48.8, 21.4; ^1H HRMS (ESI-ion trap) calcd for $\text{C}_{25}\text{H}_{20}\text{ONa}$ [$\text{M}+\text{Na}$] $^+$ 359.1406, found 359.1401.

(1*S**,2*R**)-2-Thienyl-1,2-dihydrotriphenyl-1-ol (**5e**): Colorless oil (61.0 mg, 93% yield). R_f =0.20 on silica gel (ethyl acetate/petroleum ether, $V:V$ =1:10). The *ee* was determined to be 87% using HPLC analysis on a Chiralcel AD-H column (hexane/2-propanol, $V:V$ =90:10, 1.0 mL/min, λ =254 nm). Retention times were 55.292 (major) and 58.044 (minor) min. ^1H NMR (500 MHz, CDCl_3) δ : 9.01~8.60 (m, 2H), 8.48~8.23 (m, 2H), 7.93~7.64 (m, 4H), 7.59 (dd, J =9.8, 3.3 Hz, 1H), 7.46~7.33 (m, 1H), 7.27 (d, J =3.4 Hz, 1H), 7.15 (dd, J =5.2, 3.4 Hz, 1H), 6.52~6.34 (m, 1H), 5.53 (d, J =4.2 Hz, 1H), 4.47~4.23 (m, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ : 142.7, 130.7, 130.66, 130.6, 129.7, 128.9, 128.6, 127.4, 127.1, 127.0, 126.0, 126.4, 126.1, 124.8, 124.1, 124.0, 123.9, 123.1, 123.1, 67.7, 43.5; ^1H HRMS (ESI-ion trap) calcd for $\text{C}_{22}\text{H}_{15}\text{S}$ [$\text{M}-\text{H}_2\text{O}+\text{H}$] $^+$ 311.0889, found 311.0885.

Supporting Information Copies of ^{13}C NMR and ^1H NMR and ^{19}F NMR spectra for all products, HRMS and HPLC data of all products, crystal structure and data of the product **3d**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Roberto, P.; Francesco, B.; Nicola, C. A.; Marcello, L.; Vincenzo, T.; Francesco, F.; Vincenzo, O.; Alberto, G.; Stefano, G. *J. Med. Chem.* **1995**, *38*, 942.
(b) Scott, E. S.; Felix, A. A.; Ratna, C.; David, E. N.; Val, J. W.; Richard, B. M. *J. Med. Chem.* **1995**, *38*, 2395.
(c) Lautens, M.; Rovis, T. *J. Org. Chem.* **1997**, *62*, 5246.
(d) Lautens, M.; Rovis, T. *J. Am. Chem. Soc.* **1997**, *119*, 11090.
(e) Lautens, M.; Fagnou, K.; Taylor, M.; Rovis, T. *J. Organomet. Chem.* **2001**, *624*, 259.
(f) Lautens, M.; Fagnou, K.; Zunic, V. *Org. Lett.* **2002**, *4*, 3465.
(g) Tsoung, J.; Krämer, K.; Zajdlík, A.; Liébert, C.; Lautens, M. *J. Org. Chem.* **2011**, *76*, 9031.
- [2] For reviews, see: (a) Fagnou, K.; Lautens, M. *Chem. Rev.* **2003**, *103*, 169.
(b) Hayashi, T.; Yamasaki, K. *Chem. Rev.* **2003**, *103*, 2829.
(c) Lautens, M.; Fagnou, K.; Hiebert, S. *Acc. Chem. Res.* **2003**, *36*, 48.
(d) Rayabarapu, D. K.; Cheng, C.-H. *Acc. Chem. Res.* **2007**, *40*, 971.
- [3] Nakamura, M.; Matsuo, K.; Inoue, T.; Nakamura, E. *Org. Lett.* **2003**, *5*, 1373.
- [4] (a) Lautens, M.; Ma, S. *J. Org. Chem.* **1996**, *61*, 7246.
(b) Rayabarapu, D. K.; Chiou, C.-F.; Cheng, C.-H. *Org. Lett.* **2002**, *4*, 1679.
(c) Wang, G.; Gan, Y.; Liu, Y. *Chin. J. Chem.* **2018**, *36*, 916.
(d) Fu, X.-P.; Xiao, Y.-L.; Zhang, X. *Chin. J. Chem.* **2018**, *36*, 143.
- [5] (a) Bertozzi, F.; Pineschi, M.; Macchia, F.; Arnold, L. A.; Minnaard, A. J.; Feringa, B. L. *Org. Lett.* **2002**, *4*, 2703.
(b) Zhang, W.; Wang, L.-X.; Shi, W.-J.; Zhou, Q.-L. *J. Org. Chem.* **2005**, *70*, 3734.
(c) Zhang, L.-Y.; Wang, L. *Chin. J. Chem.* **2006**, *24*, 1605.
- [6] (a) Carreras, J.; Avenoza, A.; Busto, J. H.; Peregrina, J. M. *Org. Lett.* **2007**, *9*, 1235.
(b) Cortez, G. A.; Baxter, C. A.; Schrock, R. R.; Hoveyda, A. H. *Org. Lett.* **2007**, *9*, 2871.
(c) Machin, B. P.; Howell, J.; Mandel, J.; Blanchard, N.; Tam, W. *Org. Lett.* **2009**, *11*, 2077.
- [7] (a) Lautens, M.; Fagnou, K.; Taylor, M. *Org. Lett.* **2000**, *2*, 1677.
(b) Lautens, M.; Fagnou, K. *Tetrahedron* **2001**, *57*, 5067.
(c) Lautens, M.; Fagnou, K. *J. Am. Chem. Soc.* **2001**, *123*, 7170.
(d) Preetz, A.; Kohrt, C.; Drexler, H. J.; Torrens, A.; Buschmann, H.; Lopez, M. G.; Heller, D. *Adv. Synth. Catal.* **2010**, *352*, 2073.
- [8] (a) Lautens, M.; Renaud, J.-L.; Hiebert, S. *J. Am. Chem. Soc.* **2000**, *122*, 1804.
(b) Li, L.-P.; Rayabarapu, D. K.; Nandi, M.; Cheng, C.-H. *Org. Lett.* **2003**, *5*, 1621.
(c) Cabrera, S.; Arrayás, R. G.; Carretero, J. C. *Angew. Chem., Int. Ed.* **2004**, *43*, 3944.
(d) Ogura, T.; Yoshida, K.; Yanagisawa, A.; Imamoto, T. *Org. Lett.* **2009**, *11*, 2245.
(e) Li, M.; Yan, X.-X.; Hong, W.; Zhu, X.-Z.; Cao, B.-X.; Sun, J.; Hou, X.-L. *Org. Lett.* **2004**, *6*, 2833.
- [9] (a) Pan, X.-J.; Huang, G.-B.; Long, Y.-H.; Zuo, X.-J.; Xu, X.; Gu, F.-L.; Yang, D.-Q. *J. Org. Chem.* **2014**, *79*, 187.
(b) Yang, D.-Q.; Liang, N. *Org. Biomol. Chem.* **2014**, *12*, 2080.
(c) Meng, L.; Yang, W.; Pan, X.-J.; Tao, M.; Cheng, G.; Wang, S.-Y.; Zeng, H.-P.; Long, Y.-H.; Yang, D.-Q. *J. Org. Chem.* **2015**, *80*, 2503.
- [10] (a) Yang, D.-Q.; Long, Y.-H.; Wang, H.; Zhang, Z.-M. *Org. Lett.* **2008**, *10*, 4723.
(b) Long, Y.-H.; Yang, D.-Q.; Zhang, Z.-M.; Wu, Y.-J.; Zeng, H.-P.; Chen, Y. *J. Org. Chem.* **2010**, *75*, 7291.
(c) Yang, D.-Q.; Long, Y.-H.; Zhang, J.-F.; Zeng, H.-P.; Wang, S.-Y.; Li, C.-R. *Organometallics* **2010**, *29*, 3477.
(d) Yang, D.-Q.; Long, Y.-H.; Wu, Y.-J.; Zuo, X.-J.; Tu, Q.-Q.; Fang, S.; Jiang, L.-S.; Wang, S.-Y.; Li, C.-R. *Organometallics* **2010**, *29*, 5936.
(e) Fang, S.; Liang, X.-L.; Long, Y.-H.; Li, X.-L.; Yang, D.-Q.; Wang, S.-Y.; Li, C.-R. *Organometallics* **2012**, *31*, 3113.
(f) Cheng, H.-C.; Yang, D.-Q. *J. Org. Chem.* **2012**, *77*, 9756.
(g) Yang, D.-Q.; Xia, J.-Y.; Long, Y.-H.; Zeng, Z.-Y.; Zuo, X.-J.; Wang, S.-Y.; Li, C.-R. *Org. Biomol. Chem.* **2013**, *11*, 4871.
(h) Yang, W.; Luo, R.-S.; Yang, D.-Q. *Molecules* **2015**, *20*, 21103.
- [11] (a) Feng, C.-C.; Nandi, M.; Sambaiah, T.; Cheng, C.-H. *J. Org. Chem.* **1999**, *64*, 3538.
(b) Chen, C.-L.; Martin, S. F. *Org. Lett.* **2004**, *6*, 3581.
(c) Chen, C.-L.; Martin, S.-F. *J. Org. Chem.* **2006**, *71*, 4810.
- [12] (a) Arrayas, R. G.; Cabrera, S.; Carretero, J. C. *Org. Lett.* **2003**, *5*, 1333.
(b) Zhang, W.; Zhu, S.-F.; Qiao, X.-C.; Zhou, Q.-L. *Chem.-Asian J.* **2008**, *3*, 2105.
- [13] (a) Lautens, M.; Dockendorff, C.; Fagnou, K.; Malicki, A. *Org. Lett.* **2002**, *4*, 1311.
(b) Murakami, M.; Igawa, H. *Chem. Commun.* **2002**, 390.
(c) Lautens, M.; Dockendorff, C. *Org. Lett.* **2003**, *5*, 3695.
(d) Zhang, T.-K.; Mo, D.-L.; Dai, L.-X.; Hou, X.-L. *Org. Lett.* **2008**, *10*, 3689.
(e) Luo, F.; Pan, C.; Cheng, J. *Chin. J. Org. Chem.* **2010**, *30*, 633.
- [14] (a) Li, Y.; Yang, W.; Cheng, G.; Yang, D.-Q. *J. Org. Chem.* **2016**, *81*, 4744.
(b) Wu, R. H.; Yang, W.; Chen, W. K.; Yang, D.-Q. *Org. Chem. Front.* **2017**, *4*, 1921.

- (c) Zhang, S.; Yang, S.; Huang, L.; Zhao, B.; Cheng, K.; Qi, C. *Chin. J. Org. Chem.* **2015**, *35*, 2259 (in Chinese).
(张诗浓, 杨胜虎, 黄乐浩, 赵保丽, 程凯, 齐陈泽, 有机化学, **2015**, *35*, 2259.)
- [15] (a) Seigerman, C. K.; Micyus, T. M.; Neufeldt, S. R.; Sanford, M. S. *Tetrahedron* **2013**, *69*, 5580.
(b) Dubbaka, S. R.; Narreddula, V. R.; Gadde, S.; Mathew, T. *Tetrahedron* **2014**, *70*, 9676.
- [16] Molander, G. A.; Sandrock, D. L. *Curr. Opin. Drug Discovery Dev.* **2009**, *12*, 811.
- [17] (a) Li, S. F.; Xu, J. B.; Fan, B.-M.; Lu, Z. W.; Zeng, C. Y.; Bian, Z. X.; Zhou, Y. Y.; Wang, J. *Chem.-Eur. J.* **2015**, *21*, 9003.
(b) Xu, X.; Chen, J.-C.; He, Z.-X.; Zhou, Y.-Y.; Fan, B.-M. *Org. Biomol. Chem.* **2016**, *14*, 2480.
- [18] (a) Barbeiro, C. S.; Vasconcelos, S. N. S.; de Oliveira, I. M.; Zuckerman-Schpector, J.; Caracelli, I.; Maganhi, S. H.; Stefani, H. A. *ChemistrySelect* **2017**, *2*, 8173.
(b) Liu, L.; Dong, Y.; Pang, B.; Ma, J. *J. Org. Chem.* **2014**, *79*, 7193.
- [19] Zhang, W.; Chen, J. C.; Zeng, G. Z.; Yang, F.; Xu, J. B.; Sun, W. Q.; Shinde, M. V.; Fan, B. M. *J. Org. Chem.* **2017**, *82*, 2641.
- [20] (a) Lautens, M.; Fagnou, K.; Yang, D.-Q. *J. Am. Chem. Soc.* **2003**, *125*, 14884.
(b) Villeneuve, K.; Tam, W. *Organometallics* **2007**, *26*, 6082.
- (Zhao, C.)