无催化条件下采用叔丁基过氧化氢(TBHP)选择性氧化硫醚/ 二硫醚至亚砜/单砜

蒋筱莹^a 姚传胜^b 童 踔^b 白仁仁^b 周 涛^c 谢媛媛^{*a,b}

(^a 浙江工业大学 长三角绿色制药协同创新中心 杭州 310014)

(^b 浙江工业大学药学院 杭州 310014)

(^c 浙江工商大学食品科学与生物技术学院 杭州 310018)

摘要 在常见氧化剂叔丁基过氧化氢(TBHP)的存在下,成功开发了一种选择性将硫醚氧化成亚砜,将二硫醚氧化成单砜的新方法,得到一系列亚砜和单砜类化合物,收率为67%~98%。该方法具有不使用任何催化剂和添加剂、底物适用性广、反应效率高、选择性好等优点。此外,通过控制实验,提出了一种合理的自由基反应机理。

关键词 硫醚/二硫醚;亚砜/单砜;叔丁基过氧化氢;氧化反应;无催化剂添加剂

Selective Oxidation of Sulfides/Disulfides to Sulfoxides/ Thiosulfonates Using *t*-Butyl Hydroperoxide (TBHP) without Catalyst

Jiang, Xiaoying^a Yao, Chuansheng^b Tong, Chuo^b Bai, Renren^b

Zhou, Tao^c Xie, Yuanyuan^{*a,b}

(^a Collaborative Innovation Center of Yangtze River Delta Region Green Pharmaceutical,
Zhejiang University of Technology, Hangzhou 310014)

(^b College of Pharmaceutical Sciences, Zhejiang University of Technology, Hangzhou 310014)

(^c School of Food Science and Biotechnology, Zhejiang Gongshang University, Hangzhou 310018)

Abstract A novel and simple method for the selectively oxidation of sulfides to sulfoxides and oxidation of disulfides to thiosulfonates in the presence of *t*-butyl hydroperoxide (TBHP) has been established. The oxidation system has the advantages of avoiding the use of any catalyst and additive, possessing a broad substrate scope and high efficiency. Desired sulfoxides and thiosulfonates products were obtained in moderate to good yields. Furthermore, a plausible mechanism of the reaction is proposed. The results of control experiments revealed that a radical pathway was involved in this oxidation.

Keywords sulfides/disulfides; sulfoxides/thiosulfonates; *t*-butylhydroperoxide; oxidation; catalyst/additive free

1 Introduction

Organic aromatic sulfoxides are significant skeletons in medicinal chemistry and especially in the synthesis of biologically and pharmaceutically compounds,^[1] such as anti-ulcer omeprazole^[2] and central stimulant modafinil^[3] (Figure 1). They also have other biological activities, including antigout,^[4a] antidiabetic,^[4b] anti-inflammatory,^[4c] antifungal,^[4d] anti-HIV,^[4e] antitumor^[4f] and insecticidal activities.^[4g] Furthermore, they have found extensive app-

Figure 1 Biologically active compounds containing sulfoxide moieties

lications in organic chemistry^[5~8] as versatile synthetic

* Corresponding author. E-mail: xyycz@zjut.edu.cn

Received December 26, 2019; revised March 10, 2020; published online March 31, 2020.

Project supported by the National Natural Science Foundation of China (No. 21576239).

国家自然科学基金(No. 21576239)资助项目.

intermediates, ligands and catalysts, as well as in industry as producing polymer and photographic processes materials.^[9]

In light of the importance of sulfoxides, development of the procedures for the selective oxidation of sulfides to sulfoxides becomes increasingly challenging and significant in synthetic methodology. Since the firstly reported synthesis of sulfoxides by Maercker in 1865,^[10] various reagents and oxidizing media have been employed for the conversion of sulfides to sulfoxides. Palmieri and co-workers^[11] reported the oxidation of sulfides into sulfoxides using (*n*-Bu₄N)(AuCl₄) and nitric acid as catalyst and oxidant. A number of new catalytic strategies involving the activated molecular oxygen have been developed for the oxidation of sulfides using traditional oxidants such as H₂O₂^[12] and *t*-butylhydroperoxide (TBHP), for example, which catalyzed by tetrazole amides or aluminum tri-*tert*-butoxide.^[13] Some metal or no-metal catalysts were also applied with other oxidants including hypervalent iodine reagents,^[14–16] hypervalent metal oxidants,^[17–20] Schiff bases,^[21] Oxone,^[22] ionic liquid^[23] and so on.^[24–26] In addition, new and efficient technologies have been promoted, such as porous solids,^[27] nanosphere,^[28] magnetic nanoparticle^[29] and visible light photocatalytic.^[30] Almost all these oxidation systems need catalysts or additives, their selectivity is often limited to the oxidation of sulfides to sulfoxides. Therefore, development of a new and high selective method without catalysts or additives is required (Scheme 1).

Using TBHP as an oxidant has the advantages of environmentally benign, cheap and more abundant characteristics. TBHP can be used in the oxidation of a variety of compounds including alcohols, aliphatic methylene, aromatic methylene and other compounds.^[31] Herein, a convenient protocol for the direct selective oxidation of sulfides by TBHP to give the desired products in moderate to excellent yields in the absence of catalyst and additive was

Previously reported work:

Our work:

Scheme 1 Representative of oxidation of sulfides

reported. Surprisingly, this method can also be applied to the selective oxidation of disulfides to thiosulfonates.

2 Results and discussion

Initially, the thioanisole **1a** was chosen as the model substrate for the optimization of reaction conditions in 1,2-dichloroethane at 80 °C. The desired product (methylsulfinyl)benzene (**2a**) was obtained in 50%~95% yields with the increase of TBHP amount (Table 1, Entries 1~3), while the yield of sulfone **3a** remained essentially unchanged. However, when TBHP was increased to 10.0 equiv. and reaction time was prolonged to 24 h, sulfone **3a** became the main product (Table 1, Entry 4). When H₂O₂ was used as an oxidant, the yield of sulfoxide was decreased, while the yield of byproduct sulfone was increased (Table 1, Entry 5). The reaction did not proceed in the presence of other oxidants such as K₂S₂O₈ and di-*t*-butyl peroxide (DTBP) (Table 1, Entries 6, 7). The solvents such as CHCl₃, CH₂Cl₂, MeCN, EtOAc and tetrahydrofuran (THF), were also screened for the reaction. Interestingly, a higher yield (98%) of sulfoxide was obtained in CHCl₃ with the shortest time, and almost no

Table 1 Optimization of conditions for oxidation of sulfides to sulfoxides^a

Entry	Oxidant (equiv.)	Solvent	Temp./°C	Time/h	Conversion/% of 1a	Yield ^b / % of 2a	Yield ^b / % of 3a
1 ^c	TBHP (3.0)	ClCH ₂ CH ₂ Cl	80	6	55	50	0
2 ^c	TBHP (4.0)	ClCH ₂ CH ₂ Cl	80	6	84	80	0
3	TBHP (5.0)	ClCH ₂ CH ₂ Cl	80	3	99	95	3
4	TBHP (10.0)	ClCH ₂ CH ₂ Cl	80	24	98	10	85
5	H ₂ O ₂ (5.0)	ClCH ₂ CH ₂ Cl	80	6	98	87	10
6	K ₂ S ₂ O ₈ (5.0)	ClCH ₂ CH ₂ Cl	80	6	25	N.R.	N.R.
7	DTBP (5.0)	ClCH ₂ CH ₂ Cl	80	6	30	N.R.	N.R.
8	TBHP (5.0)	CHCl₃	60	1.5	99	98	Trace
9 ^d	TBHP (5.0)	CH ₂ Cl ₂	60	10	98	90	5
10	TBHP (5.0)	CH ₃ CN	81	24	99	91	4
11	TBHP (5.0)	EtOAc	77	11	98	90	6
12	TBHP (5.0)	THF	65	7	42	N.R.	N.R.

^a Reaction conditions: **1a** (1 mmol), oxidant, solvent (2.0 mL). ^b Isolated yields based on **1a**. ^c Reaction was not complete. ^d Reaction in a sealed tube.

sulfone was detected (Table 1, Entry 8). On the contrary, the use of other solvents led to decrease in the yield of **2a**, and generation of sulfone. The best result was obtained in the presence of 5.0 equiv. of TBHP using CHCl_3 as solvent. And no better results were gained in further variations in temperature, time and so forth. Therefore, the optimal conditions for this reaction are thioanisole (1.0 mmol), TBHP (5.0 mmol) and CHCl_3 (2 mL) at 60 °C for 1.5 h.

Under the optimized reaction conditions, the scope of oxidation reaction of **1** with TBHP to **2** was next explored (Table 2). A series of sulfides with different substituents were investigated. Overall, all the substrates could be transformed into the corresponding sulfoxides smoothly in good to excellent yields ranging from 67% to 98%. The substrates with an electron-donating group on the benzene ring reacted faster than those with electron-withdrawing groups (**2b**, **2c** vs **2d**, **2e**), but they all gave the corresponding products in excellent yields (>95%). Similarly, sulfide with different R^2 also underwent efficient oxidation to afford the corresponding sulfoxide. When R^2 was ethyl or benzyl, R^2 had a little influence on the reaction and the reaction rate. The yield was slightly reduced. However, when R^2 was phenyl, the reaction rate was greatly reduced, and more oxidants (8 equiv.) were needed in order to obtain high yield of product (**2f**~**2i** vs **2j**). The difficulty of oxidation could be attributed to the steric hindrance effect and conjugative effect of benzene ring. Conversely, it had the advantage of producing byproduct more difficultly. In addition, reactions of the substrates containing sensitive groups, such as carbon-carbon double bonds, carbon-carbon triple bonds, aldehyde group and hydroxyl groups, can also proceed (**2k**~**2n**). A high yield was still achieved in the cases of allyl sulfides **1k** and propenyl sulfides **1l**. Although the aldehyde group is unstable in the presence of oxidant, oxidation of **1m** provided desired product **2m** in a moderate yield of 67% at relatively lower temperature.

Table 2 Substrate scope of sulfides to sulfoxides^a

Entry	R^1	R^2	Time/h	Product	Yield ^b /%
1	H	CH_3	1.5	2a	98
2	4- CH_3	CH_3	1.5	2b	97
3	4- OCH_3	CH_3	1	2c	98
4	4-Cl	CH_3	2	2d	96
5	4-Br	CH_3	2	2e	95
6	4- CH_3	CH_2CH_3	3	2f	93
7	4-Cl	CH_2CH_3	5	2g	92
8	4- CH_3	CH_2Ph	3	2h	96
9	4-Cl	CH_2Ph	4	2i	94
10 ^c	H	Ph	7	2j	95
11	4- CH_3	$\text{CH}_2\text{CH}=\text{CH}_2$	0.8	2k	93
12	4- CH_3	$\text{CH}_2\text{C}\equiv\text{CH}$	6	2l	87
13 ^d	4-CHO	CH_3	4	2m	67
14 ^e	4-OH	4- HOC_6H_4	9	2n	85

^a Reaction conditions: **1** (1 mmol), TBHP (5.0 mmol), CHCl_3 (2.0 mL), stirred at 60 °C unless noted. ^b Isolated yields based on **1**. ^c 8.0 equiv. TBHP were used. ^d 50 °C. ^e 2 mL CHCl_3 , 0.5 mL CH_3CN .

Product **2n** was also generated in good yield, but longer reaction time was necessary due to the steric-hindrance effect of 4-hydroxyphenyl groups. No other byproduct was observed in this reaction.

Interestingly, this system can also be applied to the selective oxidation of disulfide. Therefore, the oxidation reactions of 1,2-diphenyldisulfane (**4a**) with TBHP were discussed. Surprisingly, thiosulfonate (**5a**) was obtained rather than sulfoxide compounds. Thiosulfonate moiety exists in many compounds with various pharmacological activities including antibacterial sulfamethoxazole,^[32] antitumor probenecid,^[33] antidiabetic glibenclamide^[34] and the COX-2 inhibitor rofecoxib^[35] (Figure 2). The previously reported methods using TBHP as oxidant need other catalysts or additives.^[13,26] Therefore, oxidation of different disulfides was investigated under the same conditions for the oxidation of sulfides.

Figure 2 Biologically active compounds containing sulfone moieties

Further optimization for the reaction conditions is shown in Table 3. With the decreased of TBHP amount, the yield was reduced even the reaction proceeded for longer time. When TBHP was 3 equiv., the raw materials reacted incompletely (Table 3, Entries 1~3). It needed longer reaction time when the temperature decreased to 50 °C (Table 3, Entry 4). Ulteriorly, when the temperature dropped to 40 °C, the corresponding product **5a** was formed in trace amounts (Table 3, Entry 5). Increasing the amount of oxidant or prolonging reaction time did not increase the yield any more (Table 3, Entries 6, 7). Therefore, the optimal conditions for this reaction are ascertained as: 1,2-diphenyldisulfane (1.0 mmol), TBHP (5.0 mmol) and CHCl_3 (2 mL) at 60 °C.

With the optimized reaction condition in hand, the scope of oxidation reaction of **4** with TBHP to **5** was next explored (Table 4). Numbers of disulfides with different substituent groups on the benzene ring were investigated. In general, electron-donating groups were more tolerated in this reaction than electron-withdrawing groups (**5b**, **5c** vs **5e**, **5f**). The multi-substituted products (**5d**) were obtained in 85% yield possibly because of the steric-hindrance ef-

fect. However, the benzyl substrate (**4g**) provided the corresponding product (**5g**) in the lowest yield amongst the seven symmetric disulfides. Asymmetric disulfide (**4h**) was also applied in this system, yielding the desired product (**5h**) in 18% yield with many byproducts. It demonstrated that the oxidation of asymmetric disulfides has low selectivity.

Table 3 Optimization of conditions for oxidation of disulfides to thiosulfonates^a

Entry	Oxidant (equiv.)	Solvent	Temp./°C	Time/h	Yield ^b /%
1	TBHP (5.0)	CHCl ₃	60	2	89
2	TBHP (4.0)	CHCl ₃	60	7	88
3 ^c	TBHP (3.0)	CHCl ₃	60	14	60
4	TBHP (5.0)	CHCl ₃	50	6	88
5	TBHP (5.0)	CHCl ₃	40	8	Trace
6	TBHP (5.0)	CHCl ₃	60	10	85
7	TBHP (10.0)	CHCl ₃	60	1.5	88

^a Reaction conditions: **4a** (1 mmol), TBHP, CHCl₃ (2.0 mL). ^b Isolated yields based on **4a**. ^c Reaction was not complete.

Table 4 Substrate scope of disulfides to thiosulfonates^a

Entry	R ³	R ⁴	Time ^b /h	Product	Yield ^c /%
1	Ph	Ph	2	5a	89
2	4-CH ₃ C ₆ H ₄	4-CH ₃ C ₆ H ₄	3	5b	90
3	4-CH ₃ OC ₆ H ₄	4-CH ₃ OC ₆ H ₄	2	5c	90
4	2,4-(CH ₃) ₂ C ₆ H ₃	2,4-(CH ₃) ₂ C ₆ H ₃	5	5d	85
5	4-ClC ₆ H ₄	4-ClC ₆ H ₄	4	5e	87
6	4-BrC ₆ H ₄	4-BrC ₆ H ₄	5.5	5f	73
7	Bn	Bn	10	5g	22
8	4-CH ₃ C ₆ H ₄	4-BrC ₆ H ₄	12	5h	18

^a Reaction conditions: **4** (1 mmol), TBHP (5.0 mmol), CHCl₃ (2.0 mL), stirred at 60 °C unless noted. ^b Time taken on complete the reaction. ^c Isolated yields based on **1**.

According to the experiment results and previous reports,^[36] a plausible mechanism of this reaction is proposed (Scheme 2). Initially, *t*-butylhydroperoxide generates *t*-butyl oxygen radical and hydroxyl radical by cracking. Then, *t*-butyl peroxide radical forms through the reaction of *t*-butylhydroperoxide with *t*-butyl oxygen radical. Subsequently, thioanisole radical positive ion forms by the reaction of thioanisole **1a** and *t*-butyl peroxide radical through single electron transfer (SET) process. Finally, *t*-butyl peroxide radical attacks thioanisole radical positive ion to afford (methylsulfinyl)benzene (**2a**). For 1,2-diphenyldisulfane (**4a**), it is oxidized by TBHP to produce sulfoxide compound **A**, which is then oxidized to form sulfoxide compound **B**. The intermediate **B** is rapidly de-

composed to form sulfoxide radical intermediate **C**.^[37] Next, intermediate **C** attacks compound **A** to afford thio-sulfonate product **5a** and sulfur radical **D**. Moreover, sulfur radical **D** reacts with compound **A** to obtain raw material **4a** and sulfoxide radical intermediate **C**, forming a cycle until the termination of the reaction.

Scheme 2 A plausible reaction mechanism

Several control experiments were designed in order to gain more deep understanding about the reaction mechanism. In the first place, the reaction was performed under nitrogen atmosphere and the yield of product **2a** was the same as that obtained under air. This result revealed that oxygen was not involved in oxidation reaction (Eq. 1). Next, using TBHP (5.0~6.0 mol/L in decane) as oxidant, the result was the same as that of TBHP aqueous solution, indicating that the oxygen in the sulfoxide product doesn't come from water (Eq. 2). Finally, 2 equiv. of free radical scavenger 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) was introduced, the oxidation reaction was absolutely suppressed, and only 10% yield of product **2a** was isolated. Clearly, a radical pathway was involved (Eq. 3).

3 Conclusions

In conclusion, we have developed an efficient method for the selective oxidation of sulfides or disulfides to form sulfoxides or thiosulfonates by TBHP in the absence of catalyst and additive. This transformation proved a broad substrate scope and high efficiency. Furthermore, a plausible free radical mechanism of this reaction is proposed. The control experiments demonstrated that the oxidation performed was carried out through a radical pathway, whereas oxygen and water were not involved in oxidation.

4 Experimental section

4.1 General information

All reagents were obtained from commercial suppliers and used without further purification, unless otherwise indicated. TLC analysis was performed using pre-coated glass plates. Silica gel for column chromatography was purchased from Qingdao Haiyang Chemical Co., Ltd. Melting points were determined using a Büchi B-540 capillary melting point apparatus. ^1H NMR and ^{13}C NMR were recorded with Bruker and Varian instrument at 600, 500, 400 and 150, 125, 100 MHz, respectively, and TMS was used as internal standard. Mass spectra were measured with a Thermo Finnigan LCQ-Advantage. High resolution mass spectra (HRMS) were measured on a Bruker micrOTOF-Q II instrument using ESI techniques. The structures of known compounds were further corroborated by comparing their ^1H NMR, ^{13}C NMR and MS data with those of literature.

4.2 General procedure for the synthesis of 2a

Thioanisole **1a** (1.0 mmol), 70% TBHP (5.0 mmol, 5.0 equiv.), and CHCl_3 (2 mL) were mixed in an 25 mL round bottom flask stirred at 60 °C for 1.5 h. After the reaction was completed (monitored by TLC), the reaction mixture was cooled to room temperature, added water (5 mL) and extracted with dichloromethane (5 mL $\times$ 3). The combined organic layers were dried over Na_2SO_4 , then were concentrated under reduced pressure. The crude residue was purified by flash chromatography on silica gel using hexane/EtOAc ($V:V=20:1$) as eluent to give the product **2a**, yield 98%.

(Methylsulfinyl)benzene (**2a**): 98% yield, yellow oil. ^1H NMR (500 MHz, CDCl_3) δ : 7.64~7.62 (m, 2H), 7.53~7.46 (m, 3H), 2.70 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 145.4, 130.9, 129.2, 123.4, 44.0; ESI-MS m/z : 303 $[\text{2M} + \text{Na}]^+$; HRMS (ESI) calcd for $\text{C}_7\text{H}_7\text{N}_2\text{OS}$ $[\text{M} + \text{H}]^+$ 141.0116, found 141.0117.

(Methylsulfonyl)benzene (**3a**): 2% yield. White solid, m.p. 84~85 °C (Lit.^[38] 85~85 °C); ^1H NMR (500 MHz,

CDCl_3) δ : 7.95~7.93 (m, 2H), 7.67~7.64 (m, 1H), 7.59~7.55 (m, 2H), 3.05 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 140.7, 133.8, 129.5, 127.5, 44.6; ESI-MS m/z : 157 $[\text{M} + \text{H}]^+$.

Methyl-4-(methylsulfinyl)benzene (**2b**):^[39] 97% yield. Yellow oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.52 (d, $J=8.4$ Hz, 2H), 7.31 (d, $J=8.4$ Hz, 2H), 2.70 (s, 3H), 2.41 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 142.2, 141.3, 129.8, 123.4, 44.0, 21.5; ESI-MS m/z : 331 $[\text{2M} + \text{Na}]^+$.

Methoxy-4-(methylsulfinyl)benzene (**2c**): 98% yield. White solid, m.p. 40~42 °C (Lit.^[40] 42~45 °C); ^1H NMR (500 MHz, CDCl_3) δ : 7.56 (d, $J=9.0$ Hz, 2H), 7.00 (d, $J=9.0$ Hz, 2H), 3.82 (s, 3H), 2.67 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 162.1, 136.6, 125.5, 114.9, 55.6, 44.0; ESI-MS m/z : 171 $[\text{M} + \text{H}]^+$.

Chloro-4-(methylsulfinyl)benzene (**2d**): 96% yield. White solid, m.p. 46~48 °C (Lit.^[40] 45~47 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.58 (d, $J=8.4$ Hz, 2H), 7.49 (d, $J=8.4$ Hz, 2H), 2.72 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 144.0, 137.1, 129.5, 124.9, 44.1; ESI-MS m/z : 175 $[\text{M} + \text{H}]^+$.

Bromo-4-(methylsulfinyl)benzene (**2e**): 95% yield. White solid, m.p. 76~81 °C (Lit.^[40] 78~79 °C); ^1H NMR (500 MHz, CDCl_3) δ : 7.66 (d, $J=8.5$ Hz, 2H), 7.51 (d, $J=8.5$ Hz, 2H), 2.71 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 145.0, 132.7, 125.57, 125.2, 44.1; ESI-MS m/z : 219 $[\text{M} + \text{H}]^+$.

1-(Ethylsulfinyl)-4-methylbenzene (**2f**):^[39] 93% yield, yellow oil. ^1H NMR (500 MHz, CDCl_3) δ : 7.47 (d, $J=8.5$ Hz, 2H), 7.29 (d, $J=8.0$ Hz, 2H), 2.85 (dq, $J=13.5$, 7.5 Hz, 1H), 2.75 (dq, $J=13.0$, 7.5 Hz, 1H), 2.39 (s, 3H), 1.16 (t, $J=7.5$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 141.5, 140.0, 129.9, 124.3, 50.4, 21.5, 6.1; ESI-MS m/z : 169 $[\text{M} + \text{H}]^+$.

Chloro-4-(ethylsulfinyl)benzene (**2g**):^[41] 92% yield, yellow oil. ^1H NMR (500 MHz, CDCl_3) δ : 7.55~7.52 (m, 2H), 7.50~7.48 (m, 2H), 2.89 (dq, $J=13.5$, 7.5 Hz, 1H), 2.76 (dq, $J=13.5$, 7.5 Hz, 1H), 1.18 (t, $J=7.5$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 141.9, 137.2, 129.5, 125.7, 50.4, 5.9; ESI-MS m/z : 189 $[\text{M} + \text{H}]^+$.

1-(Benzylsulfinyl)-4-methylbenzene (**2h**): 96% yield. White solid, m.p. 138~140 °C (Lit.^[42] 140~142 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.26~7.18 (m, 7H), 6.97 (d, $J=6.4$ Hz, 2H), 4.07 (d, $J=12.4$ Hz, 1H), 3.95 (d, $J=12.4$ Hz, 1H), 2.38 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 141.5, 139.5, 130.3, 129.5, 129.3, 128.4, 128.1, 124.4, 63.8, 21.7; ESI-MS m/z : 483 $[\text{2M} + \text{Na}]^+$.

1-(Benzylsulfinyl)-4-chlorobenzene (**2i**): 94% yield. White solid, m.p. 130~131 °C (Lit.^[42] 131~133 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.38~7.34 (m, 2H), 7.28~7.21 (m, 5H), 6.95 (dd, $J=7.2$, 1.6 Hz, 2H), 4.09 (d, $J=12.8$ Hz, 1H), 3.97 (d, $J=12.4$ Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ : 141.4, 137.5, 130.5, 129.2, 128.7, 128.7, 128.5, 126.0, 63.6; ESI-MS m/z : 523 $[\text{2M} + \text{Na}]^+$.

Sulfinyldibenzene (**2j**): 95% yield. White solid, m.p. 69~70 °C (Lit.^[43] 70~71 °C); ^1H NMR (500 MHz, CDCl_3) δ : 7.65~7.62 (m, 4H), 7.47~7.41 (m, 6H); ^{13}C

NMR (125 MHz, CDCl_3) δ : 145.7, 131.2, 129.4, 124.9; ESI-MS m/z : 203 $[\text{M}+\text{H}]^+$.

1-(Allylsulfinyl)-4-methylbenzene (**2k**):^[44] 93% yield, yellow oil. ^1H NMR (500 MHz, CDCl_3) δ : 7.47 (d, $J=8.0$ Hz, 2H), 7.29 (d, $J=8.5$ Hz, 2H), 5.66~5.58 (m, 1H), 5.32~5.29 (m, 1H), 5.17 (dq, $J=17.0, 1.5$ Hz, 1H), 3.55~3.46 (m, 2H), 2.40 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 141.6, 139.8, 129.8, 125.5, 124.4, 123.8, 61.0, 21.5; ESI-MS m/z : 383 $[\text{2M}+\text{Na}]^+$.

Methyl-4-(prop-2-yn-1-ylsulfinyl)benzene (**2l**):^[45] 87% yield, yellow oil. ^1H NMR (500 MHz, CDCl_3) δ : 7.58 (d, $J=8.0$ Hz, 2H), 7.31 (d, $J=8.0$ Hz, 2H), 3.60 (qd, $J=16.0, 2.7$ Hz, 1H), 2.40 (s, 3H), 2.32 (t, $J=3.0$ Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ : 142.4, 139.7, 129.9, 124.6, 76.4, 73.0, 47.9, 21.6; ESI-MS m/z : 179 $[\text{M}+\text{H}]^+$.

4-(Methylsulfinyl)benzaldehyde (**2m**): 67% yield. White solid, m.p. 83~84 °C (Lit.^[46] 85~86 °C); ^1H NMR (500 MHz, CDCl_3) δ : 10.05 (s, 1H), 8.01 (d, $J=8.0$ Hz, 2H), 7.79 (d, $J=8.0$ Hz, 2H), 2.76 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 191.2, 152.5, 138.2, 130.5, 124.2, 43.8; ESI-MS m/z : 169 $[\text{M}+\text{H}]^+$.

4,4'-Sulfinyldiphenol (**2n**): 85% yield. White solid, m.p. 190~192 °C (Lit.^[43] 193~194 °C); ^1H NMR (500 MHz, $\text{DMSO}-d_6$) δ : 10.08 (s, 1H), 7.42 (d, $J=8.5$ Hz, 2H), 6.87 (d, $J=8.5$ Hz, 2H); ^{13}C NMR (125 MHz, $\text{DMSO}-d_6$) δ : 159.9, 135.5, 126.5, 116.1; ESI-MS m/z : 235 $[\text{M}+\text{H}]^+$.

4.3 General procedure for the synthesis of **5a**

1,2-Diphenyldisulfane (**4a**) (1.0 mmol), 70% TBHP (5.0 mmol, 5.0 equiv.), and CHCl_3 (2 mL) were mixed in an 25 mL round bottom flask stirred at 60 °C for 3.5 h. After the reaction was completed (monitored by TLC), the reaction mixture was cooled to room temperature, added water (5 mL) and extracted with dichloromethane (5 mL $\times$ 3). The combined organic layers were dried over Na_2SO_4 , then were concentrated under reduced pressure. The crude residue was purified by flash chromatography on silica gel using hexane/EtOAc ($V:V=20:1$) as eluent to give the product **5a**, 89% yield.

S-Phenyl benzenesulfonothioate (**5a**): 89% yield. White solid, m.p. 36~37 °C (Lit.^[47] 37~39 °C); ^1H NMR (500 MHz, CDCl_3) δ : 7.59~7.54 (m, 3H), 7.49~7.29 (m, 7H); ^{13}C NMR (100 MHz, CDCl_3) δ : 142.9, 136.5, 133.6, 131.4, 129.4, 128.8, 127.8, 127.5; HRMS(ESI) calcd for $\text{C}_{12}\text{H}_{10}\text{NaO}_2\text{S}_2$ $[\text{M}+\text{Na}]^+$ 273.0014, found 273.0014.

S-*p*-Tolyl 4-methylbenzenesulfonothioate (**5b**): 90% yield. White solid, m.p. 70~72 °C (Lit.^[47] 74~75 °C); ^1H NMR (500 MHz, CDCl_3) δ : 7.46 (d, $J=8.0$ Hz, 2H), 7.27~7.21 (m, 4H), 7.15 (d, $J=8.0$ Hz, 2H), 2.44 (s, 3H), 2.39 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 144.7, 142.2, 140.6, 136.6, 130.3, 129.5, 127.7, 124.7, 21.7, 21.6; ESI-MS m/z : 279 $[\text{M}+\text{H}]^+$.

S-(4-Methoxyphenyl) 4-methoxybenzenesulfonothioate (**5c**): 90% yield. White solid, m.p. 84~86 °C (Lit.^[47] 88~90 °C); ^1H NMR (500 MHz, CDCl_3) δ : 7.50 (d, $J=9.0$ Hz, 2H), 7.27 (d, $J=9.0$ Hz, 2H), 6.88 (d, $J=9.0$ Hz, 2H), 6.85 (d, $J=9.0$ Hz, 2H), 3.87 (s, 3H), 3.84 (s, 3H);

^{13}C NMR (125 MHz, CDCl_3) δ : 163.7, 162.3, 138.5, 135.0, 130.0, 119.0, 115.0, 114.0, 55.8, 55.6; ESI-MS m/z : 311 $[\text{M}+\text{H}]^+$.

S-(2,4-Dimethylphenyl) 2,4-dimethylbenzenesulfonothioate (**5d**): 85% yield. Yellow solid, m.p. 59~61 °C; ^1H NMR (500 MHz, CDCl_3) δ : 7.30 (d, $J=8.0$ Hz, 1H), 7.11 (d, $J=8.0$ Hz, 2H), 7.02 (s, 1H), 6.94~6.89 (m, 2H), 2.64 (s, 3H), 2.37 (s, 3H), 2.31 (s, 3H), 2.12 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 144.7, 144.0, 142.4, 138.7, 138.4, 137.7, 133.6, 131.8, 130.3, 127.7, 126.6, 124.0, 21.5, 21.5, 20.6, 20.5; IR (KBr) ν : 1590, 1488, 1323, 1180, 1142 cm^{-1} ; HRMS(ESI) calcd for $\text{C}_{16}\text{H}_{18}\text{NaO}_2\text{S}_2$ $[\text{M}+\text{Na}]^+$ 329.0640, found 329.0656.

S-(4-Chlorophenyl) 4-chlorobenzenesulfonothioate (**5e**): 87% yield. White solid, m.p. 131~134 °C (Lit.^[47] 135~136 °C); ^1H NMR (500 MHz, CDCl_3) δ : 7.53~7.50 (m, 2H), 7.44~7.41 (m, 2H), 7.36~7.34 (m, 2H), 7.32~7.30 (m, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ : 141.4, 140.7, 138.7, 137.8, 130.1, 129.4, 129.1, 126.2; HRMS(ESI) calcd for $\text{C}_{12}\text{H}_8\text{Cl}_2\text{NaO}_2\text{S}_2$ $[\text{M}+\text{Na}]^+$ 340.9235, found 340.9222.

S-(4-Bromophenyl) 4-bromobenzenesulfonothioate (**5f**): 73% yield. White solid, m.p. 148~152 °C (Lit.^[47] 148~149 °C); ^1H NMR (500 MHz, CDCl_3) δ : 7.61~7.58 (m, 2H), 7.53~7.50 (m, 2H), 7.45~7.42 (m, 2H), 7.25~7.22 (m, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ : 142.0, 138.0, 133.1, 132.4, 129.3, 129.1, 127.2, 126.8; ESI-MS m/z : 219.

S-Benzyl phenylmethanesulfonothioate (**5g**): 22% yield. White solid, m.p. 105~108 °C (Lit.^[48] 106~108 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.41~7.33 (m, 6H), 7.32~7.27 (m, 4H), 4.22 (s, 2H), 4.04 (s, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 135.0, 131.5, 129.5, 129.1, 128.9, 128.5, 127.9, 69.2, 41.1.

S-(4-Bromophenyl) 4-methylbenzenesulfonothioate (**5h**): 18% yield. White solid, m.p. 103~106 °C (Lit.^[49] 107 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.61~7.56 (m, 3H), 7.43 (t, $J=8.0$ Hz, 3H), 7.17 (d, $J=8.0$ Hz, 2H), 2.39 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 138.0, 136.6, 133.1, 132.5, 132.2, 130.6, 129.2, 129.1, 21.6; HRMS(ESI) calcd for $\text{C}_{13}\text{H}_{11}\text{BrNaO}_2\text{S}_2$ $[\text{M}+\text{Na}]^+$ 364.9276, found 364.9259.

Supporting Information ^1H NMR and ^{13}C NMR spectra copies of the products **2a~2n**, **3a** and **5a~5h**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Carreno, M. C. *Chem. Rev.* **1995**, *95*, 1717.
(b) Fernández, I.; Khair, N. *Chem. Rev.* **2003**, *103*, 3651.
(c) Hartz, R. A.; Arvanitis, A. G.; Arnold, C.; Rescinito, J. P.; Hung, K. L.; Zhang, G.; Wong, H.; Langley, D. R.; Gilligan, P. J.; Trainor, G. L. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 934.
(d) Weidner, J. P.; Block, S. S. *J. Med. Chem.* **1964**, *7*, 671.
(e) Block, S. S.; Weidner, J. P. *Dev. Ind. Microbiol.* **1963**, *4*, 213.
- [2] Maryam, E.; Majid, H.; Hossein, O.; Lida, F. *Iran. J. Chem. Chem. Eng.* **2017**, *36*, 21.
- [3] Lindberg, P.; Weidolf, L. *US 5877192*, **1998**.

- [4] (a) Hinsberger, S.; Hüsecken, K.; Groh, M.; Negri, M.; Hauptenthal, J.; Hartmann, R. W. *J. Med. Chem.* **2013**, *56*, 8332.
 (b) Tan, D.; Štrukil, V.; Mottillo C.; Frišić, T. *Chem. Commun.* **2014**, *50*, 5248.
 (c) Prasit, P.; Wang, Z.; Brideau, C.; Chan, C.-C.; Charleson, S.; Cromlish, W.; Ethier, D.; Evans, J. F.; Ford Hutchinson, A. W.; Gauthier, J. Y.; Gordon, R.; Guay, J.; Gresser, M.; Kargman, S.; Kennedy, B.; Leblanc, Y.; Leger, S.; Mancini, J.; O'Neill, G. P.; Ouellet, M.; Percival, M. D.; Perrier, H.; Riendeau, D.; Rodger, I.; Tagari, P.; Thérien, M.; Vickers, P.; Wong, E.; Xu, L.-J.; Young, R. N.; Zamboni, R.; Boyce, S.; Rupniak, N.; Forrest, M.; Visco, D.; Patrick, D. *Bioorg. Med. Chem. Lett.* **1999**, *9*, 1773.
 (d) Liu, F.; Luo, X. Q.; Song, B. A.; Bhadury, P. S.; Yang, S.; Jin, L. H.; Xue, W.; Hu, D. Y. *Bioorg. Med. Chem.* **2008**, *16*, 3632.
 (e) Seto, M.; Miyamoto, N.; Aikawa, K.; Aramaki, Y.; Kanzaki, N.; Iizawa, Y.; Baba, M.; Shiraishi, M. *Bioorg. Med. Chem.* **2005**, *13*, 363.
 (f) Santelli-rouvier, C.; Barrer, J. M.; Farrell, C. M.; Sharples, D.; Thill, B.; Barbe, J. *Eur. J. Med. Chem.* **2004**, *39*, 1029.
 (g) Zhao, Q. Q.; Li, Y. Q.; Xiong, L. X.; Wang, Q. M. *J. Agric. Food Chem.* **2010**, *58*, 4992.
- [5] (a) Mellah, M.; Voituriez, A.; Schulz, E. *Chem. Rev.* **2007**, *107*, 5133.
 (b) Sipos, G.; Drinkel, E. E.; Dorta, R. *Chem. Soc. Rev.* **2015**, *44*, 3834.
 (c) Jia, T.; Cao, P.; Wang, B.; Lou, Y.; Yin, X.; Wang M.; Liao, J. J. *Am. Chem. Soc.* **2015**, *137*, 13760.
 (d) Prilezhaeva, E. N. *Russ. Chem. Rev.* **2001**, *70*, 897.
 (e) Fernández I.; Khair, N. *Chem. Rev.* **2003**, *103*, 3651.
 (f) Burr S. K.; Padwa, A. *Chem. Rev.* **2004**, *104*, 2401.
 (g) Nguyen, M. H.; Imanishi, M.; Kurogi T.; Smith III, A. B. *J. Am. Chem. Soc.* **2016**, *138*, 3675.
- [6] (a) Furukawa, N.; Akutagawa, K.; Yoshiaki, T.; Oae, S. *Tetrahedron Lett.* **1981**, *22*, 2989.
 (b) Bentley, H. R.; McDermott, E. E.; Whitehead, J. K. *Nature* **1950**, *165*, 735.
 (c) Wang, Y. Y.; Hong, X. C.; Deng, Z. X. *Chin. J. Org. Chem.* **2012**, *32*, 825 (in Chinese).
 (王娅娅, 洪学传, 邓子新, 有机化学, **2012**, *32*, 825.)
 (d) Siemeister, G.; Lucking, U.; Wengner, A. M.; Lienau, P.; Steinke, W.; Schatz, C.; Mumberg, D.; Ziegelbauer, K. *Mol. Cancer Ther.* **2012**, *11*, 2265.
- [7] (a) Cross, P. E.; Arrowsmith, J. E.; Thomas, G. N.; Gwilt, M.; Burges, R. A.; Higgins, A. J. *J. Med. Chem.* **1990**, *33*, 1151.
 (b) Murai, S.; Haga, T.; Fujikawa, K.; Sakashita, N.; Shiga, K. *Symposium Series* **1991**, *443*, 98.
 (c) Tumula, V. R.; Bondwal, S.; Bisht, P.; Pendem, C.; Kumar, J. *React. Kinet., Mech. Catal.* **2012**, *107*, 449.
- [8] (a) Legros, J.; Dehli, J. R.; Bolm, C. *Adv. Synth. Catal.* **2005**, *347*, 19.
 (b) Fernandez, I.; Khair, N. *Chem. Rev.* **2003**, *103*, 3651.
 (c) Kobayashi, C.; Ogawa, C.; Konishi, H.; Sugiura, M. *J. Am. Chem. Soc.* **2003**, *125*, 6610.
 (d) Wojaczynska, E.; Wojaczynski, J. *Chem. Rev.* **2010**, *110*, 4303.
- [9] (a) Gallardo-Godoy, A.; Torres-Altoro, M. I.; White, K. J.; Barker, E. L.; Nichols, D. E. *Bioorg. Med. Chem.* **2007**, *15*, 305.
 (b) Sugata, K.; Song, L.; Nakamura, M.; Ueki, S.; Fajer, P. G.; Arata, T. *J. Mol. Biol.* **2009**, *386*, 626.
 (c) Zefirov, N. S.; Zyk, N. V.; Beloglazkina, E. K.; Kutateladze, A. *G. Sulfur Rep.* **1993**, *14*, 223.
- [10] Manesh, A. A.; Eshbala, F. H.; Hemmati, S.; Veisi, H. *RSC Adv.* **2015**, *5*, 70265.
- [11] Gasparini, F.; Giovannoli, M.; Maresca, L.; Natile, G.; Palmieri, G. *Synth. Commun.* **1984**, *14*, 1111.
- [12] (a) Scarso, A.; Strukul, G. *Adv. Synth. Catal.* **2005**, *347*, 1227.
 (b) Legros, J.; Bolm, C. *Chem.-Eur. J.* **2005**, *11*, 1086.
 (c) Haddadi, H.; Hafshejani, S. M.; Farsani, M. R. *Catal. Lett.* **2015**, *145*, 1984.
 (d) Bahrami, K.; Khodaei, M. M.; Khaledian, D. *Tetrahedron Lett.* **2012**, *53*, 354.
- [13] (a) Zhao, L. L.; Zhang, H. Y.; Wang, Y. H. *J. Org. Chem.* **2016**, *81*, 129.
 (b) Patnam, P. L.; Bhatt, M.; Singh, R.; Sandeep Saranb, S.; Jain, S. L. *RSC Adv.* **2016**, *6*, 60888.
 (c) Zaburdaeva, E. A.; Dodonov, V. A. *Russ. Chem. Bull., Int. Ed.* **2011**, *60*, 185.
- [14] (a) Chen, T. H.; Kwong, K. W.; Lee, N. F.; Ranburger, D.; Zhang, R. *Inorg. Chim. Acta* **2016**, *451*, 65.
 (b) Xie, Y. T.; Yuxin Li, Y. X.; Zhou, S.; Zhou, S.; Zhang, Y.; Chen, M. G.; Li, Z. M. *Synlett* **2018**, *29*, 340.
 (c) Dong, D. Q.; Chen, W. J.; Chen, D. M.; Li, L. X.; Li, G. H.; Wang, Z. L.; Deng, Q.; Long, S. *Chin. J. Org. Chem.* **2019**, *39*, 3190 (in Chinese).
 (董道青, 陈文静, 陈德茂, 李丽霞, 李光辉, 王祖利, 邓企, 龙妹, 有机化学, **2019**, *39*, 3190.)
- [15] Cai, M. T.; Lv, G. S.; Chen, J. X.; Gao, W. X.; Ding, J. C.; Wu, H. *Y. Chem. Lett.* **2010**, *39*, 368.
- [16] Lakouraj, M. M.; Tajbakhsh, M.; Shirini, F.; Tamami, A. *Synth. Commun.* **2005**, *35*, 775.
- [17] Sobhani, S.; Aryanejad, S.; Maleki, M. F. *Synlett* **2011**, 319.
- [18] Luu, T. X. T.; Nguyen, T. T.; Leb, T. N.; Spanget-Larsena, J.; Duus, F. *J. Sulfur Chem.* **2015**, *36*, 340.
- [19] Iranpoor, N.; Mohajer, D.; Rezaeifard, A. R. *Tetrahedron Lett.* **2004**, *45*, 3811.
- [20] (a) Bulman, P. C.; Buckley, B. R.; Elliott, C.; Chan, Y.; Dreyfus, N.; Marken, F. *Synlett* **2016**, *27*, 80.
 (b) Yue, H. L.; Bao, P. L.; Wang, L. L.; Lv, X. X.; Yang, D. S.; Wang, H.; Wei, W. *Chin. J. Org. Chem.* **2019**, *39*, 463 (in Chinese).
 (岳会兰, 鲍鹏丽, 王雷雷, 吕晓霞, 杨道山, 王桦, 魏伟, 有机化学, **2019**, *39*, 463.)
- [21] Zabardasti, A.; Shangaie, S. A. *J. Iran. Chem. Soc.* **2019**, *16*, 57.
- [22] (a) Silva, F.; Baker, A.; Stansall, J.; Michalska, W.; Yusubov, M. S.; Graz, M.; Saunders, R.; Evans, S. J. S.; Wirth, T. *Eur. J. Org. Chem.* **2018**, 2134.
 (b) Natarajan, P. *Tetrahedron Lett.* **2015**, *56*, 4131.
- [23] (a) Fareghi-Alamdari, R.; Zekri, N.; Moghadam, A. J.; Farsani, M. R. *Catal. Commun.* **2017**, *98*, 71.
 (b) Bayat, Y.; Shirini, F.; Goli-Jolodar, O. *J. Mol. Liq.* **2018**, *265*, 517.
 (c) Manesh, A. A.; Eshbala, F. H.; Hemmati, S.; Veisi, H. *RSC Adv.* **2015**, *5*, 70265.
 (d) Zhang, H.; Qi, L. L. *Tetrahedron Lett.* **2018**, *59*, 3171.
- [24] Gogoi, S. R.; Boruah, J. J.; Sengupta, G.; Saikia, G.; Ahmed, K.; Bania, K. K.; Islam, N. S. *Catal. Sci. Technol.* **2015**, *5*, 595.
- [25] Gan, S. Y.; Yin, J. J.; Yao, Y.; Liu, Y.; Chang, D. H.; Dan Zhua, D.; Shi, L. *Org. Biomol. Chem.* **2017**, *15*, 2647.
- [26] (a) Rostamnia, S.; Doustkhah, E.; Bahrami, K.; Amini, S. *J. Mol. Liq.* **2015**, *207*, 334.
 (b) Voutyritsa, E.; Triandafillidi, I.; Kokotos, C. G. *Synthesis* **2017**, *49*, 917.
 (c) Aghajania, M.; Safaei, E.; Karimi, B. *Synth. Met.* **2017**, *233*, 63.
 (d) Cravotto, G.; Garella, D.; Carnaroglio, D.; Gaudino, E. C.; Rosati, O. *Chem. Commun.* **2012**, *48*, 11632.
 (e) Liang, G. G.; Liu, M. C.; Chen, J. X.; Ding, J. C.; Gao, W. X.; Wu, H. Y. *Chin. J. Chem.* **2012**, *30*, 1611.
 (f) Peng, Z. H.; Zheng, X.; Zhang, Y. J.; An, D.; Dong, W. R. *Green Chem.* **2018**, *20*, 1760.
 (g) Billard, T.; Langlois, B. R. *J. Fluorine Chem.* **1997**, *84*, 63.
 (h) Secci, F.; Arca, M.; Frongia, A.; Piras, P. P. *Catal. Sci. Technol.* **2014**, *4*, 1407.
 (i) Iranpoor, N.; Firouzabadi, H.; Pourali, A. R. *Synlett* **2004**, 347.
 (j) Peng, S.; Song, Y. X.; He, J. Y.; Tang, S. S.; Tan, J. X.; Cao, Z.; Lin, Y. W.; He, W. M. *Chin. Chem. Lett.* **2019**, *30*, 2287.
 (k) Liu, K. J.; Deng, J. H.; Yang, J.; Gong, S. F.; Lin, Y. W.; He, J. Y.; Cao, Z.; He, W. M. *Green Chem.* **2020**, *22*, 433.
 (l) Liu, K. J.; Zeng, T. Z.; Zeng, J. L.; Gong, S. F.; He, J. Y.; Lin, Y.

- W.; Tan, J. X.; Cao, Z.; He, W. M. *Chin. Chem. Lett.* **2019**, *30*, 2304.
- [27] (a) Rostami, A.; Saedmocheshi, N.; Shirvandi, Z. *C.R. Chim.* **2018**, *21*, 835.
(b) Zhang, P. F.; Wang, Y.; Li, H. R.; Antonietti, M. *Green Chem.* **2012**, *14*, 1904.
- [28] Wang, F.; Liu, C.; Liu, G.; Li, W. X.; Liu, J. H. *Catal. Commun.* **2015**, *72*, 142.
- [29] (a) Veisi, H.; Sajjadifar, S.; Biabri, P. M.; Hemmati, S. *Polyhedron* **2018**, *153*, 240.
(b) Otokesh, S.; Kolvari, E.; Amoozadeh, A.; Koukabi, N. *RSC Adv.* **2015**, *5*, 53749.
- [30] (a) Jiang, J.; Luo, R. C.; Zhou, X. T.; Chen, Y. J.; Jia, H. B. *Adv. Synth. Catal.* **2018**, *360*, 4402.
(b) Hao, H. M.; Wang, Z.; Shi, J. L.; Li, X.; Lang, X. J. *ChemCatChem* **2018**, *10*, 4545.
(c) Xie, L. Y.; Chen, Y. L.; Qin, L.; Wen, Y.; Xie, J. W.; Tan, J. X.; Huang, Y.; Cao, Z.; He, W. M. *Org. Chem. Front.* **2019**, *6*, 3950.
(d) Xie, L. Y.; Hu, J. L.; Song, Y. X.; Jia, G. K.; Lin, Y. W.; He, J. H.; Cao, Z.; He, W. M. *ACS Sustainable Chem. Eng.* **2019**, *7*, 19993.
- [31] (a) Patnam, P. L.; Bhatt, M.; Singh, R.; Saran, S.; Jain, S. *RSC Adv.* **2016**, *6*, 60888.
(b) Li, B. D.; He, P.; Yi, G. Q.; Lin, H. Q.; Yuan, Y. Z. *Catal. Lett.* **2009**, *133*, 33.
(c) Li, Y. J.; Yan, N.; Liu, C. H.; Yu, Y.; Zhao, Y. L. *Org. Lett.* **2017**, *19*, 1160.
- [32] Bock, L.; Miller, G. H.; Schaper, K. J.; Seydel, J. K. *J. Med. Chem.* **1974**, *17*, 23.
- [33] Hinsberger, S.; Hüsecken, K.; Groh, M.; Negri, M.; Haupenthal, J.; Hartmann, R. W. *J. Med. Chem.* **2013**, *56*, 8332.
- [34] Tan, D.; Štrukil, V.; Mottillo C.; Friščić, T. *Chem. Commun.* **2014**, *50*, 5248.
- [35] Prasit, P.; Wang, Z.; Brideau, C.; Chan, C.-C.; Charleson, S.; Cromlish, W.; Ethier, D.; Evans, J. F.; Ford Hutchinson, A. W.; Gauthier, J. Y.; Gordon, R.; Guay, J.; Gresser, M.; Kargman, S.; Kennedy, B.; Leblanc, Y.; Leger, S.; Mancini, J.; O'Neill, G. P.; Ouellet, M.; Percival, M. D.; Perrier, H.; Riendeau, D.; Rodger, I.; Tagari, P.; Thérien, M.; Vickers, P.; Wong, E.; Xu, L.-J.; Young, R. N.; Zamboni, R.; Boyce, S.; Rupniak, N.; Forrest, M.; Visco, D.; Patrick, D. *Bioorg. Med. Chem. Lett.* **1999**, *9*, 1773.
- [36] Chau, M. M.; Kice, J. L. *J. Am. Chem. Soc.* **1976**, *98*, 7711.
- [37] (a) Iranpoor, N.; Firouzabadi, H.; Pourali, A. R. *Tetrahedron* **2002**, *58*, 5179.
(b) Natarajan, P. *Tetrahedron Lett.* **2015**, *56*, 4131.
- [38] Ma, B. C.; Zhao, W.; Zhang, Y. S.; Wu, S. Y.; Ding, Y. *RSC Adv.* **2014**, *4*, 30254.
- [39] Caupene, C.; Boudou, C.; Perrio, S.; Metzner, P. *J. Org. Chem.* **2005**, *70*, 2812.
- [40] Hanson, P.; Ramon, A. A.; Hendrick, J.; Smith, R. L. *Org. Biomol. Chem.* **2008**, *6*, 745.
- [41] Blakemore, P. R.; Burge, M. S.; Sephton, M. A. *Tetrahedron Lett.* **2007**, *48*, 3999.
- [42] Singh, S.; Donnell, J. S.; Schwan, A. L. *Org. Biomol. Chem.* **2010**, *8*, 1712.
- [43] Togo, H.; Nabana, T.; Yamaguchi, K. *J. Org. Chem.* **2000**, *65*, 8391.
- [44] Maitro, G.; Prestat, G.; Madec, D.; Poli, G. *J. Org. Chem.* **2006**, *71*, 7449.
- [45] Zhao, X.; Zheng, X. C.; Yang, B.; Sheng, J. Q.; Lu, K. *Org. Biomol. Chem.* **2018**, *16*, 1200.
- [46] Lowe, J. N.; Fulton, D. A.; Chiu, S. H.; Elizarow, A. M.; Cantrill, S. J.; Rowan, S. J.; Stoddart, F. J. *J. Org. Chem.* **2004**, *69*, 4390.
- [47] Zheng, Z.; Qing, F. L.; Huang, Y. G.; Xu, X. H. *Adv. Synth. Catal.* **2016**, *358*, 3477.
- [48] Wakui, T.; Nakamura, Y.; Motoki, S. *Bull. Chem. Soc. Jpn.* **1978**, *51*, 3081.
- [49] Loudon, J. D.; Livingston, A. J. *Chem. Soc.* **1935**, 896.

(Cheng, F.)