可见光促进 *N*-苯丙炔胺与磺酰肼的环化: 在无碱和催化剂下快速合成 3-磺酰基喹啉衍生物

彭美^a 郑扬帆^a 黄浩^b 叶佳^a 邓兴国^{*,c} 何纯莲^{*,a}

(^a 湖南师范大学医学院 小分子靶向药物研究与创制湖南省重点实验室 长沙 410081)

(^b 宜春学院化学与生物工程学院 江西宜春 336000)

(^c 湖南食品药品职业学院 长沙 410081)

摘要 利用可见光照射和过氧化叔丁醇作氧化剂的条件, 发展了 *N*-芳基苯丙炔胺和磺酰肼类化合物的自由基环化-氧化芳构化反应. 该反应提供了一种合成 3-磺酰基喹啉衍生物的有效方法, 显示出较好的官能团适用性、底物普适性以及很好的化学与区位选择性.

关键词 可见光; 磺酰肼; 3-磺酰基喹啉衍生物; 区位选择性

Visible-Light-Induced Cycloaddition Involving *N*-Propargylanilines with Arylsulfonylhydrazides: Rapid Access to 3-Sulfonated Quinoline Derivatives without Base and Catalyst

Peng, Mei^a Zheng, Yangfan^a Huang, Hao^b Ye, Jia^a

Deng, Xingguo^{*,c} He, Chunlian^{*,a}

(^a Key Laboratory of Study and Discovery of Small Targeted Molecules of Hunan Province, School of Medicine, Hunan Normal University, Changsha 410081)

(^b School of Chemistry and Bioengineering, Yichun University, Yichun, Jiangxi 336000)

(^c Hunan Food and Drug Vocational College, Changsha 410081)

Abstract A visible-light-induced oxidative cyclization of *N*-propargylanilines with arylsulfonylhydrazides was developed using *tert*-butyl hydroperoxide as oxidant. This transformation offers a straightforward route to 3-sulfonated quinoline derivatives with good functional group tolerance, good to excellent yields and high regio-selectivity.

Keywords visible-light; arylsulfonylhydrazides; 3-sulfonated quinoline derivatives; regio-selectivity

1 Introduction

Recently, substituted quinoline derivatives have aroused the interest and research concern of many researchers, mainly due to their fine biological activities and broad applications in medicinal chemistry.^[1] Among them, 3-substituted quinolones is one of the most important quinoline derivatives and can be found in anticancer, anti-depressant, and anti-psychotic molecules.^[2] The synthesis

of 3-substituted quinolines has been widely reported.^[3] *N*-Propargylaniline derivatives can be used for synthesizing many significant nitrogen-containing heterocyclic compounds, including 3-substituted quinolines.^[4] In recent years, visible light as a clean and renewable chemical energy has been widely utilized to promote various organic synthetic transformations.^[5] *N*-Propargylamines have been proven to be efficient radical acceptors under visible light irradiation. In 2018, Zhang *et al.*^[4b] developed an efficient

* Corresponding authors. E-mail: chunlianhe6688@163.com; 734411927@qq.com

Received February 6, 2020; revised March 25, 2020; published online April 13, 2020.

Project supported by the National Natural Science Foundation of China (Nos. 21176063, 81803720), the Hunan Provincial Natural Science Foundation (Nos. 18JJ5018, 2019JJ50383), and the Department of Science and Technology Foundation of Changsha City (Nos. kq1706047, kq1801038), and the Key Laboratory of Study and Discovery of Small Targeted Molecules of Hunan Province (No. 2019CG06).

国家自然科学基金(Nos. 21176063, 81803720)、湖南省自然科学基金(Nos. 18JJ5018, 2019JJ50383)、小分子靶向药物研究与创制湖南省重点实验室开放基金(No. 2019CG06)资助项目.

eosin Y-catalyzed three component sulfonylation of *N*-propargylanilines with DABSO (1,4-diazabicyclo-[2.2.2]octane bis(sulfur dioxide) adduct), diaryliodonium salts under green LEDs (light emitting diode, Scheme 1a). Very recently, Li's group^[4a] reported a visible-light-induced oxidative cyclization of *N*-propargyl anilines with sulfinic acids using pyridine as base (Scheme 1b). Although a few synthetic methods have been developed, stoichiometric oxidants, relatively high temperatures, additional photocatalysts or additional base would limit their applications in synthetic chemistry.

Scheme 1 Visible-light-induced methods for the synthesis of 3-sulfonated quinoline derivatives from *N*-propargylanilines

Sulfonylhydrazide is a cheap and stable sulfonyl source.^[6] It can be easily transformed into sulfonyl radicals by using of *tert*-butyl hydroperoxide (TBHP) and give a new route to produce S-containing organic compounds.^[7] It is still an attractive but challenging task to explore a more convenient and mild method to access C₃-sulfonated quinolines from more common and inexpensive starting materials.

2 Results and discussion

We began our research with the reaction of *N*-(3-phenylprop-2-yn-1-yl)aniline (**1a**) and 4-methylbenzenesulfonylhydrazide (**2a**) in 1,2-dichloroethane (DCE) by using TBHP as the oxidant under LED (435~440 nm, 7 W) irradiation in open air for 8 h without any photocatalyst and the desired product 4-phenyl-3-tosylquinoline (**3a**) was obtained in 89% isolated yield (Table 1, Entry 1). Inspired

by this result, a series of oxidants were further researched (Table 1, Entries 2~4), and TBHP was proved to be the best oxidant. The use of the other solvents resulted in lower yields (Table 1, Entries 5~10). The light sources were then studied. However, LED (455~460 nm), LED (565~570 nm), LED (390~395 nm) and white light could not improve the reaction yield (Table 1, Entries 11~14). Subsequently, the reaction atmospheres were screened (Table 1, Entries 15, 16), such as N₂ and O₂, but the yields were not improved. Finally, the reaction temperature was increased to 40 °C, and the corresponding product **3a** was isolated in 86% yield (Table 1, Entry 17). No product **3a** was generated when the reaction underwent in dark (Table 1, Entry 18). Based on all of the above results, the primary conditions were still selected as the optimized reaction conditions, namely, treatment of compound **1a** with 1.75 equiv. of **2a**, 2.5 equiv. of TBHP in DCE (0.25 mol·L⁻¹) at room temperature under blue LEDs (435~437 nm, 7 W) irradiation in open air for 8 h.

Table 1 Optimization of the reaction conditions^a

Entry	Variations from standard conditions	Yield ^b /%
1	—	90
2	DTBP instead of TBHP	Trace
3	H ₂ O ₂ instead of TBHP	Trace
4	BPO instead of TBHP	20
5	H ₂ O instead of DCE	35
6	Dioxane instead of DCE	32
7	Dichloromethane instead of DCE	50
8	CH ₃ CN instead of DCE	10
9	CH ₂ Br ₂ instead of DCE	55
10	Dimethylformamide instead of DCE	Trace
11	455~460 nm instead of 435~440 nm	65
12	565~570 nm instead of 435~440 nm	63
13	390~395 nm instead of 435~440 nm	28
14	White LED instead of 435~440 nm	30
15	N ₂ instead of air	87
16	O ₂ instead of air	85
17	40 °C instead of room temperature	86
18	In dark	Trace

^a Reaction conditions: **1a** (0.2 mmol), **2a** (0.35 mmol), TBHP (dry, 2.5 equiv.) in DCE (2 mL) and at room temperature under LEDs irradiation in open air for 8 h. ^b Yield of isolated product. DTBP = Di-*t*-butyl peroxide; TBHP = *tert*-Butyl hydroperoxide solution; BPO = Dibenzoyl peroxide.

Having the optimized reaction conditions in hand, the reactions of 4-methylbenzenesulfonylhydrazide with a variety of *N*-propargylanilines were investigated (Table 2). A series of *N*-propargylanilines (**1**) were successfully reacted with 4-methylbenzenesulfonylhydrazide (**2a**) to generate the corresponding 3-sulfonated quinoline derivatives in good to excellent yields. First, the substitution pattern on the aromatic ring of the aniline group was investi-

gated. Electron-donating groups on the aromatic ring provided the desired products in excellent yields (**3b**~**3e**). Electron-withdrawing groups on the benzene ring performed well in this transformation and smoothly produced the corresponding products in good yields (**3f**~**3j**). It is noteworthy that *N*-(3-phenylprop-2-yn-1-yl)-[1,1'-biphenyl]-2-amine performed well in this reaction, producing the product **3k** in 72% yield. Subsequently, the influence of the substituents on the phenyl ring of the aromatic alkyne moiety was also explored. Substrates bearing electron-donating (OMe) or electron-withdrawing (Cl, F, and Ac) groups on the benzene ring that is directly bound to the triple bond all provided the corresponding products in satisfactory yields. It suggested that the substituted groups did not have a transformation influence on the reaction (**3l**~**3o**).

Subsequently, the scope of sulfonyl hydrazides was also examined under the optimized reaction conditions. As shown in Table 3, various sulfonyl hydrazides including substituted groups (R=H, OMe, *t*-Bu, OCF₃, F, Cl, Br, CO₂Et and Ph) at the *para*-position of the aromatic ring reacted with **1a** to give the corresponding products (**4a**~**4i**) in 76%~87% yields. Additionally, 3-methylbenzene-sulfonylhydrazide and 2-methylbenzenesulfonylhydrazide were also effective in producing the desired 4-phenyl-3-(*m*-tolylsulfonyl)quinoline (**4j**) in 76% yield and 4-phenyl-3-(*o*-tolylsulfonyl)quinoline (**4k**) in 78% yield. Notably, naphthalene-2-sulfonylhydrazide could also perform well in this reaction, producing the product **4l** in 76% yield. Moreover, thiophene-2-sulfonylhydrazide was also evaluated, and the product **4m** was obtained in a satisfactory yield. Surprisingly, aliphatic sulfonyl hydrazides were also successfully employed under the standard conditions. Methanesulfonylhydrazide and cyclopropane-sulfonylhydrazide smoothly underwent the reaction to produce the corresponding 3-sulfonated quinolines (**4n** and **4o**) which could not be synthesized in the reported literature.^[4a-4b]

Control reactions between *N*-(3-phenylprop-2-yn-1-yl)aniline (**1a**) and 4-methylbenzenesulfonylhydrazide (**2a**) were conducted to get better understanding of the mechanistic pathway for the reaction (Scheme 2). When 2 equiv. of TEMPO was added to the reaction, the transformation was completely restrained suggesting that the reaction might proceed via radical processes (Scheme 2a).

Scheme 2 Experiments for the mechanistic study

Table 2 Scope of *N*-propargylanilines^{a,b}

^a Reaction conditions: **1** (0.2 mmol), **2a** (0.35 mmol), TBHP (dry, 2.5 equiv.) in DCE (2 mL) and at room temperature under blue LEDs (435~440 nm, 7 W) irradiation in open air for 8 h. ^b Yield of isolated product.

In the absence of 4-methylbenzenesulfonylhydrazide, no 4-phenylquinoline was observed, suggesting that the addition to the carbon-carbon triple bond might be prior to the cyclization of *N*-propargylaniline (Scheme 2b). Finally, the reaction of TsNHNH₂ with *N*-(3-phenylprop-2-yn-1-ylidene)-aniline could not produce **3a** under the optimized conditions (Scheme 2c), thus illustrating that *N*-(3-phenyl-

Table 3 Scope of arylsulfonylhydrazides^{a,b}

^a Reaction conditions: **1a** (0.2 mmol), **2** (0.35 mmol), TBHP (dry, 2.5 equiv.) in DCE (2 mL) and at room temperature under blue LEDs (435~440 nm, 7 W) irradiation in open air for 8 h. ^b Yield of isolated product.

prop-2-yn-1-ylidene)aniline might not be an intermediate in this transformation.

On the basis of the above results and previous reports,^[4] a plausible mechanism is proposed (Scheme 3). Firstly, sulfonyl radical is generated under visible-light irradiation and TBHP. Then, sulfonyl radical goes through intermolecular addition onto *N*-(3-phenylprop-2-yn-1-yl)-aniline to generate alkenyl radical **A**. Followed by the intramolecular attack of radical **A** on the pendant benzene ring subsequently produces intermediate **B**. Oxidation of **B** generates the corresponding cyclohexadienyl cation **C**, which undergoes deprotonation to provide 4-phenyl-3-tosyl-1,2-dihydroquinoline (**D**). In the end, intermediate **D** ultimately aromatizes to afford the final product **3a**.

3 Conclusions

In conclusion, we have developed a simple route to 3-sulfonylquinoline derivatives. This protocol was performed effectively at room temperature which involved cheap, readily available arylsulfonylhydrazides and low Watt blue LEDs. Moreover, this strategy exhibits good

Scheme 3 Tentative mechanistic pathway

functional group tolerance, which can be used to afford various 3-sulfonated quinoline derivatives.

4 Experimental section

4.1 Instruments and reagents

Commercially available reagents were of reagent grade (AR grade) and were used without further purification. Reactions were monitored by thin layer chromatography (TLC) using silicycle pre-coated silica gel plates. Flash column chromatography was performed over silicycle silica gel (200~300 mesh). ^1H NMR and ^{13}C NMR spectra were recorded on 400 MHz NMR plus spectrometer using residue solvent peaks as internal standards. Infrared spectra were recorded with IR spectrometer and were reported in reciprocal centimeter (cm^{-1}). High-resolution mass spectra (HRMS) were obtained with a Q-TOF Premier (ESI).

4.2 Typical experimental procedure for the preparation of 3 or 4

N-(3-Phenyl-2-propynyl)aniline derivatives **1** (0.2 mmol), sulfonylhydrazides **2** (0.4 mmol), TBHP (dry, 0.7 mmol), and DCE (2.0 mL) were sequentially added to a 10 mL oven-dried reaction vessel at room temperature. The reaction vessel was exposed to blue LED (435~440 nm, 7 W) irradiation at room temperature in open air with stirring for 8 h. After completion of the reaction, the mixture was concentrated to yield the crude product, which was further purified by flash chromatography (silica gel, petroleum ether/ethyl acetate, $V:V=10:1$) to give the desired products **3** or **4**.

4-Phenyl-3-tosylquinoline (**3a**):^[4e] Light yellow solid (64.6 mg, 90% yield), m.p. 173~175 °C (lit.^[4e] 172.6~174.3 °C); ^1H NMR (500 MHz, CDCl_3) δ : 9.77 (s, 1H), 8.20 (dd, $J=8.6, 1.5$ Hz, 1H), 7.84~7.76 (m, 1H), 7.46~7.41 (m, 2H), 7.34~7.29 (m, 3H), 7.19 (d, $J=8.5$ Hz, 2H), 7.03 (d, $J=8.1$ Hz, 2H), 6.96~6.92 (m, 2H), 2.33 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 149.89, 149.62, 147.70, 144.04, 137.91, 132.63, 132.59, 132.18, 130.00, 129.58, 129.24, 128.66, 127.92, 127.83, 127.64, 127.51, 127.42, 21.56.

6-Methyl-4-phenyl-3-tosylquinoline (**3b**):^[4e] Light yellow solid (63.4 mg, 85% yield), m.p. 199~201 °C (lit.^[4e] 200.5~202.9 °C); ^1H NMR (500 MHz, $\text{Chloroform-}d$) δ : 9.61 (s, 1H), 8.08 (d, $J=9.2$ Hz, 1H), 7.46~7.41 (m, 2H), 7.32 (t, $J=7.7$ Hz, 2H), 7.19 (d, $J=8.5$ Hz, 2H), 7.06~7.01 (m, 2H), 6.97~6.90 (m, 2H), 6.49 (d, $J=2.9$ Hz, 1H), 3.58 (s, 3H), 2.33 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 158.58, 148.03, 146.07, 145.33, 143.94, 138.06, 132.93, 132.87, 131.01, 129.91, 129.21, 128.84, 128.61, 127.91, 127.77, 124.81, 104.79, 55.34, 21.55.

6-Ethyl-4-phenyl-3-tosylquinoline (**3c**): Light yellow solid (66.6 mg, 86% yield), m.p. 215~217 °C; ^1H NMR (500 MHz, $\text{Chloroform-}d$) δ : 9.71 (s, 1H), 8.11 (d, $J=8.6$ Hz, 1H), 7.67 (dd, $J=8.6, 2.0$ Hz, 1H), 7.47~7.42 (m, 1H), 7.32 (t, $J=7.8$ Hz, 2H), 7.18 (d, $J=8.4$ Hz, 2H), 7.06~7.01 (m, 3H), 6.96~6.92 (m, 2H), 2.63 (q, $J=7.4$ Hz, 2H), 2.33 (s, 3H), 1.12 (t, $J=7.6$ Hz, 3H); ^{13}C NMR (126 MHz,

CDCl_3) δ : 149.22, 148.58, 146.89, 144.25, 143.91, 138.08, 133.38, 132.76, 132.56, 130.04, 129.47, 129.19, 128.56, 127.88, 127.61, 127.54, 124.81, 28.97, 21.55, 15.30. HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{22}\text{NO}_2\text{S}$ $[\text{M}+\text{H}]^+$ 388.1366; found, 388.1369.

7-Methyl-4-phenyl-3-tosylquinoline (**3d**):^[4a] Light yellow solid (67.1 mg, 90% yield), m.p. 205~207 °C; ^1H NMR (500 MHz, $\text{Chloroform-}d$) δ : 9.79 (s, 1H), 8.09 (dd, $J=8.5, 1.7$ Hz, 1H), 7.66 (dd, $J=8.5, 7.1$ Hz, 1H), 7.27~7.25 (m, 1H), 7.21~7.17 (m, 3H), 7.11 (d, $J=8.4$ Hz, 2H), 7.03~6.99 (m, 2H), 6.93~6.91 (m, 2H), 2.33 (s, 3H), 1.71 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 151.28, 150.36, 147.19, 143.77, 138.30, 138.08, 135.27, 133.54, 131.95, 131.72, 130.67, 129.21, 129.15, 128.62, 127.90, 127.57, 127.20, 23.95, 21.56.

8-Methyl-4-phenyl-3-tosylquinoline (**3e**):^[4e] Light yellow solid (67.9 mg, 91% yield), m.p. 172~174 °C (lit.^[4e] 168.2~174.0 °C); ^1H NMR (500 MHz, $\text{Chloroform-}d$) δ : 9.78 (s, 1H), 7.65 (d, $J=7.0$ Hz, 1H), 7.46~7.42 (m, 1H), 7.34~7.29 (m, 3H), 7.20 (d, $J=8.5$ Hz, 2H), 7.17~7.14 (m, 1H), 7.04 (d, $J=8.0$ Hz, 2H), 6.95~6.92 (m, 2H), 2.86 (s, 3H), 2.34 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 149.91, 148.68, 146.53, 143.96, 138.09, 137.57, 133.08, 132.37, 132.28, 130.06, 129.23, 128.53, 127.95, 127.57, 127.55, 127.51, 125.42, 21.57, 18.25.

6-Bromo-4-phenyl-3-tosylquinoline (**3f**):^[4e] Light yellow solid (71.7 mg, 82% yield), m.p. 199~202 °C (lit.^[4e] 201.2~203.6 °C); ^1H NMR (500 MHz, $\text{Chloroform-}d$) δ : 9.76 (s, 1H), 8.05 (d, $J=8.9$ Hz, 1H), 7.85 (dd, $J=8.9, 2.2$ Hz, 1H), 7.50~7.41 (m, 2H), 7.33 (t, $J=7.8$ Hz, 2H), 7.17 (d, $J=8.4$ Hz, 2H), 7.04 (d, $J=7.9$ Hz, 2H), 6.95~6.90 (m, 2H), 2.33 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 148.85, 148.32, 148.09, 144.27, 137.59, 135.60, 133.54, 131.87, 131.33, 129.96, 129.33, 129.01, 128.76, 127.95, 127.89, 122.28, 21.60.

1-(4-Phenyl-3-tosylquinolin-6-yl)ethanone (**3g**): Light yellow solid (62.6 mg, 78% yield), m.p. 218~220 °C; ^1H NMR (500 MHz, $\text{Chloroform-}d$) δ : 9.61 (s, 1H), 8.08 (d, $J=9.2$ Hz, 1H), 7.46~7.40 (m, 2H), 7.34~7.29 (m, 2H), 7.19 (d, $J=8.5$ Hz, 2H), 7.06~7.00 (m, 2H), 6.95~6.91 (m, 2H), 6.49 (d, $J=2.9$ Hz, 1H), 3.58 (s, 3H), 2.33 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 167.70, 158.58, 148.03, 146.07, 145.33, 143.94, 138.06, 132.93, 132.87, 131.01, 129.91, 129.21, 128.84, 128.61, 127.91, 127.77, 124.81, 104.79, 55.34, 21.55. HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{20}\text{NO}_3\text{S}$ $[\text{M}+\text{H}]^+$ 402.1158; found, 402.1155.

6-Chloro-4-phenyl-3-tosylquinoline (**3h**):^[4e] Light yellow solid (68.4 mg, 87% yield), m.p. 187~189 °C (lit.^[4e] 185.9~187.3 °C); ^1H NMR (500 MHz, $\text{Chloroform-}d$) δ : 9.75 (s, 1H), 8.13 (d, $J=9.0$ Hz, 1H), 7.72 (dd, $J=9.0, 2.3$ Hz, 1H), 7.47 (t, $J=7.6$ Hz, 1H), 7.33 (t, $J=7.8$ Hz, 2H), 7.26 (d, $J=2.3$ Hz, 1H), 7.17 (d, $J=8.5$ Hz, 2H), 7.06~7.01 (m, 2H), 6.94~6.90 (m, 2H), 2.34 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 148.99, 148.18, 148.03, 144.33, 137.66, 134.08, 133.63, 133.10, 131.99, 131.33, 130.00, 129.38, 129.05, 128.41, 128.01, 127.94, 126.06, 21.65.

6-Fluoro-4-phenyl-3-tosylquinoline (**3i**):^[4e] Light yellow

solid (68.6 mg, 91% yield), m.p. 172 ~ 174 °C (lit.^[4e] 173.3 ~ 176.7 °C); ¹H NMR (500 MHz, Chloroform-*d*) δ: 9.72 (s, 1H), 8.19 (dd, *J* = 9.3, 5.4 Hz, 1H), 7.56 (ddd, *J* = 9.3, 7.8, 2.8 Hz, 1H), 7.48 ~ 7.42 (m, 1H), 7.33 (t, *J* = 7.9 Hz, 2H), 7.18 (d, *J* = 8.4 Hz, 2H), 7.07 ~ 7.01 (m, 2H), 6.94 ~ 6.87 (m, 3H), 2.33 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ: 162.03, 160.04, 149.14, 149.09, 147.10, 147.08, 146.86, 144.22, 137.63, 133.38, 132.26, 132.19, 129.84, 129.28, 128.90, 128.69, 128.62, 127.94, 127.90, 127.84, 122.55, 122.35, 110.79, 110.60, 21.55.

7-Chloro-4-phenyl-3-tosylquinoline (**3j**): Light yellow solid (66.0 mg, 84% yield), m.p. 198 ~ 200 °C; ¹H NMR (500 MHz, Chloroform-*d*) δ: 9.84 (s, 1H), 8.16 (dd, *J* = 8.5, 1.4 Hz, 1H), 7.68 (dd, *J* = 8.5, 7.5 Hz, 1H), 7.53 (dd, *J* = 7.5, 1.4 Hz, 1H), 7.37 ~ 7.34 (m, 1H), 7.16 (dd, *J* = 8.3, 7.6 Hz, 2H), 7.10 (d, *J* = 8.5 Hz, 2H), 7.03 ~ 6.99 (m, 2H), 6.91 ~ 6.87 (m, 2H), 2.33 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ: 151.44, 149.67, 148.17, 143.99, 137.80, 134.71, 133.41, 132.62, 131.77, 131.53, 130.68, 130.02, 129.25, 128.55, 127.64, 126.96, 124.27, 21.57. HRMS (ESI) calcd for C₂₂H₁₇ClNO₂S [M + H]⁺ 394.0663; found 394.0661.

4,8-Diphenyl-3-tosylquinoline (**3k**): Light yellow solid (62.6 mg, 72% yield), m.p. 199 ~ 201 °C; ¹H NMR (500 MHz, Chloroform-*d*) δ: 9.78 (s, 1H), 7.65 (d, *J* = 7.0 Hz, 1H), 7.46 ~ 7.42 (m, 1H), 7.34 ~ 7.29 (m, 3H), 7.20 (d, *J* = 8.5 Hz, 2H), 7.17 ~ 7.14 (m, 1H), 7.04 (d, *J* = 8.0 Hz, 2H), 6.95 ~ 6.92 (m, 2H), 2.86 (s, 3H), 2.34 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ: 149.91, 148.68, 146.53, 143.96, 138.09, 137.57, 133.08, 132.37, 132.28, 130.06, 129.23, 128.53, 127.95, 127.57, 127.55, 127.51, 125.42, 21.57, 18.25. HRMS (ESI) calcd for C₂₈H₂₂NO₂S [M + H]⁺ 436.1366; found 436.1367.

4-(4-Methoxyphenyl)-3-tosylquinoline (**3l**):^[4e] Light yellow solid (58.4 mg, 75% yield), m.p. 161 ~ 164 °C; (lit.^[4e] 159.2 ~ 160.8 °C); ¹H NMR (500 MHz, Chloroform-*d*) δ: 9.75 (s, 1H), 8.18 (dt, *J* = 8.5 Hz, 1.0, 1H), 7.79 (ddd, *J* = 8.4, 6.7 Hz, 1.5, 1H), 7.44 (ddd, *J* = 8.0, 6.8, 1.3 Hz, 1H), 7.38 (ddd, *J* = 8.6, 1.6, 0.7 Hz, 1H), 7.21 (d, *J* = 8.5 Hz, 2H), 7.07 ~ 7.02 (m, 2H), 6.88 ~ 6.83 (m, 4H), 3.89 (s, 3H), 2.33 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ: 159.93, 149.99, 149.66, 147.74, 143.96, 137.97, 133.02, 132.09, 131.39, 129.58, 129.15, 127.92, 127.86, 127.74, 127.42, 124.58, 113.10, 55.40, 21.55.

4-(4-Chlorophenyl)-3-tosylquinoline (**3m**):^[4a] Light yellow solid (60.5 mg, 77% yield), m.p. 192 ~ 194 °C; ¹H NMR (500 MHz, Chloroform-*d*) δ: 9.75 (s, 1H), 8.13 (d, *J* = 9.0 Hz, 1H), 7.72 (dd, *J* = 9.0, 2.3 Hz, 1H), 7.47 (t, *J* = 7.6, 1H), 7.33 (t, *J* = 7.8 Hz, 2H), 7.26 (d, *J* = 2.3 Hz, 1H), 7.17 (d, *J* = 8.5 Hz, 2H), 7.06 ~ 7.01 (m, 2H), 6.94 ~ 6.90 (m, 2H), 2.34 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ: 148.99, 148.18, 148.03, 144.33, 137.66, 134.08, 133.63, 133.10, 131.99, 131.33, 130.00, 129.38, 129.05, 128.41, 128.01, 127.94, 126.06, 21.65.

4-(4-Bromophenyl)-3-tosylquinoline (**3n**):^[4a] Light yellow solid (76.9 mg, 88% yield), m.p. 198 ~ 200 °C; ¹H NMR (500 MHz, Chloroform-*d*) δ: 9.76 (s, 1H), 8.05 (d, *J* = 8.9 Hz, 1H), 7.85 (dd, *J* = 8.9 Hz, 2.2, 1H), 7.50 ~ 7.41

(m, 2H), 7.33 (t, *J* = 7.8 Hz, 2H), 7.17 (d, *J* = 8.4 Hz, 2H), 7.04 (d, *J* = 7.9 Hz, 2H), 6.95 ~ 6.90 (m, 2H), 2.33 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ: 148.85, 148.32, 148.09, 144.27, 137.59, 135.60, 133.54, 131.87, 131.33, 129.96, 129.33, 129.01, 128.76, 127.95, 127.89, 122.28, 21.60.

1-(4-(3-Tosylquinolin-4-yl)phenyl)ethanone (**3o**):^[4e] Light yellow solid (73.0 mg, 91% yield), m.p. 201 ~ 203 °C (lit.^[4e] 202.5 ~ 204.6 °C); ¹H NMR (500 MHz, Chloroform-*d*) δ: 9.71 (s, 1H), 8.11 (d, *J* = 8.6 Hz, 1H), 7.67 (dd, *J* = 8.6, 2.0 Hz, 1H), 7.47 ~ 7.42 (m, 1H), 7.32 (t, *J* = 7.8 Hz, 2H), 7.18 (d, *J* = 8.4 Hz, 2H), 7.06 ~ 7.01 (m, 3H), 6.96 ~ 6.92 (m, 2H), 2.63 (q, *J* = 7.4 Hz, 2H), 2.33 (s, 3H), 1.12 (t, *J* = 7.6 Hz, 3H); ¹³C NMR (126 MHz, CDCl₃) δ: 149.22, 148.58, 146.89, 144.25, 143.91, 138.08, 133.38, 132.76, 132.56, 130.04, 129.47, 129.19, 128.56, 127.88, 127.61, 127.54, 124.81, 28.97, 21.55, 15.30.

4-Phenyl-3-(phenylsulfonyl)quinoline (**4a**):^[4e] Light yellow solid (56.6 mg, 82% yield), m.p. 187 ~ 189 °C (lit.^[4e] 187.8 ~ 189.4 °C); ¹H NMR (400 MHz, Chloroform-*d*) δ: 9.72 (s, 1H), 8.15 (d, *J* = 8.1 Hz, 1H), 7.78 ~ 7.69 (m, 1H), 7.37 ~ 7.32 (m, 3H), 7.24 ~ 7.13 (m, 7H), 6.83 (d, *J* = 8.1 Hz, 2H); ¹³C NMR (101 MHz, CDCl₃) δ: 150.29, 149.58, 147.53, 140.77, 134.24, 133.06, 132.44, 130.07, 130.02, 128.75, 128.67, 128.00, 127.89, 127.76, 127.50.

3-((4-Methoxyphenyl)sulfonyl)-4-phenylquinoline (**4b**):^[4e] Light yellow solid (60.0 mg, 80% yield), m.p. 170 ~ 172 °C (lit.^[4e] 167.1 ~ 168.9 °C); ¹H NMR (500 MHz, Chloroform-*d*) δ: 9.77 (s, 1H), 8.19 (dt, *J* = 8.5, 1.0 Hz, 1H), 7.79 (ddd, *J* = 8.4, 6.8, 1.4 Hz, 1H), 7.46 ~ 7.40 (m, 2H), 7.37 ~ 7.30 (m, 3H), 7.22 (d, *J* = 8.9 Hz, 2H), 6.99 ~ 6.95 (m, 2H), 6.70 (d, *J* = 9.0 Hz, 2H), 3.79 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ: 163.27, 149.64, 147.74, 132.93, 132.74, 132.45, 132.08, 130.15, 130.07, 129.63, 128.67, 127.80, 127.70, 127.54, 127.40, 113.88, 55.66.

3-((4-(Tert-butyl)phenyl)sulfonyl)-4-phenylquinoline (**4c**): Light yellow solid (61.8 mg, 77% yield), m.p. 168 ~ 170 °C; ¹H NMR (500 MHz, Chloroform-*d*) δ: 9.77 (s, 1H), 8.19 (dt, *J* = 8.5, 1.0 Hz, 1H), 7.79 (ddd, *J* = 8.4, 6.8, 1.4 Hz, 1H), 7.46 ~ 7.40 (m, 2H), 7.37 ~ 7.30 (m, 3H), 7.22 (d, *J* = 8.9 Hz, 2H), 6.99 ~ 6.95 (m, 2H), 6.70 (d, *J* = 9.0 Hz, 2H), 3.79 (s, 3H); ¹³C NMR (126 MHz, CDCl₃) δ: 163.27, 149.64, 147.74, 132.93, 132.74, 132.45, 132.08, 130.15, 130.07, 129.63, 128.67, 127.80, 127.70, 127.54, 127.40, 113.88, 55.66. HRMS (ESI) calcd for C₂₅H₂₄NO₂S [M + H]⁺ 402.1522; found 402.1518.

4-Phenyl-3-((4-(trifluoromethoxy)phenyl)sulfonyl)quinoline (**4d**):^[4e] Light yellow solid (66.9 mg, 78% yield), m.p. 161 ~ 165 °C (lit.^[4e] 162.2 ~ 165.7 °C); ¹H NMR (400 MHz, Chloroform-*d*) δ: 9.80 (s, 1H), 8.23 (d, *J* = 8.4 Hz, 1H), 7.89 ~ 7.80 (m, 1H), 7.50 ~ 7.43 (m, 2H), 7.35 ~ 7.29 (m, 5H), 7.07 ~ 7.02 (m, 2H), 6.96 ~ 6.91 (m, 2H); ¹³C NMR (101 MHz, CDCl₃) δ: 152.30, 152.29, 150.13, 149.77, 147.24, 138.95, 132.61, 132.35, 132.06, 130.09, 130.06, 129.64, 128.93, 128.14, 127.85, 127.42, 127.37, 120.57.

3-((4-Fluorophenyl)sulfonyl)-4-phenylquinoline (**4e**):^[4e] Light yellow solid (63.2 mg, 87% yield), m.p. 181 ~ 183 °C (lit.^[4e] 180 ~ 182.3 °C); ¹H NMR (400 MHz, Chloroform-*d*)

δ : 9.79 (s, 1H), 8.28~8.19 (m, 1H), 7.89~7.77 (m, 1H), 7.51~7.44 (m, 2H), 7.38~7.27 (m, 5H), 7.00~6.87 (m, 4H); ^{13}C NMR (101 MHz, CDCl_3) δ : 166.57, 164.02, 150.13, 149.64, 147.37, 136.73, 132.54, 132.44, 132.29, 130.81, 130.71, 130.09, 129.59, 128.91, 128.09, 127.85, 127.44, 116.05, 115.82.

3-((4-Chlorophenyl)sulfonyl)-4-phenylquinoline (**4f**):^[4e] Light yellow solid (65.2 mg, 86% yield), m.p. 168~170 °C (lit.^[4e] 167.1~170.2 °C); ^1H NMR (400 MHz, Chloroform-*d*) δ : 9.78 (s, 1H), 8.27~8.21 (m, 1H), 7.86~7.82 (m, 1H), 7.51~7.45 (m, 2H), 7.38~7.32 (m, 3H), 7.21 (s, 4H), 6.98~6.92 (m, 2H); ^{13}C NMR (101 MHz, CDCl_3) δ : 150.25, 149.63, 147.33, 139.81, 139.19, 132.61, 132.37, 132.12, 130.09, 129.56, 129.34, 128.95, 128.92, 128.13, 127.85, 127.45.

3-((4-Bromophenyl)sulfonyl)-4-phenylquinoline (**4g**):^[4e] Light yellow solid (71.1 mg, 84% yield), m.p. 250~253 °C (lit.^[4e] 252.2~253.8 °C); ^1H NMR (400 MHz, Chloroform-*d*) δ : 9.79 (s, 1H), 8.28~8.19 (m, 1H), 7.89~7.77 (m, 1H), 7.51~7.44 (m, 2H), 7.38~7.27 (m, 5H), 7.00~6.87 (m, 4H); ^{13}C NMR (101 MHz, CDCl_3) δ : 166.57, 164.02, 150.13, 149.64, 147.37, 136.73, 132.54, 132.44, 132.29, 130.81, 130.71, 130.09, 129.59, 128.91, 128.09, 127.85, 127.44, 116.05, 115.82.

Ethyl 4-((4-phenylquinolin-3-yl)sulfonyl)benzoate (**4h**): Light yellow solid (63.4 mg, 76% yield), m.p. 265~267 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 9.80 (s, 1H), 8.24 (d, $J=8.6$ Hz, 1H), 7.92~7.80 (m, 3H), 7.49~7.45 (m, 2H), 7.37~7.29 (m, 5H), 6.96~6.88 (m, 2H), 4.38 (q, $J=7.1$ Hz, 2H), 1.38 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 165.00, 150.47, 149.73, 147.41, 144.47, 134.36, 132.66, 132.23, 131.84, 130.06, 129.67, 129.60, 128.97, 128.14, 127.87, 127.83, 127.47, 127.42, 61.81, 14.24. HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{20}\text{NO}_4\text{S}$ [$\text{M} + \text{H}$]⁺ 418.1108; found 418.1109.

3-([1,1'-Biphenyl]-4-ylsulfonyl)-4-phenylquinoline (**4i**): Light yellow solid (65.7 mg, 78% yield), m.p. 187~189 °C; ^1H NMR (500 MHz, Chloroform-*d*) δ : 9.83 (s, 1H), 8.21 (d, $J=1.5$ Hz, 1H), 7.86~7.78 (m, 1H), 7.53~7.50 (m, 2H), 7.48~7.43 (m, 6H), 7.42~7.39 (m, 1H), 7.38~7.30 (m, 5H), 7.00~6.95 (m, 2H); ^{13}C NMR (126 MHz, CDCl_3) δ : 149.90, 149.79, 147.65, 145.91, 139.31, 139.14, 132.62, 132.55, 132.26, 130.09, 129.69, 129.08, 128.71, 128.65, 128.39, 127.90, 127.73, 127.49, 127.44, 127.28, 127.23. HRMS (ESI) calcd for $\text{C}_{27}\text{H}_{20}\text{NO}_2\text{S}$ [$\text{M} + \text{H}$]⁺ 422.1209; found, 422.1205.

4-Phenyl-3-(*m*-tolylsulfonyl)quinoline (**4j**):^[4a] Light yellow solid (54.6 mg, 76% yield), m.p. 156~159 °C; ^1H NMR (500 MHz, Chloroform-*d*) δ : 9.83 (s, 1H), 8.21 (d, $J=1.5$ Hz, 1H), 7.86~7.78 (m, 1H), 7.53~7.50 (m, 2H), 7.48~7.43 (m, 6H), 7.42~7.39 (m, 1H), 7.38~7.30 (m, 5H), 7.00~6.95 (m, 2H); ^{13}C NMR (126 MHz, CDCl_3) δ : 149.90, 149.79, 147.65, 145.91, 139.31, 139.14, 132.62, 132.55, 132.26, 130.09, 129.69, 129.08, 128.71, 128.65, 128.39, 127.90, 127.73, 127.49, 127.44, 127.28, 127.23.

4-Phenyl-3-(*o*-tolylsulfonyl)quinoline (**4k**): Light yellow solid (56.0 mg, 78% yield), m.p. 152~154 °C; ^1H NMR

(400 MHz, Chloroform-*d*) δ : 9.82 (s, 1H), 8.24 (d, $J=8.4$ Hz, 1H), 7.87~7.78 (m, 1H), 7.46 (ddd, $J=8.2, 6.8, 1.2$ Hz, 1H), 7.34~7.27 (m, 3H), 7.19 (t, $J=7.9$ Hz, 2H), 7.14 (dd, $J=8.0, 1.5$ Hz, 1H), 7.08~7.05 (m, 1H), 6.93~6.87 (m, 1H), 6.87~6.82 (m, 2H), 2.24 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 149.79, 149.50, 147.91, 138.53, 136.70, 133.18, 132.30, 132.20, 132.09, 131.87, 129.66, 129.63, 129.31, 128.57, 127.98, 127.63, 127.41, 127.37, 126.03, 20.00. HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{18}\text{NO}_2\text{S}$ [$\text{M} + \text{H}$]⁺ 360.1053; found 360.1051.

3-(Naphthalen-2-ylsulfonyl)-4-phenylquinoline (**4l**):^[4a] Light yellow solid (60.0 mg, 76% yield), m.p. 155~157 °C; ^1H NMR (500 MHz, Chloroform-*d*) δ : 9.87 (s, 1H), 8.25~8.19 (m, 1H), 7.83~7.78 (m, 2H), 7.74~7.66 (m, 3H), 7.62~7.59 (m, 1H), 7.55 (ddd, $J=8.1, 6.9, 1.3$ Hz, 1H), 7.45~7.35 (m, 3H), 7.29 (dd, $J=8.5, 1.9$ Hz, 1H), 7.20~7.14 (m, 2H), 6.90~6.84 (m, 2H); ^{13}C NMR (126 MHz, CDCl_3) δ : 150.18, 149.83, 147.80, 137.35, 134.87, 132.39, 132.30, 132.23, 131.73, 130.23, 130.02, 129.69, 129.54, 129.19, 129.01, 128.85, 127.88, 127.80, 127.60, 127.50, 127.43, 127.41, 122.45.

4-Phenyl-3-(thiophen-2-ylsulfonyl)quinoline (**4m**):^[4a] Light yellow solid (55.5 mg, 79% yield), m.p. 193~195 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 7.66 (d, $J=8.3$ Hz, 2H), 7.32 (s, 1H), 7.23 (d, $J=8.2$ Hz, 2H), 7.18 (d, $J=8.8$ Hz, 2H), 7.03 (d, $J=8.8$ Hz, 2H), 2.38 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 144.33, 135.67, 135.25, 130.92, 129.91, 129.51, 127.37, 122.94, 21.70.

3-(Methylsulfonyl)-4-phenylquinoline (**4n**): White solid (37.9 mg, 67% yield), m.p. 88~90 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 9.59 (s, 1H), 8.27~8.21 (m, 1H), 7.92~7.83 (m, 1H), 7.59~7.52 (m, 4H), 7.48 (dd, $J=8.5$ Hz, 1.9, 1H), 7.44~7.40 (m, 2H), 2.77 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 149.80, 149.68, 147.45, 132.93, 132.47, 131.67, 130.05, 129.72, 129.52, 128.28, 128.16, 127.69, 127.23, 44.53. HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{13}\text{NO}_2\text{S}$ [$\text{M} + \text{H}$]⁺ 283.0667; found 283.0669.

3-(Cyclopropylsulfonyl)-4-phenylquinoline (**4o**): White solid (48.2 mg, 78% yield), m.p. 95~97 °C; ^1H NMR (500 MHz, Chloroform-*d*) δ : 9.49 (s, 1H), 8.23 (d, $J=8.5$ Hz, 1H), 7.89~7.82 (m, 1H), 7.58~7.50 (m, 4H), 7.49~7.42 (m, 3H), 2.04 (tt, $J=8.0, 4.8$ Hz, 1H), 1.18~1.12 (m, 2H), 0.89~0.81 (m, 2H); ^{13}C NMR (126 MHz, CDCl_3) δ : 149.65, 149.41, 147.68, 133.50, 132.18, 132.05, 130.20, 129.72, 129.30, 128.04, 127.98, 127.71, 127.50, 32.95, 6.33. HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{15}\text{NO}_2\text{S}$ [$\text{M} + \text{H}$]⁺ 309.0823; found 309.0826.

Supporting Information ^1H NMR and ^{13}C NMR spectra of new compounds. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Michael, J. P. *Nat. Prod. Rep.* **2000**, *17*, 603.
(b) Strekowski, L.; Say, M.; Henary, M.; Ruiz, P.; Manzel, L. Macfarlane, D. E.; Bojarski, A. J. *J. Med. Chem.* **2003**, *46*, 1242.
(c) Twin, H.; Batey, R. A. *Org. Lett.* **2004**, *6*, 4913.

- (d) Michael, J. P. *Nat. Prod. Rep.* **2008**, *25*, 166.
- (e) Hayat, F.; Moseley, E.; Salahuddin, A.; Van Zyl, R. L.; Azam, A. *Eur. J. Med. Chem.* **2011**, *46*, 1897.
- (f) Chung, P.-Y.; Bian, Z.-X.; Pun, H.-Y.; Chan, D.; Chan, S.-C.; Chui, C.-H.; Tang, J. C.-O.; Lam, K.-H. *Future Med. Chem.* **2015**, *7*, 947.
- (g) Andries, K.; Verhasselt, P.; Guillemont, J.; Göhlmann, H. W. H.; Neefs, J.-M.; Winkler, H.; Van Gestel, J.; Timmerman, P.; Zhu, M.; Lee, E.; Williams, P.; de Chaffoy, D.; Huitric, E.; Hoffner, S.; Cambau, E.; Truffot-Pernot, C.; Lounis, N.; Jarlier, V. *Science* **2005**, *307*, 223.
- (h) Sun, M.; Jiang, J.; Chen, J.; Yang, Q.; Yu, X. Y. *Tetrahedron* **2019**, *75*, 130456.
- (i) Xie, L.-Y.; Fang, T.-G.; Tan, J.-X.; Zhang, B.; Cao, Z.; Yang, L.-H.; He, W.-M. *Green Chem.* **2019**, *21*, 3858.
- (j) Cao, Z.; Zhu, Q.; Lin, Y.-W.; He, W.-M. *Chin. Chem. Lett.* **2019**, *30*, 2132.
- (k) Peng, S.; Song, Y.-X.; He, J.-Y.; Tang, S.-S.; Tan, J.-X.; Cao, Z.; Lin, Y.-W.; He, W.-M. *Chin. Chem. Lett.* **2019**, *30*, 2287.
- (l) Wang, L. L.; Bao, P. L.; Liu, W. W.; Liu, S. T.; Hu, C. S.; Yue, H. L.; Yang, D. S.; Wei, W. *Chin. J. Org. Chem.* **2018**, *38*, 3189 (in Chinese).
- (王雷雷, 鲍鹏丽, 刘维伟, 刘思彤, 胡昌松, 岳会兰, 杨道山, 魏伟, 有机化学, **2018**, *38*, 3189.)
- (m) Ye, J.-L.; Zhu, Y.-N.; Geng, H.; Huang, P.-Q. *Sci. China Chem.* **2018**, *61*, 687.
- (n) Bao, P.; Wang, L.; Liu, Q.; Yang, D.; Wang, H.; Zhao, X.; Yue, H.; Wei, W. *Tetrahedron Lett.* **2019**, *60*, 214.
- (o) Wang, L.; Zhang, M.; Zhang, Y.; Liu, Q.; Zhao, X.; Li, J.-S.; Luo, Z.; Wei, W. *Chin. Chem. Lett.* **2020**, *31*, 67.
- [2] (a) Galambos, J.; Domány, G.; Nógrádi, K.; Wágner, G.; Keserü, G. M.; Bobok, A.; Kolok, S.; Mikó-Bakk, M. L.; Vastag, M.; Sághy, K.; Kóti, J.; Szakács, Z.; Béni, Z.; Gál, K.; Szombathelyi, Z.; Greiner, I. *Biorg. Med. Chem. Lett.* **2016**, *26*, 1249.
- (b) Codony, X.; Vela, J. M.; Ramírez, M. J. *Curr. Opin. Pharm.* **2011**, *11*, 94.
- (c) Zajdel, P.; Marciniak, K.; Maślankiewicz, A.; Paluchowska, M.; Satała, H. G.; Partyka, A.; Jastrzębska-Więsek, M.; Wróbel, D.; Wesolowska, A.; Duszyńska, B.; Bojarski, A. J.; Pawłowski, M. *Biorg. Med. Chem.* **2011**, *19*, 6750.
- (d) Xie, L.-Y.; Peng, S.; Tan, J.-X.; Sun, R.-X.; Yu, X.; Dai, N.-N.; Tang, Z.-L.; Xu, X.; He, W.-M. *ACS Sustainable Chem. Eng.* **2018**, *6*, 16976.
- [3] (a) Kong, L.; Yu, S.; Zhou, X.; Li, X. *Org. Lett.* **2016**, *18*, 588.
- (b) Zheng, J.; Li, Z.; Huang, L.; Wu, W.; Li, J.; Jiang, H. *Org. Lett.* **2016**, *18*, 3514.
- (c) An, X.-D.; Yu, S. *Org. Lett.* **2015**, *17*, 2692.
- (d) Kong, L.; Zhou, Y.; Huang, H.; Yang, Y.; Liu, Y.; Li, Y. *J. Org. Chem.* **2015**, *80*, 1275.
- (e) Xie, L.-Y.; Peng, S.; Fan, T.-G.; Liu, Y.-F.; Sun, M.; Jiang, L.-L.; Wang, X.-X.; Cao, Z.; He, W.-M. *Sci. China Chem.* **2019**, *62*, 460.
- (f) Liu, B.; Gao, H.; Yu, Y.; Wu, W.; Jiang, H. *J. Org. Chem.* **2013**, *78*, 10319.
- [4] (a) Zhang, Y.; Chen, W.; Jia, X.; Wang, L.; Li, P. *Chem. Commun.* **2019**, *55*, 2785.
- (b) Sun, D.; Yin, K.; Zhang, R. *Chem. Commun.* **2018**, *54*, 1335.
- (c) Deng, Q.; Xu, Y.; Liu, P.; Tan, L.; Sun, P. *Org. Chem. Front.* **2018**, *5*, 19.
- (d) Wu, Z.-G.; Liang, X.; Zhou, J.; Yu, L.; Wang, Y.; Zheng, Y.-X.; Li, Y.-F.; Zuo, J.-L.; Pan, Y. *Chem. Commun.* **2017**, *53*, 6637.
- (e) Zhang, L.; Chen, S.; Gao, Y.; Zhang, P.; Wu, Y.; Tang, G.; Zhao, Y. *Org. Lett.* **2016**, *18*, 1286.
- [5] (a) Wei, Y.; Zhou, Q.-Q.; Tan, F.; Lu, L.-Q.; Xiao, W.-J. *Synthesis* **2019**, *51*, 3021.
- (b) Xie, L.-Y.; Bai, Y.-S.; Xu, X.-Q.; Peng, X.; Tang, H.-S.; Huang, Y.; Lin, Y.-W.; Cao, Z.; He, W.-M. *Green Chem.* **2020**, *22*, 1720.
- (c) Peng, S.; Lin, Y.-W.; He, W.-M. *Chin. J. Org. Chem.* **2020**, *40*, 541 (in Chinese).
- (彭莎, 林英武, 何卫民, 有机化学, **2020**, *40*, 541.)
- (d) Xie, L.-Y.; Chen, Y.-L.; Qin, L.; Wen, Y.; Xie, J.-W.; Tan, J.-X.; Huang, Y.; Cao, Z.; He, W.-M. *Org. Chem. Front.* **2019**, *6*, 3950.
- [6] (a) Yang, F.-L.; Wang, F.-X.; Wang, T.-T.; Wang, Y.-J.; Tian, S.-K. *Chem. Commun.* **2014**, *50*, 2111.
- (b) Bao, W.-H.; Wang, Z.; Tang, X.; Zhang, Y.-F.; Tan, J.-X.; Zhu, Q.; Cao, Z.; Lin, Y.-W.; He, W.-M. *Chin. Chem. Lett.* **2019**, *30*, 2259.
- (c) Wei, W.; Liu, C.; Yang, D.; Wen, J.; You, J.; Suo, Y.; Wang, H. *Chem. Commun.* **2013**, *49*, 10239.
- [7] (a) Taniguchi, T.; Idota, A.; Ishibashi, H. *Org. Biomol. Chem.* **2011**, *9*, 3151.
- (b) Zhang, M.; Xie, P.; Zhao, W.; Niu, B.; Wu, W.; Bian, Z.; Pittman, C. U.; Zhou, A. *J. Org. Chem.* **2015**, *80*, 4176.
- (c) Li, X.; Xu, X.; Hu, P.; Xiao, X.; Zhou, C. *J. Org. Chem.* **2013**, *78*, 7343.
- (d) Wei, W.; Wen, J.; Yang, D.; Guo, M.; Wang, Y.; You, J.; Wang, H. *Chem. Commun.* **2015**, *51*, 768.
- (e) Li, X.; Xu, X.; Zhou, C. *Chem. Commun.* **2012**, *48*, 12240.

(Li, L.; Fan, Y.)