四氢呋喃螺氧化吲哚衍生物的一锅法高效合成

郭欣 郭亚军 孔德志 卢会杰 华远照* 王敏灿*

(郑州大学化学学院 绿色催化研究所 郑州 450000)

摘要 报道了 α -羟基芳基酮和 β,γ -不饱和- α -酮酰胺发生的 Michael/半缩酮化和傅-克(Friedel-Crafts)反应的两步一锅反应。该方法利用不包含氧化吲哚和四氢呋喃结构的链状底物, 高效构建出包含螺碳原子、氧化吲哚环和四氢呋喃环的四氢呋喃螺氧化吲哚衍生物。

关键词 一锅法; 串联反应; 四氢呋喃螺氧化吲哚

Efficient Synthesis of Tetrahydrofuran Spirooxindoles via One-Pot Reaction

Guo, Xin Guo, Yajun Kong, Dezhi Lu, Huijie Hua, Yuanzhao* Wang, Mincan*

(Institute of Green Catalysis, College of Chemistry, Zhengzhou University, Zhengzhou 450000)

Abstract A one-pot reaction of Michael/hemiketalization and Friedel-Crafts reaction of α -hydroxy aryl ketones and β,γ -unsaturated α -ketoamides has been developed. The process enables efficient synthesis of tetrahydrofuran spirooxindoles using chain substrates that do not contain oxindole and tetrahydrofuran skeletons. A spiro-carbon center, an oxindole ring and a tetrahydrofuran ring, are constructed in this process.

Keywords one-pot; cascade reaction; tetrahydrofuran spirooxindoles

1 Introduction

Spirocyclic oxindoles are privileged structural motifs existing in a broad range of natural products and pharmaceuticals, which possess significant biological activities.^[1] It has been shown that substituents and spatial structures greatly influence the properties of those compounds.^[2] And potentially promising skeletons are those bearing a heteroatom (especially N or O) in the spiro stereocenter at the C(3) position of the oxindole.^[3] Among various spirocyclic oxindoles, tetrahydrofuran spirooxindole derivatives have also attracted much attention of organic chemists for their potential biological activities. For example, spiro[furo-2,3'-oxindole]s could inhibit the growth of lung adenocarcinoma (A549) cells and hepatocellular carcinoma (HepG2) cells.^[4] Consequently, a lot of elegant methods for their generation have been developed.^[5]

Among these documented methods, there are two pathways to access tetrahydrofuran spirooxindoles. One method for the synthesis of tetrahydrofuran spirooxindoles starts with isatins or 3-hydroxy-2-oxindoles containing

ring A, followed by the construction of the spiro-carbon center and the formation of ring B (Pathway a, Scheme 1).^[6] The other needs readily available B-ring-containing amides to construct spiro-carbon and form ring A (Pathway b, Scheme 1).^[7] Only one ring is built and the other has already existed in one of the substrates in both above strategies. Meanwhile, sometimes these strategies have some inherent drawbacks, such as harsh reaction conditions, multiple operation procedures, poor compatibility of functional groups, and poor availability of the starting material to diversify the molecular structure. It is highly desirable and challenging to develop simple and integrated methods for the construction of a spiro-carbon center, ring A, and ring B in one-pot reaction.

Recently, our group has endeavored to develop the applications of metal-catalysts.^[8] And we have successfully realized some dinuclear metal-catalytic asymmetric cascade reactions for the synthesis of structurally novel and functionalized compounds,^[9] such as Michael/hemiketalization to construct five- or six-membered oxygenated het-

* Corresponding authors. E-mail: hyzhao@zzu.edu.cn; wangmincan@zzu.edu.cn

Received March 11, 2020; revised April 20, 2020; published online April 30, 2020.

Project supported by the National Natural Science Foundation of China (No. 21871237), the China Postdoctoral Science Foundation (No. 2017M622361) and the Education Department of Henan Province (Nos. 17B150014, 18B150028).

国家自然科学基金(No. 21272216)、中国博士后科学基金(No. 2017M622361)和河南省教育厅(Nos. 17B150014, 18B150028)资助项目。

Scheme 1 Construction of tetrahydrofuran spirooxindoles

erocycles,^[9a-9b] phospho-Michael/Michael cascade reaction for the synthesis of 1,2,3-trisubstituted indanes,^[9c] Knoevenagel/Michael/cyclization for the synthesis of dihydrofuran spirooxindoles,^[9d] Michael/transesterification for the formation of spiro-indanone γ -butyrolactones,^[9e] and asymmetric *exo'*-selective [3+2] cycloaddition to construct trifluoromethyl-substituted 2,3-pyrrolidinyl dispirooxindoles.^[9f] Particularly, we recently demonstrated that β,γ -unsaturated α -ketoamides could serve as especially appropriate Michael acceptors for the enantioselective construction of dispirocyclic oxindoles through cascade Michael/cyclization/Friedel-Crafts reaction.^[9g-9h]

As a continuation, we envisioned that the reaction of α -hydroxy aryl ketones **1** with β,γ -unsaturated α -ketoamides **2** may provide a straightforward approach to construct multisubstituted tetrahydrofuran spirooxindoles in one pot reaction. Herein we wish to report our progress with acceptable results in this area.^[10]

1 Results and discussion

1.1 Catalyst screening and optimization

Initially, α -hydroxy aryl ketone **1a** and β,γ -unsaturated α -ketoamide **2a** were chosen as model substrates for optimization studies (Table 1). Reactions were conducted in dichloromethane (DCM) at room temperature for 24 h. However, no obvious reaction occurred when ZnCl_2 (10 mol%) was used as catalyst (Table 1, Entry 1). A number of Lewis acids, such as FeCl_3 , $\text{Zn}(\text{OTf})_2$, $\text{Zn}(\text{OAc})_2$, $\text{Cu}(\text{OTf})_2$, $\text{Cu}(\text{OAc})_2$, $\text{Mn}(\text{OAc})_2$, $\text{La}(\text{OAc})_2$, and so on, were also tested, and none of them could catalyze the model reaction (Table 1, Entries 1~7). Then, some bases were employed as catalysts in this model reaction. Fortunately, the product **3** was obtained in 49% isolated yield when NaOH was used (Table 1, Entry 8). Encouraged by this, KOH and CH_3ONa were examined subsequently, and the yields of product **3** were improved to 49% and 70%, respectively (Table 1, Entries 9 and 10). A Brønsted base ZnEt_2 was also evaluated, and gave the product **3** in 81%

yield (Table 1, Entry 11). However, no reaction occurred when Lewis base NEt_3 was used as catalyst (Table 1, Entry 12).

Table 1 Optimization of the reaction conditions^a

Entry	Catalyst	x/mol%	Solvent	Yield/% of 3	Yield/% of 4/4' (4/4')
1	ZnCl_2	10	CH_2Cl_2	Trace	—
2	FeCl_3	10	CH_2Cl_2	NR	—
3	$\text{Zn}(\text{OTf})_2$	10	CH_2Cl_2	NR	—
4	$\text{Cu}(\text{OTf})_2$	10	CH_2Cl_2	NR	—
5	$\text{La}(\text{OAc})_3$	10	CH_2Cl_2	NR	—
6	$\text{Co}(\text{OAc})_2$	10	CH_2Cl_2	NR	—
7	$\text{Zn}(\text{OAc})_2$	10	CH_2Cl_2	NR	—
8	NaOH	10	CH_2Cl_2	45	—
9	KOH	10	CH_2Cl_2	49	—
10	CH_3ONa	10	CH_2Cl_2	70	—
11	ZnEt_2	10	CH_2Cl_2	81	—
12	NEt_3	10	CH_2Cl_2	NR	—
13	ZnEt_2	10	THF	62	—
14	ZnEt_2	10	Toluene	64	—
15	ZnEt_2	10	Benzene	74	—
16	ZnEt_2	10	CHCl_3	76	—
17	ZnEt_2	10	CH_3CN	79	—
18 ^b	ZnEt_2	5	CH_2Cl_2	76	—
19 ^c	ZnEt_2	20	CH_2Cl_2	88	—
20 ^c	ZnEt_2	30	CH_2Cl_2	88	—
21 ^{d,e}	ZnEt_2	20	CH_2Cl_2	—	60 (32/28)
22 ^{d,f}	ZnEt_2	20	CH_2Cl_2	—	71 (39/32)
23 ^{d,g}	ZnEt_2	20	CH_2Cl_2	—	84 (46/38)

^a Unless otherwise noted, all the reactions were conducted with 10 mol% catalyst, 0.20 mmol of **1**, and 0.21 mmol of **2** in 2 mL of solvent at 20 °C for 24 h. ^b Reaction time was 40 h. ^c Reaction time was 6 h. ^d 6 h later, acid was added at 0 °C and the mixture was stirred for another 1 h. ^e Two drops of H_2SO_4 was added. ^f 0.5 mL of HCl was added. ^g 0.5 mL of CF_3COOH (TFA) was added.

Next, the evaluation of different solvents, including tetrahydrofuran (THF), toluene, benzene, CHCl_3 , and CH_3CN indicated that CH_2Cl_2 was the optimal solvent which gave the product **3** in 81% yield (Table 1, Entries 11, 13~17). When the catalyst loading amount was lowered to 5 mol%,

the yield of product **3** was decreased to 76% and the reaction time was prolonged to 48 h. Once increasing the catalyst loading amount to 20 mol%, the yield of product **3** was increased to 88% in 6 h. While the yield was not further improved when 30 mol% catalyst loading was employed, which indicated that 20 mol% of ZnEt_2 was the best choice (Table 1, Entries 18~20).

The ^1H NMR and ^{13}C NMR spectra revealed that product **3** was obtained as a mixture of the chain structure (Michael addition product) and the cyclic structure (Michael/hemiketalization product).^[10] There was a dynamic balance between the chain structure and the cyclic structure, which led to the difficulty to purify and characterize. It was found that this mixture could give the title compound tetrahydrofuran spirooxindole **4/4'** through an intramolecular Friedel-Crafts reaction, when it was treated with H_2SO_4 (Table 1, Entry 21). Then, concentrated hydrochloric acid (HCl) and CF_3COOH (TFA) were evaluated, and TFA was the optimal choice giving product **4/4'** in 84% total yields through 2 steps one-pot reaction (Table 1, Entries 22 and 23). Although the diastereoselectivity was low, the two isomers of *anti*-**4aa** and *syn*-**4aa'** could be separated by flash column chromatography on silica gel.

Thus the optimal conditions were determined as follows (Table 1, Entry 23): 20 mol% ZnEt_2 , 2 mL of CH_2Cl_2 , α -hydroxy aryl ketones **1** (0.20 mmol), β,γ -unsaturated α -ketoamide **2** (0.21 mmol) at 20 °C for 6 h. Then 0.5 mL of TFA was dropwise added at 0 °C and stirred for another 1 h.

1.2 Substrate scope

The substrate scope of this multisubstituted tetrahydrofuran spirooxindoles forming reaction was explored under the optimal conditions. Using **1a** as a model substrate, the *N*-protecting groups R^1 of β,γ -unsaturated- α -ketoamides **2** was first examined. The experiment results revealed that the steric bulk of the substituent at *N* atom in **2a**~**2c** had an obvious influence on the yield in this reaction. When substituent R^1 was changed from Me to Et or Bn, the total yields of products **4/4'** decreased to 53% and 51% (Table 2, Entries 2 and 3). The reason might owe to the larger steric hindrance of Et or Bn.

Then a series of *N*-methyl β,γ -unsaturated- α -ketoamides **2** bearing various substituents attached on the phenyl ring were investigated. Substrates **2d**~**2h** with electron-withdrawing halogen groups (F, Cl or Br) or electron-donating groups (Me or MeO) at the *para*-position of the *N*-aromatic ring reacted with **1a** efficiently under the standard conditions. And the corresponding products **4/4'** were obtained in 62%~87% total yields (Table 2, Entries 4~8).

The effect of the *para*-substituent of the phenyl ring (Ar^2) was also explored. Various functional groups, both the electron-withdrawing groups (NO_2 , F, Cl or Br) and the electron-donating groups (Me or MeO), were well tolerated (Table 2, Entries 9~14). Meanwhile, it was also observed that nitro group of Ar^2 affected the yield obviously (51%, Table 2, Entry 9). Substrates **2o** and **2p** with Br substituent

at the *meta*- or *ortho*-position of Ar^2 provided the corresponding products **4/4'** in 72% and 79% total yields, respectively (Table 2, Entries 15, 16). Notably, 1-naphthyl and 2-naphthyl substituted ketoamides **2q**~**2r** took part in this reaction successfully, giving corresponding products **4/4'** in total yields of 75% and 84%, respectively. Further study showed that 2-furyl substituted ketoamide **2s** was also a suitable substrate to react with **1a** in 85% total yields.

The reaction of ketoamide **2a** with various nucleophiles was next investigated. A series of nucleophiles **1b**~**1i** bearing either electron-withdrawing groups (F, Cl and Br) or electron-donating groups (Me and MeO) at aromatic ring Ar^1 underwent the reaction efficiently, giving 71%~95% total yields (Table 2, Entries 20~27).

To determine the relative configuration of the products, product **4al'** was treated with hydroxylamine hydrochloride and NaOAc in CH_3OH , which led to the corresponding product of oxime **5'** in 96% yield (Scheme 2). A single crystal of compound **5'** was obtained for X-ray crystallographic analysis (Figure 1).^[11] Thus, the relative stereochemistry of **4al'** could be deduced.

Scheme 2 Derivatization of product **4al'**

Figure 1 X-ray structure of compound **5'**

2 Conclusions

In conclusion, we have developed an efficient method for the direct formation of tetrahydrofuran spirooxindoles,

Table 2 Substrate scope of the reaction^a

Entry	Ar ¹	R ¹	Ar ²	R ²	Product	Yield/% (4/4')
1	Ph	Me	Ph	H	4aa/4aa'	84 (46/38)
2	Ph	Et	Ph	H	4ab/4ab'	53 (22/31)
3	Ph	Bn	Ph	H	4ac/4ac'	51 (25/26)
4	Ph	Me	Ph	F	4ad/4ad'	86 (56/30)
5	Ph	Me	Ph	Cl	4ae/4ae'	87 (57/30)
6	Ph	Me	Ph	Br	4af/4af'	61 (39/22)
7	Ph	Me	Ph	Me	4ag/4ag'	74 (44/30)
8	Ph	Me	Ph	MeO	4ah/4ah'	76 (37/39)
9	Ph	Me	4-O ₂ NC ₆ H ₄	H	4ai/4ai'	51 (24/27)
10	Ph	Me	4-FC ₆ H ₄	H	4aj/4aj'	77 (39/38)
11	Ph	Me	4-ClC ₆ H ₄	H	4ak/4ak'	88 (44/44)
12	Ph	Me	4-BrC ₆ H ₄	H	4al/4al'	94 (61/33)
13	Ph	Me	4-MeC ₆ H ₄	H	4am/4am'	79 (41/38)
14	Ph	Me	4-MeOC ₆ H ₄	H	4an/4an'	72 (39/33)
15	Ph	Me	3-BrC ₆ H ₄	H	4ao/4ao'	72 (49/23)
16	Ph	Me	2-BrC ₆ H ₄	H	4ap/4ap'	79 (50/29)
17	Ph	Me	1-Naphthyl	H	4aq/4aq'	75 (37/38)
18	Ph	Me	2-Naphthyl	H	4ar/4ar'	84 (58/26)
19	Ph	Me	2-Furyl	H	4as/4as'	85 (43/42)
20	4-FC ₆ H ₄	Me	Ph	H	4ba/4ba'	87 (47/40)
21	4-ClC ₆ H ₄	Me	Ph	H	4ca/4ca'	92 (48/44)
22	4-BrC ₆ H ₄	Me	Ph	H	4da/4da'	94 (59/36)
23	4-MeC ₆ H ₄	Me	Ph	H	4ea/4ea'	98 (46/42)
24	4-MeOC ₆ H ₄	Me	Ph	H	4fa/4fa'	91 (62/29)
25	3-BrC ₆ H ₄	Me	Ph	H	4ga/4ga'	80 (48/42)
26	3-MeOC ₆ H ₄	Me	Ph	H	4ha/4ha'	78 (41/37)
27	2-MeOC ₆ H ₄	Me	Ph	H	4ia/4ia'	71 (42/29)

^a Unless otherwise noted, all the reactions were conducted with 20 mol% catalyst, 0.20 mmol of **1**, and 0.21 mmol of **2** in 2 mL of CH₂Cl₂ at 20 °C. 6 h later, 0.5 mL of TFA was dropwise added at 0 °C.

including the constructions of a spiro-carbon center, an oxindole ring, and a tetrahydrofuran ring in one-pot reaction. This chemistry is based on Michael/hemiketalization and Friedel-Crafts reaction of α -hydroxy aryl ketones and β,γ -unsaturated α -ketoamides. Using ZnEt₂ and TFA as catalysts, the one-pot reaction proceeded efficiently and delivered a wide range of tetrahydrofuran spirooxindoles in moderate to good yields. We expect that this novel protocol would provide a convenient and versatile synthetic approach toward structurally diverse tetrahydrofuran spirooxindoles using simple chain substrates.

3 Experimental section

3.1 General procedures

The melting point (m.p.) was determined on an electro-thermal digital melting point apparatus and uncorrected.

NMR spectra were recorded on an NMR spectrometer using CDCl₃ or DMSO as the solvents and TMS as an internal standard (400 MHz for ¹H NMR, 100 MHz for ¹³C NMR). HRMS was performed on a Q-TOF Micro LC/MS System ESI spectrometer.

3.2 General procedure for the catalytic reaction

In a flame-dried Schlenk tube, a solution of diethylzinc (0.04 mL, 1.0 mol/L in hexane, 0.04 mmol) was added to a solution of compound **1** (0.2 mmol) and substrate **2** (0.21 mmol) in dry CH₂Cl₂ (2 mL) under nitrogen at 20 °C. The mixture was stirred for 6 h to complete the Michael/hemiketalization reaction (step 1, detected by TLC). Then, 0.5 mL of TFA was added dropwise to the mixture at 0 °C, and stirred for another 1 h at room temperature. Finally, the solution was diluted with a saturated solution of Na₂CO₃ and extracted with CH₂Cl₂ (10 mL $\times$ 3). The combined

organic layer was dried over MgSO_4 and concentrated *in vacuo*. The residue was purified by flash column chromatography on silica gel to give products **4aa**^[10] and **4aa'**, respectively.

5-Benzoyl-1'-methyl-4-phenyl-4,5-dihydro-3*H*-spiro[furan-2,3'-indolin]-2'-one (**4aa'**): White solid (38% yield), m.p. 209~211 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.70 (d, $J=7.6$ Hz, 2H), 7.51 (d, $J=7.3$ Hz, 1H), 7.45~7.25 (m, 6H), 7.18~7.00 (m, 4H), 6.86 (d, $J=7.8$ Hz, 1H), 5.97 (d, $J=8.2$ Hz, 1H), 4.38 (q, $J=8.3$ Hz, 1H), 3.23 (s, 3H), 3.08 (dd, $J=13.0$, 8.7 Hz, 1H), 2.69 (dd, $J=12.9$, 8.2 Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 195.9, 174.7, 143.55, 138.6, 136.5, 132.8, 131.5, 130.1, 128.7, 128.4, 128.3, 128.2, 127.1, 123.4, 123.0, 108.7, 85.2, 83.6, 49.0, 41.9, 26.5; HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{22}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 384.1594, found 384.1596.

5-Benzoyl-1'-ethyl-4-phenyl-4,5-dihydro-3*H*-spiro[furan-2,3'-indolin]-2'-one (**4ab'**): White solid (31% yield), m.p. 200~202 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.69 (d, $J=7.7$ Hz, 2H), 7.52 (d, $J=7.3$ Hz, 1H), 7.45~7.22 (m, 6H), 7.17~6.97 (m, 4H), 6.87 (d, $J=7.8$ Hz, 1H), 5.98 (d, $J=8.2$ Hz, 1H), 4.37 (q, $J=8.3$ Hz, 1H), 3.78 (q, $J=7.1$ Hz, 2H), 3.08 (dd, $J=12.8$, 8.7 Hz, 1H), 2.68 (dd, $J=12.9$, 8.3 Hz, 1H), 1.29 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 196.3, 174.7, 143.0, 138.9, 137.0, 133.1, 132.2, 130.3, 129.0, 128.8, 128.6, 128.6, 127.4, 123.9, 123.2, 109.2, 85.6, 84.0, 49.3, 42.5, 35.5, 13.0; HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{24}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 398.1751, found 398.1754.

5-Benzoyl-1'-benzyl-4-phenyl-4,5-dihydro-3*H*-spiro[furan-2,3'-indolin]-2'-one (**4ac'**): White solid (26% yield), m.p. 175~177 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.73 (d, $J=7.6$ Hz, 2H), 7.54 (d, $J=7.2$ Hz, 1H), 7.48~7.41 (m, 1H), 7.38~7.21 (m, 10H), 7.19~7.04 (m, 4H), 6.77 (d, $J=7.8$ Hz, 1H), 6.03 (d, $J=8.3$ Hz, 1H), 4.95 (dd, $J=54.7$, 15.7 Hz, 2H), 4.43 (q, $J=8.4$ Hz, 1H), 3.19 (dd, $J=12.8$, 9.0 Hz, 1H), 2.74 (dd, $J=12.8$, 8.2 Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 196.3, 175.4, 142.9, 138.7, 136.9, 135.9, 133.1, 131.9, 130.3, 129.2, 129.0, 128.8, 128.6, 128.5, 128.1, 127.8, 127.5, 123.8, 123.4, 110.1, 85.5, 84.1, 49.4, 44.5, 42.4; HRMS (ESI) calcd for $\text{C}_{31}\text{H}_{26}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 460.1907, found 460.1909.

5-Benzoyl-5'-fluoro-1'-methyl-4-phenyl-4,5-dihydro-3*H*-spiro[furan-2,3'-indolin]-2'-one (**4ad'**): White solid (30% yield), m.p. 196~198 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.67 (d, $J=7.7$ Hz, 2H), 7.50~7.36 (m, 1H), 7.35~7.21 (m, 5H), 7.19~7.01 (m, 4H), 6.88~6.71 (m, 1H), 5.94 (d, $J=8.3$ Hz, 1H), 4.33 (dd, $J=16.8$, 8.4 Hz, 1H), 3.21 (s, 3H), 3.10 (dd, $J=12.8$, 9.1 Hz, 1H), 2.65 (dd, $J=12.9$, 8.1 Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 195.6, 174.5, 159.3 (d, $J=240.3$ Hz), 139.48 (d, $J=1.9$ Hz), 138.0, 136.5, 132.8, 128.6, 128.4, 128.3, 128.2, 127.2, 116.1, (d, $J=23.2$ Hz), 111.7 (d, $J=24.7$ Hz), 109.2 (d, $J=7.87$ Hz), 84.98, 83.7, 48.9, 41.8, 26.6; HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{21}\text{FNO}_3$ $[\text{M}+\text{H}]^+$ 402.1500, found 402.1503.

5-Benzoyl-5'-chloro-1'-methyl-4-phenyl-4,5-dihydro-3*H*-spiro[furan-2,3'-indolin]-2'-one (**4ae'**): White solid (30% yield), m.p. 180~182 °C; ^1H NMR (600 MHz, CDCl_3) δ :

7.71~7.70 (m, 2H), 7.56 (d, $J=7.7$ Hz, 2H), 7.41~7.34 (m, 2H), 7.31~7.26 (m, 3H), 7.21~7.12 (m, 2H), 6.96~6.82 (m, 2H), 6.16 (d, $J=8.5$ Hz, 1H), 4.89 (dt, $J=10.5$, 8.2 Hz, 1H), 3.23 (s, 3H), 3.16 (dd, $J=20.6$, 8.3 Hz, 2H), 2.52 (dd, $J=12.4$, 7.3 Hz, 1H); ^{13}C NMR (151 MHz, CDCl_3) δ : 196.4, 174.7, 143.4, 136.7, 136.2, 132.8, 132.3, 131.2, 130.0, 129.7, 128.7, 128.3, 128.2, 127.7, 125.3, 123.4, 123.1, 108.6, 83.4, 81.9, 47.4, 40.6, 29.7, 26.5; HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{21}\text{ClNO}_3$ $[\text{M}+\text{H}]^+$ 418.1204, found 418.1205.

5-Benzoyl-5'-bromo-1'-methyl-4-phenyl-4,5-dihydro-3*H*-spiro[furan-2,3'-indolin]-2'-one (**4af'**): White solid (23% yield), m.p. 203~205 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.57~7.40 (m, 3H), 7.36~7.28 (m, 1H), 7.28~7.18 (m, 1H), 7.17~7.03 (m, 4H), 6.95~6.80 (m, 3H), 6.62 (d, $J=8.3$ Hz, 1H), 5.84 (d, $J=8.4$ Hz, 1H), 4.31~4.16 (m, 1H), 3.03 (s, 3H), 2.85 (dd, $J=12.9$, 9.1 Hz, 1H), 2.49 (dd, $J=13.0$, 8.3 Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 196.2, 174.5, 142.2, 137.97, 136.3, 133.3, 132.7, 132.6, 128.4, 128.1, 127.97, 127.95, 126.9, 126.5, 115.5, 110.1, 84.8, 83.5, 48.6, 41.9, 26.1; HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{21}\text{BrNO}_3$ $[\text{M}+\text{H}]^+$ 462.0699, found 462.0700.

5-Benzoyl-1',5'-dimethyl-4-phenyl-4,5-dihydro-3*H*-spiro[furan-2,3'-indolin]-2'-one (**4ag'**): White solid (30% yield), m.p. 201~203 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.71 (d, $J=7.6$ Hz, 2H), 7.47~7.36 (m, 1H), 7.35~7.22 (m, 5H), 7.20~7.02 (m, 4H), 6.73 (d, $J=7.9$ Hz, 1H), 5.95 (d, $J=8.2$ Hz, 1H), 4.38 (q, $J=8.4$ Hz, 1H), 3.19 (s, 3H), 3.07 (dd, $J=12.9$, 8.7 Hz, 1H), 2.67 (dd, $J=12.9$, 8.3 Hz, 1H), 2.38 (s, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 195.98, 174.7, 141.2, 138.6, 136.6, 132.7, 132.6, 131.5, 130.2, 128.7, 128.5, 128.2, 128.2, 127.0, 124.2, 108.4, 85.3, 83.8, 48.99, 41.9, 26.5, 21.2; HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{24}\text{NO}_3$ $[\text{M}+\text{H}]^+$ 398.1751, found 398.1753.

5-Benzoyl-5'-methoxy-1'-methyl-4-phenyl-4,5-dihydro-3*H*-spiro[furan-2,3'-indolin]-2'-one (**4ah'**): White solid (39% yield), m.p. 215~217 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.69 (d, $J=7.6$ Hz, 2H), 7.45~7.35 (m, 1H), 7.34~7.22 (m, 4H), 7.16~6.99 (m, 4H), 6.91~6.78 (m, 1H), 6.79~6.70 (m, 1H), 5.93 (d, $J=8.3$ Hz, 1H), 4.34 (q, $J=8.4$ Hz, 1H), 3.82 (s, 3H), 3.18 (s, 3H), 3.07 (dd, $J=12.9$, 8.7 Hz, 1H), 2.66 (dd, $J=12.9$, 8.2 Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 195.8, 174.4, 156.3, 138.5, 136.99, 136.5, 132.7, 128.6, 128.4, 128.2, 128.2, 127.0, 113.2, 111.8, 108.9, 85.2, 83.9, 56.1, 48.9, 41.9, 26.6; HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{24}\text{NO}_4$ $[\text{M}+\text{H}]^+$ 414.1700, found 414.1701.

5-Benzoyl-1'-methyl-4-(4-nitrophenyl)-4,5-dihydro-3*H*-spiro[furan-2,3'-indolin]-2'-one (**4ai'**): Yellow solid (27% yield), m.p. 135~137 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.89~7.87 (m, 2H), 7.47~7.44 (m, 2H), 7.36~7.33 (m, 3H), 7.29~7.27 (m, 1H), 7.12~7.08 (m, 1H), 7.04~7.03 (m, 1H), 6.84 (d, $J=7.8$ Hz, 1H), 6.26 (d, $J=3.3$ Hz, 1H), 6.11~6.10 (m, 1H), 5.87 (d, $J=8.5$ Hz, 1H), 4.56~4.49 (m, 1H), 3.22 (s, 3H), 3.09 (dd, $J=12.9$, 8.7 Hz, 1H), 2.52 (dd, $J=12.9$, 8.2 Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 196.1, 174.4, 150.7, 143.4, 141.5, 135.8, 132.8, 130.99,

130.1, 129.4, 128.9, 128.7, 128.6, 128.1, 125.8, 123.2, 123.0, 110.5, 108.6, 107.9, 83.6, 83.1, 42.7, 39.5, 26.5; HRMS (ESI) calcd for $C_{25}H_{21}N_2O_5$ $[M+H]^+$ 429.1445, found 429.1445.

5-Benzoyl-4-(4-fluorophenyl)-1'-methyl-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4aj'**): White solid (38% yield), m.p. 220~222 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.95 (d, $J=7.7$ Hz, 2H), 7.52~7.42 (m, 4H), 7.39~7.26 (m, 3H), 7.08 (t, $J=7.5$ Hz, 1H), 7.00 (t, $J=8.5$ Hz, 2H), 6.80 (d, $J=7.8$ Hz, 1H), 5.71 (d, $J=9.6$ Hz, 1H), 4.31 (dd, $J=20.0, 9.5$ Hz, 1H), 3.21 (s, 3H), 2.89~2.79 (m, 1H), 2.66 (dd, $J=12.8, 8.7$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 195.2, 177.1, 162.1 (d, $J=244.3$ Hz), 143.4, 135.5, 134.2 (d, $J=3.2$ Hz), 133.7, 130.1, 130.0, 129.9, 129.8, 129.5, 128.6, 124.3, 123.5, 115.7 (d, $J=21.2$ Hz), 108.6, 86.6, 83.9, 46.6, 43.96, 26.4; HRMS (ESI) calcd for $C_{25}H_{21}FNO_3$ $[M+H]^+$ 402.1500, found 402.1503.

5-Benzoyl-4-(4-chlorophenyl)-1'-methyl-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4ak'**): White solid (44% yield), m.p. 221~223 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.72~7.71 (m, 2H), 7.50~7.44 (m, 2H), 7.38~7.26 (m, 5H), 7.14~7.08 (m, 3H), 6.86 (d, $J=7.8$ Hz, 1H), 5.94 (d, $J=7.9$ Hz, 1H), 4.31 (q, $J=7.9$ Hz, 1H), 3.23 (s, 3H), 2.98 (dd, $J=13.2, 7.5$ Hz, 1H), 2.74 (dd, $J=13.2, 8.5$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 195.5, 174.6, 143.5, 137.5, 136.4, 132.96, 132.9, 131.4, 130.1, 130.0, 128.4, 128.3, 123.3, 123.1, 108.7, 85.5, 83.5, 48.3, 42.1, 26.5; HRMS (ESI) calcd for $C_{25}H_{21}ClNO_3$ $[M+H]^+$ 418.1204, found 418.1205.

5-Benzoyl-4-(4-bromophenyl)-1'-methyl-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4al'**): White solid (33% yield), m.p. 180~182 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.69 (d, $J=7.0$ Hz, 2H), 7.49~7.44 (m, 2H), 7.38~7.30 (m, 3H), 7.24~7.20 (m, 4H), 7.14~7.10 (m, 1H), 6.86 (d, $J=7.6$ Hz, 1H), 5.96 (d, $J=7.7$ Hz, 1H), 4.31 (dd, $J=14.8, 7.2$ Hz, 1H), 3.22 (m, 3H), 3.00~2.95 (m, 1H), 2.71 (dd, $J=12.2, 8.7$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 195.9, 174.9, 143.4, 137.7, 136.2, 133.1, 131.3, 130.3, 130.2, 128.4, 128.3, 123.3, 123.2, 121.1, 108.8, 85.2, 83.6, 48.4, 42.0, 26.5; HRMS (ESI) calcd for $C_{25}H_{21}BrNO_3$ $[M+H]^+$ 462.0699, found 462.0700.

5-Benzoyl-1'-methyl-4-(*p*-tolyl)-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4am'**): White solid (38% yield), m.p. 197~199 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.72 (d, $J=7.7$ Hz, 2H), 7.50 (d, $J=7.3$ Hz, 1H), 7.45~7.38 (m, 1H), 7.38~7.32 (m, 1H), 7.31~7.26 (m, 2H), 7.22~7.15 (m, 2H), 7.14~7.07 (m, 1H), 6.88 (dd, $J=28.0, 7.8$ Hz, 3H), 5.93 (d, $J=8.3$ Hz, 1H), 4.36 (q, $J=8.4$ Hz, 1H), 3.22 (s, 3H), 3.07 (dd, $J=12.8, 9.0$ Hz, 1H), 2.65 (dd, $J=12.9, 8.1$ Hz, 1H), 2.19 (s, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 196.3, 175.1, 143.97, 137.2, 136.9, 135.7, 132.8, 132.0, 130.3, 129.2, 128.9, 128.9, 128.4, 123.7, 123.3, 108.9, 85.7, 84.0, 49.1, 42.3, 26.7, 21.2; HRMS (ESI) calcd for $C_{26}H_{24}NO_3$ $[M+H]^+$ 398.1751, found 398.1753.

5-Benzoyl-4-(4-methoxyphenyl)-1'-methyl-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4an'**): White solid (33% yield), m.p. 189~191 °C; 1H NMR (600 MHz,

$CDCl_3$) δ : 7.70 (d, $J=7.4$ Hz, 2H), 7.50 (d, $J=7.1$ Hz, 1H), 7.45~7.39 (m, 1H), 7.37~7.32 (m, 1H), 7.32~7.27 (m, 2H), 7.25~7.20 (m, 2H), 7.11 (t, $J=7.5$ Hz, 1H), 6.85 (d, $J=7.7$ Hz, 1H), 6.65 (d, $J=8.6$ Hz, 2H), 5.93 (d, $J=8.3$ Hz, 1H), 4.34 (q, $J=8.3$ Hz, 1H), 3.69 (s, 3H), 3.23 (s, 3H), 3.03 (dd, $J=12.9, 8.5$ Hz, 1H), 2.67 (dd, $J=13.0, 8.4$ Hz, 1H); ^{13}C NMR (151 MHz, $CDCl_3$) δ : 196.1, 174.8, 158.5, 143.5, 136.6, 132.7, 131.6, 130.5, 129.97, 129.7, 128.4, 128.1, 123.3, 122.97, 113.6, 108.6, 85.3, 83.5, 55.2, 48.3, 42.2, 26.5; HRMS (ESI) calcd for $C_{26}H_{24}NO_4$ $[M+H]^+$ 414.1700, found 414.1701.

5-Benzoyl-4-(3-bromophenyl)-1'-methyl-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4ao'**): White solid (23% yield), m.p. 206~208 °C; 1H NMR (600 MHz, $CDCl_3$) δ : 7.71 (d, $J=7.4$ Hz, 2H), 7.51~7.43 (m, 2H), 7.41~7.29 (m, 5H), 7.20 (d, $J=7.8$ Hz, 1H), 7.11 (t, $J=7.3$ Hz, 1H), 7.02 (t, $J=7.8$ Hz, 1H), 6.85 (d, $J=7.7$ Hz, 1H), 5.94 (d, $J=7.9$ Hz, 1H), 4.28 (q, $J=8.0$ Hz, 1H), 3.22 (s, 3H), 3.01 (dd, $J=13.0, 8.1$ Hz, 1H), 2.70 (dd, $J=13.1, 8.3$ Hz, 1H); ^{13}C NMR (151 MHz, $CDCl_3$) δ : 195.5, 174.6, 143.5, 141.1, 136.4, 132.98, 131.8, 131.3, 130.1, 130.1, 129.9, 128.4, 128.3, 127.2, 123.3, 123.1, 122.1, 108.7, 85.3, 83.5, 48.5, 41.6, 26.5; HRMS (ESI) calcd for $C_{25}H_{21}BrNO_3$ $[M+H]^+$ 462.0699, found 462.0700.

5-Benzoyl-4-(2-bromophenyl)-1'-methyl-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4ap'**): White solid (29% yield), m.p. 203~205 °C; 1H NMR (600 MHz, $CDCl_3$) δ : 7.71 (d, $J=7.5$ Hz, 2H), 7.56 (d, $J=7.6$ Hz, 2H), 7.43~7.34 (m, 2H), 7.30~7.26 (m, 3H), 7.22~7.11 (m, 2H), 6.96~6.90 (m, 1H), 6.86 (d, $J=7.7$ Hz, 1H), 6.16 (d, $J=8.5$ Hz, 1H), 4.89 (dd, $J=18.4, 8.1$ Hz, 1H), 3.23 (s, 3H), 3.19~3.13 (m, 1H), 2.52 (dd, $J=12.4, 7.3$ Hz, 1H); ^{13}C NMR (151 MHz, $CDCl_3$) δ : 196.3, 174.7, 143.4, 136.7, 136.2, 132.8, 132.3, 131.2, 130.1, 129.7, 128.7, 128.3, 128.2, 127.7, 125.3, 123.4, 123.1, 108.6, 83.4, 81.9, 47.4, 40.7, 26.5; HRMS (ESI) calcd for $C_{25}H_{21}BrNO_3$ $[M+H]^+$ 462.0699, found 462.0700.

5-Benzoyl-1'-methyl-4-(naphthalen-1-yl)-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4aq'**): White solid (38% yield), m.p. 197~199 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.71~7.60 (m, 6H), 7.54~7.39 (m, 2H), 7.35~7.34 (m, 4H), 7.21~7.12 (m, 3H), 6.87 (d, $J=7.8$ Hz, 1H), 6.04 (d, $J=8.2$ Hz, 1H), 4.54 (q, $J=8.3$ Hz, 1H), 3.25 (s, 3H), 3.20 (dd, $J=12.9, 8.7$ Hz, 1H), 2.76 (dd, $J=13.0, 8.2$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 195.9, 174.8, 143.5, 136.5, 135.97, 133.1, 132.6, 132.4, 131.5, 130.0, 128.4, 128.1, 127.97, 127.8, 127.5, 126.6, 125.9, 125.7, 123.3, 123.0, 108.7, 85.3, 83.7, 49.1, 41.9, 26.5; HRMS (ESI) calcd for $C_{29}H_{24}NO_3$ $[M+H]^+$ 434.1751, found 434.1753.

5-Benzoyl-1'-methyl-4-(naphthalen-2-yl)-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4ar'**): White solid (26% yield), m.p. 233~235 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.71~7.60 (m, 6H), 7.56~7.49 (m, 2H), 7.38~7.31 (m, 4H), 7.21~7.11 (m, 3H), 6.87 (d, $J=7.8$ Hz, 1H), 6.04 (d, $J=8.2$ Hz, 1H), 4.54 (q, $J=8.4$ Hz, 1H), 3.25 (s, 3H), 3.20 (dd, $J=12.9, 8.7$ Hz, 1H), 2.76 (dd, $J=12.9, 8.2$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 195.9, 174.8, 143.5, 136.5,

135.99, 133.1, 132.6, 132.4, 131.5, 130.0, 128.4, 128.1, 127.96, 127.8, 127.5, 126.6, 125.9, 125.7, 123.3, 123.0, 108.7, 85.3, 83.7, 49.1, 26.5; HRMS (ESI) calcd for $C_{29}H_{24}NO_3$ $[M+H]^+$ 434.1751, found 434.1753.

5-Benzoyl-4-(furan-2-yl)-1'-methyl-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4as'**): White solid (42% yield), m.p. 207~209 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.02~8.00 (m, 2H), 7.75 (dd, $J=8.3, 1.1$ Hz, 2H), 7.58~7.56 (m, 2H), 7.50~7.47 (m, 2H), 7.41~7.32 (m, 3H), 7.15~7.11 (m, 1H), 6.88 (d, $J=7.8$ Hz, 1H), 5.98 (d, $J=7.4$ Hz, 1H), 4.39 (dd, $J=15.6, 7.1$ Hz, 1H), 3.23 (s, 3H), 2.99 (dd, $J=13.3, 6.7$ Hz, 1H), 2.85 (dd, $J=13.3, 8.7$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 194.8, 174.6, 147.3, 146.8, 143.6, 135.96, 133.4, 131.1, 130.4, 129.6, 128.51, 128.45, 123.4, 123.3, 123.2, 108.9, 85.8, 83.5, 48.5, 41.9, 26.5; HRMS (ESI) calcd for $C_{23}H_{20}NO_4$ $[M+H]^+$ 374.1387, found 374.1388.

5-(4-Fluorobenzoyl)-1'-methyl-4-phenyl-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4ba'**): White solid (40% yield), m.p. 197~199 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.81~7.77 (m, 2H), 7.50 (d, $J=7.2$ Hz, 1H), 7.38~7.31 (m, 3H), 7.15~7.06 (m, 4H), 6.97~6.92 (m, 2H), 6.86 (d, $J=7.8$ Hz, 1H), 5.88 (d, $J=8.3$ Hz, 1H), 4.38 (q, $J=8.4$ Hz, 1H), 3.23 (s, 3H), 3.09 (dd, $J=12.9, 8.9$ Hz, 1H), 2.73~2.61 (m, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 194.7, 174.8, 165.4 (d, $J=253.2$ Hz), 143.5, 138.2, 132.93 (d, $J=3.1$ Hz), 131.3 (d, $J=9.3$ Hz), 131.33, 130.1, 128.6, 128.3, 127.1, 123.3, 123.1, 115.2 (d, $J=21.8$ Hz), 108.7, 85.5, 83.8, 48.97, 41.5, 26.5; HRMS (ESI) calcd for $C_{25}H_{21}FN-O_3$ $[M+H]^+$ 402.1500, found 402.1503.

5-(4-chlorobenzoyl)-1'-methyl-4-phenyl-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4ca'**): White solid (44% yield), m.p. 178~180 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.69 (d, $J=8.6$ Hz, 2H), 7.51~7.48 (m, 1H), 7.39~7.30 (m, 3H), 7.27~7.23 (m, 2H), 7.17~7.08 (m, 4H), 6.86 (d, $J=7.8$ Hz, 1H), 5.87 (t, $J=6.8$ Hz, 1H), 4.49~4.27 (m, 1H), 3.23 (s, 3H), 3.08 (dd, $J=12.9, 9.0$ Hz, 1H), 2.67 (dd, $J=12.9, 8.2$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 195.2, 174.8, 143.4, 139.1, 138.1, 134.8, 131.3, 130.1, 130.0, 129.8, 129.6, 129.4, 129.1, 128.6, 128.4, 128.3, 127.2, 123.3, 123.1, 108.7, 85.5, 83.8, 48.95, 41.5, 26.5; HRMS (ESI) calcd for $C_{25}H_{21}ClNO_3$ $[M+H]^+$ 418.1204, found 418.1205.

5-(4-Bromobenzoyl)-1'-methyl-4-phenyl-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4da'**): White solid (36% yield), m.p. 218~220 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.60 (t, $J=5.4$ Hz, 2H), 7.49 (d, $J=7.2$ Hz, 1H), 7.45~7.28 (m, 5H), 7.19~7.04 (m, 4H), 6.86 (d, $J=7.8$ Hz, 1H), 5.86 (d, $J=8.3$ Hz, 1H), 4.38 (q, $J=8.4$ Hz, 1H), 3.23 (s, 3H), 3.08 (dd, $J=12.9, 9.0$ Hz, 1H), 2.67 (dd, $J=12.9, 8.2$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 195.4, 174.7, 143.5, 138.1, 135.2, 132.6, 131.4, 131.3, 130.1, 130.1, 129.8, 128.6, 128.3, 127.9, 127.2, 123.3, 123.1, 108.7, 85.5, 83.8, 48.9, 41.5, 26.5; HRMS (ESI) calcd for $C_{25}H_{21}BrNO_3$ $[M+H]^+$ 462.0699, found 462.0700.

1'-Methyl-5-(4-methylbenzoyl)-4-phenyl-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4ea'**): White solid (42%

yield), m.p. 240~242 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.63 (d, $J=8.2$ Hz, 2H), 7.54~7.46 (m, 1H), 7.38~7.30 (m, 3H), 7.16~7.03 (m, 6H), 6.85 (d, $J=7.8$ Hz, 1H), 5.93 (d, $J=8.2$ Hz, 1H), 4.37 (q, $J=8.4$ Hz, 1H), 3.23 (s, 3H), 3.14~3.03 (m, 1H), 2.68 (dt, $J=23.0, 11.5$ Hz, 1H), 2.32 (s, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 195.4, 174.7, 143.5, 143.5, 138.6, 133.99, 131.5, 129.97, 128.9, 128.6, 128.6, 128.2, 126.99, 123.3, 122.96, 108.6, 85.1, 83.5, 48.99, 41.96, 26.5, 21.6; HRMS (ESI) calcd for $C_{26}H_{24}NO_3$ $[M+H]^+$ 398.1751, found 398.1753.

5-(4-Methoxybenzoyl)-1'-methyl-4-phenyl-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4fa'**): White solid (29% yield), m.p. 205~207 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.74~7.68 (m, 2H), 7.50 (d, $J=6.9$ Hz, 1H), 7.45~7.39 (m, 1H), 7.38~7.32 (m, 1H), 7.32~7.26 (m, 2H), 7.25~7.21 (m, 2H), 7.11 (dd, $J=11.0, 4.0$ Hz, 1H), 6.85 (d, $J=7.8$ Hz, 1H), 6.68~6.60 (m, 2H), 5.93 (d, $J=8.2$ Hz, 1H), 4.34 (q, $J=8.3$ Hz, 1H), 3.68 (s, 3H), 3.23 (s, 3H), 3.02 (dd, $J=13.0, 8.4$ Hz, 1H), 2.68 (dd, $J=13.0, 8.3$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 196.1, 174.7, 158.5, 143.5, 136.5, 132.7, 131.5, 130.5, 129.98, 129.7, 128.4, 128.2, 123.3, 122.98, 113.6, 108.6, 85.3, 83.5, 55.2, 48.3, 42.2, 26.5; HRMS (ESI) calcd for $C_{26}H_{24}NO_4$ $[M+H]^+$ 414.1700, found 414.1701.

5-(3-Bromobenzoyl)-1'-methyl-4-phenyl-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4ga'**): White solid (42% yield), m.p. 219~221 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.77~7.72 (m, 1H), 7.67 (d, $J=7.8$ Hz, 1H), 7.55~7.47 (m, 2H), 7.39~7.33 (m, 1H), 7.32~7.27 (m, 2H), 7.19~7.06 (m, 5H), 6.86 (d, $J=7.8$ Hz, 1H), 5.88 (d, $J=8.3$ Hz, 1H), 4.45~4.32 (m, 1H), 3.23 (s, 3H), 3.08 (dt, $J=13.0, 6.5$ Hz, 1H), 2.66 (dd, $J=12.9, 8.1$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 194.9, 174.7, 143.4, 138.2, 137.9, 135.5, 131.4, 131.2, 130.1, 129.7, 128.5, 128.4, 127.3, 127.0, 123.3, 123.1, 122.4, 108.7, 85.1, 83.8, 48.9, 41.4, 26.5; HRMS (ESI) calcd for $C_{25}H_{21}BrNO_3$ $[M+H]^+$ 462.0699, found 462.0700.

5-(3-Methoxybenzoyl)-1'-methyl-4-phenyl-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4ha'**): White solid (37% yield), m.p. 170~172 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.53~7.47 (m, 1H), 7.40~7.28 (m, 4H), 7.25~7.04 (m, 6H), 7.03~6.93 (m, 1H), 6.91~6.78 (m, 1H), 5.84 (dd, $J=87.2, 8.9$ Hz, 1H), 4.44~4.23 (m, 1H), 3.74 (d, $J=6.7$ Hz, 3H), 3.23 (s, 3H), 3.09 (dd, $J=12.9, 8.7$ Hz, 1H), 2.69 (dd, $J=12.9, 8.2$ Hz, 1H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 195.8, 174.7, 159.4, 143.5, 138.5, 137.9, 131.4, 130.0, 129.1, 128.8, 128.7, 128.2, 127.0, 123.3, 123.0, 120.9, 119.6, 112.4, 108.7, 85.2, 83.6, 55.4, 48.99, 41.8, 26.5; HRMS (ESI) calcd for $C_{26}H_{24}NO_4$ $[M+H]^+$ 414.1700, found 414.1701.

5-(2-Methoxybenzoyl)-1'-methyl-4-phenyl-4,5-dihydro-3H-spiro[furan-2,3'-indolin]-2'-one (**4ia'**): White solid (29% yield), m.p. 158~160 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.47 (d, $J=7.1$ Hz, 1H), 7.38~7.23 (m, 4H), 7.15~6.99 (m, 5H), 6.89~6.76 (m, 2H), 6.76~6.68 (m, 1H), 6.16 (d, $J=8.2$ Hz, 1H), 4.27 (q, $J=8.2$ Hz, 1H), 3.90 (d, $J=8.9$ Hz, 3H), 3.23 (s, 3H), 2.94 (dd, $J=13.0, 7.9$ Hz,

1H), 2.67 (dd, $J=13.0, 8.6$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 196.9, 174.7, 157.3, 143.6, 139.9, 133.3, 131.8, 130.8, 129.8, 128.5, 128.0, 127.2, 126.8, 123.3, 122.8, 120.8, 110.8, 108.6, 88.1, 82.8, 55.7, 47.8, 42.9, 26.4; HRMS (ESI) calcd for $\text{C}_{26}\text{H}_{24}\text{NO}_4$ $[\text{M} + \text{H}]^+$ 414.1700, found 414.1701.

3.3 General procedure for the synthesis of 5'

Hydroxylamine hydrochloride (106 mg, 1.5 mmol) and sodium acetate (100 mg, 1.2 mmol) were added to a solution of **4a1'** (92 mg, 0.20 mmol) in methanol (5 mL). Then, the reaction mixture was refluxed in a heating mantle for 10 h. After cooling to room temperature, water (3 mL) was added to quench the reaction. The solution was extracted with CH_2Cl_2 (10 mL $\times$ 3). The combined organic layer was washed with brine, dried over Na_2SO_4 and concentrated under reduced pressure. The crude reaction mixture was purified via column chromatography on silica gel to afford pure product **5'** as colorless solid in 96% yield.

4-(4-Bromophenyl)-5-((*E*)-(hydroxyimino)(phenyl)methyl)-1'-methyl-4,5-dihydro-3*H*-spiro[furan-2,3'-indolin]-2'-one (**5'**): White solid (96% yield), m.p. 183~184 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 11.71 (s, 1H), 7.65 (d, $J=7.2$ Hz, 1H), 7.52 (d, $J=8.5$ Hz, 2H), 7.41~7.36 (m, 3H), 7.26~7.22 (m, 1H), 7.17~7.11 (m, 5H), 7.07 (d, $J=7.8$ Hz, 1H), 5.92 (d, $J=7.2$ Hz, 1H), 4.54~4.49 (m, 1H), 3.17 (s, 3H), 2.82 (dd, $J=13.7, 9.6$ Hz, 1H), 2.58 (dd, $J=13.7, 4.7$ Hz, 1H), 2.03~1.96 (m, 1H); ^{13}C NMR (101 MHz, DMSO) δ : 174.8, 156.4, 143.5, 141.7, 134.5, 131.95, 131.5, 130.8, 130.3, 130.1, 129.0, 128.6, 127.6, 124.3, 123.5, 120.2, 109.6, 82.6, 81.4, 46.5, 41.96, 26.7; HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{22}\text{BrN}_2\text{O}_3$ $[\text{M} + \text{H}]^+$ 477.0808, found 477.0809.

Supporting information ^1H NMR and ^{13}C NMR spectra of **4'** and **5'**. The Supporting Information is available free of charge via the internet at <http://sioc-journal.cn/>.

References and note

- [1] (a) Rottmann, M.; McNamara, C.; Yeung, B. K. S.; Lee, M. C. S.; Zou, B.; Russell, B.; Seitz, P.; Plouffe, D. M.; Dharia, N. V.; Tan, J.; Cohen, S. B.; Spencer, K. R.; Gonzalez-Paez, G. E.; Lakshminarayana, S. B.; Goh, A.; Suwanarusk, R.; Jegla, T.; Schmitt, E. K.; Beck, H. P.; Brun, R.; Nosten, F.; Renia, L.; Dartois, V.; Keller, T. H.; Fidock, D. A.; Winzeler, E. A.; Diagana, T. T. *Science* **2010**, 329, 1175.
(b) Singh, G. S.; Desta, Z. Y. *Chem. Rev.* **2012**, 112, 6104.
(c) Zheng, Y. J.; Tice, C. M.; Singh, S. B. *Bioorg. Med. Chem. Lett.* **2014**, 24, 3673.
(d) Yu, B.; Yu, D. Q.; Liu, H. M. *Eur. J. Med. Chem.* **2015**, 97, 673.
(e) Ye, N.; Chen, H.; Wold, E. A.; Shi, P.-Y.; Zhou, J. *ACS Infect. Dis.* **2016**, 2, 382.
(f) Mei, G. J.; Shi, F. *Chem. Commun.* **2018**, 54, 6607.
- [2] (a) Peddibhotla, S. *Curr. Bioact. Compd.* **2009**, 5, 20.
(b) Cui, B.-D.; Zuo, J.; Zhao, J.-Q.; Zhou, M.-Q.; Wu, Z.-J.; Zhang, X.-M.; Yuan, W.-C. *J. Org. Chem.* **2014**, 79, 5305.
- [3] (a) Vintonyak, V. V.; Warburg, K.; Kruse, H.; Grimme, S.; Hübel, K.; Rauh, D.; Waldmann, H. *Angew. Chem., Int. Ed.* **2010**, 49, 5902.
(b) Jiang, X.; Cao, Y.; Wang, Y.; Liu, L.; Shen, F.; Wang, R. *J. Am. Chem. Soc.* **2010**, 132, 15328.
(c) Crosignani, S.; Jorand-Lebrun, C.; Page, P.; Campbell, G.; Colovray, V.; Missotten, M.; Humbert, Y.; Cleva, C.; Arrighi, J.-F.; Gaudet, M.; Johnson, Z.; Ferro, P.; Chollet, A. *ACS Med. Chem. Lett.* **2011**, 2, 644.
(d) Rana, S.; Natarajan, A. *Org. Biomol. Chem.* **2013**, 11, 244.
- [4] Franz, A. K.; Dreyfuss, P. D.; Schreiber, S. L. *J. Am. Chem. Soc.* **2007**, 129, 1020.
- [5] Ghosh, A.; Carter, R. G. *Angew. Chem., Int. Ed.* **2019**, 58, 681.
- [6] (a) Alcaide, B.; Almendros, P.; Rodriguez-Acebes, R. *J. Org. Chem.* **2006**, 71, 2346.
(b) Li, J.; Liu, Y. J.; Li, C. J.; Jia, X. *Chem.-Eur. J.* **2011**, 17, 7409.
(c) Hanhan, N. V.; Jones, N. R.; Tran, N. T.; Franz, A. K. *Angew. Chem., Int. Ed.* **2012**, 51, 989.
(d) Yang, L.; Xie, P.; Li, E.; Li, X.; Huang, Y.; Chen, R. *Org. Biomol. Chem.* **2012**, 10, 7628.
(e) Lian, Z.; Shi, M. *Eur. J. Org. Chem.* **2012**, 2012, 581.
(f) Mei, L.-Y.; Wei, Y.; Xu, Q.; Shi, M. *Organometallics* **2013**, 32, 3544.
(g) Tang, Z.; Liu, Z.; An, Y.; Jiang, R.; Zhang, X.; Li, C.; Jia, X.; Li, J. *J. Org. Chem.* **2016**, 81, 9158.
(h) Silvi, M.; Chatterjee, I.; Liu, Y.-K.; Melchiorre, P. *Angew. Chem., Int. Ed.* **2013**, 52, 10780.
(i) Zhao, B.-L.; Du, D.-M. *Adv. Synth. Catal.* **2016**, 358, 3992.
(j) Zhu, Y.-S.; Wang, W.-B.; Yuan, B.-B.; Li, Y.-N.; Wang, Q.-L.; Bu, Z.-W. *Org. Biomol. Chem.* **2017**, 15, 984.
(k) Zhang, J.-H.; Wang, R.-B.; Li, D.-F.; Zhao, L.-M. *ACS Omega* **2017**, 2, 7022.
- [7] (a) Lee, S.; Hartwig, J. F. *J. Org. Chem.* **2001**, 66, 3402.
(b) Jones, K.; Wilkinson, J. *Chem. Commun.* **1992**, 1767.
(c) Overman, L. E.; Rosen, M. D. *Angew. Chem., Int. Ed.* **2000**, 39, 4596.
(d) Kyei, A. S.; Tchabanenko, K.; Baldwin, J. E.; Adlington, R. M. *Tetrahedron Lett.* **2004**, 45, 8931.
(e) Jaegli, S.; Dufour, J.; Wei, H.-L.; Piou, T.; Duan, X.-H.; Vors, J.-P.; Neuville, L.; Zhu, J.-P. *Org. Lett.* **2010**, 12, 4498.
(f) Yin, B.-L.; Lai, J.-Q.; Zhang, Z.-R.; Jiang, H.-F. *Adv. Synth. Catal.* **2011**, 353, 1961.
- [8] (a) Hua, Y.-Z.; Lu, L.-J.; Huang, P.-J.; Wei, D.-H.; Tang, M.-S.; Wang, M.-C.; Chang, J.-B. *Chem.-Eur. J.* **2014**, 20, 12394.
(b) Hua, Y.-Z.; Han, X.-W.; Yang, X.-C.; Song, X.; Wang, M.-C.; Chang, J.-B. *J. Org. Chem.* **2014**, 79, 11690.
(c) Hua, Y.-Z.; Yang, X.-C.; Liu, M.-M.; Song, X.; Wang, M.-C.; Chang, J.-B. *Macromolecules* **2015**, 48, 1651.
(d) Wang, X.-W.; Hua, Y.-Z.; Wang, M.-C. *J. Org. Chem.* **2016**, 81, 9227.
(e) Shao, N.; Luo, Y.-Y.; Lu, H.-J.; Hua, Y.-Z.; Wang, M.-C. *Tetrahedron* **2018**, 74, 2130.
(f) Hua, Y.-Z.; Han, X.-W.; Huang, L.-H.; Wang, M.-C. *Chin. J. Org. Chem.* **2018**, 38, 237 (in Chinese).
(g) 华远照, 韩兴旺, 黄利华, 王敏灿, 有机化学, **2018**, 38, 237.
(h) Hua, Y.-Z.; Chen, J.-W.; Yang, H.; Wang, M.-C. *J. Org. Chem.* **2018**, 83, 1160.
(i) Zhang, Z.-F.; Yang, X.-C.; Lu, H.-J.; Wang, M.-C. *Eur. J. Org. Chem.* **2018**, 785.
(j) Gao, Y.-Y.; Hua, Y.-Z.; Wang, M.-C. *Adv. Synth. Catal.* **2018**, 360, 80.
(k) Liu, S.; Gao, W.-C.; Miao, Y.-H.; Wang, M.-C. *J. Org. Chem.* **2019**, 84, 2652.
- [9] (a) Song, X.; Liu, J.; Liu, M.-M.; Wang, X.; Zhang, Z.-F.; Wang, M.-C.; Chang, J. *Tetrahedron* **2014**, 70, 5468.
(b) Hua, Y.-Z.; Liu, M.-M.; Huang, P.-J.; Song, X.; Wang, M.-C.; Chang, J.-B. *Chem.-Eur. J.* **2015**, 21, 11994.
(c) Tao, B.-K.; Yang, H.; Hua, Y.-Z.; Wang, M.-C. *Org. Biomol. Chem.* **2019**, 17, 4301.
(d) Miao, Y.-H.; Hua, Y.-Z.; Wang, M.-C. *Org. Biomol. Chem.* **2019**, 17, 7172.
(e) Liu, M.-M.; Yang, X.-C.; Hua, Y.-Z.; Chang, J.-B.; Wang, M.-C.

Org. Lett. **2019**, *21*, 7089.

(f) Yi, Y.; Hua, Y.-Z.; Lu, H.-J.; Liu, L.-T.; Wang, M.-C. *Org. Lett.* **2020**, *22*, 2527.

(g) Liu, M.-M.; Yang, X.-C.; Hua, Y.-Z.; Chang, J.-B.; Wang, M.-C. *Org. Lett.* **2019**, *21*, 2111.

(h) Yang, X.-C.; Liu, M.-M.; Mathey, F.; Yang, H.; Hua, Y.-Z.; Wang, M.-C. *J. Org. Chem.* **2019**, *84*, 7762.

[10] Guo, Y.-J.; Guo, X.; Kong, D.-Z.; Lu, H.-J.; Liu, L.-T.; Hua, Y.-Z.; Wang, M.-C. *J. Org. Chem.*, **2020**, *85*, 4195. The minor diastereomers in this article are obtained in 23%~44% isolated yields with about 1 : 1 *dr* value in the current study.

[11] CCDC 1987133 (**5'**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from the Cambridge Crystallographic Data Centre.

(Li, L.; Fan, Y.)