Indan-f-amphox: 可用于铱催化 β -芳基 β -酮酸酯 不对称加氢反应的 PNN 配体

古国贤^a 胡 杨^b 刘绍东^c 董秀琴^{*,b} 张绪穆^{*,d}

(^a 河北工业大学化工学院 天津市本质安全化工技术重点实验室 天津 300130)

(^b 武汉大学化学与分子科学学院 武汉 430072)

(^c 美国新泽西州立大学罗格斯分校化学系 新泽西州皮斯卡塔维 08854 美国)

(^d 南方科技大学化学系 广东深圳 518055)

摘要 从 Ug's 胺出发通过 4 步化学反应合成了一种新的手性 PNN 三齿配体 indan-f-amphox (该配体为 f-amphox 的姊妹配体), 并将其用于铱催化的 β -芳基 β -酮酸酯的不对称氢化反应, 结果显示 Ir(III)-indan-f-amphox 催化剂对该类反应具有非常高的反应活性和立体选择性 (对应选择性最高 99%, 转化数高达 10000), 适用于多种 β -芳基 β -酮酸酯。

关键词 PNN 配体; 不对称氢化; β -酮酸酯; f-amphox

Indan-f-amphox: An Efficient PNN Ligand for Iridium-Catalyzed Asymmetric Hydrogenation of β -Aryl β -Ketoesters

Gu, Guoxian^a Hu, Yang^b Liu, Shaodong^c Dong, Xiu-Qin^{*,b} Zhang, Xumu^{*,d}

(^a School of Chemical Engineering and Technology, Tianjin Key Laboratory of Chemical Process Safety, Hebei University of Technology, Tianjin 300130, China)

(^b College of Chemistry and Molecular Science, Wuhan University, Wuhan 430072, China)

(^c Department of Chemistry, Rutgers, The State University of New Jersey, Piscataway, NJ, 08854, USA)

(^d Department of Chemistry, Southern University of Science and Technology, Shenzhen, Guangdong 518055, China)

Abstract A new chiral tridentate PNN ligand, indan-f-amphox (a sister ligand of f-amphox), was synthesized and applied in iridium-catalyzed asymmetric hydrogenation of β -aryl β -ketoesters. A wide range of β -aryl β -ketoesters are reduced using an Ir(III)-indan-f-amphox catalyst with excellent enantioselectivities and reactivities (up to 99% ee, TON=10000).

Keywords PNN ligand; asymmetric hydrogenation; β -ketoesters; f-amphox

1 Introduction

Enantiopure compounds are important building blocks in the fields of pharmaceutical and agrochemical industries due to their distinct biological activities.^[1] In principle, the preparation of enantiopure compounds through catalytic asymmetric hydrogenation (AH) of readily available unsaturated compounds such as olefins, ketones and imines is one of the most straightforward and operationally simple approaches.^[2] Although many chiral catalysts have been developed in this field, there is no universal catalyst suitable for many substrates.^[3] Development of readily available new ligand systems is the key factor for asymmetric cata-

lysis.

In many catalysts, the M-NH system has proved to be a key moiety for achieving high catalytic activities and excellent enantioselectivities in the hydrogenation of ketones and imines.^[4] In 1995, Noyori and co-workers^[5] reported ruthenium bisphosphine/diamine system, which showed high catalytic activity for hydrogenation of simple ketones. In this system, a proton from the NH₂ ligand and a hydride from the Ru center interact with the C=O bond of ketone via a six-membered pericyclic Ru-NH transition state, and this "NH effect" has resulted in the high catalytic efficiency of simple aromatic ketones.^[6] In 2011, Zhou and co-workers^[7] have made a breakthrough using catalysts with

* Corresponding authors. E-mail: zhangxm@sustc.edu.cn; xiuqindong@whu.edu.cn

Received October 28, 2019; revised November 28, 2019; published online December 11, 2019.

Project supported by the Shenzhen Free Exploration Fund (No. JCYJ20170817104853520).

深圳市科创委自由探索基金(No. JCYJ20170817104853520)资助项目.

Ir-NH species for hydrogenation of simple ketones. Introducing a coordination pyridine group to the SpiroAP ligand can inhibit dimerization, which is crucial important for the exceptionally high stability and activity in hydrogenation reaction. The resulting modified ligand SpiroPAP afforded chiral alcohols products in the hydrogenation of ketones with up to 99.9% *ee* and TONs as high as 4550000. In 1998, Zhang and co-workers^[8] developed a new tridentate phosphine free ligand, bis(oxazolinylmethyl)amine (ambox), and successfully applied the *in situ* generated Ru(II)-ambox complex in the transfer hydrogenation of simple ketones. M-NH was demonstrated to be an important element for the hydrogenation of aromatic ketones. In 2007, indan-ambox was successfully prepared and applied in direct AH of simple ketones achieving high enantioselectivities and decent activity.^[9] Based on our success with chiral ambox systems, more electron donating ligands were designed and synthesized to achieve higher activity. Lately, (*S_C,S_C,R_{FC}*)-f-amphox was reported by our group and it was also successfully applied in AH of simple and functionalized ketones (Scheme 1).^[10]

Scheme 1 Evolution of NH-tridentate ligands in Zhang's group

In Ru-NH catalyzed hydrogenation of β -alkyl β -ketoesters, Ru(II)-BINAP/diamine system offered excellent *ee* and turnovers under an acidic condition. However, for hydrogenation of β -aryl β -ketoesters, an important reaction for synthesizing key drug intermediates such as duloxetine and atomoxetine, much lower *ee*'s and turnovers were observed.^[2c,11] In 2012, Zhou *et al.*^[12] reported an Ir-NH species for AH of β -aryl β -ketoesters in the Ir-SpiroPAP system with excellent *ee*'s. A slight drawback is that the ligand synthesis is tedious. Considering the success of these Ir-NH catalyzed systems, we envisioned that Ir/f-amphox catalyst could also promote the AH of β -ketoesters.

Here we report our achievements in the synthesis of another steric hindered and electron-rich tridentate ligand, [(*R_C,S_C*)-*S_C,R_{FC}*]-indan-f-amphox, which has a deeper chi-

ral concave pocket around the metal center than its sister ligand of (*S_C,S_C,R_{FC}*)-f-amphox. As shown in Scheme 2, the synthesis of indan-f-amphox started from functionalization of (*S*)-Ugi's amine **1** to obtain (*S_C,R_{FC}*)-**2** based on previous literature.^[13] Compound (*S_C,R_{FC}*)-**2** reacted with 2-chloromethyloxazoline, which was obtained from chiral *cis*-amino indanol and 2-chloroacetonitrile, to afford indan-f-amphox with an overall yield of 23%. This short synthetic route with four steps allows easy modification and both enantiomers are readily to be prepared.

Scheme 2 Ligand design and synthesis

2 Results and discussion

Our initial study began with ethyl 3-oxo-3-phenylpropanoate **3a** as the model substrate. Under 2 MPa of hydrogen with 0.1 mol% Ir-ligand complex generated *in situ*, a brief screening of ligands in *i*-PrOH was conducted and the results is shown in Table 1. Preliminary results displayed that indan-f-amphox (**L1**) provided moderate *ee* value (Entry 1) but with only 15% conversion. Then a screening of different solvents with Ir/indan-f-amphox was carried out and methanol was turned out to be the best solvent for this reaction, affording 96% *ee* and >99% conversion (Entry 5). It is interesting that the *i*-PrOH, which had been proved to be beneficial for Ir/f-amphox in the AH of other functionalized ketones,^[10] was proved to be negative for Ir/indan-f-amphox in this reaction (Entry 1). Non-polar or polar aprotic solvents gave low reactivities or low enantioselectivities (Entries 7~9). If the oxazoline arm is replaced by pyridine (**L3**), which is planar,^[14] *ee* value dropped dramatically to 33% (Entry 9). It is inexplicable that (*S_C,S_C,R_{FC}*)-f-amphox (**L2**), the sister ligand of [(*R_C,S_C*)-*S_C,R_{FC}*]-indan-f-amphox, did not work in either *i*-PrOH or MeOH in this reaction (Entries 2 and 10) with 0.1 mol% catalyst loading. The other Ir-NH ligand, f-ampha, which works well in AH of α -ketoesters^[15] performed poorly in this reaction (Entry 4). When the catalyst loading decreased from 0.1 to 0.01 mol%, the Ir/indan-f-amphox

also provided 99% conversion and 96% *ee* (Entry 11, S/C = 10000).

Table 1 AH of ethyl 3-oxo-3-phenylpropanoate^a

Entry	Ligand	Solvent	Conv. ^b /%	<i>ee</i> ^c /%
1	L1	<i>i</i> -PrOH	15	70
2	L2	<i>i</i> -PrOH	NR	NA
3	L3	<i>i</i> -PrOH	>99	33
4	L4	<i>i</i> -PrOH	<5%	NA
5	L1	MeOH	>99	96
6	L1	EtOH	>99	64
7	L1	CH ₂ Cl ₂	>99	10
8	L1	Toluene	NR	NA
9	L1	THF	5	60
10	L2	MeOH	NR	NA
11d	L1	MeOH	>99	96

^a Reaction conditions: 0.2 mmol **3a**, 0.05 mol% [Ir(COD)Cl]₂, 0.1 mol% ligand, 1 mol% ^tBuOK, 2.0 mL solvent, reaction time 16 h. ^b Determined by ¹H NMR analysis. ^c *ee* was determined by high performance liquid chromatography (HPLC) analysis; the absolute configuration was assigned by the comparison of the optical rotation with reported standard.^[16] ^d S/C = 10000, 24 h.

With the optimized condition in hand, a variety of methyl 3-oxo-3-arylpropanoates (**3a**~**3o**) have been reduced to provide the corresponding chiral secondary alcohols (Table 2) with Ir/indan-f-amphox at S/C = 1000. High yields (95%~99%) and excellent enantioselectivities (88%~99% *ee*) were obtained. Change of substituents on substrates has little effect on the reactivities (>99% conv.). Substrates with electron-donating groups at *meta*- and *para*- or *ortho*-position, provided the hydrogenated products with excellent *ee*'s (**4b**, **4c**, **4d**, **4k** and **4l**). In contrast, substrates with a electron-withdrawing substituent on *para*-position (**4f** and **4g**) gave product in diminished *ee*'s. The substrates with naphthyl (**4m** and **4n**) were hydrogenated smoothly with both reactivity (both >99% conv.) and enantioselectivities (96% *ee* for **4m** and 94% *ee* for **4n**). Particularly, the heterocyclic substrate (**4o**) also

proceed smoothly affording the desired hydrogenation product with 90% *ee*. Interestingly, transesterification occurs under this reaction condition, giving methyl ester products exclusively (**3a'** and **3m'**).

Table 2 Iridium-catalyzed AH of β -ketoesters^a

Entry	3	Ar	R	Product	Yield ^b /%	<i>ee</i> ^c /%
1	3a	C ₆ H ₅	Me	4a	99	96
2	3a'	C ₆ H ₅	Et	4a	98	96
3	3b	4-MeC ₆ H ₄	Me	4b	99	99
4	3c	4-EtC ₆ H ₄	Me	4c	99	96
5	3d	4-MeOC ₆ H ₄	Me	4d	99	95
6	3e	4-FC ₆ H ₄	Me	4e	96	96
7	3f	4-F ₃ CC ₆ H ₄	Me	4f	96	88
8	3g	4-NO ₂ C ₆ H ₄	Me	4g	97	90
9	3h	4-ClC ₆ H ₄	Me	4h	96	93
10	3i	2-ClC ₆ H ₄	Me	4i	99	88
11	3j	3-ClC ₆ H ₄	Me	4j	95	91
12	3k	2-MeC ₆ H ₄	Me	4k	98	95
13	3l	3-MeC ₆ H ₄	Me	4l	99	96
14	3m	1-Naphthyl	Me	4m	97	96
15	3m'	1-Naphthyl	Et	4m	99	96
16	3n	2-Naphthyl	Me	4n	97	94
17	3o	2-Thienyl	Me	4o	96	90

^a Reaction conditions: 0.2 mmol of substrate **3**, 0.05 mol% [Ir(COD)Cl]₂, 0.1 mol% ligand, 2.0 mL of MeOH, 0.1 mol% ^tBuOK, 20 h. ^b Isolated yield. ^c *ee* was determined by chiral HPLC analysis.

3 Conclusions

A new chiral tridentate ligand, indan-f-amphox, was synthesized. In contrast to its sister ligand f-amphox, indan-f-amphox showed highly reactivity and enantioselectivity in iridium catalyzed AH of β -aryl β -ketoesters. By using an Ir/indan-f-amphox complex, a variety of β -ketoesters were reduced efficiently to chiral alcohols with high optical purities (up to 96% *ee* at S/C = 10000).

4 Experimental section

4.1 General

¹H NMR (400 MHz), ¹³C NMR (101 MHz) and ³¹P NMR (162 MHz) spectra were recorded on a Bruker ADVANCE III spectrometer with CDCl₃ as solvent and tetramethylsilane (TMS) as internal standard. Chemical shifts (δ) were reported up field to TMS for ¹H NMR and relative to CDCl₃ (δ 77.0) for ¹³C NMR. HRMS was recorded on APEXII and ZAB-HS spectrometer. HPLC analyses were

performed using an Agilent 1260 Infinity II instrument. Column chromatography was performed with silica gel Merck 60 (300~400 mesh). All reactions and manipulations, which are sensitive to moisture or air, were performed in an argon-filled glove box or using standard schlenk techniques. Aromatic ketones were purchased from commercial suppliers and purified by simple distillation or flash column chromatography prior to use. Anhydrous MeOH, EtOH, DCM, ⁱPrOH, THF and toluene were purchased from Sigma-Adrich. [Ir(COD)Cl]₂ was prepared according to the literature.^[17]

2-Chloromethylloxazoline was synthesized following the procedure reported by Ref. [18] and (*S,R*_{FC})-**2** was synthesized by the method reported in Ref. [11].

4.2 Synthesis of ((*R*_C,*S*_C)-*S*_C,*R*_{FC})-indan-f-amphox

(*S,R*)-**3** (500 mg, 1.20 mmol), K₂CO₃ (1.66 g, 12.0 mmol) and oxazoline (274 mg, 1.33 mmol) were placed in a round bottom flask, and purged by nitrogen twice. 15 mL of anhydrous acetonitrile was added by using syringe. Then the reaction was refluxed overnight. The solvent was removed under vacuum. Crude product was purified by column chromatography with DCM/MeOH (*V* : *V* = 99 : 1) as eluent to give 230 mg of pure product as yellow solid (33% yield). m.p. 210.7~212.4 °C; ¹H NMR (400 MHz, CDCl₃) δ: 1.40 (d, *J* = 6.6 Hz, 3H), 3.07~2.99 (m, 2H), 3.18 (d, *J* = 17.9 Hz, 1H), 3.32 (dd, *J* = 17.9, 6.9 Hz, 1H), 3.74 (d, *J* = 10.0 Hz, 1H), 4.02~3.90 (m, 7H), 4.26 (t, *J* = 2.4 Hz, 1H), 4.43 (s, 1H), 5.10~5.06 (m, 1H), 5.36 (d, *J* = 7.9 Hz, 1H), 7.58~7.05 (m, 14H); ¹³C NMR (101 MHz, CDCl₃) δ: 19.15, 39.62, 43.17, 50.65, 50.74, 69.02, 69.66, 69.72, 71.24, 75.20, 75.28, 76.22, 83.00, 96.87, 97.11, 125.18, 125.50, 127.31, 128.07, 128.15, 128.20, 128.28, 128.37, 129.04, 132.58, 132.77, 134.76, 134.97, 137.10, 139.77, 142.07, 166.02; ³¹P NMR (162 MHz, CDCl₃) δ: -25.07; HRMS calcd for C₃₅H₃₄N₂OPFe 585.1753, found 585.1750.

4.3 Procedure for asymmetric hydrogenation of β-ketoesters

4.3.1 General procedure with S/C = 1000

In an argon-filled glove box, a 4.0 mL vial was charged with the metal precursor [Ir(COD)Cl]₂ (3.4 mg, 5.0 × 10⁻³ mmol), indan-f-amphox (6.1 mg, 10.5 × 10⁻³ mmol) and anhydrous MeOH (1.0 mL). The mixture was stirred for 1.0 h at 25 °C, giving an orange red solution. The resulting solution (20 μL) and solution of ^tBuOK in MeOH (20 μL, *c* = 0.1 mol/L) were transferred by a syringe into a 5.0 mL vial charged with β-keto esters (0.2 mmol) in 1.0 mL of anhydrous MeOH. The vial was transferred to an autoclave, which was then charged with 2 MPa of H₂ and stirred at room temperature for 16 h. The hydrogen gas was released slowly in a well-ventilated hood and the solution was passed through a short column of silica gel to remove the metal complex. The crude product was analyzed by ¹H NMR for conversion and chiral HPLC for *ee* values.

Methyl (*R*)-3-hydroxy-3-phenylpropanoate (**4a**).^[19] Colorless oil, 36 mg, >99% conv., 99% yield, 96% *ee*. [α]_D²² +31.4 (*c* 1.0, MeOH); ¹H NMR (400 MHz, CDCl₃) δ:

7.39~7.36 (m, 3H), 7.35~7.24 (m, 2H), 5.14~5.11 (m, 1H), 3.71 (s, 3H), 3.34 (d, *J* = 3.2 Hz, 1H), 2.80~2.67 (m, 2H); ¹³C NMR (126 MHz, CDCl₃) δ: 172.79, 142.55, 128.58, 127.85, 125.68, 70.32, 51.94, 43.22; The enantiomeric excess was determined by HPLC on Chiracel OD-H column, 210 nm, 27 °C, *n*-hexane/*i*-PrOH (*V* : *V* = 95 : 5); flow 1.0 mL/min; *t*_{R1} = 16.2 min (*S*), *t*_{R2} = 26.1 min (*R*).

Methyl (*R*)-3-hydroxy-3-(*p*-tolyl)propanoate (**4b**).^[20] Colorless oil, 39 mg, >99% conv., 99% yield, 99% *ee*. [α]_D²² +27.3 (*c* 1.0, MeOH); ¹H NMR (400 MHz, CDCl₃) δ: 7.26 (d, *J* = 7.9 Hz, 2H), 7.16 (d, *J* = 7.8 Hz, 2H), 5.10 (dd, *J* = 9.1, 3.8 Hz, 1H), 3.72 (s, 3H), 3.17 (s, 1H), 2.77 (dd, *J* = 16.3, 9.2 Hz, 1H), 2.69 (dd, *J* = 16.3, 3.8 Hz, 1H), 2.34 (s, 3H); ¹³C NMR (101 MHz, CDCl₃) δ: 172.83, 139.56, 137.57, 129.25, 125.61, 70.20, 51.90, 43.18, 21.14. The enantiomeric excess was determined by HPLC on Chiracel AS-3 column, 210 nm, 27 °C, *n*-hexane/*i*-PrOH (*V* : *V* = 99 : 1); flow 1.0 mL/min; *t*_{R1} = 23.9 min (*S*), *t*_{R2} = 27.8 min (*R*).

Methyl (*R*)-3-(4-ethylphenyl)-3-hydroxypropanoate (**4c**): Colorless oil, 41 mg, >99% conv., 98% yield, 96% *ee*. [α]_D²² +17.9 (*c* 1.6, MeOH); ¹H NMR (400 MHz, CDCl₃) δ: 7.29 (d, *J* = 7.9 Hz, 2H), 7.19 (d, *J* = 7.9 Hz, 2H), 5.11 (dd, *J* = 9.3, 3.6 Hz, 1H), 3.72 (s, 3H), 3.18 (s, 1H), 2.8~2.61 (m, 4H), 1.23 (t, *J* = 7.6 Hz, 3H); ¹³C NMR (126 MHz, CDCl₃) δ: 172.85, 143.95, 139.80, 128.07, 125.70, 70.23, 51.91, 43.18, 28.55, 15.60. The enantiomeric excess was determined by HPLC on Chiracel AS-H column, 210 nm, 40 °C, *n*-hexane/*i*-PrOH (*V* : *V* = 95 : 5); flow 0.8 mL/min; *t*_{R1} = 12.5 min (*R*), *t*_{R2} = 14.6 min (*S*).

Methyl (*R*)-3-hydroxy-3-(4-methoxyphenyl)propanoate (**4d**): Colorless oil, 42 mg, >99% conv., 99% yield, 96% *ee*. [α]_D²² +19.3 (*c* 1.3, MeOH); ¹H NMR (400 MHz, CDCl₃) δ: 7.31~7.28 (m, 2H), 6.90~6.87 (m, 2H), 5.09 (dd, *J* = 9.3, 3.7 Hz, 1H), 3.80 (d, *J* = 0.8 Hz, 3H), 3.72 (d, *J* = 0.9 Hz, 3H), 3.17 (s, 1H), 2.80~2.65 (m, 2H); ¹³C NMR (126 MHz, CDCl₃) δ: 172.80, 159.20, 134.73, 126.96, 113.92, 69.96, 55.29, 51.89, 43.20. The enantiomeric excess was determined by HPLC on Chiracel OD-H column, 210 nm, 40 °C, *n*-hexane/*i*-PrOH (*V* : *V* = 95 : 5); flow 0.8 mL/min; *t*_{R1} = 19.5 min (*R*), *t*_{R2} = 20.7 min (*S*).

Methyl (*R*)-3-(4-fluorophenyl)-3-hydroxypropanoate (**4e**): Colorless oil, 38 mg, >99% conv., 96% yield, 96% *ee*. [α]_D²² +12.8 (*c* 1.3, MeOH); ¹H NMR (400 MHz, CDCl₃) δ: 7.36~7.33 (m, 2H), 7.06~7.01 (m, 2H), 5.14~5.10 (m, 1H), 3.72 (s, 3H), 3.36 (d, *J* = 3.2 Hz, 1H), 2.78~2.66 (m, 2H); ¹³C NMR (126 MHz, CDCl₃) δ: 172.72, 162.29 (d, *J* = 245.8 Hz), 138.24 (d, *J* = 3.1 Hz), 127.38 (d, *J* = 8.2 Hz), 115.41 (d, *J* = 21.5 Hz), 69.68, 51.98, 43.16. The enantiomeric excess was determined by HPLC on Chiracel OD-H column, 210 nm, 40 °C, *n*-hexane/*i*-PrOH (*V* : *V* = 95 : 5); flow 0.8 mL/min; *t*_{R1} = 13.2 min (*S*), *t*_{R2} = 15.3 min (*R*).

Methyl (*R*)-3-hydroxy-3-(4-(trifluoromethyl)phenyl)propanoate (**4f**): White solid, 48 mg, >99% conv., 97% yield, 88% *ee*. m.p. 42.3~44.2 °C (lit.^[20] 42.2~43.7 °C); [α]_D²² +11.9 (*c* 1.6, MeOH); ¹H NMR (400 MHz, CDCl₃) δ:

7.63~7.61 (m, 2H), 7.51~7.49 (m, 2H), 5.20 (t, $J=6.3$ Hz, 1H), 3.74 (s, 3H), 2.74 (d, $J=6.4$ Hz, 2H); ^{13}C NMR (126 MHz, CDCl_3) δ : 172.61, 146.31, 130.02 (d, $J=32.5$ Hz), 125.97, 125.55 (q, $J=3.8$ Hz), 69.65, 52.09, 42.89. The enantiomeric excess was determined by HPLC on Chiralcel AS-H column, 210 nm, 40 °C, *n*-hexane/*i*-PrOH ($V:V=95:5$); flow 0.8 mL/min; $t_{\text{R}1}=11.9$ min (R), $t_{\text{R}2}=13.3$ min (S).

Methyl (R)-3-hydroxy-3-(4-nitrophenyl)propanoate (**4g**):^[21] Colorless oil, 44 mg, >99% conv., 97% yield, 88% ee. $[\alpha]_{\text{D}}^{22}+17.1$ (c 1.5, MeOH); ^1H NMR (400 MHz, CDCl_3) δ : 8.24~8.20 (m, 2H), 7.59~7.55 (m, 2H), 5.25 (dd, $J=8.1, 4.6$ Hz, 1H), 3.75 (s, 3H), 2.80~2.70 (m, 2H); ^{13}C NMR (126 MHz, CDCl_3) δ : 172.41, 149.58, 147.46, 126.50, 123.83, 69.35, 52.19, 42.73. The enantiomeric excess was determined by HPLC on Chiralcel ID column, 210 nm, 40 °C, *n*-hexane/*i*-PrOH ($V:V=90:10$); flow 0.8 mL/min; $t_{\text{R}1}=23.4$ min (R), $t_{\text{R}2}=29.4$ min (S).

Methyl (R)-3-(4-chlorophenyl)-3-hydroxypropanoate (**4h**):^[20] Colorless oil, 41 mg, >99% conv., 96% yield, 93% ee. $[\alpha]_{\text{D}}^{22}+13.0$ (c 1.3, MeOH); ^1H NMR (400 MHz, CDCl_3) δ : 7.34~7.30 (m, 4H), 5.11 (dd, $J=8.2, 4.6$ Hz, 1H), 3.73 (s, 3H), 3.35 (s, 1H), 2.77~2.66 (m, 2H); ^{13}C NMR (126 MHz, CDCl_3) δ : 142.85, 133.13, 128.61, 127.03, 73.62, 61.33, 40.42. The enantiomeric excess was determined by HPLC on Chiralcel AS-H column, 210 nm, 40 °C, *n*-hexane/*i*-PrOH ($V:V=95:5$); flow 0.8 mL/min; $t_{\text{R}1}=16.6$ min (R), $t_{\text{R}2}=18.3$ min (S).

Methyl (R)-3-(2-chlorophenyl)-3-hydroxypropanoate (**4i**):^[22] Colorless oil, 42 mg, >99% conv., 99% yield, 88% ee. $[\alpha]_{\text{D}}^{22}+24.3$ (c 0.9, MeOH). ^1H NMR (400 MHz, CDCl_3) δ : 7.62 (dd, $J=7.6, 1.7$ Hz, 1H), 7.33~7.26 (m, 2H), 7.24~7.10 (m, 1H), 5.49 (dd, $J=9.8, 2.6$ Hz, 1H), 3.73 (s, 3H), 3.59 (s, 1H), 2.86 (dd, $J=16.6, 2.7$ Hz, 1H), 2.59 (dd, $J=16.6, 9.8$ Hz, 1H); ^{13}C NMR (126 MHz, CDCl_3) δ : 172.91, 139.84, 131.34, 129.38, 128.78, 127.24, 127.02, 67.02, 52.02, 41.28. The enantiomeric excess was determined by HPLC on Chiralcel AS-H column, 210 nm, 40 °C, *n*-hexane/*i*-PrOH ($V:V=95:5$); flow 0.8 mL/min; $t_{\text{R}1}=16.6$ min (R), $t_{\text{R}2}=18.3$ min (S).

Methyl (R)-3-(3-chlorophenyl)-3-hydroxypropanoate (**4j**):^[22] Colorless oil, 41 mg, >99% conv., 95% yield, 91% ee. $[\alpha]_{\text{D}}^{22}+14.4$ (c 1.6, MeOH); ^1H NMR (400 MHz, CDCl_3) δ : 7.40 (d, $J=2.2$ Hz, 1H), 7.32~7.23 (m, 3H), 5.11 (dd, $J=7.9, 4.9$ Hz, 1H), 3.74 (s, 3H), 3.51 (s, 1H), 2.78~2.69 (m, 2H); ^{13}C NMR (126 MHz, CDCl_3) δ : 172.60, 144.56, 134.46, 129.86, 127.93, 125.92, 123.81, 69.63, 52.04, 43.02. The enantiomeric excess was determined by HPLC on Chiralcel OJ-H column, 210 nm, 40 °C, *n*-hexane/*i*-PrOH ($V:V=90:10$); flow 0.8 mL/min; $t_{\text{R}1}=11.3$ min (S), $t_{\text{R}2}=12.4$ min (R).

Methyl (R)-3-hydroxy-3-(*o*-tolyl)propanoate (**4k**):^[23] Colorless oil, 38 mg, >99% conv., 98% yield, 95% ee. $[\alpha]_{\text{D}}^{22}+9.6$ (c 1.4, MeOH); ^1H NMR (500 MHz, CDCl_3) δ : 7.50 (dd, $J=7.6, 1.4$ Hz, 1H), 7.25~7.14 (m, 3H), 5.36 (dd, $J=9.2, 3.5$ Hz, 1H), 3.75 (s, 3H), 3.15 (s, 1H), 2.74~2.64 (m, 2H), 2.35 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ :

173.01, 140.43, 134.27, 130.49, 127.64, 126.45, 125.20, 66.98, 51.98, 41.96, 19.02. The enantiomeric excess was determined by HPLC on Chiralcel OD-H column, 210 nm, 40 °C, *n*-hexane/*i*-PrOH ($V:V=90:10$); flow 0.8 mL/min; $t_{\text{R}1}=10.7$ min (S), $t_{\text{R}2}=15.7$ min (R).

Methyl (R)-3-hydroxy-3-(*m*-tolyl)propanoate (**4l**): Colorless oil, 38 mg, >99% conv., 98% yield, 96% ee. $[\alpha]_{\text{D}}^{22}+22.5$ (c 1.4, MeOH); ^1H NMR (500 MHz, CDCl_3) δ : 7.25~7.17 (m, 2H), 7.16~7.11 (m, 1H), 7.10~7.09 (m, 1H), 5.11~5.08 (m, 1H), 3.72 (s, 3H), 3.24~3.23 (m, 1H), 2.79~2.68 (m, 2H), 2.35 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 172.88, 142.44, 138.29, 128.61, 128.49, 126.36, 122.71, 70.35, 51.93, 43.19, 21.47. The enantiomeric excess was determined by HPLC on Chiralcel OD-H column, 210 nm, 40 °C, *n*-hexane/*i*-PrOH ($V:V=90:10$); flow 0.8 mL/min; $t_{\text{R}1}=9.2$ min (S), $t_{\text{R}2}=11.8$ min (R).

Methyl (R)-3-hydroxy-3-(naphthalen-1-yl)propanoate (**4m**): Colorless oil, 41 mg, >99% conv., 95% yield, 91% ee. $[\alpha]_{\text{D}}^{22}+17.2$ (c 1.1, MeOH); ^1H NMR (400 MHz, CDCl_3) δ : 8.07~8.04 (m, 1H), 7.89~7.87 (m, 1H), 7.81~7.79 (m, 1H), 7.72~7.70 (m, 1H), 7.54~7.47 (m, 3H), 5.93 (dd, $J=9.6, 3.1$ Hz, 1H), 3.77 (s, 3H), 3.34 (s, 1H), 2.96~2.82 (m, 2H); ^{13}C NMR (126 MHz, CDCl_3) δ : 173.09, 138.02, 133.76, 129.94, 129.03, 128.32, 126.32, 125.64, 125.54, 122.94, 122.76, 67.33, 52.04, 42.59. The enantiomeric excess was determined by HPLC on Chiralcel AS-3 column, 210 nm, 40 °C, *n*-hexane/*i*-PrOH ($V:V=95:5$); flow 0.5 mL/min; $t_{\text{R}1}=30.8$ min (S), $t_{\text{R}2}=32.6$ min (R).

Methyl (R)-3-hydroxy-3-(naphthalen-2-yl)propanoate (**4n**):^[20] Colorless oil, 45 mg, >99% conv., 97% yield, 94% ee. $[\alpha]_{\text{D}}^{22}+21.0$ (c 1.0, MeOH); ^1H NMR (400 MHz, CDCl_3) δ : 7.85~7.82 (m, 4H), 7.51~7.45 (m, 3H), 5.33~5.29 (m, 1H), 3.74 (s, 3H), 3.34 (t, $J=3.1$ Hz, 1H), 2.89~2.78 (m, 2H); ^{13}C NMR (126 MHz, CDCl_3) δ : 172.83, 139.86, 139.84, 133.27, 133.03, 128.44, 128.04, 127.70, 126.27, 126.04, 124.47, 123.70, 70.45, 52.00, 43.14. The enantiomeric excess was determined by HPLC on Chiralcel AS-H column, 210 nm, 40 °C, *n*-hexane/*i*-PrOH ($V:V=90:10$); flow 0.8 mL/min; $t_{\text{R}1}=12.6$ min (R), $t_{\text{R}2}=13.9$ min (S).

Methyl (R)-3-hydroxy-3-(thiophen-2-yl)propanoate (**4o**):^[20] Yellow oil, 36 mg, >99% conv., 96% yield, 90% ee. $[\alpha]_{\text{D}}^{22}+9.6$ (c 1.2, MeOH); ^1H NMR (400 MHz, CDCl_3) δ : 7.27~7.25 (m, 1H), 6.99~6.96 (m, 2H), 5.40~5.37 (m, 1H), 3.74 (s, 3H), 3.41 (s, 1H), 2.94~2.83 (m, 2H); ^{13}C NMR (126 MHz, CDCl_3) δ : 172.37, 146.20, 126.76, 124.94, 123.69, 66.54, 52.04, 42.98. The enantiomeric excess was determined by HPLC on Chiralcel AS-H column, 210 nm, 40 °C, *n*-hexane/*i*-PrOH ($V:V=90:10$); flow 0.8 mL/min; $t_{\text{R}1}=7.3$ min (R), $t_{\text{R}2}=5.0$ min (S).

4.3.2 Procedure with S/C=10000

In an argon-filled glove box, a 4.0 mL vial was charged with the metal precursor $[\text{Ir}(\text{COD})\text{Cl}]_2$ (3.4 mg, 5.0×10^{-3} mmol), indan-*f*-amphox (6.1 mg, 10.5×10^{-3} mmol) and anhydrous MeOH (1.0 mL). The mixture was stirred for 1.0 h at 25 °C, giving an orange red solution. The resulting

solution (2 μ L) and *t*-BuOK (1 mg) were transferred by a syringe into a 5.0 mL vial charged with **3a** (0.2 mmol) in 1.0 mL of anhydrous MeOH. The vial was transferred to an autoclave, which was then charged with 1 MPa of H₂ and stirred at room temperature for 24 h. The work-up was identical to that described for the asymmetric hydrogenation at S/C=1000. (*R*)-3-Hydroxy-3-phenylpropanoate (**4a**): > 99% conversion, 96% *ee*.

Supporting Information ¹H NMR and ¹³C NMR spectra of all compounds, and chromatography of **4a**~**4o**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] Johnson, R. A.; Sharpless, K. B. In *Catalytic Asymmetric Synthesis*, Ed.: Ojima, I., Wiley-VCH, New York, **2000**, p. 13.
- [2] (a) Noyori R.; Kitamura M. In *Modern Synthetic Methods*, Ed.: Scheffold, R., Springer, Berlin, **1989**, Vol. 5, p. 115.
(b) Takaya, H.; Ohta, T.; Noyori, R. In *Catalytic Asymmetric Synthesis*, Ed.: Ojima, I., VCH, New York, **1993**, p. 1.
(c) Tang, W.; Zhang, X. *Chem. Rev.* **2003**, *103*, 3029.
- [3] (a) Chen, S.; Yang, W.; Yao, Y.; Yang, X.; Deng, Y.; Yang, D. *Chin. J. Org. Chem.* **2018**, *38*, 2534 (in Chinese).
(陈姝琪, 杨文, 姚永祺, 杨新, 邓颖颖, 杨定乔, 有机化学, **2018**, *38*, 2534.)
(b) Shan, H.; Ling, L.; Hu, J.; Zhang, H. *Chin. J. Org. Chem.* **2019**, *39*, 1548 (in Chinese).
(单贺, 伶俐, 胡剑锋, 张浩, 有机化学, **2019**, *39*, 1548.)
(c) Yuan, Q.; Zhang, W. *Chin. J. Org. Chem.* **2016**, *36*, 274 (in Chinese).
(袁乾家, 张万斌, 有机化学, **2016**, *36*, 274.)
- [4] For M-NH catalysis see review: (a) Zhao, B.; Han, Z.; Ding, K. *Angew. Chem., Int. Ed.* **2013**, *52*, 4744.
The recent advances in M-NH ligands see: (b) Li, Y.; Yu, S.; Wu, X.; Xiao, J.; Shen, W.; Dong, Z.; Gao, J. *J. Am. Chem. Soc.* **2014**, *136*, 4031.
(c) Bao, D.-H.; Wu, H.-L.; Liu, C.-L.; Xie, J.-H.; Zhou, Q.-L. *Angew. Chem., Int. Ed.* **2015**, *54*, 8791.
- [5] Ohkuma, T.; Ooka, H.; Hashiguchi, S.; Ikariya, T.; Noyori, R. *J. Am. Chem. Soc.* **1995**, *117*, 2675.
- [6] (a) Hashiguchi, S.; Fujii, A.; Takehara, J.; Ikariya, T.; Noyori, R. *J. Am. Chem. Soc.* **1995**, *117*, 7562.
(b) Gao, J.; Ikariya, T.; Noyori, R. *Organometallics* **1996**, *15*, 1087.
(c) Gamez, P.; Fache, F.; Lemaire, M. *Tetrahedron: Asymmetry* **1995**, *6*, 705.
- [7] Xie, J.-H.; Liu, X.-Y.; Xie, J.-B.; Wang, L.-X.; Zhou, Q.-L. *Angew. Chem., Int. Ed.* **2011**, *50*, 7329.
- [8] Jiang, Y.; Jiang, Q.; Zhang, X. *J. Am. Chem. Soc.* **1998**, *120*, 3817.
- [9] Li, W.; Hou, G.; Wang, C.; Jiang, Y.; Zhang, X. *Chem. Commun.* **2010**, *46*, 3979.
- [10] (a) Wu, W.; Liu, S.; Duan, M.; Tan, X.; Chen, C.; Xie, Y.; Lan, Y.; Dong, X.-Q.; Zhang, X. *Org. Lett.* **2016**, *18*, 2938.
(b) Wu, W.; You, C.; Yin, C.; Liu, Y.; Dong, X.-Q.; Zhang, X. *Org. Chem. Front.* **2017**, *4*, 1499.
(c) Wu, W.; You, C.; Yin, C.; Liu, Y.; Dong, X.-Q.; Zhang, X. *Org. Lett.* **2017**, *19*, 2548.
(d) Wu, W.; Xie, Y.; Li, P.; Li, X.; Liu, Y.; Dong, X.-Q.; Zhang, X. *Org. Chem. Front.* **2017**, *4*, 555.
(e) Gu, G.; Yang, T.; Yu, O.; Qian, H.; Wang, J.; Dang, L.; Zhang, X. *Org. Lett.* **2017**, *19*, 5920.
- [11] Noyori, R. *Angew. Chem., Int. Ed.* **2002**, *41*, 2008.
- [12] Xie, J.-H.; Liu, X.-Y.; Yang, X.-H.; Xie, J.-B.; Wang, L.-X.; Zhou, Q.-L. *Angew. Chem., Int. Ed.* **2012**, *51*, 201.
- [13] Hayashi, T.; Mise, T.; Fukushima, M.; Nagashima, N.; Hamada, Y.; Matsumoto, A.; Kawakami, S.; Konishi, M.; Yamamoto, K. *Bull. Chem. Soc. Jpn.* **1980**, *53*, 1138.
- [14] Nie, H.; Zhou, G.; Wang, Q.; Chen, W.; Zhang, S. *Tetrahedron: Asymmetry* **2013**, *24*, 1567.
- [15] Gu, G.; Yang, T.; Lu, J.; Wen, J.; Dang, L.; Zhang, X. *Org. Chem. Front.* **2018**, *5*, 1209.
- [16] Jiang, W.; Zhao, Q.; Tan, W. *Chin. J. Chem.* **2018**, *36*, 153.
- [17] (a) Schenck, T. G.; Downes, J. M.; C. Milne, R. C.; Mackenzie, P. B.; Boucher, H.; Whelan, J.; Bosnich, B. *Inorg. Chem.* **1985**, *24*, 2334.
(b) Pohl, M.; Lyon, D. K.; Mizuno, N.; Nomiya, K.; Finke, R. G. *Inorg. Chem.* **1995**, *34*, 1413.
- [18] Ye, M.-C.; Li, B.; Zhou, J.; Sun, X.-L.; Tang, Y. *J. Org. Chem.* **2005**, *70*, 6108.
- [19] Jiang, W.; Zhao, Q.; Tang, W. *Chin. J. Chem.* **2018**, *36*, 153.
- [20] Zhang, Y.; Yu, D.; Huang, Z.; Xu, J.; Wu, X.; Wang, Y.; Wang, M.; Yang, S.; Webster, R. D.; Chi, Y. R. *J. Am. Chem. Soc.* **2015**, *137*, 2416.
- [21] White, N. A.; Rovis, T. *J. Am. Chem. Soc.* **2014**, *136*, 14674.
- [22] Chen, X.-S.; Hou, C.-J.; Qin, C.; Liu, H.; Liu, Y.-J.; Huang, D.-Z.; Hu, X.-P. *RSC Adv.* **2017**, *7*, 12871.
- [23] Ohtani, I. I.; Hotta, K.; Ichikawa, Y.; Isobe, M. *Chem. Lett.* **1995**, *7*, 513.
- [24] Andrea, K.; Michel, S.; Barbare, M. *Org. Lett.* **2013**, *15*, 1614.

(Zhao, C.)