基于 δ -腈基取代对亚甲基苯醌 1,6-氮杂共轭加成的大位阻 α -氰胺合成研究

王琳^{†,a} 王楠^{†,c} 齐越^a 孙书涛^b 刘希功^{*,b}
李伟^{*,a} 刘磊^{*,b}

(^a 山东中医药大学药学院 济南 250355)

(^b 山东大学化学与化工学院 济南 250100)

(^c 山东省肿瘤防治研究院(山东省肿瘤医院) 山东第一医科大学(山东省医学科学院) 济南 250117)

摘要 目前对亚甲基苯醌参与的 1,6-共轭加成反应的研究主要集中在对预先合成的 δ -单取代底物进行三级碳中心的构建。报道了一级芳胺对预先制备的 δ -氰基- δ -芳基二取代对亚甲基苯醌的 1,6-氮杂共轭加成反应, 高效地合成了一系列含有四级碳手性中心的大位阻胺类化合物。该方法条件温和、反应迅速, 同时芳胺和对亚甲基苯醌的适用范围广泛。此外, 环状二级胺例如吗啉和咪唑也是合适的亲核试剂。

关键词 对亚甲基苯醌; δ -氰基- δ -芳基二取代; 大位阻胺; 氮杂四级碳中心; 1,6-氮杂共轭加成

Synthesis of Sterically Hindered α -Aminonitriles through 1,6-Aza-conjugate Addition of Anilines to δ -Cyano Substituted *para*-Quinone Methides

Wang, Lin^{†,a} Wang, Nan^{†,c} Qi, Yue^a Sun, Shutao^b Liu, Xigong^{*,b}
Li, Wei^{*,a} Liu, Lei^{*,b}

(^a Department of Pharmaceutical Analysis, School of Pharmacy, Shandong University of Traditional Chinese Medicine, Jinan 250355)

(^b School of Chemistry and Chemical Engineering, Shandong University, Jinan 250100)

(^c The First Ward of Intervention Department, Shandong Cancer Hospital and Institute, Shandong First Medical University and Shandong Academy of Medical Sciences, Jinan 250117)

Abstract Current 1,6-conjugate addition typically focused on pre-synthesized *para*-quinone methides bearing a δ -mono substituent for tertiary stereocenter formation. Here, an efficient 1,6-aza-conjugate addition of primary anilines to pre-prepared δ -CN- δ -aryl disubstituted *para*-quinone methides for facile access to sterically hindered amines with a fully substituted α -carbon center has been described. The mild and expeditious method exhibited broad scopes of both aniline and *para*-quinone methide components. The generality of the method in modular preparation of medicinally valuable, sterically hindered amines was further demonstrated by using cyclic secondary amines like morpholine and imidazole as nucleophilic components.

Keywords *para*-quinone methide; δ -CN- δ -aryl disubstitution; sterically hindered amine; aza-quaternary carbon; 1,6-aza-conjugate addition

1 Introduction

para-Quinone methides (*p*-QMs), structurally charac-

terized by the assembly of carbonyl and olefinic moieties, have been widely employed as prominent acceptors for 1,6-conjugate addition reactions.^[1-3] However, existing

* Corresponding authors. E-mail: 201990000024@sdu.edu.cn; liwei6911@163.com; lei.liu@sdu.edu.cn

Received April 17, 2020; revised May 14, 2020; published online May 19, 2020.

Dedicated to the 40th anniversary of Chinese Journal of Organic Chemistry.

Project supported by the National Natural Science Foundation of China (Nos. 21722204, 21971148).

国家自然科学基金(Nos. 21722204, 21971148)资助项目。

[†] 共同第一作者(These authors contributed equally to this work).

methods suffered from notable structural restriction on *p*-QMs, probably due to the stability issue.^[4] Current studies predominantly concentrate on *p*-QMs having two bulky α -substituents (e.g. ^tBu). Additionally, *p*-QM substrates typically contain a mono-aryl substitution at the δ -position for tertiary stereocenter construction. Sun and co-workers developed a delicate protocol to address the structural limitations by *in situ* generating *p*-QM intermediates bearing two electron-donating groups (EDGs) at the δ -position via Brønsted acid catalyzed dehydration of *p*-hydroxybenzyl alcohols.^[5] Despite great innovation, *p*-QM intermediates bearing a δ -electron-withdrawing group (EWG) might not be accessible through the protocol. In addition, the electrophilicity of *p*-QMs bearing two EDGs at δ -position towards 1,6-conjugate addition might be impaired. Recently, our group employed an oxidative C—H cleavage strategy to *in situ* generate δ -EWG substituted *p*-QMs.^[6] However, diverse nucleophiles might not be compatible with the strongly oxidative conditions, and therefore the utilities of the protocol would be restricted. Therefore, further expanding the scope of *p*-QMs and applying them to address existing synthetic challenges would be highly desired. Using pre-prepared, isolable, and storable *p*-QMs as substrates would be an attractive solution. Placing an electron-withdrawing cyano group at the exocyclic methylene δ -position would tune the molecular electron-density distribution, and the stability of resulting *p*-QMs might be improved by lowering their polymerizability.^[7]

Sterically hindered amines with a fully substituted α carbon center are important structural motifs in various natural products and pharmaceuticals.^[8] C—N bond forming reactions represent practical and robust approaches for their synthesis. In this context, 1,6-aza-conjugate addition of amines to *p*-QMs has received considerable attentions.^[9] However, existing methods typically involved intra- or inter-molecular addition of cyclic secondary amines to *p*-QMs providing corresponding adducts with a tertiary α carbon center. To our knowledge, 1,6-aza-conjugate addition for the synthesis of sterically hindered amines with an acyclic, fully substituted α carbon center has never been reported. In addition, the use of primary amines as nucleophiles for 1,6-conjugate addition proved to be challenging due to the instability of the resulting adducts.^[2c] Given the significance of α -amino-nitriles in organic synthesis and pharmaceutical science,^[10] we herein reported a 1,6-aza-conjugate addition of a range of primary anilines to δ -CN- δ -aryl substituted *p*-QMs for efficient synthesis of sterically hindered amines with a fully substituted α carbon center. The generality of the method was further demonstrated by 1,6-aza-conjugate addition with valuable secondary amines.

2 Results and discussion

Initially, 1,6-aza-conjugate addition of aniline **2a** to δ -CN- δ -phenyl substituted *p*-QM **1a** in CH₂Cl₂ was selected as the model reaction for optimization (Table 1).

When diphenyl phosphate was used as a Brønsted acid catalyst, the reaction proceeded smoothly, affording the expected sterically hindered amine **3a** in 52% yield together with undesired triarylmethane **3a'** in 16% yield (Entry 1, Table 1). The solvent effect was then carefully investigated, and MeOH was identified to be optimal with respect to the chemoselectivity and efficiency (Entries 1~7, Table 1). Notably, no undesired **3a'** was detected (Entry 7, Table 1). An extensive screen of Brønsted and Lewis acid additives revealed that the 1,6-aza-conjugate addition proceeded most efficiently when *p*-toluenesulfonic acid (PTSA) was used as the catalyst, furnishing desired **3a** in 91% yield within 0.5 h (Entries 7~11, Table 1).

Table 1 Reaction condition optimization^a

Entry	Catalyst	Solvent	Time/h	Yield ^b /%	
				3a	3a'
1	DPP	CH ₂ Cl ₂	6	52	16
2	DPP	Toluene	6	57	8
3	DPP	EtOAc	6	60	10
4	DPP	THF	2	66	18
5	DPP	Acetone- <i>d</i> ₆	0.5	69	15
6	DPP	CH ₃ CN	0.5	71	13
7	DPP	MeOH	0.5	75	<5
8	AcOH	MeOH	0.5	82	<5
9	PTSA	MeOH	0.5	91	<5
10	BF ₃ ·OEt ₂	MeOH	1	77	<5
11	AgOTf	MeOH	1	68	<5

^a Reaction conditions, unless otherwise specified: a solution of **1a** (0.1 mmol), **2a** (0.105 mmol), catalyst (0.01 mmol) in solvent (1.0 mL) at room temperature for indicated time period. ^b Isolated yield. DPP=diphenyl phosphate. PTSA=*p*-toluenesulfonic acid, Tf=trifluoromethanesulfonyl.

With the optimized conditions in hand, the scope of δ -CN substituted *p*-QMs was examined (Scheme 1). A wide variety of *p*-QMs **1** bearing electronically varied aryl groups with diverse substituent patterns at the δ -position joined in the 1,6-aza-conjugate addition reactions smoothly, furnishing corresponding sterically hindered amines **3a**~**3o** in 85%~94% yields (Scheme 1A). δ -Polyarene naphthalene substituted *p*-QMs were also suitable substrates, as demonstrated by the formation of respective **3p** and **3q** in 94% and 90% yields. The substituent pattern on quinone moiety was then explored (Scheme 1B). *p*-QMs **1r**~**1u** bearing diverse electron-donating and electron-withdrawing small substituents at α -position were well tolerated, giving respective **3r**~**3u** in 90%~94% yields.

The scope of aniline components was next evaluated (Scheme 2). A broad range of anilines bearing diverse electron-donating and electron-withdrawing substituents at

Scheme 1 Scope of *p*-QM components

para-position were competent nucleophilic partners, as illustrated by the generation of respective sterically hindered amines **4a**~**4d** in high efficiency. Anilines **2e**~**2g** bearing substituents at *meta*- or *ortho*-position were also well tolerated, and respective **4e**~**4g** were isolated in 88%~90% yields. 2-Naphthalene substituted **2h** was suitable component for the 1,6-aza-conjugate addition process.

The success in amination with primary anilines prompted us to further explore 1,6-aza-conjugate addition with secondary amines (Scheme 3). Triarylmethane **7** bearing a morpholine moiety has been identified as a potent inhibitor of apolipoprotein E production (Scheme 3A).^[11] Under the standard conditions, morpholine (**5**) reacted with **1a** giving diarylmethane **6** bearing a fully substituted stereocenter in 85% yield. Clotrimazole (**10**) is a broad-spectrum antifun-

gal drug that is used for the treatment of dermal infections caused by various species of pathogenic dermatophytes, yeasts, and *Malassezia furfur* (Scheme 3B).^[12] Imidazole (**8**) participated in 1,6-aza-conjugate addition of **1a** smoothly, affording diarylmethane **9** bearing a fully substituted stereocenter in 90% yield. These results suggested that the method would provide a robust handle to facilitate prepare structurally diverse and sterically hindered amines for further medicinal study.

3 Conclusions

In summary, an efficient 1,6-aza-conjugate addition of primary anilines to δ -CN- δ -aryl disubstituted *p*-QMs for facile access to sterically hindered amines with a fully substituted α carbon center has been described. The mild and

Scheme 2 Scope of aniline components

Scheme 3 Secondary amine components

expeditious method exhibited broad scopes of both aniline and *p*-QM components. The generality of the method in modular preparation of medicinally valuable, sterically hindered amines was further demonstrated by using cyclic secondary amines like morpholine and imidazole as nucleophilic components. Ongoing studies focus on applying the method in medicinal chemistry study.

4 Experimental section

4.1 General information

Proton nuclear magnetic resonance (^1H NMR) spectra were recorded at 500 MHz respectively. Carbon nuclear magnetic resonance (^{13}C NMR) spectra were recorded at 126 MHz respectively. The solvent peak was used as a reference value for ^1H NMR (CDCl_3 δ 7.26, $(\text{CD}_3)_2\text{CO}$ δ 2.05) and for ^{13}C NMR (CDCl_3 δ 77.23, $(\text{CD}_3)_2\text{CO}$ δ 206.23/29.82). Analytical thin-layer chromatography (TLC) was performed on precoated silica gel GF254 plates. Column chromatography was carried out on silica gel (200~300 mesh). HRMS were carried out on an Orbitrap analyzer.

4.1 Preparation of substrate 1

To a solution of **S-1** (10 mmol, 1.0 equiv.) in anhydrous tetrahydrofuran (THF) at 0 °C was added PhMgBr (1.0 mol·L $^{-1}$, 10 mL) in THF dropwise under N_2 (Scheme 4). After stirred at the same temperature for 0.5 h, the reaction was warmed up to room temperature slowly and stirred for 2 h. Upon completion, the mixture was quenched by a saturated aqueous NH_4Cl solution (15 mL). The organic layer was extracted with ethyl acetate, and the combined organic layers were washed with saturated brine, and dried over anhydrous MgSO_4 . The solvent was removed under vacuum to afford crude alcohol for next step.

Reagent and conditions: (a) PhMgBr (1.0 equiv.), THF, 0 °C; (b) InCl_3 (0.1 equiv.), TMSCN (1.3 equiv.), CH_2Cl_2 , 0 °C; (c) Pd/C (10%), MeOH, hydrogen balloon, r.t.; (d) DDQ (1.0 equiv.), CH_2Cl_2 , r.t.

Scheme 4 Preparation of substrate 1

To a mixture of InCl_3 (1.0 mmol, 0.1 equiv.) and TMSCN (13 mmol, 1.3 equiv.) in anhydrous CH_2Cl_2 (20 mL) was added a solution of crude alcohol in anhydrous CH_2Cl_2 (10 mL) at 0 °C. The reaction was stirred at the same temperature and monitored by TLC until the complete conversion. Then it was quenched by saturated aqueous NaHCO_3 solution (20 mL) and extracted with CH_2Cl_2 . The combined organic layers were washed with saturated brine, and dried over anhydrous MgSO_4 . The solvent was removed under vacuum to provide the crude product, which was used as such for next step.

The crude product was dissolved in 20 mL of MeOH solution, and 10% Pd/C was added. The mixture was

stirred overnight under a hydrogen balloon at room temperature before it was filtered through a Celite pad. The solvent was removed under vacuum and the residue was purified by a column chromatography on silica gel using ethyl acetate/petroleum ether as eluent to give **S-2**.

To a solution of **S-4** in CH_2Cl_2 (20.0 mL) was added DDQ (1.0 equiv.) and it was stirred at room temperature for 1 h. Then the solvent was removed under vacuum and the residue was purified by silica gel chromatography using ethyl acetate/petroleum ether as eluent to afford expected **1a**.

(4-Oxocyclohexa-2,5-dien-1-ylidene)-2-phenylacetonitrile (**1a**): ^1H NMR (500 MHz, CDCl_3) δ : 7.84 (dd, $J=10.0, 2.7$ Hz, 1H), 7.56~7.50 (m, 5H), 7.48 (dd, $J=10.1, 2.7$ Hz, 1H), 6.61 (dd, $J=10.0, 1.9$ Hz, 1H), 6.48 (dd, $J=10.1, 1.9$ Hz, 1H); ^{13}C NMR (126 MHz, CDCl_3) δ : 186.6, 139.8, 137.1, 134.6, 132.1, 131.6, 131.5, 131.5, 130.6, 129.5, 124.2, 117.3; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{10}\text{NO}$ $[\text{M}+\text{H}]^+$ 208.0757, found 208.0760.

([1,1'-Biphenyl]-4-yl)-2-(4-oxocyclohexa-2,5-dien-1-ylidene)acetonitrile (**1b**): ^1H NMR (500 MHz, CDCl_3) δ : 7.87 (dd, $J=10.0, 2.7$ Hz, 1H), 7.76 (d, $J=8.4$ Hz, 2H), 7.66~7.60 (m, 4H), 7.58 (dd, $J=10.1, 2.7$ Hz, 1H), 7.53~7.48 (m, 2H), 7.46~7.41 (m, 1H), 6.63 (dd, $J=10.0, 1.8$ Hz, 1H), 6.52 (dd, $J=10.1, 1.8$ Hz, 1H); ^{13}C NMR (126 MHz, CDCl_3) δ : 186.7, 144.5, 139.6, 139.5, 137.2, 134.7, 131.5, 131.3, 131.0, 129.3, 128.7, 128.1, 127.4, 124.0, 117.3; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{14}\text{NO}$ $[\text{M}+\text{H}]^+$ 284.1070, found 284.1065.

(4-Oxocyclohexa-2,5-dien-1-ylidene)-2-(*p*-tolyl)acetonitrile (**1c**): ^1H NMR (500 MHz, CDCl_3) δ : 7.83 (dd, $J=10.0, 2.7$ Hz, 1H), 7.51 (dd, $J=10.1, 2.7$ Hz, 1H), 7.43 (d, $J=8.2$ Hz, 2H), 7.34 (d, $J=8.0$ Hz, 2H), 6.60 (dd, $J=10.0, 1.9$ Hz, 1H), 6.48 (dd, $J=10.1, 1.9$ Hz, 1H), 2.45 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 186.8, 142.5, 139.2, 137.2, 134.9, 131.4, 131.4, 130.7, 130.3, 129.4, 124.5, 117.4, 21.8; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{12}\text{NO}$ $[\text{M}+\text{H}]^+$ 222.0913, found 222.0915.

(4-Isopropylphenyl)-2-(4-oxocyclohexa-2,5-dien-1-ylidene)acetonitrile (**1d**): ^1H NMR (500 MHz, CDCl_3) δ : 7.83 (dd, $J=10.0, 2.7$ Hz, 1H), 7.53 (dd, $J=10.1, 2.7$ Hz, 1H), 7.48~7.43 (m, 2H), 7.40~7.36 (m, 2H), 6.60 (dd, $J=10.0, 1.9$ Hz, 1H), 6.48 (dd, $J=10.1, 1.9$ Hz, 1H), 3.04~2.94 (m, 1H), 1.30 (s, 3H), 1.29 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 186.7, 153.2, 139.2, 137.2, 134.9, 131.4, 131.3, 130.9, 129.7, 127.7, 124.5, 117.4, 34.4, 23.9; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{16}\text{NO}$ $[\text{M}+\text{H}]^+$ 250.1226, found 250.1229.

(4-*tert*-Butylphenyl)-2-(4-oxocyclohexa-2,5-dien-1-ylidene)acetonitrile (**1e**): ^1H NMR (500 MHz, CDCl_3) δ : 7.84 (dd, $J=10.0, 2.7$ Hz, 1H), 7.56~7.51 (m, 3H), 7.50~7.44 (m, 2H), 6.60 (dd, $J=10.0, 1.9$ Hz, 1H), 6.49 (dd, $J=10.1, 1.9$ Hz, 1H), 1.37 (s, 9H); ^{13}C NMR (126 MHz, CDCl_3) δ : 186.8, 155.5, 139.3, 137.3, 135.0, 131.4, 131.4, 130.7, 129.4, 126.6, 124.5, 117.4, 35.4, 31.3; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{18}\text{NO}$ $[\text{M}+\text{H}]^+$ 264.1383, found 264.1386.

(4-Bromophenyl)-2-(4-oxocyclohexa-2,5-dien-1-ylidene)acetonitrile (**1f**): ^1H NMR (500 MHz, CDCl_3) δ : 7.82 (dd, $J=10.0, 2.6$ Hz, 1H), 7.68 (d, $J=8.4$ Hz, 2H), 7.45~7.37 (m, 3H), 6.62 (dd, $J=10.0, 1.5$ Hz, 1H), 6.50 (dd, $J=10.1, 1.6$ Hz, 1H); ^{13}C NMR (126 MHz, CDCl_3) δ : 186.5, 140.2, 136.9, 134.1, 132.9, 132.0, 131.8, 130.9, 126.5, 122.7, 116.9; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_9\text{BrNO}$ $[\text{M}+\text{H}]^+$ 285.9862, found 285.9858.

(4-Chlorophenyl)-2-(4-oxocyclohexa-2,5-dien-1-ylidene)acetonitrile (**1g**): ^1H NMR (500 MHz, CDCl_3) δ : 7.82 (dd, $J=10.0, 2.7$ Hz, 1H), 7.55~7.50 (m, 2H), 7.49~7.45 (m, 2H), 7.43 (dd, $J=10.1, 2.7$ Hz, 1H), 6.62 (dd, $J=10.0, 1.9$ Hz, 1H), 6.50 (dd, $J=10.1, 1.9$ Hz, 1H); ^{13}C NMR (126 MHz, CDCl_3) δ : 186.5, 140.2, 138.1, 136.9, 134.1, 131.9, 131.8, 130.5, 129.9, 122.6, 117.0; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_9\text{ClNO}$ $[\text{M}+\text{H}]^+$ 242.0367, found 242.0364.

(4-Fluorophenyl)-2-(4-oxocyclohexa-2,5-dien-1-ylidene)acetonitrile (**1h**): ^1H NMR (500 MHz, CDCl_3) δ : 7.84 (dd, $J=10.0, 2.7$ Hz, 1H), 7.58~7.52 (m, 2H), 7.46 (dd, $J=10.1, 2.7$ Hz, 1H), 7.29~7.23 (m, 2H), 6.63 (dd, $J=10.0, 1.8$ Hz, 1H), 6.51 (dd, $J=10.1, 1.8$ Hz, 1H); ^{13}C NMR (126 MHz, CDCl_3) δ : 186.6, 165.5, 163.4, 139.9, 137.0, 134.3, 132.9, 132.8, 131.7, 131.7, 128.2, 128.2, 122.8, 117.1, 117.0, 116.9; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_9\text{FNO}$ $[\text{M}+\text{H}]^+$ 226.0663, found 226.0659.

(4-Oxocyclohexa-2,5-dien-1-ylidene)-2-(4-(trifluoromethyl)phenyl)acetonitrile (**1i**): ^1H NMR (500 MHz, CDCl_3) δ : 7.85~7.77 (m, 3H), 7.65 (d, $J=8.2$ Hz, 2H), 7.38 (dd, $J=10.1, 2.7$ Hz, 1H), 6.62 (dd, $J=10.0, 1.8$ Hz, 1H), 6.48 (dd, $J=10.1, 1.9$ Hz, 1H); ^{13}C NMR (126 MHz, CDCl_3) δ : 186.3, 141.1, 136.7, 135.4, 133.8, 133.1, 132.9, 132.1, 132.0, 130.9, 126.5, 126.5, 126.5, 126.4, 124.6, 122.5, 121.9, 116.8; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_9\text{F}_3\text{NO}$ $[\text{M}+\text{H}]^+$ 276.0631, found 276.0635.

(4-Oxocyclohexa-2,5-dien-1-ylidene)-2-(*m*-tolyl)acetonitrile (**1j**): ^1H NMR (500 MHz, CDCl_3) δ : 7.82 (dd, $J=10.0, 2.7$ Hz, 1H), 7.48 (dd, $J=10.1, 2.7$ Hz, 1H), 7.40 (t, $J=7.6$ Hz, 1H), 7.36~7.27 (m, 3H), 6.59 (dd, $J=10.0, 1.8$ Hz, 1H), 6.47 (dd, $J=10.1, 1.9$ Hz, 1H), 2.43 (s, 3H); ^{13}C NMR (126 MHz, CDCl_3) δ : 186.6, 139.6, 139.5, 137.1, 134.8, 132.3, 132.0, 131.5, 131.4, 131.0, 129.3, 127.8, 124.4, 117.3, 21.5; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{12}\text{NO}$ $[\text{M}+\text{H}]^+$ 222.0913, found 222.0915.

(3-Bromophenyl)-2-(4-oxocyclohexa-2,5-dien-1-ylidene)acetonitrile (**1k**): ^1H NMR (500 MHz, CDCl_3) δ : 7.81 (dd, $J=10.0, 2.7$ Hz, 1H), 7.71~7.64 (m, 2H), 7.46~7.39 (m, 3H), 6.62 (dd, $J=10.0, 1.8$ Hz, 1H), 6.50 (dd, $J=10.1, 1.8$ Hz, 1H); ^{13}C NMR (126 MHz, CDCl_3) δ : 186.4, 140.7, 136.7, 134.4, 134.0, 133.8, 133.1, 132.0, 131.9, 131.0, 129.1, 123.6, 122.0, 116.9; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_9\text{BrNO}$ $[\text{M}+\text{H}]^+$ 285.9862, found 285.9859.

(3-Chlorophenyl)-2-(4-oxocyclohexa-2,5-dien-1-ylidene)acetonitrile (**1l**): ^1H NMR (500 MHz, CDCl_3) δ : 7.82 (dd, $J=10.0, 2.7$ Hz, 1H), 7.55~7.46 (m, 3H), 7.45~7.38 (m, 2H), 6.63 (dd, $J=10.0, 1.9$ Hz, 1H), 6.51 (dd, $J=10.1, 1.9$ Hz, 1H); ^{13}C NMR (126 MHz, CDCl_3) δ : 186.5, 140.7,

136.8, 135.7, 134.0, 133.6, 132.0, 132.0, 131.5, 130.8, 130.3, 128.7, 122.1, 116.9; HRMS (ESI) calcd for $C_{14}H_9ClNO$ $[M+H]^+$ 242.0367, found 242.0363.

(3-Fluorophenyl)-2-(4-oxocyclohexa-2,5-dien-1-ylidene)acetonitrile (**1m**): 1H NMR (500 MHz, $CDCl_3$) δ : 7.85 (dd, $J=10.0$, 2.7 Hz, 1H), 7.81 (d, $J=8.2$ Hz, 2H), 7.65 (d, $J=8.1$ Hz, 2H), 7.38 (dd, $J=10.1$, 2.7 Hz, 1H), 6.64 (dd, $J=10.0$, 1.9 Hz, 1H), 6.50 (dd, $J=10.1$, 1.9 Hz, 1H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 186.6, 163.9, 161.9, 140.6, 136.9, 134.2, 133.8, 133.7, 133.1, 131.9, 131.9, 131.3, 131.3, 126.4, 126.4, 122.4, 118.6, 118.5, 117.6, 117.4, 116.9, 115.6; HRMS (ESI) calcd for $C_{14}H_9FNO$ $[M+H]^+$ 226.0663, found 226.0661.

(4-Oxocyclohexa-2,5-dien-1-ylidene)-2-(o-tolyl)acetonitrile (**1n**): 1H NMR (500 MHz, $CDCl_3$) δ : 7.85 (dd, $J=10.0$, 2.6 Hz, 1H), 7.45~7.39 (m, 1H), 7.36~7.29 (m, 2H), 7.18 (dd, $J=7.6$, 0.9 Hz, 1H), 7.08 (dd, $J=10.1$, 2.6 Hz, 1H), 6.61 (dd, $J=10.0$, 1.9 Hz, 1H), 6.39 (dd, $J=10.1$, 1.9 Hz, 1H), 2.37 (s, 3H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 186.7, 141.6, 137.4, 136.4, 134.5, 131.9, 131.5, 131.4, 131.0, 130.9, 130.7, 126.7, 123.4, 116.4, 20.2; HRMS (ESI) calcd for $C_{15}H_{12}NO$ $[M+H]^+$ 222.0913, found 222.0917.

(3,5-Dimethylphenyl)-2-(4-oxocyclohexa-2,5-dien-1-ylidene)acetonitrile (**1o**): 1H NMR (500 MHz, $CDCl_3$) δ : 7.82 (dd, $J=10.0$, 2.7 Hz, 1H), 7.50 (dd, $J=10.1$, 2.7 Hz, 1H), 7.17 (s, 1H), 7.12 (s, 2H), 6.60 (dd, $J=10.0$, 1.9 Hz, 1H), 6.48 (dd, $J=10.1$, 1.9 Hz, 1H), 2.39 (s, 6H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 186.8, 139.6, 139.4, 137.2, 135.0, 133.3, 132.1, 131.5, 131.2, 128.4, 124.8, 118.5, 117.4, 21.5; HRMS (ESI) calcd for $C_{16}H_{14}NO$ $[M+H]^+$ 236.1070, found 236.1073.

(Naphthalen-2-yl)-2-(4-oxocyclohexa-2,5-dien-1-ylidene)acetonitrile (**1p**): 1H NMR (500 MHz, $CDCl_3$) δ : 8.02 (s, 1H), 7.99 (d, $J=8.5$ Hz, 1H), 7.96~7.87 (m, 3H), 7.68~7.54 (m, 4H), 6.64 (dd, $J=10.0$, 1.8 Hz, 1H), 6.52 (dd, $J=10.1$, 1.8 Hz, 1H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 186.7, 139.8, 137.2, 134.9, 134.2, 132.9, 131.7, 131.7, 131.6, 129.5, 129.0, 128.8, 128.1, 127.8, 126.5, 124.3, 117.4; HRMS (ESI) calcd for $C_{18}H_{12}NO$ $[M+H]^+$ 258.0913, found 258.0918.

(Naphthalen-1-yl)-2-(4-oxocyclohexa-2,5-dien-1-ylidene)acetonitrile (**1q**): 1H NMR (500 MHz, $CDCl_3$) δ : 8.03 (d, $J=8.3$ Hz, 1H), 8.00~7.93 (m, 2H), 7.88~7.82 (m, 1H), 7.66~7.55 (m, 3H), 7.47 (dd, $J=7.1$, 0.9 Hz, 1H), 7.08 (dd, $J=10.1$, 2.6 Hz, 1H), 6.67 (dd, $J=10.0$, 1.9 Hz, 1H), 6.34 (dd, $J=10.1$, 1.9 Hz, 1H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 186.8, 142.5, 136.4, 134.8, 133.9, 132.0, 131.8, 131.3, 131.1, 129.6, 129.1, 128.8, 128.2, 127.4, 125.4, 124.7, 122.2, 117.0; HRMS (ESI) calcd for $C_{18}H_{12}NO$ $[M+H]^+$ 258.0913, found 258.0916.

2-(3-Methyl-4-oxocyclohexa-2,5-dien-1-ylidene)-2-phenylacetonitrile (**1r**): **1r** was obtained as a 1 : 1 mixture of the *Z* : *E* isomers. 1H NMR (500 MHz, $CDCl_3$) δ : 7.78 (dd, $J=9.9$, 2.7 Hz, 1H), 7.68~7.65 (m, 1H), 7.56~7.48 (m, 10H), 7.40 (dd, $J=10.0$, 2.7 Hz, 1H), 7.31~7.27 (m, 1H), 6.60 (d, $J=9.9$ Hz, 1H), 6.47 (d, $J=10.0$ Hz, 1H), 2.14 (d,

$J=1.4$ Hz, 3H), 2.03 (d, $J=1.4$ Hz, 3H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 187.0, 140.5, 140.3, 140.3, 136.8, 134.3, 133.7, 132.3, 131.4, 131.3, 131.1, 130.7, 130.6, 129.4, 129.4, 122.2, 122.1, 117.7, 16.8, 16.5; HRMS (ESI) calcd for $C_{15}H_{12}NO$ $[M+H]^+$ 222.0913, found 222.0916.

(3-Bromo-4-oxocyclohexa-2,5-dien-1-ylidene)-2-phenylacetonitrile (**1s**): **1s** was obtained as a 1 : 1 mixture of the *Z* : *E* isomers. 1H NMR (500 MHz, $CDCl_3$) δ : 8.30 (d, $J=2.5$ Hz, 1H), 7.92 (d, $J=2.5$ Hz, 1H), 7.85 (dd, $J=9.8$, 2.5 Hz, 1H), 7.61~7.51 (m, 10H), 7.51~7.48 (m, 1H), 6.75 (d, $J=9.8$ Hz, 1H), 6.63 (d, $J=10.0$ Hz, 1H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 179.2, 179.1, 139.6, 139.4, 138.4, 137.1, 136.6, 136.2, 134.8, 133.3, 132.0, 131.9, 131.9, 130.8, 130.8, 130.7, 130.7, 130.6, 130.2, 129.8, 129.7, 124.7, 117.1, 117.1; HRMS (ESI) calcd for $C_{14}H_9BrNO$ $[M+H]^+$ 285.9862, found 285.9866.

2-(3-Chloro-4-oxocyclohexa-2,5-dien-1-ylidene)-2-phenylacetonitrile (**1t**): **1t** was obtained as a 1 : 1 mixture of the *Z* : *E* isomers. 1H NMR (500 MHz, $CDCl_3$) δ : 8.02 (d, $J=2.5$ Hz, 1H), 7.83 (dd, $J=9.9$, 2.5 Hz, 1H), 7.65 (d, $J=2.5$ Hz, 1H), 7.59~7.50 (m, 10H), 7.50~7.45 (m, 1H), 6.71 (d, $J=9.9$ Hz, 1H), 6.59 (d, $J=10.0$ Hz, 1H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 179.1, 179.0, 139.2, 139.1, 137.5, 137.3, 136.9, 134.7, 134.3, 132.1, 131.9, 131.8, 131.1, 131.1, 130.7, 130.6, 129.7, 129.6, 124.9, 124.8, 117.0; HRMS (ESI) calcd for $C_{14}H_9ClNO$ $[M+H]^+$ 242.0367, found 242.0369.

2-(3-Fluoro-4-oxocyclohexa-2,5-dien-1-ylidene)-2-phenylacetonitrile (**1u**): **1u** was obtained as a 2 : 1 mixture of the *Z* : *E* isomers. 1H NMR (500 MHz, $CDCl_3$) δ : 7.78 (dd, $J=9.9$, 2.5 Hz, 1H), 7.56~7.37 (m, 8H), 7.09 (dd, $J=11.1$, 2.5 Hz, 1H), 6.63 (dd, $J=9.9$, 7.4 Hz, 1H), 6.50 (dd, $J=9.9$, 7.3 Hz, 0.5 H); ^{13}C NMR (126 MHz, $CDCl_3$) δ : 178.4, 178.2, 157.2, 157.1, 155.0, 154.9, 139.8, 139.7, 136.7, 134.5, 132.0, 132.0, 131.8, 131.7, 131.2, 131.2, 131.2, 130.8, 130.5, 129.7, 129.6, 125.2, 125.1, 117.1, 117.1, 117.0, 116.7, 116.6, 114.6, 114.5; HRMS (ESI) calcd for $C_{14}H_9FNO$ $[M+H]^+$ 226.0663, found 226.0659.

4.3 General procedure for 1,6-aza-conjugate addition of anilines to δ -cyano substituted *para*-quinone methides

To a solution of **1** (0.1 mmol, 1.0 equiv.) in MeOH (1.0 mL) was successively added **2** (0.105 mmol, 1.05 equiv.), PTSA (0.01 mmol, 10 mol%) at room temperature. The mixture was stirred at the same temperature for 30 min~6 h. The solvent was removed and the residue was purified by flash column chromatography to give the desired product.

2-(4-Hydroxyphenyl)-2-phenyl-2-(phenylamino)acetonitrile (**3a**): It was purified by flash chromatography on silica gel using acetone/petroleum ether (V : $V=1$: 9~1 : 4) as eluent to afford **3a** 27.2 mg, 91% yield. 1H NMR (500 MHz, Acetone- d_6) δ : 8.65 (s, 1H), 7.67~7.56 (m, 2H), 7.49~7.29 (m, 5H), 7.18~7.04 (m, 2H), 6.94~6.83 (m, 2H), 6.75 (t, $J=7.9$ Hz, 3H), 6.02 (s, 1H); ^{13}C NMR (126

MHz, Acetone- d_6) δ : 158.1, 145.2, 140.8, 131.4, 129.4, 128.9, 128.1, 126.6, 120.6, 116.4, 65.0; HRMS (ESI) calcd for $C_{20}H_{15}N_2O$ $[M-H]^-$ 299.1190, found 299.1195.

2-([1,1'-Biphenyl]-4-yl)-2-(4-hydroxyphenyl)-2-(phenylamino)acetonitrile (**3b**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9 \sim 1:4$) as eluent to afford **3b** 31.9 mg, 85% yield. 1H NMR (500 MHz, Acetone- d_6) δ : 8.67 (s, 1H), 7.76~7.65 (m, 6H), 7.49~7.44 (m, 4H), 7.39 (d, $J=7.5$ Hz, 1H), 7.13 (dd, $J=8.5, 7.4$ Hz, 2H), 6.90 (d, $J=8.7$ Hz, 2H), 6.83~6.73 (m, 3H), 6.07 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.4, 145.4, 141.7, 140.4, 140.1, 131.5, 129.5, 129.1, 128.3, 128.2, 128.0, 127.4, 127.4, 120.7, 119.6, 116.6, 116.3, 65.0; HRMS (ESI) calcd for $C_{26}H_{19}N_2O$ $[M-H]^-$ 375.1503, found 375.1506.

2-(4-Hydroxyphenyl)-2-(phenylamino)-2-(*p*-tolyl)acetonitrile (**3c**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9 \sim 1:4$) as eluent to afford **3c** 27.2 mg, 87% yield. 1H NMR (500 MHz, Acetone- d_6) δ : 8.62 (d, $J=12.2$ Hz, 1H), 7.47 (d, $J=8.3$ Hz, 2H), 7.41~7.37 (m, 2H), 7.24 (d, $J=8.0$ Hz, 2H), 7.10 (dd, $J=8.6, 7.3$ Hz, 2H), 6.89~6.85 (m, 2H), 6.78~6.71 (m, 3H), 5.95 (s, 1H), 2.33 (s, 3H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.3, 145.5, 138.9, 138.2, 131.8, 130.1, 129.0, 128.3, 126.8, 120.9, 119.5, 116.5, 116.2, 64.9, 20.6; HRMS (ESI) calcd for $C_{21}H_{17}N_2O$ $[M-H]^-$ 313.1346, found 313.1342.

2-(4-Hydroxyphenyl)-2-(4-isopropylphenyl)-2-(phenylamino)acetonitrile (**3d**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9 \sim 1:4$) as eluent to afford **3d** 30.7 mg, 90% yield. 1H NMR (500 MHz, Acetone- d_6) δ : 8.64 (s, 1H), 7.51 (d, $J=8.2$ Hz, 2H), 7.42 (d, $J=8.5$ Hz, 2H), 7.31 (d, $J=8.3$ Hz, 2H), 7.10 (dd, $J=8.4, 7.5$ Hz, 2H), 6.88 (d, $J=8.7$ Hz, 2H), 6.79~6.71 (m, 3H), 2.95~2.90 (m, 1H), 1.23 (s, 3H), 1.22 (s, 3H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.2, 149.7, 145.4, 138.6, 131.6, 129.0, 128.2, 127.4, 126.8, 120.8, 119.4, 116.5, 116.2, 64.9, 34.0, 23.8; HRMS (ESI) calcd for $C_{23}H_{21}N_2O$ $[M-H]^-$ 341.1659, found 341.1655.

2-(4-(tert-Butyl)phenyl)-2-(4-hydroxyphenyl)-2-(phenylamino)acetonitrile (**3e**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9 \sim 1:4$) as eluent to afford **3e** (31.3 mg, 88% yield). 1H NMR (500 MHz, Acetone- d_6) δ : 8.64 (s, 1H), 7.53~7.50 (m, 2H), 7.49~7.45 (m, 2H), 7.44~7.41 (m, 2H), 7.17~7.05 (m, 2H), 6.91~6.85 (m, 2H), 6.82~6.68 (m, 3H), 5.98 (s, 1H), 1.30 (s, 9H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.2, 152.0, 145.5, 138.3, 131.5, 129.0, 128.2, 126.5, 126.4, 120.8, 119.4, 116.5, 116.2, 64.9, 34.8, 31.2; HRMS (ESI) calcd for $C_{24}H_{23}N_2O$ $[M-H]^-$ 355.1816, found 355.1814.

2-(4-Bromophenyl)-2-(4-hydroxyphenyl)-2-(phenylamino)acetonitrile (**3f**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9 \sim 1:4$) as eluent to afford **3f** 33.9 mg, 90% yield. 1H NMR (500 MHz, Acetone- d_6) δ : 8.72 (s, 1H),

7.63 (d, $J=8.5$ Hz, 2H), 7.59~7.54 (m, 2H), 7.42 (d, $J=8.7$ Hz, 2H), 7.16~7.09 (m, 2H), 6.93~6.87 (m, 2H), 6.80~6.72 (m, 3H), 6.07 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.2, 144.9, 140.2, 132.4, 130.9, 128.9, 128.7, 128.1, 122.4, 120.2, 119.6, 116.4, 116.1, 64.6; HRMS (ESI) calcd for $C_{20}H_{14}BrN_2O$ $[M-H]^-$ 377.0295, found 377.0291.

2-(4-Chlorophenyl)-2-(4-hydroxyphenyl)-2-(phenylamino)acetonitrile (**3g**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9 \sim 1:4$) as eluent to afford **3g** 29.6 mg, 89% yield. 1H NMR (500 MHz, Acetone- d_6) δ : 8.70 (s, 1H), 7.66~7.59 (m, 2H), 7.51~7.45 (m, 2H), 7.43~7.38 (m, 2H), 7.12 (dd, $J=8.5, 7.5$ Hz, 2H), 6.91~6.85 (m, 2H), 6.79~6.72 (m, 3H), 6.07 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.5, 145.1, 139.9, 134.5, 131.2, 129.6, 129.1, 128.7, 128.3, 120.4, 119.7, 116.6, 116.3; HRMS (ESI) calcd for $C_{20}H_{14}ClN_2O$ $[M-H]^-$ 333.0800, found 333.0803.

2-(4-Forophenyl)-2-(4-hydroxyphenyl)-2-(phenylamino)acetonitrile (**3h**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9 \sim 1:4$) as eluent to afford **3h** 29.2 mg, 92% yield. 1H NMR (500 MHz, Acetone- d_6) δ : 8.69 (s, 1H), 7.70~7.59 (m, 2H), 7.41 (d, $J=8.6$ Hz, 2H), 7.24~7.17 (m, 2H), 7.15~7.07 (m, 2H), 6.89 (d, $J=8.6$ Hz, 2H), 6.79~6.71 (m, 3H), 6.06 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 163.5, 161.6, 157.9, 144.7, 136.6, 136.6, 130.9, 128.5, 127.8, 120.1, 119.1, 116.0, 115.8, 115.8, 115.6, 64.1; HRMS (ESI) calcd for $C_{20}H_{14}FN_2O$ $[M-H]^-$ 317.1096, found 317.1099.

2-(4-Hydroxyphenyl)-2-(phenylamino)-2-(4-(trifluoromethyl)phenyl)acetonitrile (**3i**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9 \sim 1:4$) as eluent to afford **3i** 34.5 mg, 94% yield. 1H NMR (500 MHz, Acetone- d_6) δ : 8.75 (s, 1H), 7.88 (d, $J=8.1$ Hz, 2H), 7.81 (d, $J=8.4$ Hz, 2H), 7.45 (d, $J=7.4$ Hz, 2H), 7.14 (t, $J=7.9$ Hz, 2H), 6.95~6.85 (m, 2H), 6.84~6.72 (m, 3H), 6.18 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.1, 158.0, 144.8, 144.5, 144.5, 130.4, 130.2, 130.0, 128.6, 127.8, 127.1, 126.1, 126.1, 125.2, 123.1, 119.7, 119.4, 116.1, 115.9, 115.8, 64.5; HRMS (ESI) calcd for $C_{21}H_{14}F_3N_2O$ $[M-H]^-$ 367.1064, found 367.1069.

2-(4-Hydroxyphenyl)-2-(phenylamino)-2-(*m*-tolyl)acetonitrile (**3j**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9 \sim 1:4$) as eluent to afford **3j** 27.6 mg, 88% yield. 1H NMR (500 MHz, Acetone- d_6) δ : 8.65 (s, 1H), 7.46 (s, 1H), 7.41 (dd, $J=11.9, 5.1$ Hz, 3H), 7.31 (t, $J=7.7$ Hz, 1H), 7.19 (d, $J=7.5$ Hz, 1H), 7.14~7.06 (m, 2H), 6.90~6.85 (m, 2H), 6.76 (d, $J=8.6$ Hz, 3H), 5.97 (s, 1H), 2.34 (s, 3H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.2, 145.4, 141.0, 139.2, 131.6, 129.7, 129.3, 129.0, 128.3, 127.2, 123.8, 120.8, 119.4, 116.5, 116.1, 65.1, 21.1; HRMS (ESI) calcd for $C_{21}H_{17}N_2O$ $[M-H]^-$ 313.1346, found 313.1343.

2-(3-Bromophenyl)-2-(4-hydroxyphenyl)-2-(phenylami-

no)acetonitrile (**3k**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **3k** 34.7 mg, 92% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.73 (s, 1H), 7.78 (s, 1H), 7.65 (dd, $J=8.0, 0.7$ Hz, 1H), 7.59~7.55 (m, 1H), 7.47~7.40 (m, 3H), 7.14 (dd, $J=8.4, 7.5$ Hz, 2H), 6.91 (d, $J=8.6$ Hz, 2H), 6.79~6.74 (m, 3H), 6.13 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.4, 144.9, 143.4, 132.0, 131.5, 130.8, 129.4, 129.0, 128.1, 125.7, 122.9, 120.1, 119.7, 116.5, 116.2, 64.6; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{14}\text{BrN}_2\text{O} [\text{M}-\text{H}]^-$ 377.0295, found 377.0293.

2-(3-Chlorophenyl)-2-(4-hydroxyphenyl)-2-(phenylamino)acetonitrile (**3l**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **3l** 30.3 mg, 91% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.75 (d, $J=1.8$ Hz, 1H), 7.66~7.56 (m, 2H), 7.53~7.38 (m, 4H), 7.13 (dd, $J=8.5, 7.5$ Hz, 2H), 6.90 (d, $J=8.5$ Hz, 2H), 6.82~6.71 (m, 3H), 6.14 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.4, 144.9, 143.2, 134.8, 131.2, 130.9, 129.1, 129.0, 128.1, 126.5, 125.3, 120.1, 119.7, 116.5, 116.2, 64.7; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{14}\text{ClN}_2\text{O} [\text{M}-\text{H}]^-$ 333.0800, found 333.0797.

2-(3-Fluorophenyl)-2-(4-hydroxyphenyl)-2-(phenylamino)acetonitrile (**3m**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **3m** 29.8 mg, 94% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.71 (s, 1H), 7.53~7.48 (m, 2H), 7.45 (dd, $J=6.9, 1.9$ Hz, 2H), 7.36 (dd, $J=10.3, 1.0$ Hz, 1H), 7.19~7.08 (m, 3H), 6.90 (dd, $J=7.0, 1.7$ Hz, 2H), 6.80~6.72 (m, 3H), 6.11 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 164.4, 162.5, 158.5, 145.04, 143.8, 143.7, 131.6, 131.6, 131.1, 129.1, 128.2, 122.7, 122.7, 120.3, 119.8, 116.6, 116.3, 116.0, 115.8, 113.8, 113.6, 64.8; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{14}\text{FN}_2\text{O} [\text{M}-\text{H}]^-$ 317.1096, found 317.1091.

2-(4-Hydroxyphenyl)-2-(phenylamino)-2-(*o*-tolyl)acetonitrile (**3n**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **3n** 28.2 mg, 90% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.73 (s, 1H), 7.47 (d, $J=7.7$ Hz, 1H), 7.37~7.22 (m, 5H), 7.12 (dd, $J=8.4, 7.5$ Hz, 2H), 6.89 (d, $J=8.7$ Hz, 2H), 6.80~6.69 (m, 3H), 5.86 (s, 1H), 2.36 (s, 3H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.0, 144.8, 136.5, 136.3, 132.8, 129.8, 128.8, 128.4, 128.0, 126.0, 120.3, 119.4, 118.7, 115.8, 115.6, 63.1, 20.3; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{17}\text{N}_2\text{O} [\text{M}-\text{H}]^-$ 313.1346, found 313.1343.

2-(3,5-Dimethylphenyl)-2-(4-hydroxyphenyl)-2-(phenylamino)acetonitrile (**3o**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **3o** 28.1 mg, 86% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.64 (s, 1H), 7.41 (d, $J=8.7$ Hz, 2H), 7.24 (s, 2H), 7.10 (dd, $J=8.6, 7.4$ Hz, 2H), 7.00 (s, 1H), 6.87 (dd, $J=6.9, 4.9$ Hz, 2H), 6.75 (t, $J=8.2$ Hz, 3H), 5.93 (s, 1H), 2.29 (s, 6H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 157.7, 145.0, 140.6, 138.5,

131.2, 130.0, 128.4, 127.7, 123.9, 120.3, 118.9, 116.0, 115.6, 64.6, 20.5; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{19}\text{N}_2\text{O} [\text{M}-\text{H}]^-$ 327.1503, found 327.1507.

2-(4-Hydroxyphenyl)-2-(naphthalen-2-yl)-2-(phenylamino)acetonitrile (**3p**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **3p** 32.8 mg, 94% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.67 (s, 1H), 8.18 (s, 1H), 8.00~7.91 (m, 3H), 7.70 (d, $J=8.7$ Hz, 1H), 7.58~7.53 (m, 2H), 7.52~7.43 (m, 2H), 7.11 (t, $J=7.9$ Hz, 2H), 6.88 (d, $J=7.2$ Hz, 2H), 6.81 (d, $J=7.9$ Hz, 2H), 6.75 (t, $J=7.4$ Hz, 1H), 6.13 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.3, 145.4, 138.3, 133.7, 131.5, 129.5, 129.0, 128.9, 128.4, 128.1, 127.4, 127.3, 125.8, 124.4, 120.7, 119.6, 65.3; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{17}\text{N}_2\text{O} [\text{M}-\text{H}]^-$ 349.1346, found 349.1343.

2-(4-Hydroxyphenyl)-2-(naphthalen-1-yl)-2-(phenylamino)acetonitrile (**3q**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **3q** 31.4 mg, 90% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.76 (d, $J=16.6$ Hz, 1H), 8.42~8.30 (m, 1H), 8.04~7.97 (m, 2H), 7.62~7.46 (m, 4H), 7.45~7.39 (m, 2H), 7.19~7.06 (m, 2H), 6.91 (d, $J=8.8$ Hz, 2H), 6.85~6.79 (m, 2H), 6.79~6.69 (m, 1H), 6.07 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.5, 145.2, 135.5, 134.2, 130.9, 130.6, 130.3, 129.7, 129.4, 129.0, 127.5, 126.7, 126.4, 125.9, 125.3, 120.4, 120.3, 119.3, 116.4, 116.3, 63.8; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{17}\text{N}_2\text{O} [\text{M}-\text{H}]^-$ 349.1346, found 349.1348.

2-(4-Hydroxy-3-methylphenyl)-2-phenyl-2-(phenylamino)acetonitrile (**3r**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **3r** 28.2 mg, yield 90%. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.54 (s, 1H), 7.62 (d, $J=7.4$ Hz, 2H), 7.43 (t, $J=7.6$ Hz, 2H), 7.39~7.32 (m, 2H), 7.21 (d, $J=8.3$ Hz, 1H), 7.14~7.06 (m, 2H), 6.84 (d, $J=8.4$ Hz, 1H), 6.79~6.72 (m, 3H), 5.98 (s, 1H), 2.19 (s, 3H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 156.2, 145.4, 141.1, 131.6, 129.5, 129.3, 129.0, 126.8, 125.5, 125.5, 120.8, 119.4, 116.5, 115.3, 65.1, 16.0; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{17}\text{N}_2\text{O} [\text{M}-\text{H}]^-$ 313.1346, found 313.1342.

2-(3-Bromo-4-hydroxyphenyl)-2-phenyl-2-(phenylamino)acetonitrile (**3s**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **3s** 34.7 mg, 92% yield. ^1H NMR (500 MHz, CDCl_3) δ : 7.72 (d, $J=2.3$ Hz, 1H), 7.61~7.57 (m, 2H), 7.47~7.37 (m, 4H), 7.17 (t, $J=7.9$ Hz, 2H), 7.02 (d, $J=8.6$ Hz, 1H), 6.87 (t, $J=7.4$ Hz, 1H), 6.63 (d, $J=7.9$ Hz, 2H), 4.41 (s, 1H); ^{13}C NMR (126 MHz, CDCl_3) δ : 153.1, 143.6, 139.6, 133.3, 130.3, 129.6, 129.4, 129.3, 127.5, 126.4, 120.6, 119.8, 116.9, 116.4, 111.0, 64.7; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{14}\text{BrN}_2\text{O} [\text{M}-\text{H}]^-$ 377.0295, found 377.0299.

2-(3-Chloro-4-hydroxyphenyl)-2-phenyl-2-(phenylamino)acetonitrile (**3t**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V:$

$V=1:9\sim 1:4$) as eluent to afford **3t** 30.3 mg, 91% yield. ^1H NMR (500 MHz, CDCl_3) δ : 7.65~7.56 (m, 3H), 7.46~7.37 (m, 4H), 7.20~7.14 (m, 2H), 7.06~7.01 (m, 1H), 6.87 (t, $J=7.4$ Hz, 1H), 6.63 (d, $J=7.7$ Hz, 2H), 5.72 (s, 1H), 4.37 (s, 1H); ^{13}C NMR (126 MHz, CDCl_3) δ : 152.1, 143.5, 139.6, 133.0, 129.6, 129.4, 129.3, 127.3, 126.8, 126.4, 120.8, 120.6, 119.7, 117.1, 116.4, 64.8; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{14}\text{ClN}_2\text{O}$ $[\text{M}-\text{H}]^-$ 333.0800, found 333.0803.

2-(3-Fluoro-4-hydroxyphenyl)-2-phenyl-2-(phenylamino)acetonitrile (**3u**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **3u** 29.8 mg, 94% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 9.00 (s, 1H), 7.66~7.61 (m, 2H), 7.48~7.43 (m, 2H), 7.42~7.36 (m, 1H), 7.35~7.26 (m, 2H), 7.15~7.09 (m, 2H), 7.06 (t, $J=8.8$ Hz, 1H), 6.77 (t, $J=8.1$ Hz, 3H), 6.12 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 152.8, 150.8, 145.8, 145.7, 145.1, 140.6, 132.6, 132.6, 129.6, 129.3, 129.0, 126.7, 123.2, 123.2, 120.3, 119.7, 118.7, 118.7, 116.6, 114.9, 114.7, 64.8; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{14}\text{FN}_2\text{O}$ $[\text{M}-\text{H}]^-$ 317.1096, found 317.1091.

2-(4-Hydroxyphenyl)-2-phenyl-2-(*p*-tolylamino)acetonitrile (**4a**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **4a** 28.8 mg, 92% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.67 (s, 1H), 7.63 (d, $J=7.6$ Hz, 2H), 7.42 (t, $J=7.9$ Hz, 4H), 7.36 (t, $J=7.3$ Hz, 1H), 6.90 (dd, $J=19.7$, 8.3 Hz, 4H), 6.68 (d, $J=8.2$ Hz, 2H), 5.84 (s, 1H), 2.18 (s, 3H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 157.9, 142.7, 140.9, 131.5, 129.2, 129.2, 128.7, 128.2, 128.0, 126.5, 120.6, 116.5, 115.9, 65.1, 19.9; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{17}\text{N}_2\text{O}$ $[\text{M}-\text{H}]^-$ 313.1346, found 313.1350.

2-((4-Bromophenyl)amino)-2-(4-hydroxyphenyl)-2-phenylacetonitrile (**4b**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **4b** 34.3 mg, 91% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.71 (s, 1H), 7.60 (d, $J=7.6$ Hz, 2H), 7.47~7.37 (m, 5H), 7.27 (d, $J=8.8$ Hz, 2H), 6.91~6.86 (m, 2H), 6.75~6.68 (m, 2H), 6.26 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.4, 144.6, 140.4, 131.8, 131.0, 129.6, 129.2, 128.3, 126.7, 120.4, 118.3, 116.2, 111.1, 65.0; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{14}\text{BrN}_2\text{O}$ $[\text{M}-\text{H}]^-$ 377.0295, found 377.0298.

2-((4-Chlorophenyl)amino)-2-(4-hydroxyphenyl)-2-phenylacetonitrile (**4c**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **4c** 30.6 mg, 92% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.72 (s, 1H), 7.61 (d, $J=7.6$ Hz, 2H), 7.50~7.33 (m, 5H), 7.14 (d, $J=8.8$ Hz, 2H), 6.90 (d, $J=8.7$ Hz, 2H), 6.77 (d, $J=8.8$ Hz, 2H), 6.23 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.1, 143.9, 140.2, 130.8, 129.3, 128.9, 128.6, 128.0, 126.5, 123.6, 120.2, 117.6, 116.0, 64.9; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{14}\text{ClN}_2\text{O}$ $[\text{M}-\text{H}]^-$ 333.0800, found 333.0797.

2-(4-Hydroxyphenyl)-2-phenyl-2-((4-(trifluoromethoxy)phenyl)amino)acetonitrile (**4d**): It was purified by flash

chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **4d** 36.4 mg, 95% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.73 (s, 1H), 7.66~7.58 (m, 2H), 7.48~7.35 (m, 5H), 7.12 (d, $J=8.5$ Hz, 2H), 6.94~6.87 (m, 2H), 6.87~6.80 (m, 2H), 6.34 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.2, 144.3, 141.7, 140.2, 130.8, 129.4, 129.0, 128.1, 126.5, 122.0, 120.2, 120.0, 116.9, 116.1, 64.9; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{14}\text{F}_3\text{N}_2\text{O}_2$ $[\text{M}-\text{H}]^-$ 383.1013, found 383.1009.

2-(4-Hydroxyphenyl)-2-phenyl-2-(*m*-tolylamino)acetonitrile (**4e**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **4e** 27.8 mg, 89% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.68 (s, 1H), 7.64~7.58 (m, 2H), 7.46~7.39 (m, 4H), 7.36 (t, $J=4.6$ Hz, 1H), 6.97 (t, $J=7.8$ Hz, 1H), 6.90~6.84 (m, 2H), 6.65 (s, 1H), 6.58 (d, $J=7.5$ Hz, 1H), 6.51~6.45 (m, 1H), 5.92 (s, 1H), 2.16 (s, 3H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.1, 145.2, 141.0, 138.3, 131.6, 129.4, 128.9, 128.7, 128.2, 126.7, 120.7, 120.3, 117.3, 116.1, 113.4, 65.0, 21.2; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{17}\text{N}_2\text{O}$ $[\text{M}-\text{H}]^-$ 313.1346, found 313.1343.

2-((3-Chlorophenyl)amino)-2-(4-hydroxyphenyl)-2-phenylacetonitrile (**4f**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **4f** 30.0 mg, 90% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.71 (s, 1H), 7.61 (d, $J=7.6$ Hz, 2H), 7.46 (t, $J=7.6$ Hz, 2H), 7.43~7.37 (m, 3H), 7.11 (t, $J=8.1$ Hz, 1H), 6.90 (d, $J=8.7$ Hz, 2H), 6.82 (d, $J=1.7$ Hz, 1H), 6.77 (d, $J=7.9$ Hz, 1H), 6.68 (dd, $J=8.2$, 1.6 Hz, 1H), 6.34 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.4, 146.7, 140.3, 134.2, 130.9, 130.4, 129.6, 129.2, 128.2, 126.7, 120.3, 119.1, 116.2, 116.1, 114.7, 64.9; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{14}\text{ClN}_2\text{O}$ $[\text{M}-\text{H}]^-$ 333.0800, found 333.0805.

2-(4-Hydroxyphenyl)-2-phenyl-2-(*o*-tolylamino)acetonitrile (**4g**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **4g** 27.6 mg, 88% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.69 (s, 1H), 7.69~7.62 (m, 2H), 7.48~7.41 (m, 4H), 7.38~7.34 (m, 1H), 7.13 (d, $J=7.2$ Hz, 1H), 6.93~6.86 (m, 3H), 6.74~6.69 (m, 1H), 6.49 (d, $J=8.1$ Hz, 1H), 5.03 (s, 1H), 2.32 (s, 3H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.3, 142.6, 141.1, 131.8, 130.7, 129.6, 129.0, 128.0, 126.5, 126.3, 125.5, 120.8, 119.7, 116.3, 115.4, 64.7, 17.6; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{17}\text{N}_2\text{O}$ $[\text{M}-\text{H}]^-$ 313.1346, found 313.1348.

2-(4-Hydroxyphenyl)-2-(naphthalen-2-ylamino)-2-phenylacetonitrile (**4h**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V: V=1:9\sim 1:4$) as eluent to afford **4h** 32.2 mg, 92% yield. ^1H NMR (500 MHz, Acetone- d_6) δ : 8.77 (s, 1H), 7.69 (dd, $J=21.8$, 8.1 Hz, 4H), 7.49~7.36 (m, 6H), 7.33~7.25 (m, 2H), 7.22 (t, $J=7.3$ Hz, 1H), 6.90 (d, $J=8.5$ Hz, 2H), 6.82 (s, 1H), 6.31 (s, 1H); ^{13}C NMR (126 MHz, Acetone- d_6) δ : 158.3, 142.8, 140.6, 134.7, 131.2, 129.5, 129.3, 129.0, 128.9, 128.6, 128.3, 127.9, 126.8, 126.6, 123.2, 120.5, 119.8, 116.2, 110.0, 65.1; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{17}$

$\text{N}_2\text{O} [\text{M}-\text{H}]^-$ 349.1346, found 349.1343.

2-(4-Hydroxyphenyl)-2-morpholino-2-phenylacetonitrile (**6**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V:V=1:9\sim 1:4$) as eluent to afford **6** 25.0 mg, 85% yield. ^1H NMR (500 MHz, CDCl_3) δ : 7.67 (d, $J=7.6$ Hz, 2H), 7.53 (d, $J=8.6$ Hz, 2H), 7.39~7.33 (m, 2H), 7.29~7.27 (m, 1H), 6.81 (d, $J=8.7$ Hz, 2H), 5.69 (s, 1H), 3.85~3.77 (m, 4H), 2.60 (d, $J=4.1$ Hz, 4H); ^{13}C NMR (126 MHz, CDCl_3) δ : 156.1, 139.3, 130.9, 129.3, 128.6, 128.0, 126.4, 117.0, 116.1, 75.5, 67.2, 49.6; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{17}\text{N}_2\text{O}_2$ $[\text{M}-\text{H}]^-$ 293.1296, found 293.1291.

2-(4-Hydroxyphenyl)-2-(1H-imidazol-1-yl)-2-phenylacetonitrile (**9**): It was purified by flash chromatography on silica gel using acetone/petroleum ether ($V:V=1:9\sim 1:4$) as eluent to afford **9** 24.7 mg, 90% yield. ^1H NMR (400 MHz, CD_3OD) δ : 7.52~7.46 (m, 3H), 7.43 (s, 1H), 7.22~7.09 (m, 4H), 7.06 (d, $J=8.7$ Hz, 2H), 6.87 (d, $J=8.7$ Hz, 2H); ^{13}C NMR (101 MHz, CD_3OD) δ : 160.6, 138.4, 131.4, 131.1, 130.6, 130.5, 130.4, 128.3, 128.1, 127.6, 120.3, 117.1, 67.9; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{12}\text{N}_3\text{O}$ $[\text{M}-\text{H}]^-$ 274.0986, found 274.0990.

Supporting Information Copies of ^1H NMR and ^{13}C NMR spectra. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Turner, A. B. *Q. Rev., Chem. Soc.* **1964**, 18, 347.
(b) Peter, M. G. *Angew. Chem. Int. Ed.* **1989**, 28, 555.
(c) Itoh, T. *Prog. Polym. Sci.* **2001**, 26, 1019.
(d) Parra, A.; Tortosa, M. *ChemCatChem* **2015**, 7, 1524.
(e) Li, W.; Xu, X.; Zhang, P.; Li, P. *Chem. Asian J.* **2018**, 13, 2350.
- [2] (a) Angle, S. R.; Turnbull, K. D. *J. Am. Chem. Soc.* **1989**, 111, 1136.
(b) Angle, S. R.; Arnaiz, D. O. *J. Org. Chem.* **1990**, 55, 3708.
(c) Baik, W.; Lee, H. J.; Jang, J. M.; Koo, S.; Kim, B. H. *J. Org. Chem.* **2000**, 65, 108.
(d) Reddy, V.; Anand, R. V. *Org. Lett.* **2015**, 17, 3390.
(e) Ramanjaneyulu, B. T.; Mahesh, S.; Anand, R. V. *Org. Lett.* **2015**, 17, 3952.
(f) Shen, Y.; Qi, J.; Mao, Z.; Cui, S. *Org. Lett.* **2016**, 18, 2722.
(g) Huang, X. Y.; Ding, R.; Mo, Z. Y.; Xu, Y. L.; Tang, H. T.; Wang, H. S.; Chen, Y. Y.; Pan, Y. M. *Org. Lett.* **2018**, 20, 4819.
(h) Wu, Q. Y.; Ao, G. Z.; Liu, F. *Org. Chem. Front.* **2018**, 5, 2061.
(i) Ke, M.; Song, Q. *Adv. Synth. Catal.* **2017**, 359, 384.
- [3] (a) Chu, W. D.; Zhang, L. F.; Bao, X.; Zhao, X. H.; Zeng, C.; Du, J. Y.; Zhang, G. B.; Wang, F. X.; Ma, X. Y.; Fan, C. A. *Angew. Chem. Int. Ed.* **2013**, 52, 9229.
(b) Caruana, L.; Kniep, F.; Johansen, T. K.; Poulsen, P. H.; Jørgensen, K. A. *J. Am. Chem. Soc.* **2014**, 136, 15929.
(c) Lou, Y.; Cao, P.; Jia, T.; Zhang, Y.; Wang, M.; Liao, J. *Angew. Chem. Int. Ed.* **2015**, 54, 12134.
(d) Dong, N.; Zhang, Z. P.; Xue, X. S.; Li, X.; Cheng, J. P. *Angew. Chem. Int. Ed.* **2016**, 55, 1460.
(e) Li, X.; Xu, X.; Wei, W.; Lin, A.; Yao, H. *Org. Lett.* **2016**, 18, 428.
(f) Ge, L.; Lu, X.; Cheng, C.; Chen, J.; Cao, W.; Wu, X.; Zhao, G. J. *Org. Chem.* **2016**, 81, 9315.
(g) Ma, C.; Huang, Y.; Zhao, Y. *ACS Catal.* **2016**, 6, 6408.
(h) He, F. S.; Jin, J. H.; Yang, Z. T.; Yu, X.; Fossey, J. S.; Deng, W. P. *ACS Catal.* **2016**, 6, 652.
(i) Jarava-Barrera, C.; Parra, A.; López, A.; Cruz-Acosta, F.; Collado-Sanz, D.; Cárdenas, D. J.; Tortosa, M. *ACS Catal.* **2016**, 6, 442.
(j) Li, S.; Liu, Y.; Huang, B.; Zhou, T.; Tao, H.; Xiao, Y.; Liu, L.; Zhang, J. *ACS Catal.* **2017**, 7, 2805.
(k) Huang, G. B.; Huang, W. H.; Guo, J.; Xu, D. L.; Qu, X. C.; Zhai, P. H.; Zheng, X. H.; Weng, J.; Lu, G. *Adv. Synth. Catal.* **2019**, 361, 1241.
- [4] Errede, L. A.; Szwarc, M. *Q. Rev., Chem. Soc.* **1958**, 12, 301.
- [5] (a) Wang, Z.; Wong, Y. F.; Sun, J. *Angew. Chem. Int. Ed.* **2015**, 54, 13711.
(b) Chen, M.; Sun, J. *Angew. Chem. Int. Ed.* **2017**, 56, 11966.
- [6] (a) Wang, Z.; Zhu, Y.; Pan, X.; Wang, G.; Liu, L. *Angew. Chem. Int. Ed.* **2020**, 59, 3053.
(b) Pan, X.; Wang, Z.; Kan, L.; Mao, Y.; Zhu, Y.; Liu, L. *Chem. Sci.* **2020**, 11, 2414.
- [7] Qi, Y.; Zhang, F.; Wang, L.; Feng, A.; Zhu, R.; Sun, S.; Li, W.; Liu, L. *Org. Biomol. Chem.* **2020**, 18, 3522.
- [8] (a) Beatty, J. W.; Stephenson, C. R. J. *Acc. Chem. Res.* **2015**, 48, 1474.
(b) Enders, D.; Reinhold, U. *Tetrahedron: Asymmetry* **1997**, 8, 1895.
(c) Lu, Z.; Ma, S. *Angew. Chem. Int. Ed.* **2008**, 47, 258.
(d) Legnani, L.; Bhawal, B. N.; Morandi, B. *Synthesis* **2017**, 49, 776.
(e) Ellman, J. A. *Pure Appl. Chem.* **2003**, 75, 39.
- [9] (a) Jadhav, A. S.; Pankhade, Y. A.; Anand, R. V. *J. Org. Chem.* **2018**, 83, 8596.
(b) Jiang, F.; Yuan, F. R.; Jin, L. W.; Mei, G. J.; Shi, F. *ACS Catal.* **2018**, 8, 10234.
(c) Feng, Z.; Yuan, Z.; Zhao, X.; Huang, Y.; Yao, H. *Org. Chem. Front.* **2019**, 6, 3535.
(d) Roy, D.; Panda, G. *Synthesis* **2019**, 51, 4434.
(e) Torán, R.; Vila, C.; Sanz-Marco, A.; Muñoz, M.; Pedro, J.; Blay, G. *Eur. J. Org. Chem.* **2020**, 5, 627.
(f) Zhang, J.-R.; Jin, H.-S.; Wang, R.-B.; Zhao, L.-M. *Adv. Synth. Catal.* **2019**, 361, 4811.
- [10] (a) Otto, N.; Opatz, T. *Chem. Eur. J.* **2014**, 20, 13064.
(b) Huang, P. Q. *Acta Chim. Sinica* **2018**, 76, 357 (in Chinese). (黄培强, 化学学报, **2018**, 76, 357.)
(c) Gao, Y. J.; Xiao, Z. H.; Liu, L. X.; Huang, P. Q. *Chin. J. Org. Chem.* **2017**, 37, 1189.
- [11] Singh, M.; Schott, J. T.; Leon, M. A.; Granata, R. T.; Dhaha, H. K.; Welles, J. A.; Boyce, M. A.; Oseni-Olalemi, H. S.; Mordaunt, C. E.; Vargas, A. J.; Patel, N. V.; Maitra, S. *Bioorg. Med. Chem. Lett.* **2012**, 22, 6252.
- [12] (a) Tiffert, T.; Ginsburg, H.; Krugliak, M.; Elford, B. C.; Lew, V. L. *Proc. Natl. Acad. Sci. U. S. A.* **2000**, 97, 331.
(b) Saliba, K. J.; Kirk, K. *Trans. R. Soc. Trop. Med. Hyg.* **1998**, 92, 666.

(Li, L.; Fan, Y.)