中心手性金属铑配合物催化的不对称光诱导 Giese 自由基加成反应

陈亮^{a,b} 胡良建^{a,c} 杜宇^a 苏伟平^{*,a} 康强^{*,a}^(a) 中国科学院福建物质结构研究所 结构化学国家重点实验室 福州 350002)^(b) 中国科学院大学 北京 100049)^(c) 福建农林大学材料工程学院 福州 350002)

摘要 报道了一种可见光驱动的双功能中心手性金属铑配合物催化的不对称自由基加成反应, 由对氨基苄乙酸生成的对氨基苄基自由基对 α,β -不饱和酰基咪唑进行立体选择性的 1,4-加成, 以中高收率和高对映选择性合成了一系列加成产物. 该反应具有对映选择性高、反应条件温和、操作简单等特点. 根据对双功能中心手性金属配合物催化的研究和文献报道, 提出了一个合理的反应机理.

关键词 可见光催化; 双功能手性铑配合物; Giese 自由基加成反应; 对氨基苄基自由基

Asymmetric Photoinduced Giese Radical Addition Enabled by a Single Chiral-at-Metal Rhodium Complex

Chen, Liang^{a,b} Hu, Liangjian^{a,c} Du, Yu^a Su, Weiping^{*,a} Kang, Qiang^{*,a}^(a) State Key Laboratory of Structural Chemistry, Fujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences, Fuzhou 350002)^(b) University of Chinese Academy of Sciences, Beijing 100049)^(c) College of Material Engineering, Fujian Agriculture and Forestry University, Fuzhou 350002)

Abstract A visible-light driven asymmetric Giese radical addition catalyzed by a bifunctional chiral-at-metal rhodium complex has been developed. *para*-Aminobenzyl radicals generated from *para*-aminophenylacetic acids could undergo a 1,4-addition in a highly enantioselective approach to α,β -unsaturated 2-acyl imidazoles, and a variety of corresponding adducts were obtained in moderate to high yields with excellent enantioselectivities. This reaction features high enantioselectivity, mild reaction conditions and an operationally simple procedure. A plausible reaction mechanism is proposed based on our research and literature precedents on dual functional chiral-at-metal catalysis.

Keywords visible-light photoredox catalysis; bifunctional chiral-at-metal rhodium complex; Giese radical addition; *para*-aminobenzyl radical

1 Introduction

Visible-light photoredox catalysis which produces a various range of radical intermediates in a mild and convenient process has been identified as a vigorous tool in organic synthesis.^[1] However, due to the involvement of open-shell radical species and quick subsequent reactions featuring low energetic barriers, the development of catalytic asymmetric photoredox reactions remains a challenging task in controlling the stereoselectivity of highly reactive intermediates.^[2-3] Currently the major solutions for asymmetric catalysis promoted by visible light are typi-

cally relying on two strategies: (1) dual catalysis systems^[4] involving two catalysts; (2) asymmetric/photoredox bifunctional photocatalyst.^[5] The dual catalysis systems usually contain one photocatalyst for the radical generation and an additional chiral co-catalyst for controlling the enantioselectivity. On the other hand, single bifunctional photocatalyst integrates visible light harvest for activation of substrates and stereocontrol during the chemical bond formation together. In a word, both strategies show the potential to overcome the inherent difficulties of visible light driven asymmetric catalysis. For instance, the Giese-

* Corresponding authors. E-mail: wpsu@fjirsm.ac.cn, kangq@fjirsm.ac.cn

Received April 25, 2020; revised May 14, 2020; published online May 19, 2020.

Dedicated to the 40th anniversary of Chinese Journal of Organic Chemistry.

Project supported by the National Natural Science Foundation of China (No. 21871261).

国家自然科学基金面上(No. 21871261)资助项目.

type reaction^[6] that is the addition of nucleophilic alkyl radicals to electron-poor alkenes has long been considered as one of the most established bimolecular radical reactions. As a paradigm, Sibi, Porter and other researchers^[7] have revealed that chiral Lewis acid catalysts are capable of dominating the stereoselective conjugate addition of radicals. Inspired by these pioneering achievements, Yoon and co-workers^[8] developed the first highly enantioselective intermolecular conjugate addition of α -aminoalkyl radicals generated from α -silylalkyl anilines by merging visible-light photoredox and Lewis acid catalysis. Recently, Gong group^[5g] utilized a chiral nickel DBFOX complex as a single catalyst for a catalytic asymmetric conjugate radical addition.

Oxidative decarboxylation of carboxylic acids is a substantial process in nature. Since from the seminal work by MacMillan,^[9] the decarboxylation of α -amino acids via visible-light photoredox catalysis has become a powerful method to generate α -amino alkyl radicals.^[10] As the ‘phenylogue’ of the α -amino acids, synthetic utilization of the aminoarylacetic acids with visible light has been quite limited.^[11] In 2003, Miyake and co-workers^[11d] succeeded in utilizing *para*-aminophenylacetic acids as *para*-aminobenzyl radical precursors in addition reactions with electron-deficient alkenes, but no enantioselectivity was observed (Scheme 1, a). As a continuation of our interest in the development of chiral-at-metal complexes^[12-13] as bifunctional photocatalysts in catalytic asymmetric reactions, herein, we report our recent findings on a catalytic asymmetric conjugate addition of *para*-aminobenzyl radicals generated from *para*-aminophenylacetic acids to α,β -unsaturated carbonyl compounds catalyzed by a chiral-at-metal rhodium complex under visible light irradiation (Scheme 1, b).

2 Results and discussion

We initiated our studies with α,β -unsaturated 2-acyl imidazole **1a** and 4-(*N*-methyl-*N*-phenylamino)-phenylacetic acid **2a** as model substrates in the presence of 2 mol% of

chiral-at-metal Rh(III) complex **Λ -Rh1** developed by Meggers’ group under the irradiation with blue LEDs. To our delight, the reaction proceeded smoothly in degassed 1,2-dichloroethane (DCE) to afford the desired product **3a** in 77% yield with 82% *ee* (Table 1, Entry 1). Control

Table 1 Optimization of reaction conditions^a

Entry	Λ -M	Solvent	h ν	Yield ^b /%	<i>ee</i> ^c /%
1	Λ -Rh1	DCE	✓	77	82
2	Λ -Rh1	DCE	—	—	—
3	None	DCE	✓	—	—
4	Λ -Ir	DCE	✓	53	89
5	Λ -Rh2	DCE	✓	70	88
6	Λ -Rh2	MeOH	✓	79	96
7	Λ -Rh2	Toluene	✓	46	93
8	Λ -Rh2	DMSO	✓	31	60
9	Λ -Rh2	Acetone	✓	75	91
10	Λ -Rh2	CH ₃ CN	✓	65	87

^a Unless otherwise noted, reactions were carried out by using **1a** (0.1 mmol), **2a** (0.15 mmol) and Λ -M (0.002 mmol, 2 mol%) in degassed solvent (0.3 mL) at room temperature under irradiation with 20 W blue LEDs. ^b Isolated yields based on **1a**. ^c Determined by chiral HPLC analysis.

(a) Miyake, Nishibayashi (2013)

(b) This work

Scheme 1 Photoredox catalyzed addition of *para*-aminobenzyl radicals to electron-deficient alkenes

experiments devoid of catalyst (no reaction) or performed in the dark (no reaction) reveal that it is the combination of chiral rhodium complex **A-Rh1** and visible light that is required for an efficient reaction (Entries 2, 3). Encouraged by these promising results, other chiral-at-metal complexes were examined. **A-Ir** could lead to the formation of the target molecule **3a** with an increased enantioselectivity (89% *ee*) but with a diminished yield (53%, Entry 4). Gratifyingly, **A-Rh2** developed by our group gave the desired product in 70% yield with 88% *ee* (Entry 5). Further screening of solvents revealed that methanol was the superior one (Entries 6~10), leading to the formation of product **3a** in 79% yield with 96% *ee* (Entry 6). These reactions could be finished in 12 h.

With the optimal conditions established for **3a** (Table 1, Entry 6), the scope of this photoinduced conjugate addition reaction was then examined. The reaction between *para*-aminobenzyl radical precursor **2a** and α,β -unsaturated 2-acyl imidazoles **1** bearing a β -aryl substituent was first studied using the optimal conditions developed above. The results are summarized in Table 2. The introduction of electron-donating and electron-withdrawing groups on the phenyl ring had little influence on enantioselectivities. The

Table 2 Substrate scope of α,β -unsaturated 2-acyl imidazoles^a

^a Unless otherwise noted, reactions were carried out by using **1** (0.1 mmol), **2a** (0.15 mmol) and **A-Rh2** (2 mol%) in degassed MeOH (0.3 mL) at room temperature under irradiation with 20 W blue LEDs. All isolated yields were based on substrate **1**, *ee* values were determined by chiral HPLC analysis.

desired products **3b**~**3h** were obtained in moderate to high yields (58%~95%) with high enantioselectivities (91%~96% *ee*). Naphthyl-based substrates **1i**, **1j** and heteroaromatic (eg. thienyl) **1k** were also well tolerated, affording **3i**~**3k** in moderate yields (54%~72%) with high enantioselectivities (90%~95% *ee*). α,β -Unsaturated 2-acyl imidazole with a β -alkyl substituent (eg. isopropyl) also worked well under the optimal reaction conditions, although the conversion was incomplete even after 48 h, affording product **3l** in moderate yield and high enantioselectivity (50% yield, 91% *ee*). The relative configuration of compound **3h** was confirmed by X-ray crystallographic studies.^[14]

Further examination of the substrate scope of arylacetic acids **2** was also conducted using the optimal conditions with **1a** (Table 3). Phenylacetic acids bearing a tertiary amino group such as *N*-dimethylamino, *N*-methyl-*N*-benzylamino, *N*-dibenzylamino moieties at the *para*-position of the aryl substituent were applicable, giving the corresponding products **4a**~**4c** in moderate yields (up to 83%) with good enantioselectivities (up to 91% *ee*). The phenylacetic acid bearing a secondary amino group (eg., *N*-benzylamino) at the *para*-position also reacted smoothly to give **4d** in 70% yield with 95% *ee*. *para*-Primary amino phenylacetic acid also worked for this reaction, although a low yield and enantioselectivity was obtained (**4e**, 38% yield, 58% *ee*). In addition, consistent with the report of Miyake *et al.*,^[11d] 4-methoxy-substituted arylacetic acid and phenylacetic acids with an amino group at the *meta*- or *ortho*-position of the aryl substituent (**2g**~**2i**) could not give any products under the optimal reaction conditions, probably because the radical character of their one-electron oxidized intermediate is too small to induce decarboxylation, or an intramolecular hydrogen bonding between the amino group and carboxylic acid inhibits the electron transfer oxidation.

Table 3 Substrate scope of arylacetic acids^a

Reaction scheme for Table 3: $\text{1a} + \text{2b} \sim \text{2i} \xrightarrow[\text{r.t., 12 h}]{\text{A-Rh2 (2.0 mol\%), hv, MeOH}} \text{4}$

Entry	Ar (2)	4	Yield ^b /%	<i>ee</i> ^c /%
1	4-Me ₂ NC ₆ H ₄ (2b)	4a	36	90
2	4-BnMeNC ₆ H ₄ (2c)	4b	83	91
3	4-Bn ₂ NC ₆ H ₄ (2d)	4c	60	80
4	4-BnHNC ₆ H ₄ (2e)	4d	70	95
5 ^d	4-H ₂ NC ₆ H ₄ (2f)	4e	38	58
6	4-MeOC ₆ H ₄ (2g)	—	—	—
7	3-PhMeNC ₆ H ₄ (2h)	—	—	—
8	2-PhMeNC ₆ H ₄ (2i)	—	—	—

^a Unless otherwise noted, reactions were carried out by using **1a** (0.1 mmol), **2** (0.15 mmol) and **A-Rh2** (2 mol%) in degassed MeOH (0.3 mL) at room temperature under irradiation with 20 W blue LEDs. ^b Isolated yields based on **1a**. ^c Determined by chiral HPLC analysis. ^d DMSO was used in place of MeOH.

To illustrate the potential application of current protocol, a large-scale reaction of α,β -unsaturated 2-acyl imidazole **1h** (0.52 g, 2.0 mmol) with arylacetic acid **2a** (0.72 g, 3.0 mmol) was conducted in the presence of 2 mol% of Δ -Rh2 (Scheme 2, a). Gratifyingly, the reaction proceeded smoothly to afford **3h** in 54% yield (0.49 g) with 93% *ee*. Moreover, the imidazole moiety of products could be easily transferred to other functional groups.^[15] For example, the removal of the imidazole moiety of **3h** worked smoothly to afford ester **5** in good yield without any loss in enantiomeric excess (Scheme 2, b).

Based on control experiments and literature precedents

on dual functional chiral-at-metal catalysis,^[5g,13,16] a plausible reaction pathway is proposed in Scheme 3. Substrate **1a** first coordinates with the rhodium complex Δ -Rh2 in a bidentate fashion to generate a *N,O*-coordinated intermediate **I**, which could be photoexcited (**I**^{*}) and then oxidizes aminoarylacetic acid **2a** to generate a benzyl radical **II** and the reduced **I**[−] via a single electron transfer (SET) process. The benzyl radical **II** then trapped by the intermediate **I** in a highly stereoselective approach to form a radical intermediate **III**. This intermediate **III** undergoes another SET process with strong reductant **I**[−] to regenerate **I** and furnish an anion **IV**, followed by protonation to afford a neutral

Scheme 2 Large-scale experiment and synthetic transformation of product **3h**

Scheme 3 Proposed mechanism and transition-state model

complex **V**. The substitution of complex **V** with **1a** eventually releases the desired product **3a**, and a new catalytic cycle is initiated.

3 Conclusions

In conclusion, a highly efficient and practical catalytic enantioselective Giese radical addition of *para*-aminobenzyl radicals generated from *para*-aminophenylacetic acids to electron-deficient alkenes catalyzed by a bifunctional chiral-at-metal rhodium complex has been developed, affording the desired adducts in moderate to high yields with excellent enantioselectivities. Here the rhodium complex works not only as a chiral Lewis acid to improve the electrophilicity of substrates and control the stereoselectivity during the chemical bond formation, but also a photocatalyst to form photosensitive substrate-coordinated complex. This reaction features high enantioselectivity, mild reaction conditions and an operationally simple procedure. Further research on the development of new types of bifunctional chiral-at-metal complexes and their application in asymmetric reactions are ongoing in our laboratory.

4 Experimental section

4.1 Experimental details

All reactions were performed in Schlenk tubes under an atmosphere of argon using oven-dried glassware. Commercially obtained reagents were used without further purification, unless otherwise noted. Dry DCE, toluene, CH₃CN and tetrahydrofuran (THF) were obtained from solvent distillation machine (Vigor VSPS-5). Dimethyl sulfoxide (DMSO) was distilled from CaH₂. Methanol and acetone (IPA) were used without further purification. Reactions were checked by thin-layer chromatography (TLC) analysis and plates were visualized with short-wave UV light (254 nm). The ¹H NMR and ¹³C NMR and ¹⁹F NMR spectra were obtained in CDCl₃ using a Bruker-BioSpin AVANCE III HD NMR spectrometer at 400, 100 and 376.4 MHz, respectively. HPLC analyses of the compounds were done using chiralcel IA-IF columns and chiralcel AD-H, AS-H, OJ-H and OD-H columns using hexane and isopropanol as eluent. The infrared spectra were recorded on a Bruker VERTEX 70 IR spectrometer as KBr pellets. High-resolution mass spectra were recorded on a Bruker Impact II UHR TOF LC/MS mass spectrometry. Chiral-at-metal complexes **A-Rh1**,^[17] **A-Rh2**^[13] and **A-Ir**^[18] were prepared according to reported procedure. α,β -Unsaturated 2-acyl imidazoles **1**^[13] and arylacetic acid (**2a**~**2e**, **2h** and **2i**)^[11d-11e] were synthesized according to reported procedures.

4.2 General procedure for chiral Rhodium(III) complex catalyzed enantioselective Giese addition reaction

To an oven-dried 25 mL Schlenk tube equipped with a stir bar, **A-Rh2** (2.5 mg, 2 mol%) was added along with

α,β -unsaturated 2-acyl imidazoles **1** (0.1 mmol, 1.0 equiv.), 4-amino phenylacetic acid **2** (1.5 equiv.) and MeOH (0.3 mL). After degassing via three freeze-pump-thaw cycles, the Schlenk tube was sealed and positioned at a distance of 5~10 cm from 20 W blue LEDs. The reaction was stirred at room temperature for 12~48 h under argon, and then the mixture directly purified by silica gel column chromatography (EtOAc/petroleum ether, *V* : *V* = 1 : 10 to 1 : 5) to afford the title products **3** or **4**.

1-(1-Isopropyl-1*H*-imidazol-2-yl)-4-(4-(methyl(phenyl)-amino)phenyl)-3-phenylbutan-1-one (**3a**): Pale yellow oil, 35 mg, 79% yield, 96% *ee* [HPLC: chiralpak AD-H column, 254 nm, hexane/isopropanol (*V* : *V* = 90 : 10), flow rate 1 mL/min, 40 °C, *t*_R(minor) = 8.42 min, *t*_R(major) = 9.18 min]. [α]_D²⁵ + 32.919 (*c* 0.5, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ : 7.39~7.13 (m, 9H), 7.07 (d, *J* = 8.3 Hz, 1H), 7.01~6.87 (m, 5H), 5.49~5.35 (m, 1H), 3.85~3.63 (m, 2H), 3.55~3.41 (m, 1H), 3.30 (s, 3H), 3.03~2.88 (m, 2H), 1.38 (d, *J* = 6.7 Hz, 3H), 1.33 (d, *J* = 6.7 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 191.8, 149.2, 147.0, 144.4, 142.6, 133.6, 130.2, 129.4, 129.1, 128.3, 127.8, 126.2, 121.3, 120.9, 120.4, 119.2, 49.0, 44.8, 43.3, 43.0, 40.2, 23.6, 23.5; IR (KBr) ν : 2926, 2361, 2343, 1678, 1595, 1512, 1497, 1457, 1397, 1346, 1258, 982, 760, 704 cm⁻¹; HRMS (ESI) calcd for C₂₉H₃₁N₃NaO [M + Na]⁺ 460.2365, found 460.2359.

1-(1-Isopropyl-1*H*-imidazol-2-yl)-4-(4-(methyl(phenyl)-amino)phenyl)-3-(*p*-tolyl)butan-1-one (**3b**): Pale yellow oil, 31 mg, 68% yield, 95% *ee* [HPLC: chiralpak AD-H column, 254 nm, hexane/isopropanol (*V* : *V* = 90 : 10), flow rate 1.0 mL/min, 40 °C, *t*_R(minor) = 7.58 min, *t*_R(major) = 8.49 min]. [α]_D²⁵ + 46.781 (*c* 0.5, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ : 7.27~7.19 (m, 2H), 7.19~7.09 (m, 4H), 7.19~7.09 (m, 4H), 7.06~6.99 (m, 4H), 5.42~5.30 (m, 1H), 3.72~3.54 (m, 2H), 3.40 (dd, *J* = 15.8, 4.7 Hz, 1H), 3.26 (s, 3H), 2.95~2.81 (m, 2H), 2.27 (s, 3H), 1.33 (d, *J* = 6.6 Hz, 3H), 1.30 (d, *J* = 6.7 Hz, 3H); ¹³C NMR (101 MHz, CDCl₃) δ : 191.9, 149.3, 147.0, 142.5, 141.5, 135.7, 133.8, 130.2, 129.2, 129.1, 127.7, 121.5, 121.0, 120.4, 119.1, 49.2, 44.9, 43.1, 42.8, 40.3, 23.7, 23.5, 21.2; IR (KBr) ν : 2922, 2855, 2361, 2345, 1676, 1597, 1512, 1498, 1395, 1340, 1255, 981, 916, 822, 695 cm⁻¹; HRMS (ESI) calcd for C₃₀H₃₃N₃NaO [M + Na]⁺ 474.2521, found 474.2515.

1-(1-Isopropyl-1*H*-imidazol-2-yl)-3-(4-methoxyphenyl)-4-(4-(methyl(phenyl)amino)phenyl)butan-1-one (**3c**): Pale yellow oil, 27 mg, 58% yield, 92% *ee* [HPLC: chiralpak IC column, 254 nm, hexane/isopropanol (*V* : *V* = 75 : 25), flow rate 1.0 mL/min, 40 °C, *t*_R(minor) = 6.02 min, *t*_R(major) = 6.77 min]. [α]_D²⁵ + 37.251 (*c* 0.5, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ : 7.25~7.19 (m, 1H), 7.20~7.11 (m, 4H), 7.01 (d, *J* = 8.4 Hz, 1H), 6.95~6.84 (m, 3H), 6.77 (d, *J* = 8.7 Hz, 1H), 5.43~5.31 (m, 1H), 3.75 (s, 1H), 3.71~3.54 (m, 2H), 3.39 (dd, *J* = 15.9, 5.0 Hz, 1H), 3.26 (s, 3H), 2.87 (d, *J* = 7.2 Hz, 1H), 1.34 (d, *J* = 6.6 Hz, 1H), 1.31 (d, *J* = 6.7 Hz, 1H); ¹³C NMR (101 MHz, CDCl₃) δ : 191.9, 158.0, 149.3, 147.0, 142.5, 136.5, 133.8, 130.2,

129.2, 128.8, 121.5, 121.0, 120.4, 119.2, 113.7, 55.3, 49.2, 45.1, 43.2, 42.5, 40.4, 23.7, 23.6; IR (KBr) ν : 2925, 2851, 2362, 2342, 1678, 1598, 1512, 1497, 1395, 1343, 1250, 1039, 986, 830, 697 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{30}\text{H}_{33}\text{N}_3\text{NaO}_2$ $[\text{M}+\text{Na}]^+$ 490.2470, found 490.2464.

3-(4-Fluorophenyl)-1-(1-isopropyl-1*H*-imidazol-2-yl)-4-(4-(methyl(phenyl)amino)phenyl)butan-1-one (**3d**): Pale yellow oil, 37.1 mg, 81% yield, 91% *ee* [HPLC: chiralpak AD-H column, 254 nm, hexane/isopropanol ($V:V=90:10$), flow rate 1.0 mL/min, 40 $^{\circ}\text{C}$, $t_{\text{R}}(\text{minor})=8.29$ min, $t_{\text{R}}(\text{major})=8.97$ min]. $[\alpha]_{\text{D}}^{25}+32.119$ (c 0.5, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 7.21~7.03 (m, 6H), 6.95~6.73 (m, 9H), 5.36~5.24 (m, 1H), 3.65~3.49 (m, 2H), 3.38~3.27 (m, 1H), 3.18 (s, 3H), 2.86~2.71 (m, 2H), 1.27 (d, $J=6.6$ Hz, 3H), 1.23 (d, $J=6.7$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 191.7, 161.5 (d, $J=244.6$ Hz), 149.3, 147.2, 142.6, 140.0 (d, $J=3.0$ Hz), 133.3, 130.2, 129.5, 129.3 (d, $J=7.7$ Hz), 129.2, 121.3, 121.2, 120.6, 119.4, 115.1 (d, $J=21.1$ Hz), 49.2, 45.0, 43.1, 42.7, 40.3, 23.7, 23.6; ^{19}F NMR (376 MHz, CDCl_3) δ : -117.12; IR (KBr) ν : 2923, 2367, 2340, 1676, 1595, 1512, 1498, 1456, 1396, 1348, 1257, 1225, 984, 917, 834, 698 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{29}\text{H}_{30}\text{FN}_3\text{NaO}$ $[\text{M}+\text{Na}]^+$ 478.2271, found 478.2263.

3-(4-Chlorophenyl)-1-(1-isopropyl-1*H*-imidazol-2-yl)-4-(4-(methyl(phenyl)amino)phenyl)butan-1-one (**3e**): Pale yellow oil, 33 mg, 70% yield, 96% *ee* [HPLC: chiralpak AD-H column, 254 nm, hexane/isopropanol ($V:V=92:8$), flow rate 0.6 mL/min, 40 $^{\circ}\text{C}$, $t_{\text{R}}(\text{minor})=17.35$ min, $t_{\text{R}}(\text{major})=18.17$ min]. $[\alpha]_{\text{D}}^{25}+67.172$ (c 0.5, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 7.28~7.10 (m, 8H), 6.98 (d, $J=8.4$ Hz, 2H), 6.96~6.85 (m, 5H), 5.43~5.30 (m, 1H), 3.76~3.56 (m, 2H), 3.40 (dd, $J=16.0, 4.8$ Hz, 1H), 3.26 (s, 3H), 2.98~2.78 (m, 2H), 1.35 (d, $J=6.6$ Hz, 3H), 1.31 (d, $J=6.7$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 191.4, 149.2, 147.2, 142.9, 142.4, 133.0, 131.9, 130.1, 129.4, 129.3, 129.2, 128.4, 121.2, 121.1, 120.7, 119.5, 49.2, 44.8, 42.8, 42.8, 40.3, 23.7, 23.5; IR (KBr) ν : 2934, 2363, 2336, 1676, 1596, 1518, 1503, 1346, 1259, 992, 828, 762, 697 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{29}\text{H}_{30}\text{ClN}_3\text{NaO}$ $[\text{M}+\text{Na}]^+$ 494.1975, found 494.1969.

1-(1-Isopropyl-1*H*-imidazol-2-yl)-4-(4-(methyl(phenyl)amino)phenyl)-3-(*m*-tolyl)butan-1-one (**3f**): Colorless oil, 43 mg, 95% yield, 93% *ee* [HPLC: chiralpak IA column, 254 nm, hexane/isopropanol ($V:V=94:6$), flow rate 0.4 mL/min, 40 $^{\circ}\text{C}$, $t_{\text{R}}(\text{minor})=21.32$ min, $t_{\text{R}}(\text{major})=22.55$ min]. $[\alpha]_{\text{D}}^{25}+30.054$ (c 0.5, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 7.18~7.09 (m, 3H), 7.07~7.00 (m, 2H), 6.98~6.791 (m, 4H), 6.89~6.76 (m, 6H), 5.36~5.23 (m, 1H), 3.64~3.47 (m, 2H), 3.33 (dd, $J=15.6, 4.8$ Hz, 1H), 3.18 (s, 1H), 2.87~2.73 (m, 2H), 2.20 (s, 3H), 1.26 (d, $J=6.6$ Hz, 3H), 1.22 (d, $J=6.6$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 191.9, 149.3, 147.1, 144.5, 142.7, 137.8, 133.9, 130.2, 129.4, 129.2, 128.7, 128.2, 127.1, 124.8, 121.6, 121.0, 120.4, 119.1, 49.2, 44.7, 43.2, 43.1, 40.3, 23.7, 23.5, 21.6; IR (KBr) ν : 2926, 2362, 2345, 1676, 1599, 1512, 1498, 1456, 1396, 1342, 1257, 984, 702 cm^{-1} ; HRMS

(ESI) calcd for $\text{C}_{30}\text{H}_{33}\text{N}_3\text{NaO}$ $[\text{M}+\text{Na}]^+$ 474.2521, found 474.2514.

1-(1-Isopropyl-1*H*-imidazol-2-yl)-4-(4-(methyl(phenyl)amino)phenyl)-3-(3-(trifluoromethyl)phenyl)butan-1-one (**3g**): Pale yellow oil, 40 mg, 79% yield, 95% *ee* [HPLC: chiralpak AD-H column, 254 nm, hexane/isopropanol ($V:V=75:25$), flow rate 1.0 mL/min, 40 $^{\circ}\text{C}$, $t_{\text{R}}(\text{minor})=4.25$ min, $t_{\text{R}}(\text{major})=4.62$ min]. $[\alpha]_{\text{D}}^{25}+43.915$ (c 0.5, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 7.46~7.30 (m, 4H), 7.27~7.12 (m, 4H), 7.00~6.84 (m, 7H), 5.44~5.30 (m, 1H), 3.75~3.63 (m, 2H), 3.54~3.41 (m, 1H), 3.25 (s, 3H), 2.95 (dd, $J=13.6, 6.6$ Hz, 1H), 2.86 (dd, $J=13.5, 6.7$ Hz, 1H), 1.35 (d, $J=6.7$ Hz, 3H), 1.30 (d, $J=6.7$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 191.2, 149.2, 147.3, 145.2, 142.4, 132.7, 131.2, 130.4 (q, $J=31.3$ Hz), 130.1, 129.5, 129.5, 128.7, 124.9 (q, $J=3.6$ Hz), 124.3 (q, $J=273.7$ Hz), 123.2 (q, $J=3.7$ Hz), 121.3, 121.3, 120.6, 119.4, 49.3, 44.5, 43.3, 42.8, 40.3, 23.6, 23.5; ^{19}F NMR (376 MHz, CDCl_3) δ : -62.47; IR (KBr) ν : 2929, 2361, 2344, 1676, 1595, 1512, 1498, 1455, 1408, 1397, 1330, 1256, 1176, 1127, 982, 919, 707 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{30}\text{H}_{30}\text{F}_3\text{N}_3\text{NaO}$ $[\text{M}+\text{Na}]^+$ 528.2239, found 528.2230.

3-(2-Fluorophenyl)-1-(1-isopropyl-1*H*-imidazol-2-yl)-4-(4-(methyl(phenyl)amino)phenyl)butan-1-one (**3h**): Pale yellow oil, 31 mg, 68% yield, 94% *ee* [HPLC: chiralpak IC column, 254 nm, hexane/isopropanol ($V:V=70:30$), flow rate 1.0 mL/min, 40 $^{\circ}\text{C}$, $t_{\text{R}}(\text{minor})=4.67$ min, $t_{\text{R}}(\text{major})=5.04$ min]. $[\alpha]_{\text{D}}^{25}+49.312$ (c 0.5, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 7.28~7.19 (m, 4H), 7.17~7.07 (m, 2H), 7.05 (d, $J=8.5$ Hz, 2H), 7.02~6.95 (m, 2H), 6.95~6.83 (m, 5H), 5.45~5.32 (m, 1H), 4.03~3.89 (m, 1H), 3.73 (dd, $J=17.2, 8.8$ Hz, 1H), 3.48 (dd, $J=17.2, 5.9$ Hz, 1H), 3.24 (s, 3H), 3.01~2.84 (m, 2H), 1.34 (d, $J=6.7$ Hz, 3H), 1.31 (d, $J=6.7$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 191.4, 161.0 (d, $J=246.4$ Hz), 149.3, 147.1, 142.4, 133.4, 131.2, 131.0, 130.1, 129.4, 129.1, 129.07, 127.8 (d, $J=8.4$ Hz), 124.0 (d, $J=3.3$ Hz), 121.4, 121.1, 120.4, 119.2, 115.5 (d, $J=22.8$ Hz), 49.2, 43.6, 41.4, 40.3, 36.7, 23.7, 23.6; ^{19}F NMR (376 MHz, CDCl_3) δ : -117.41; IR (KBr) ν : 2921, 2363, 1676, 1595, 1512, 1497, 1457, 1409, 1395, 1346, 1258, 982, 919, 759, 704 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{29}\text{H}_{30}\text{FN}_3\text{NaO}$ $[\text{M}+\text{Na}]^+$ 478.2271, found 478.2265.

1-(1-Isopropyl-1*H*-imidazol-2-yl)-4-(4-(methyl(phenyl)amino)phenyl)-3-(naphthalen-2-yl)butan-1-one (**3i**): Colorless oil, 34 mg, 72% yield, 90% *ee* [HPLC: chiralpak OD-H column, 254 nm, hexane/isopropanol ($V:V=95:5$), flow rate 0.8 mL/min, 40 $^{\circ}\text{C}$, $t_{\text{R}}(\text{minor})=13.79$ min, $t_{\text{R}}(\text{major})=18.65$ min]. $[\alpha]_{\text{D}}^{25}+62.240$ (c 0.5, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 8.23 (d, $J=8.4$ Hz, 1H), 7.81 (d, $J=7.9$ Hz, 1H), 7.67 (d, $J=8.1$ Hz, 1H), 7.56~7.36 (m, 4H), 7.24~7.12 (m, 4H), 7.06 (d, $J=8.1$ Hz, 2H), 6.91~6.78 (m, 5H), 5.32~5.19 (m, 1H), 4.72~4.51 (m, 1H), 3.83 (dd, $J=16.9, 8.0$ Hz, 1H), 3.65 (dd, $J=16.9, 6.5$ Hz, 1H), 3.22 (s, 3H), 3.11 (dd, $J=13.9, 6.6$ Hz, 1H), 2.98 (dd, $J=13.6, 8.2$ Hz, 1H), 1.24 (t, $J=6.0$ Hz, 6H); ^{13}C NMR (101 MHz, CDCl_3) δ : 191.8, 149.3, 147.1, 142.7,

142.0, 133.6, 133.6, 132.4, 130.2, 129.5, 129.2, 128.0, 127.8, 127.7, 126.5, 126.4, 125.8, 125.3, 121.5, 121.1, 120.4, 119.2, 49.2, 44.9, 43.5, 43.0, 40.3, 23.7, 23.5; IR (KBr) ν : 2924, 2363, 1675, 1597, 1510, 1497, 1457, 1396, 1340, 1255, 984, 918, 822, 750, 698 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{33}\text{H}_{33}\text{N}_3\text{NaO}$ $[\text{M} + \text{Na}]^+$ 510.2521, found 510.2512.

1-(1-Isopropyl-1*H*-imidazol-2-yl)-4-(4-(methyl(phenyl)-amino)phenyl)-3-(naphthalen-1-yl)butan-1-one (**3j**): Colorless oil, 25 mg, 54% yield, 95% *ee* [HPLC: chiralpak OD-H column, 254 nm, hexane/isopropanol ($V : V = 95 : 5$), flow rate 0.8 mL/min, 40 $^{\circ}\text{C}$, $t_{\text{R}}(\text{minor}) = 14.09$ min, $t_{\text{R}}(\text{major}) = 15.41$ min]. $[\alpha]_{\text{D}}^{25} + 30.454$ (c 0.5, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 7.83~7.69 (m, 3H), 7.64 (s, 1H), 7.47~7.34 (m, 3H), 7.29~7.11 (m, 4H), 7.03 (d, $J = 7.4$ Hz, 2H), 6.96~6.81 (m, 5H), 5.40~5.29 (m, 1H), 3.89~3.74 (m, 2H), 3.57~3.43 (m, 1H), 3.24 (d, $J = 2.1$ Hz, 3H), 3.00 (d, $J = 6.5$ Hz, 2H), 1.30 (d, $J = 6.6$ Hz, 3H), 1.24 (d, $J = 6.6$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 191.9, 149.3, 147.1, 142.7, 140.7, 134.0, 133.8, 131.9, 130.2, 129.5, 129.1, 128.9, 126.8, 125.9, 125.6, 125.4, 123.6, 123.4, 121.7, 121.0, 120.3, 119.0, 49.1, 44.7, 42.6, 40.3, 23.6, 23.5; IR (KBr) ν : 2928, 2365, 1676, 1595, 1510, 1497, 1457, 1395, 1342, 1258, 979, 919, 779, 696 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{33}\text{H}_{33}\text{N}_3\text{NaO}$ $[\text{M} + \text{Na}]^+$ 510.2521, found 510.2516.

1-(1-Isopropyl-1*H*-imidazol-2-yl)-4-(4-(methyl(phenyl)-amino)phenyl)-3-(thiophen-3-yl)butan-1-one (**3k**): Yellow oil, 32.2 mg, 72% yield, 93% *ee* [HPLC: chiralpak AD-H column, 254 nm, hexane/isopropanol ($V : V = 90 : 10$), flow rate 1.0 mL/min, 40 $^{\circ}\text{C}$, $t_{\text{R}}(\text{minor}) = 9.68$ min, $t_{\text{R}}(\text{major}) = 10.67$ min]. $[\alpha]_{\text{D}}^{25} + 23.790$ (c 0.5, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 7.28~7.14 (m, 4H), 7.10~7.03 (m, 3H), 6.99~6.87 (m, 3H), 6.86~6.76 (m, 2H), 5.49~5.36 (m, 1H), 4.06~3.90 (m, 1H), 3.68 (dd, $J = 17.0$, 8.7 Hz, 1H), 3.45 (dd, $J = 16.9$, 5.7 Hz, 1H), 3.27 (s, 3H), 3.00 (dd, $J = 13.6$, 7.1 Hz, 1H), 2.94 (dd, $J = 13.6$, 7.9 Hz, 1H), 1.36 (t, $J = 6.5$ Hz, 1H); ^{13}C NMR (101 MHz, CDCl_3) δ : 191.2, 149.2, 148.3, 147.3, 142.3, 133.0, 130.2, 129.3, 129.2, 126.6, 124.1, 123.1, 121.2, 121.1, 120.7, 119.5, 49.3, 46.0, 43.7, 40.4, 38.5, 23.7, 23.6; IR (KBr) ν : 2925, 2369, 2333, 1676, 1597, 1518, 1503, 1346, 1259, 979, 917, 695 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{27}\text{H}_{29}\text{N}_3\text{NaOS}$ $[\text{M} + \text{Na}]^+$ 466.1929, found 466.1924.

1-(1-Isopropyl-1*H*-imidazol-2-yl)-4-methyl-3-(4-(methyl(phenyl)amino)benzyl)pentan-1-one (**3l**): Colorless oil, 20 mg, 50% yield, 91% *ee* [HPLC: chiralpak AD-H column, 254 nm, hexane/isopropanol ($V : V = 92 : 8$), flow rate 0.6 mL/min, 40 $^{\circ}\text{C}$, $t_{\text{R}}(\text{minor}) = 9.06$ min, $t_{\text{R}}(\text{major}) = 9.81$ min]; $[\alpha]_{\text{D}}^{25} - 20.658$ (c 0.5, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 7.25~7.18 (m, 3H), 7.17~7.06 (m, 3H), 6.93 (dd, $J = 8.5$, 2.6 Hz, 4H), 6.87 (t, $J = 7.3$ Hz, 1H), 5.52~5.40 (m, 1H), 3.26 (s, 3H), 3.16 (dd, $J = 17.1$, 5.5 Hz, 1H), 3.06 (dd, $J = 17.1$, 6.8 Hz, 1H), 2.72~2.56 (m, 1H), 2.51~2.33 (m, 2H), 1.84~1.72 (m, 1H), 1.40 (dd, $J = 8.7$, 6.7 Hz, 6H), 0.97 (d, $J = 6.8$ Hz, 3H), 0.93 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 193.4, 149.3,

146.8, 142.9, 135.1, 130.2, 129.4, 129.2, 121.7, 121.0, 120.2, 119.0, 49.3, 41.9, 40.4, 40.2, 37.1, 29.7, 23.8, 23.7, 19.9, 18.7; IR (KBr) ν : 2958, 2925, 2855, 2365, 1675, 1597, 1512, 1500, 1457, 1340, 1257, 870, 704 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{30}\text{H}_{33}\text{N}_3\text{NaO}$ $[\text{M} + \text{Na}]^+$ 426.2521, found 426.2516.

4-(4-(Dimethylamino)phenyl)-1-(1-isopropyl-1*H*-imidazol-2-yl)-3-phenylbutan-1-one (**4a**): Colorless oil, 36% yield, 90% *ee* [HPLC: chiralpak AD-H column, 254 nm, hexane/isopropanol ($V : V = 90 : 10$), flow rate 1.0 mL/min, 40 $^{\circ}\text{C}$, $t_{\text{R}}(\text{minor}) = 7.92$ min, $t_{\text{R}}(\text{major}) = 9.18$ min]. $[\alpha]_{\text{D}}^{25} + 27.322$ (c 0.5, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 7.22~7.00 (m, 7H), 6.91 (d, $J = 8.6$ Hz, 2H), 6.53 (d, $J = 8.6$ Hz, 2H), 5.32~5.26 (m, 1H), 3.64~3.46 (m, 2H), 3.30 (dd, $J = 15.0$, 3.9 Hz, 1H), 2.86~2.68 (m, 2H), 2.80 (s, 6H), 1.24 (d, $J = 6.6$ Hz, 3H), 1.21 (d, $J = 6.6$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 192.1, 149.2, 144.9, 142.8, 130.1, 129.4, 128.3, 127.9, 126.2, 120.9, 112.9, 49.1, 44.8, 43.5, 42.9, 41.0, 23.7, 23.6; IR (KBr) ν : 2926, 2854, 2364, 1676, 1522, 1455, 1396, 1340, 1257, 984, 948, 917, 810, 703 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{29}\text{N}_3\text{NaO}$ $[\text{M} + \text{Na}]^+$ 398.2208, found 398.2203.

4-(4-(Benzyl(methyl)amino)phenyl)-1-(1-isopropyl-1*H*-imidazol-2-yl)-3-phenylbutan-1-one (**4b**): Colorless oil, 83% yield, 91% *ee* [HPLC: chiralpak AD-H column, 254 nm, hexane/isopropanol ($V : V = 75 : 25$), flow rate 1.0 mL/min, 40 $^{\circ}\text{C}$, $t_{\text{R}}(\text{minor}) = 5.83$ min, $t_{\text{R}}(\text{major}) = 7.11$ min]. $[\alpha]_{\text{D}}^{25} + 32.986$ (c 0.5, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 7.26~6.98 (m, 12H), 6.88 (d, $J = 8.6$ Hz, 2H), 6.54 (d, $J = 8.6$ Hz, 2H), 5.33~5.21 (m, 1H), 4.38 (s, 1H), 3.67~3.47 (m, 2H), 3.29 (dd, $J = 15.7$, 4.6 Hz, 1H), 2.85 (s, 3H), 2.81 (dd, $J = 13.7$, 6.4 Hz, 1H), 2.73 (dd, $J = 13.7$, 7.8 Hz, 1H), 1.24 (d, $J = 6.7$ Hz, 3H), 1.20 (d, $J = 6.7$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 192.1, 148.3, 144.9, 142.8, 139.3, 130.1, 129.4, 128.6, 128.3, 128.1, 127.9, 127.0, 126.9, 126.2, 120.9, 112.5, 57.0, 49.1, 44.8, 43.5, 42.8, 38.6, 23.7, 23.5; IR (KBr) ν : 2923, 2853, 2367, 2344, 1676, 1522, 1455, 1373, 1257, 1225, 984, 917, 809, 700 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{30}\text{H}_{33}\text{N}_3\text{NaO}$ $[\text{M} + \text{Na}]^+$ 474.2521, found 474.2516.

4-(4-(Dibenzylamino)phenyl)-1-(1-isopropyl-1*H*-imidazol-2-yl)-3-phenylbutan-1-one (**4c**): Colorless oil, 60% yield, 80% *ee* [HPLC: chiralpak AD-H column, 254 nm, hexane/isopropanol ($V : V = 75 : 25$), flow rate 1.0 mL/min, 40 $^{\circ}\text{C}$, $t_{\text{R}}(\text{minor}) = 7.05$ min, $t_{\text{R}}(\text{major}) = 13.24$ min]. $[\alpha]_{\text{D}}^{25} + 37.784$ (c 0.5, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 7.34~7.26 (m, 4H), 7.27~7.17 (m, 10H), 7.17~7.07 (m, 3H), 6.92 (d, $J = 8.3$ Hz, 2H), 6.60 (d, $J = 8.1$ Hz, 2H), 5.41~5.28 (m, 1H), 4.58 (s, 4H), 3.69 (dd, $J = 16.5$, 9.3 Hz, 1H), 3.63~3.51 (m, 1H), 3.33 (dd, $J = 16.5$, 5.1 Hz, 1H), 2.87 (dd, $J = 13.7$, 6.5 Hz, 1H), 2.78 (dd, $J = 13.7$, 8.3 Hz, 1H), 1.31 (d, $J = 6.7$ Hz, 3H), 1.27 (d, $J = 6.7$ Hz, 3H); ^{13}C NMR (101 MHz, CDCl_3) δ : 192.1, 147.7, 144.8, 142.7, 138.9, 130.1, 129.3, 128.7, 128.3, 128.2, 127.9, 126.9, 126.9, 126.2, 120.9, 112.5, 54.3, 49.1, 44.7, 43.3, 42.7, 23.7, 23.5; IR (KBr) ν : 2929, 2854, 2363, 1676, 1524, 1497, 1457, 1395, 1363, 1258, 959, 813, 700 cm^{-1} ;

HRMS (ESI) calcd for $C_{36}H_{37}N_3NaO$ $[M + Na]^+$ 550.2834, found 550.2829.

4-(4-(Benzylamino)phenyl)-1-(1-isopropyl-1*H*-imidazol-2-yl)-3-phenylbutan-1-one (**4d**) Pale yellow oil, 33 mg, 70% yield, 95% *ee* [HPLC: chiralpak IC column, 254 nm, hexane/isopropanol (*V* : *V* = 70 : 30), flow rate 1.0 mL/min, 40 °C, t_R (major) = 6.77 min, t_R (minor) = 7.38 min]. $[\alpha]_D^{25} + 34.052$ (*c* 0.5, $CHCl_3$); 1H NMR (400 MHz, $CDCl_3$) δ : 7.37~7.05 (m, 12H), 6.91 (d, *J* = 8.1 Hz, 2H), 6.51 (d, *J* = 8.2 Hz, 2H), 5.41~5.28 (m, 1H), 4.26 (s, 2H), 3.66 (dd, *J* = 16.0, 8.9 Hz, 1H), 3.63~3.52 (m, 1H), 3.37 (dd, *J* = 16.0, 4.9 Hz, 1H), 2.86 (dd, *J* = 13.6, 6.7 Hz, 1H), 2.80 (dd, *J* = 13.6, 7.7 Hz, 1H), 1.32 (d, *J* = 6.6 Hz, 3H), 1.28 (d, *J* = 6.6 Hz, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 192.0, 146.0, 144.7, 142.7, 139.3, 130.2, 129.5, 129.3, 128.7, 128.3, 127.9, 127.8, 127.4, 126.2, 120.9, 113.1, 49.1, 48.9, 44.8, 43.5, 43.0, 23.7, 23.5; IR (KBr) ν : 2925, 2855, 2363, 2342, 1676, 1616, 1519, 1497, 1455, 1409, 1395, 1324, 1256, 979, 917, 702 cm^{-1} ; HRMS (ESI) calcd for $C_{29}H_{31}N_3NaO$ $[M + Na]^+$ 460.2365, found 460.2359.

4-(4-Aminophenyl)-1-(1-isopropyl-1*H*-imidazol-2-yl)-3-phenylbutan-1-one (**4e**) Pale Yellow oil, 38% yield, 58% *ee* [HPLC: chiralpak IC column, 254 nm, hexane/isopropanol (*V* : *V* = 75 : 25), flow rate 1.0 mL/min, 40 °C, t_R (major) = 11.49 min, t_R (minor) = 16.95 min]. $[\alpha]_D^{25} + 20.859$ (*c* 0.5, $CHCl_3$); 1H NMR (400 MHz, $CDCl_3$) δ : 7.24~7.16 (m, 5H), 7.15~7.08 (m, 2H), 6.87 (d, *J* = 8.2 Hz, 2H), 6.53 (d, *J* = 8.3 Hz, 2H), 5.41~5.28 (m, 1H), 3.69~3.50 (m, 2H), 3.39 (dd, *J* = 15.6, 4.8 Hz, 1H), 2.89~2.76 (m, 2H), 1.33 (d, *J* = 6.6 Hz, 3H), 1.29 (d, *J* = 6.6 Hz, 3H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 192.0, 144.6, 144.3, 142.7, 130.3, 129.7, 129.4, 128.3, 127.9, 126.2, 120.9, 115.2, 49.1, 44.9, 43.5, 43.0, 23.7, 23.6; IR (KBr) ν : 2923, 2852, 2362, 2344, 1673, 1519, 1458, 1396, 1395, 981, 919, 830, 703 cm^{-1} ; HRMS (ESI) calcd for $C_{22}H_{25}N_3NaO$ $[M + Na]^+$ 370.1895, found 370.1890.

Methyl-3-(2-fluorophenyl)-4-(4-(methyl(phenyl)amino)-phenyl)butanoate (**5**) Colorless oil, 60% yield, 97% *ee* [HPLC: chiralpak IC column, 254 nm, hexane/isopropanol (*V* : *V* = 97.5 : 2.5), flow rate 0.6 mL/min, 40 °C, t_R (major) = 11.23 min, t_R (minor) = 11.88 min]. $[\alpha]_D^{25} + 5.888$ (*c* 0.5, $CHCl_3$); 1H NMR (400 MHz, $CDCl_3$) δ : 7.32~6.79 (m, 13H), 3.76~3.61 (m, 1H), 3.53 (s, 3H), 3.25 (s, 3H), 2.99~2.80 (m, 2H), 2.78~2.61 (m, 2H); ^{13}C NMR (101 MHz, $CDCl_3$) δ : 172.7, 161.0 (d, *J* = 244.4 Hz), 149.2, 147.3, 132.7, 130.3, 130.2, 130.0, 129.1, 128.1 (d, *J* = 8.4 Hz), 124.0 (d, *J* = 3.2 Hz), 121.2, 120.7, 119.5, 115.6 (d, *J* = 22.7 Hz), 51.6, 40.7, 40.3, 38.4, 38.3; ^{19}F NMR (376 MHz, $CDCl_3$) δ : -117.48; IR (KBr) ν : 2363, 2345, 1736, 1637, 1595, 1510, 1492, 1384, 1341, 1254, 1152, 755, 695 cm^{-1} ; HRMS (ESI) calcd for $C_{24}H_{24}NFNaO_2$ $[M + Na]^+$ 400.1689, found 400.1684.

4.3 General procedure for synthetic transformation

To a solution of **3h** (227 mg, 0.5 mmol) in *N,N*-dimethylformamide (DMF) (5.0 mL) was added MeI (710 mg, 5.0 mmol, 10.0 equiv.). The mixture was stirred at 80 °C

for 6 h (monitored by TLC) under argon atmosphere, followed by addition of MeOH (2.0 mL) and DBU (76 mg, 2.5 mmol, 5.0 equiv.). After stirring at room temperature for 20 h, the reaction mixture was diluted with EtOAc (25 mL) and H_2O (25 mL). The aqueous layer was separated and extracted with EtOAc (25 mL $\times$ 2). The combined organic layers were washed with brine (10 mL), dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (EtOAc/petroleum ether, *V* : *V* = 2 : 8) to provide the compound **5** as pale yellow oil (113 mg, 60% yield).

Supporting Information X-ray data for compound **3h**, 1H NMR, ^{13}C NMR spectra, and HPLC chromatograms for compounds **3**, **4** and **5**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] For selected reviews on visible-light photoredox catalysis, see: (a) Xuan, J.; Xiao, W. J. *Angew. Chem., Int. Ed.* **2012**, *51*, 6828. (b) Prier, C. K.; Rankic, D. A.; MacMillan, D. W. C. *Chem. Rev.* **2013**, *113*, 5322. (c) Douglas, J. J.; Sevrin, M. J.; Stephenson, C. R. J. *Org. Process Res. Dev.* **2016**, *20*, 1134. (d) Gentry, E. C.; Knowles, R. R. *Acc. Chem. Res.* **2016**, *49*, 1546. (e) Romero, N. A.; Nicewicz, D. A. *Chem. Rev.* **2016**, *116*, 10075. (f) Shaw, M. H.; Twilton, J.; MacMillan, D. W. C. *J. Org. Chem.* **2016**, *81*, 6898. (g) Skubi, K. L.; Blum, T. R.; Yoon, T. P. *Chem. Rev.* **2016**, *116*, 10035. (h) Parasram, M.; Gevorgyan, V. *Chem. Soc. Rev.* **2017**, *46*, 6227. (i) Larsen, C. B.; Wenger, O. S. *Chem.-Eur. J.* **2018**, *24*, 2039. (j) Marzo, L.; Pagire, S. K.; Reiser, O.; König, B. *Angew. Chem., Int. Ed.* **2018**, *57*, 10034. (k) Zhao, Y. T.; Xia, W. J. *Chem. Soc. Rev.* **2018**, *47*, 2591. (l) Chuentragool, P.; Kurandina, D.; Gevorgyan, V. *Angew. Chem., Int. Ed.* **2019**, *58*, 11586. (m) Zhou, Q. Q.; Zou, Y. Q.; Lu, L. Q.; Xiao, W. J. *Angew. Chem., Int. Ed.* **2019**, *58*, 1586. (n) Xiao, L.; Li, J. H.; Wang, T. *Acta Chim. Sinica* **2019**, *77*, 841 (in Chinese). (肖丽, 李嘉恒, 王挺, 化学学报, **2019**, *77*, 841.) (o) Chen, Y. L.; Chang, L.; Zuo, Z. W. *Acta Chim. Sinica* **2019**, *77*, 794 (in Chinese). (陈奕霖, 常亮, 左智伟, 化学学报, **2019**, *77*, 794.) (p) Chen, D.; Liu, J. C.; Zhang, X. Y.; Jiang, H. Z.; Li, J. H. *Chin. J. Org. Chem.* **2019**, *39*, 3353 (in Chinese). (陈丹, 刘剑沉, 张馨元, 蒋合众, 李加洪, 有机化学, **2019**, *39*, 3353.) (q) Zhang, H.; Yu, S. Y. *Chin. J. Org. Chem.* **2019**, *39*, 95 (in Chinese). (张昊, 俞寿云, 有机化学, **2019**, *39*, 95.) [2] For recent reviews on asymmetric photoredox catalysis, see: (a) Meggers, E. *Chem. Commun.* **2015**, *51*, 3290. (b) Wang, C. F.; Lu, Z. *Org. Chem. Front.* **2015**, *2*, 179. (c) Yoon, T. P. *Acc. Chem. Res.* **2016**, *49*, 2307. (d) Wang, D. H.; Zhang, L.; Luo, S. Z. *Acta Chim. Sinica* **2017**, *75*, 22 (in Chinese). (王德红, 张龙, 罗三中, 化学学报, **2017**, *75*, 22.) (e) Brenninger, C.; Jolliffe, J. D.; Bach, T. *Angew. Chem., Int. Ed.* **2018**, *57*, 14338.

- (f) Garrido-Castro, A. F.; Maestro, M. C.; Aleman, J. *Tetrahedron Lett.* **2018**, 59, 1286.
- (g) Silvi, M.; Melchiorre, P. *Nature* **2018**, 554, 41.
- (h) Zou, Y. Q.; Hormann, F. M.; Bach, T. *Chem. Soc. Rev.* **2018**, 47, 278.
- (i) Jiang, C.; Chen, W.; Zheng, W. H.; Lu, H. *Org. Biomol. Chem.* **2019**, 17, 867.
- [3] For examples of deracemization enabled by visible-light photocatalysis, see:
- (a) Hoelzl-Hobmeier, A.; Bauer, A.; Silva, A. V.; Huber, S. M.; Bannwarth, C.; Bach, T. *Nature* **2018**, 564, 240.
- (b) Shin, N. Y.; Ryss, J. M.; Zhang, X.; Miller, S. J.; Knowles, R. R. *Science* **2019**, 366, 364.
- (c) Troster, A.; Bauer, A.; Jandl, C.; Bach, T. *Angew. Chem., Int. Ed.* **2019**, 58, 3538.
- (d) Shi, Q.; Ye, J. *Angew. Chem., Int. Ed.* **2020**, 59, 4998.
- [4] For selected examples of dual catalysis systems, see:
- (a) Blum, T. R.; Miller, Z. D.; Bates, D. M.; Guzei, I. A.; Yoon, T. P. *Science* **2016**, 354, 1391.
- (b) Capacci, A. G.; Malinowski, J. T.; McAlpine, N. J.; Kuhne, J.; MacMillan, D. W. C. *Nat. Chem.* **2017**, 9, 1073.
- (c) Lin, L.; Bai, X.; Ye, X.; Zhao, X.; Tan, C. H.; Jiang, Z. *Angew. Chem., Int. Ed.* **2017**, 56, 13842.
- (d) Yang, Q.; Zhang, L.; Ye, C.; Luo, S. Z.; Wu, L. Z.; Tung, C. H. *Angew. Chem., Int. Ed.* **2017**, 56, 3694.
- (e) Proctor, R. S. J.; Davis, H. J.; Phipps, R. J. *Science* **2018**, 360, 419.
- (f) Ye, C.-X.; Melcamu, Y. Y.; Li, H.-H.; Cheng, J.-T.; Zhang, T.-T.; Ruan, Y.-P.; Zheng, X.; Lu, X.; Huang, P.-Q. *Nat. Commun.* **2018**, 9, 410.
- (g) Zhang, H. H.; Zhao, J. J.; Yu, S. J. *Am. Chem. Soc.* **2018**, 140, 16914.
- (h) Cheng, Y. Z.; Zhao, Q. R.; Zhang, X.; You, S. L. *Angew. Chem., Int. Ed.* **2019**, 58, 18069.
- (i) Li, Y.; Lei, M.; Gong, L. *Nat. Catal.* **2019**, 2, 1016.
- (j) Zhang, K.; Lu, L. Q.; Jia, Y.; Wang, Y.; Lu, F. D.; Pan, F.; Xiao, W. J. *Angew. Chem., Int. Ed.* **2019**, 58, 13375.
- (k) Cheng, Z. M.; Chen, P. H.; Liu, G. S. *Acta Chim. Sinica* **2019**, 77, 856 (in Chinese).
- (成忠明, 陈品红, 刘国生, 化学学报, **2019**, 77, 856.)
- [5] For selected examples of bifunctional photocatalysts, see:
- (a) Arceo, E.; Jurberg, I. D.; Alvarez-Fernandez, A.; Melchiorre, P. *Nat. Chem.* **2013**, 5, 750.
- (b) Ding, W.; Lu, L. Q.; Zhou, Q. Q.; Wei, Y.; Chen, J. R.; Xiao, W. J. *Am. Chem. Soc.* **2017**, 139, 63.
- (c) Silvi, M.; Verrier, C.; Rey, Y. P.; Buzzetti, L.; Melchiorre, P. *Nat. Chem.* **2017**, 9, 868.
- (d) Skubi, K. L.; Kidd, J. B.; Jung, H.; Guzei, I. A.; Baik, M. H.; Yoon, T. P. *J. Am. Chem. Soc.* **2017**, 139, 17186.
- (e) Li, Y.; Zhou, K.; Wen, Z.; Cao, S.; Shen, X.; Lei, M.; Gong, L. *J. Am. Chem. Soc.* **2018**, 140, 15850.
- (f) Rigotti, T.; Casado-Sanchez, A.; Cabrera, S.; Perez-Ruiz, R.; Liras, M.; O'Shea, V. A. D.; Aleman, J. *ACS Catal.* **2018**, 8, 5928.
- (g) Shen, X.; Li, Y.; Wen, Z.; Cao, S.; Hou, X.; Gong, L. *Chem. Sci.* **2018**, 9, 4562.
- (h) Stegbauer, S.; Jandl, C.; Bach, T. *Angew. Chem., Int. Ed.* **2018**, 57, 14593.
- (i) Guo, Q.; Wang, M.; Peng, Q.; Huo, Y.; Liu, Q.; Wang, R.; Xu, Z. *ACS Catal.* **2019**, 9, 4470.
- [6] (a) Giese, B. *Angew. Chem., Int. Ed.* **1983**, 22, 753.
- (b) Giese, B.; González-Gómez, J. A.; Witzel, T. *Angew. Chem., Int. Ed.* **1984**, 23, 69.
- [7] (a) Wu, J. H.; Radinov, R.; Porter, N. A. *J. Am. Chem. Soc.* **1995**, 117, 11029.
- (b) Sibi, M. P.; Ji, J.; Wu, J. H.; Gürtler, S.; Porter, N. A. *J. Am. Chem. Soc.* **1996**, 118, 9200.
- (c) Sibi, M. P.; Porter, N. A. *Acc. Chem. Res.* **1999**, 32, 163.
- [8] Espelt, L. R.; McPherson, I. S.; Wiensch, E. M.; Yoon, T. P. *J. Am. Chem. Soc.* **2015**, 137, 2452.
- [9] (a) Zuo, Z. W.; Ahneman, D. T.; Chu, L. L.; Terrett, J. A.; Doyle, A. G.; MacMillan, D. W. C. *Science* **2014**, 345, 437.
- (b) Zuo, Z. W.; Gong, H.; Li, W.; Choi, J.; Fu, G. C.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2016**, 138, 1832.
- [10] For accounts on synthetic utilization of α -aminoalkyl radicals in visible-light photoredox catalysis, see:
- (a) Cho, D. W.; Yoon, U. C.; Mariano, P. S. *Acc. Chem. Res.* **2011**, 44, 204.
- (b) Nakajima, K.; Miyake, Y.; Nishibayashi, Y. *Acc. Chem. Res.* **2016**, 49, 1946.
- [11] (a) Cermenati, L.; Mella, M.; Albin, A. *Tetrahedron* **1998**, 54, 2575.
- (b) Smitha, M. A.; Prasad, E.; Gopidas, K. R. *J. Am. Chem. Soc.* **2001**, 123, 1159.
- (c) Yoshimi, Y.; Kobayashi, K.; Kamakura, H.; Nishikawa, K.; Haga, Y.; Maeda, K.; Morita, T.; Itou, T.; Okada, Y.; Hatanaka, M. *Tetrahedron Lett.* **2010**, 51, 2332.
- (d) Miyake, Y.; Nakajima, K.; Nishibayashi, Y. *Chem. Commun.* **2013**, 49, 7854.
- (e) Lang, S. B.; O'Nele, K. M.; Tunge, J. A. *J. Am. Chem. Soc.* **2014**, 136, 13606.
- [12] (a) Bauer, E. B. *Chem. Soc. Rev.* **2012**, 41, 3153.
- (b) Cao, Z.-Y.; Brittain, W. D. G.; Fossey, J. S.; Zhou, F. *Catal. Sci. Technol.* **2015**, 5, 3441.
- (c) Zhang, L.; Meggers, E. *Chem.-Asian. J.* **2017**, 12, 2335.
- (d) Cruchter, T.; Larionov, V. A. *Coord. Chem. Rev.* **2018**, 376, 95.
- [13] Lin, S. X.; Sun, G. J.; Kang, Q. *Chem. Commun.* **2017**, 53, 7665.
- [14] CCDC 1998446 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [15] (a) Ohta, S.; Hayakawa, S.; Nishimura, K.; Okamoto, M. *Chem. Pharm. Bull.* **1987**, 35, 1058.
- (b) Miyashita, A.; Suzuki, Y.; Nagasaki, I.; Ishiguro, C.; Iwamoto, K.-I.; Higashino, T. *Chem. Pharm. Bull.* **1997**, 45, 1254.
- [16] (a) de Assis, F. F.; Huang, X.; Akiyama, M.; Pilli, R. A.; Meggers, E. *J. Org. Chem.* **2018**, 83, 10922.
- (b) Ma, J.; Lin, J.; Zhao, L.; Harms, K.; Marsch, M.; Xie, X.; Meggers, E. *Angew. Chem., Int. Ed.* **2018**, 57, 11193.
- [17] Wang, C.; Chen, L. A.; Huo, H.; Shen, X.; Harms, K.; Gong, L.; Meggers, E. *Chem. Sci.* **2015**, 6, 1094.
- [18] Huo, H.; Shen, X.; Wang, C.; Zhang, L.; Rose, P.; Chen, L. A.; Harms, K.; Marsch, M.; Hilt, G.; Meggers, E. *Nature* **2014**, 515, 100.

(Zhao, C.)