铜催化邻烯基芳基异硫氰酸酯与叠氮化钠串联双环化反应合成 5H-苯并四氮唑并噻嗪化合物

张亚辉 刘 洋 缪建康 郝文燕*

(江西师范大学化学化工学院 功能有机小分子教育部重点实验室 南昌 330022)

摘要 以低毒、廉价的铜盐作为催化剂,实现了邻烯基芳基异硫氰酸酯与叠氮化钠的串联双环化反应,简单有效地合成了一系列 5H-苯并四氮唑并噻嗪衍生物. 在该催化体系下,反应底物显示出良好的官能团兼容性和反应性,并以中高产率制备了 5H-苯并四氮唑并噻嗪类化合物. 该策略的提出为含氮、含硫小分子化合物的合成提供了一条方便的路径.
关键词 铜催化; 串联双环化; 5H-苯并四氮唑并噻嗪; 邻烯基芳基异硫氰酸酯

Copper-Catalyzed Cascade Bicyclization of *o*-Alkenylphenyl Isothiocyanates with Sodium Azide Leading to the 5H-Benzo[*d*]tetrazolo[5,1-*b*][1,3]thiazines

Zhang, Yahui Liu, Yang Miao, Jiankang Hao, Wenyan*

(Key Laboratory of Functional Small Organic Molecules, Ministry of Education, College of Chemistry & Chemical Engineering, Jiangxi Normal University, Nanchang 330022)

Abstract A simple and efficient method for the preparation of 5H-benzo[*d*]tetrazolo[5,1-*b*][1,3]thiazines has been developed. The transformation involved the copper(I)-catalyzed cascade bicyclization of *o*-alkenylphenyl isothiocyanates with sodium azide to afford corresponding products in moderate to good yields. This present strategy provides an effective way to construct small molecular *N*-, and *S*-heterocycles.

Keywords copper-catalysis; cascade bicyclization; 5H-benzo[*d*]tetrazolo[5,1-*b*][1,3]thiazine; *o*-alkenylphenyl isothiocyanate

1 Introduction

The synthesis of tetrazole has become one of the most important research topics in heterocyclic chemistry due to their important biological activities displayed in medicinal chemistry such as antituberculosis,^[1] anti-allergy,^[2] anti-inflammatory,^[3] antitumor^[4] and so on. Furthermore, as the isostere of the carboxyl acid group,^[5] tetrazole derivatives have attracted increasing attention and are often used as testing materials, explosives, metal coordination compounds, chemical catalysts and other materials.^[6] By far, the addition of azides to isothiocyanates^[7] or the [3+2] cycloaddition of an azide to a nitrile by using a Lewis acid as the catalyst have been recognized as the powerful approaches for constructing tetrazole derivatives.^[8] For instance, Hantzsch and Vagt^[8a] disclosed the first example of

Lewis acid catalyzed addition reaction of hydrazoic acid to the cyanide group to synthesis of 5-substituted tetrazole derivatives. In 2001, Sharpless and co-workers^[8b] developed this method to synthesize a variety of 5-substituted 1H-tetrazoles from sodium azide to nitriles with water as solvent. However, to our surprise, the synthesis of the polycyclic skeleton 5H-benzo[*d*]tetrazolo[5,1-*b*][1,3]thiazine is quite rare, only two published example are known.^[8c,9] Recently, a silver-catalyzed cascade bicyclization of *o*-alkynylphenyl isothiocyanates with sodium azide for the synthesis of 5H-benzo[*d*]tetrazolo[5,1-*b*][1,3]thiazines has been reported by our group (Scheme 1).^[9] However, with the urgent demand for green and sustainable chemistry, developing a new catalytic system in place of the noble catalyst like silver or palladium would be highly desired. Of the available cheap metals, copper salt is the repre-

* Corresponding author. E-mail: wenyanhao@jxnu.edu.cn

Received December 25, 2019; revised May 12, 2020; published online May 25, 2020.

Project supported by the National Natural Science Foundation of China (No. 21762023), the Natural Science Foundation of Jiangxi Province (No. 2018BAB203006) and the Research Fund of Jiangxi provincial Education Department (No. GJJ160285).

国家自然科学基金(No. 21762023)、江西省自然科学基金(No. 2018BAB203006)和江西省教育厅(No. GJJ160285)资助项目.

sentative example. It is inexpensive, low toxic and environmentally friendly. The catalytic activities of copper salts are more remarkable, even comparable to those of noble metals such as palladium catalysts in cross coupling reactions.^[10] Numerous catalytic systems have been developed for C—O,^[11] C—N,^[12] C—S^[13] and C—C^[14] bond formation. Meanwhile, the applications of copper catalysts in heterocycle synthesis have also been explored by many organic chemists. In continuation of our interest in the synthesis of biologically relevant heterocyclic compounds by using copper salts as the catalyst,^[15] we envisioned that *o*-alkenylphenyl isothiocyanate could also be applied in the reaction with sodium azide by using tandem bicyclization strategy.^[16] Herein, we wish to introduce our recent work for the generation of 5*H*-benzo[*d*]tetrazolo[5,1-*b*][1,3]-thiazines via copper catalyzed cascade bicyclization of *o*-alkenylphenyl isothiocyanate with sodium azide.

Our previous work:

This work:

Scheme 1 Comparison between the previous and present work

2 Results and discussion

The starting material *o*-alkenylphenyl isothiocyanates were easily obtained from Heck coupling of 2-iodoanilines with acrylates, acrylonitrile or *N,N*-dimethyl acrylamide,^[17] followed by the treatment with thiophosgene according to the literature procedure.^[18] In order to find an optimal reaction condition, methyl (*E*)-3-(2-isothiocyanatophenyl)-acrylate (**1a**) and sodium azide (**2**) were used as the model substrates (Table 1). Reaction of **1a** (0.2 mmol) with sodium azide **2** (2.0 equiv.) using 10 mol% CuI and NaHCO₃ (2.0 equiv.) in 2.0 mL of MeCN at 80 °C for 20 h afforded the desired product **3a** in 25% yield (Table 1, Entry 1). Other copper salts used as catalysts in the reaction were then screened (Table 1, Entries 2~7). CuCl was found to be more effective and provided the desired product **3a** in 54% yield (Table 1, Entry 3). A screen of different protonic acid effect on the reaction indicated that CH₃CO₂H was the best choice, leading to the desired product **3a** in 89% yield (Table 1, Entry 9). Lower yields were observed when other protonic acids such as HCl, TsOH, PivOH were used as the additive (Table 1, Entries 8~11). It is worth noting that only 15% yield of product was obtained in the absence of acid (Table 1, Entry 12). When different solvents such as *N,N*-dimethylformamide (DMF), dimethyl sulfoxide (DMSO), toluene, tetrahydrofuran (THF), dichloromethane

(DCM) and 1,2-dichloroethane (DCE) were examined, it was observed that no better result emerged (Table 1, Entries 13~18). The temperature also has great impact on the reaction, and the yield of **3a** was decreased to 45% when the temperature was lowered to 60 °C (Table 1, Entry 19). Raising the reaction temperature to 100 °C also resulted in a diminished yield (Table 1, Entry 20).

Table 1 Initial studies for the tandem reaction of *o*-alkenylphenyl isothiocyanate (**1a**) with sodium azide **2**^a

Entry	Catalyst	Additive	Solvent	Temp./°C	Yield ^b /%
1	CuI	NaHCO ₃	MeCN	80	25
2	CuBr	NaHCO ₃	MeCN	80	28
3	CuCl	NaHCO ₃	MeCN	80	54
4	Cu(OAc) ₂	NaHCO ₃	MeCN	80	53
5	CuO	NaHCO ₃	MeCN	80	52
6	CuCl ₂	NaHCO ₃	MeCN	80	35
7	Cu(OTf) ₂	NaHCO ₃	MeCN	80	26
8	CuCl	TsOH	MeCN	80	25
9	CuCl	CH ₃ CO ₂ H	MeCN	80	89
10	CuCl	PivOH	MeCN	80	45
11	CuCl	HCl	MeCN	80	80
12	CuCl	—	MeCN	80	23
13	CuCl	CH ₃ CO ₂ H	DMF	80	NR
14	CuCl	CH ₃ CO ₂ H	DMSO	80	NR
15	CuCl	CH ₃ CO ₂ H	Toluene	80	35
16	CuCl	CH ₃ CO ₂ H	THF	80	60
17	CuCl	CH ₃ CO ₂ H	DCM	80	27
18	CuCl	CH ₃ CO ₂ H	DCE	80	29
19	CuCl	CH ₃ CO ₂ H	MeCN	60	45
20	CuCl	CH ₃ CO ₂ H	MeCN	100	82

^a Reaction was performed with **1a** (0.2 mmol), **2** (0.4 mmol), catalyst (0.02 mmol), acid (0.4 mmol) in solvent (2 mL) for 20 h. ^b Isolated yield based on *o*-alkenylphenyl isothiocyanate (**1a**).

With the optimized conditions in hand, the reaction scope of different *o*-alkenylphenyl isothiocyanates was investigated. The results are summarized in Table 2. All reactions proceeded smoothly, leading to the desired 5*H*-benzo[*d*]tetrazolo[5,1-*b*][1,3]thiazines in moderate to good yields. As shown in Table 2, substrates **1** bearing different ester groups including COOMe, COOEt, COO^{*n*}Bu, COO^{*i*}Bu, COO^{*t*}Bu were all tolerated in the reaction to afford the corresponding products **3a**~**3e** in good

yields. Electronic properties and substitution position on the benzene ring of substrate **1** did not hamper the reaction process. With both of electron withdrawing groups, such as F, Cl, Br, CF₃, and electron-donating group CH₃ on the benzene ring, the reactions could afford the desired products in moderate yields **3f**~**3l**. To further demonstrate the substrate scope, the reactions of (*E*)-3-(2-isothiocyanatophenyl)acrylonitrile (**1m**) and (*E*)-3-(2-isothiocyanatophenyl)-*N,N*-dimethylacrylamides (**1p**) with sodium azide **2** were next investigated. As shown in Table 2, the reactions proceeded well under the optimized conditions to afford the desired products **3m** and **3p** in 80% and 84% yield, respectively. Compared with the electron-donating group (CH₃) on the benzene ring (**3o** and **3r**), the reaction of (*E*)-3-(2-isothiocyanatophenyl)acrylonitriles or (*E*)-3-(2-isothiocyanatophenyl)-*N,N*-dimethylacrylamides bearing electron-withdrawing group (Cl or F) could lead to the desired products (**3n** and **3q**) in lower yields.

The structure of **3a** was corroborated by X-ray diffraction analysis of the crystal structure of **3a**, and its ORTEP diagram is displayed in Figure 1.

Figure 1 Single-crystal X-ray diffraction structure of **3a**
The thermal ellipsoids are at a 30% probability level, the CCDC number is 1959528

Interestingly, when DMF was used as the solvent and trimethylchlorosilane as the additive, the reaction between **1a** and sodium azide **2** still could happen at room temperature and a sulfur-containing five-membered ring **4** (the

Table 2 Synthesis of 5*H*-benzo[*d*]tetrazolo[5,1-*b*][1,3]thiazines via the copper(I)-promoted tandem reaction of *o*-alkenylphenyl isothiocyanates^a

	3a 89% yield		3b 88% yield		3c 90% yield
	3d 91% yield		3e 91% yield		3f 60% yield
	3g 76% yield		3h 75% yield		3i 76% yield
	3j 50% yield		3k 76% yield		3l 74% yield
	3m 80% yield		3n 52% yield		3o 84% yield
	3p 84% yield		3q 66% yield		3r 88% yield

^a Reaction was performed with *o*-alkenylphenyl isothiocyanates **1** (0.2 mmol), sodium azide **2** (0.4 mmol), CH₃COOH (0.4 mmol), CuCl (0.02 mmol) in MeCN (2 mL) under 80 °C for 20 h.

structure of **4** was corroborated by X-ray diffraction analysis) can be obtained in 97% yield. Then intermediate **4** were treated with iodine in the presence of sodium hydrogencarbonate to give methyl 1-(1,2,3,4-thiazol-5-yl)-1*H*-indole-2-carboxylate **5** in 45% yield (Scheme 2).

Scheme 2 Synthesis of indole from *o*-alkenylphenyl isothiocyanate

A possible mechanism for this transformation was proposed in Scheme 3. Firstly, a [3+2] cycloaddition of azido anion **2** to an isothiocyanate moiety in substrate **1** would occur to produce the intermediate **A**. Then, intermediate **A** could undergo isomerization to afford intermediate **B**. Next, copper-catalyzed intramolecular Michael addition occurred to give the intermediate **C**. Finally, the protonation of intermediate **C** would happen to give the target product **3**.

Scheme 3 Proposed mechanistic pathway

3 Conclusions

In summary, a new method for the synthesis of various valuable 5*H*-benzo[*d*]tetrazolo[5,1-*b*][1,3] thiazines from easily available *o*-alkenylphenyl isothiocyanates and sodium azide in one pot has been developed. Mechanistically, stepwise [3+2] cycloaddition-Michael addition-proto-

nation sequence, is proposed. This present cascade bicyclization strategy provided an effective route for the generation of small molecular *N*-, and *S*-heterocycles.

4 Experimental section

4.1 Instruments and reagents

¹H NMR and ¹³C NMR spectra were recorded in 400 MHz apparatus and the frequencies for ¹H NMR and ¹³C NMR test are 400 MHz and 100 MHz, respectively. The chemical shifts were reported with TMS as internal standard. Melting points were tested in an X-4A instrument without correcting temperature and the HRMS were obtained under ESI model in a mass spectrometer equipped with TOF analyzer. All other chemicals and solvents used in the experiments were obtained from commercial sources and used directly without further treatment. *o*-Alkenylphenyl isothiocyanates **1** were synthesized following literature processes.^[17-18]

4.2 General procedure for the synthesis of product 3

A mixture of *o*-alkenylphenyl isothiocyanate **1** (0.20 mmol), sodium azide **2** (0.4 mmol), CuCl (10 mol%) and CH₃COOH (0.40 mmol) was added into a tube. Subsequently MeCN (2 mL) was added. Then, the sealed tube was heated at 80 °C for 20 h. After completion of reaction as indicated by thin-layer chromatography (TLC), the mixture was concentrated and directly purified by flash column chromatography (EtOAc/petroleum ether, *V*: *V* = 1:2) to give the desired product **3**.

Methyl 2-(5*H*-benzo[*d*]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetate (**3a**): Yellow solid, 46.7 mg, 89% yield. m.p. 140.1~141.8 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.00 (d, *J*=8.0 Hz, 1H), 7.47~7.52 (m, 1H), 7.40 (d, *J*=4.0 Hz, 2H), 4.71 (d, *J*=8.4 Hz, 1H), 3.60 (s, 3H), 2.61~2.75 (dq, *J*=8.8, 16.8 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 169.5, 148.9, 130.9, 130.2, 129.4, 128.0, 124.9, 119.2, 52.3, 42.2, 39.8; HRMS calcd for C₁₁H₁₁N₄O₂S (M+H⁺) 263.0597, found 263.0600.

Ethyl 2-(5*H*-benzo[*d*]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetate (**3b**): Yellow solid, 48.6 mg, 88% yield. m.p. 156.6~158.4 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.07 (d, *J*=8.0 Hz, 1H), 7.56~7.60 (m, 1H), 7.49 (d, *J*=3.6 Hz, 2H), 4.81 (dd, *J*=6.6, 8.4 Hz, 1H), 4.15 (dq, *J*=2.4, 7.2 Hz, 2H), 2.75 (dq, *J*=8.8, 16.8 Hz, 2H), 1.22 (t, *J*=6.8 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 169.0, 149.0, 130.9, 130.2, 129.4, 128.1, 125.0, 119.2, 61.5, 42.5, 42.2, 14.1; HRMS calcd for C₁₂H₁₃N₄O₂S (M+H⁺) 277.0754, found 277.0761.

Butyl 2-(5*H*-benzo[*d*]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetate (**3c**): Yellow solid, 54.8 mg, 90% yield. m.p. 98.4~99.8 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.08 (d, *J*=8.0 Hz, 1H), 7.56~7.60 (m, 1H), 7.49 (d, *J*=4.0 Hz, 2H), 4.80 (dd, *J*=6.8, 8.8 Hz, 1H), 4.09 (t, *J*=6.8 Hz, 2H), 2.75 (dq, *J*=8.8, 16.8 Hz, 2H), 1.53~1.60 (m, 2H), 1.26~1.36 (m, 2H), 0.90 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 169.1, 149.0, 130.9, 130.2, 129.4, 128.1,

125.0, 119.2, 65.3, 42.5, 39.8, 30.4, 19.0, 13.6; HRMS calcd for $C_{14}H_{17}N_4O_2S$ ($M + H^+$) 305.1067, found 305.1068.

Isobutyl 2-(5*H*-benzo[d]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetate (**3d**): Yellow solid, 55.4 mg, 91% yield. m.p. 102.6~103.8 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.99 (d, $J=8.0$ Hz, 1H), 7.45~7.51 (m, 1H), 7.39 (d, $J=4.0$ Hz, 2H), 4.72 (dd, $J=6.8, 8.8$ Hz, 1H), 3.79 (dq, $J=2.4, 6.8$ Hz, 2H), 2.69 (dq, $J=8.8, 16.4$ Hz, 2H), 1.74~1.84 (m, 1H), 0.79 (dd, $J=0.8, 6.8$ Hz, 6H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 169.1, 148.9, 130.9, 130.2, 129.4, 128.1, 125.0, 119.2, 71.5, 42.4, 39.8, 27.5, 19.0; HRMS calcd for $C_{14}H_{17}N_4O_2S$ ($M + H^+$) 305.1067, found 305.1061.

tert-Butyl 2-(5*H*-benzo[d]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetate (**3e**): Yellow solid, 55.4 mg, 91% yield. m.p. 110.2~111.8 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.07 (d, $J=8.0$ Hz, 1H), 7.54~7.60 (m, 1H), 7.48 (d, $J=4.0$ Hz, 2H), 4.75 (dd, $J=6.8, 8.8$ Hz, 1H), 2.66 (dq, $J=8.4, 16.4$ Hz, 2H), 1.42 (s, 9H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 168.2, 149.1, 130.9, 130.1, 129.3, 128.1, 125.1, 119.1, 82.4, 43.5, 40.1, 28.0; HRMS calcd for $C_{14}H_{17}N_4O_2S$ ($M + H^+$) 305.1067, found 305.1077.

Methyl 2-(7-fluoro-5*H*-benzo[d]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetate (**3f**): Yellow solid, 33.6 mg, 60% yield. m.p. 100.6~100.8 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.04 (d, $J=8.4$ Hz, 1H), 7.56 (dd, $J=2.0, 8.8$ Hz, 1H), 7.48 (d, $J=2.0$ Hz, 1H), 4.73 (dd, $J=6.8, 8.4$ Hz, 1H), 3.71 (s, 3H), 2.76 (dq, $J=8.4, 16.8$ Hz, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 169.3, 162.1 (d, $^1J_{CF}=250$ Hz), 148.5, 139.1, 127.2, 121.4 (d, $^3J_{CF}=9$ Hz), 117.4 (d, $^2J_{CF}=23$ Hz), 115.3 (d, $^2J_{CF}=24$ Hz), 52.4, 41.9, 39.4; HRMS calcd for $C_{11}H_{10}FN_4O_2S$ ($M + H^+$) 281.0503, found 281.0517.

Methyl 2-(7-chloro-5*H*-benzo[d]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetate (**3g**): Yellow solid, 45.1 mg, 76% yield. m.p. 111.3~112.5 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.03 (q, $J=8.4$ Hz, 1H), 7.55 (dd, $J=2.0, 8.6$ Hz, 1H), 7.49 (d, $J=2.0$ Hz, 1H), 4.72~4.76 (m, 1H), 3.71 (s, 3H), 2.76 (dq, $J=8.0, 16.8$ Hz, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 169.3, 148.7, 135.1, 130.4, 129.5, 128.1, 126.4, 120.6, 52.4, 42.0, 39.4; HRMS calcd for $C_{11}H_{10}ClN_4O_2S$ ($M + H^+$) 297.0208, found 297.0207.

Methyl 2-(8-chloro-5*H*-benzo[d]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetate (**3h**): Yellow solid, 44.5 mg, 75% yield. m.p. 89.4~89.6 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.09 (s, 1H), 7.46 (d, $J=8.0$ Hz, 2H), 4.80 (dd, $J=6.4, 8.0$ Hz, 1H), 3.69 (s, 3H), 2.76 (dq, $J=8.4, 16.8$ Hz, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 169.3, 149.1, 136.2, 131.6, 129.5, 129.4, 123.2, 119.5, 52.4, 42.2, 39.4; HRMS calcd for $C_{11}H_{10}ClN_4O_2S$ ($M + H^+$) 297.0208, found 297.0202.

Methyl 2-(7-bromo-5*H*-benzo[d]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetate (**3i**): Yellow solid, 51.9 mg, 76% yield. m.p. 121.0~121.2 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.97 (d, $J=8.4$ Hz, 1H), 7.71 (d, $J=8.8$ Hz, 1H), 7.65 (s, 1H), 4.74 (t, $J=8.0$ Hz, 1H), 3.71 (s, 3H), 2.76 (dq, $J=8.8, 16.8$ Hz, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 169.3, 148.8, 133.4, 131.0, 129.9, 126.6, 122.9, 120.8, 52.5, 42.1,

39.3; HRMS calcd for $C_{11}H_{10}BrN_4O_2S$ ($M + H^+$) 340.9702, found 340.9691.

Methyl 2-(7-(trifluoromethyl)-5*H*-benzo[d]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetate (**3j**): Yellow solid, 33.0 mg, 50% yield. m.p. 148.6~149.8 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.24 (d, $J=8.4$ Hz, 1H), 7.86 (d, $J=7.6$ Hz, 1H), 7.78 (s, 1H), 4.86 (t, $J=8.0$ Hz, 1H), 3.70 (s, 3H), 2.80 (dq, $J=8.4, 16.8$ Hz, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 169.1, 149.4, 133.3, 131.5 (q, $J_{CF3}=33$ Hz), 127.4 (d, $^4J_{CF}=4$ Hz), 125.6 (d, $^4J_{CF}=3$ Hz), 125.5, 124.4, 119.9, 52.5, 42.1, 39.5; HRMS calcd for $C_{12}H_{10}F_3N_4O_2S$ ($M + H^+$) 331.0471, found 331.0480.

Methyl 2-(7-methyl-5*H*-benzo[d]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetate (**3k**): Yellow solid, 42.0 mg, 76% yield. m.p. 105.6~105.8 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.95 (d, $J=8.4$ Hz, 1H), 7.36 (d, $J=8.0$ Hz, 1H), 7.26 (s, 1H), 4.72 (dd, $J=6.4, 8.8$ Hz, 1H), 3.69 (s, 3H), 2.73 (dq, $J=8.8, 16.8$ Hz, 2H), 2.44 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 169.6, 148.5, 139.9, 130.8, 128.7, 128.3, 124.8, 119.1, 52.3, 42.3, 39.9, 21.2; HRMS calcd for $C_{12}H_{13}N_4O_2S$ ($M + H^+$) 277.0754, found 277.0751.

Methyl 2-(8-methyl-5*H*-benzo[d]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetate (**3l**): Yellow solid, 40.9 mg, 74% yield. m.p. 193.6~193.8 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.88 (s, 1H), 7.36 (d, $J=8.0$ Hz, 1H), 7.28 (d, $J=8.0$ Hz, 1H), 4.76 (t, $J=8.0$ Hz, 1H), 3.68 (s, 3H), 2.68~3.82 (m, 2H), 2.48 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 169.5, 148.9, 140.8, 130.7, 130.1, 127.8, 122.0, 119.5, 52.2, 42.4, 39.6, 21.3; HRMS calcd for $C_{12}H_{13}N_4O_2S$ ($M + H^+$) 277.0754, found 277.0757.

2-(5*H*-Benzo[d]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetonitrile (**3m**): Yellow solid, 36.7 mg, 80% yield. m.p. 178.6~179.8 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.00 (d, $J=8.0$ Hz, 1H), 7.62 (t, $J=7.6$ Hz, 2H), 7.54 (t, $J=7.6$ Hz, 1H), 5.17 (t, $J=6.4$ Hz, 1H), 3.16 (dq, $J=7.2, 17.2$ Hz, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 148.5, 131.1, 130.8, 129.9, 129.6, 124.3, 119.0, 117.7, 39.1, 27.1; HRMS calcd for $C_{10}H_8N_5S$ ($M + H^+$) 230.0495, found 230.0505.

2-(7-Chloro-5*H*-benzo[d]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetonitrile (**3n**): Yellow solid, 27.4 mg, 52% yield. m.p. 188.6~189.8 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.10 (d, $J=8.8$ Hz, 1H), 7.64 (dd, $J=2.0, 8.8$ Hz, 1H), 7.54 (d, $J=2.0$ Hz, 1H), 4.56 (t, $J=7.6$ Hz, 1H), 2.85 (dq, $J=6.4, 17.2$ Hz, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 148.9, 135.6, 131.4, 128.4, 124.2, 121.0, 119.5, 114.9, 39.5, 27.3; HRMS calcd for $C_{10}H_7ClN_5S$ ($M + H^+$) 264.0105, found 264.0114.

2-(7-Methyl-5*H*-benzo[d]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetonitrile (**3o**): Yellow solid, 40.9 mg, 84% yield. m.p. 168.0~168.8 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.95 (d, $J=8.8$ Hz, 1H), 7.50 (d, $J=2.8$ Hz, 2H), 5.17 (d, $J=6.8$ Hz, 1H), 3.21 (dq, $J=6.8, 17.2$ Hz, 2H), 2.41 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 148.1, 139.8, 131.6, 129.7, 128.6, 124.2, 118.9, 117.7, 27.2, 21.2; HRMS calcd for $C_{11}H_{10}N_5S$ ($M + H^+$) 244.0651, found 244.0656.

2-(5*H*-Benzo[d]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)-*N,N*-

dimethylacetamide (**3p**): Yellow solid, 46.3 mg, 84% yield. m.p. 140.6~140.8 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.98 (d, $J=8.0$ Hz, 1H), 7.38~7.50 (m, 3H), 4.91 (q, $J=4.8$ Hz, 1H), 2.87 (s, 3H), 2.80 (dd, $J=9.2$, 16.4 Hz, 1H), 2.73 (s, 3H), 2.59 (dd, $J=4.8$, 16.4 Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 167.9, 149.6, 131.0, 129.9, 129.4, 128.4, 125.8, 119.0, 41.0, 40.2, 36.9, 35.6; HRMS calcd for $\text{C}_{12}\text{H}_{14}\text{N}_5\text{OS}$ ($\text{M} + \text{H}^+$) 276.0914, found 276.0918.

2-(7-Fluoro-5*H*-benzo[*d*]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)-*N,N*-dimethylacetamide (**3q**): Yellow solid, 38.7 mg, 66% yield. m.p. 182.3~182.5 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.98 (q, $J=4.8$ Hz, 1H), 7.17~7.24 (m, 2H), 4.90 (dd, $J=5.6$, 8.4 Hz, 1H), 2.88 (s, 3H), 2.81 (dd, $J=8.8$, 16.8 Hz, 1H), 2.76 (s, 3H), 2.62 (dd, $J=5.2$, 16.4 Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 167.7, 162.1 (d, $^1J_{\text{CF}}=250$ Hz), 149.2, 128.2 (d, $^3J_{\text{CF}}=8$ Hz), 127.4 (d, $^4J_{\text{CF}}=3$ Hz), 121.1 (d, $^3J_{\text{CF}}=9$ Hz), 117.1 (d, $^2J_{\text{CF}}=23$ Hz), 115.6 (d, $^2J_{\text{CF}}=24$ Hz), 41.2, 39.8, 36.9, 35.7; HRMS calcd for $\text{C}_{12}\text{H}_{13}\text{FN}_5\text{OS}$ ($\text{M} + \text{H}^+$) 294.0819, found 294.0814.

N,N-Dimethyl-2-(7-methyl-5*H*-benzo[*d*]tetrazolo[5,1-*b*][1,3]thiazin-5-yl)acetamide (**3r**): Yellow solid, 50.9 mg, 88% yield. m.p. 174.7~174.9 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.87 (q, $J=8.4$ Hz, 1H), 7.45 (s, 1H), 7.40 (d, $J=8.4$ Hz, 1H), 5.00 (dd, $J=4.4$, 10.0 Hz, 1H), 3.03 (dd, $J=10.0$, 16.8 Hz, 1H), 2.81 (s, 3H), 2.77 (s, 3H), 2.75 (dd, $J=4.8$, 16.8 Hz, 1H), 2.37 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 168.6, 149.4, 139.6, 130.7, 129.6, 128.8, 126.0, 118.5, 41.2, 36.7, 35.4, 21.2; HRMS calcd for $\text{C}_{13}\text{H}_{16}\text{N}_5\text{OS}$ ($\text{M} + \text{H}^+$) 290.1070, found 290.1084.

4.3 General procedure for the synthesis of product 4

A mixture of *o*-alkenylphenyl isothiocyanate **1** (0.20 mmol), sodium azide **2** (0.4 mmol), CuCl (10 mol%), $(\text{CH}_3)_3\text{SiCl}$ (3.0 equiv.) was added into a tube. Subsequently DMF (2 mL) was added. Then, the reaction was carried out at room temperature for 16 h. After completion of reaction as indicated by TLC, the mixture was concentrated and directly purified by flash column chromatography (EtOAc /petroleum ether, $V:V=1:2$) to give the desired product methyl (*E*)-3-2-((1,2,3,4-thiatriazol-5-yl)-amino)phenyl)acrylate (**4**): Yellow solid, 50.9 mg, 97% yield. m.p. 155.0~157.0 °C; ^1H NMR (400 MHz, CDCl_3) δ : 10.47 (s, 1H), 7.90 (d, $J=15.6$ Hz, 1H), 7.43~7.61 (m, 3H), 7.28 (s, 1H), 6.81 (s, 1H), 3.87 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 178.5, 167.0, 139.4, 138.7, 131.8, 128.9, 128.1, 127.6, 122.4, 121.3, 51.9; HRMS calcd for $\text{C}_{13}\text{H}_{16}\text{N}_5\text{OS}$ ($\text{M} + \text{H}^+$) 263.0597, found 263.0610.

4.4 General procedure for the synthesis of product 5

A mixture of product **4** (0.20 mmol), I_2 (3.0 equiv.) and NaHCO_3 (2.0 equiv.) was added into a tube. Subsequently DMF (2 mL) was added. Then, the reaction was cooled at 0 °C for 6 h. After completion of reaction as indicated by TLC, the mixture was concentrated and directly purified by flash column chromatography (EtOAc /petroleum ether, $V:V=1:2$) to give the desired product methyl 1-(1,2,3,

4-thiatriazol-5-yl)-1*H*-indole-2-carboxylate (**5**): Yellow solid, 23.4 mg, 45% yield. m.p. 177.6~178.8 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.37 (d, $J=8.0$ Hz, 1H), 7.91 (d, $J=8.0$ Hz, 1H), 7.75 (t, $J=7.6$ Hz, 1H), 7.55 (t, $J=7.6$ Hz, 1H), 6.81 (s, 1H), 3.87 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 166.1, 148.1, 141.1, 133.4, 130.9, 129.3, 126.1, 119.5, 118.9, 113.0, 52.2; HRMS calcd for $\text{C}_{13}\text{H}_{16}\text{N}_5\text{OS}$ ($\text{M} + \text{H}^+$) 261.0441, found 261.0442.

Supporting Information ^1H NMR and ^{13}C NMR spectra for compounds **3a**~**3r**, **4**, **5**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Ren, Q. C.; Zhang, S.; Gao, C.; Xu, Z.; Ding, J. W.; Huang, L.; Feng, L. S. *World Notes Antibiot.* **2017**, *38*, 238.
(b) Xie, M. S.; Cheng, X.; Chen, Y. G.; Wu, X. X.; Qua, G. R.; Guo, H. M. *Org. Biomol. Chem.* **2018**, *16*, 6890.
- [2] (a) Peet, N. P.; Baugh, L. E.; Sunder, S.; Lewis, J. E.; Matthews, E. H.; Olberding, E. L.; Shah, D. N. *J. Med. Chem.* **1986**, *29*, 2403.
(b) Li, G. Q.; Li, Z.; Lu, Y. H. *Chin. J. Med. Chem.* **1996**, *6*, 50.
- [3] Poonian, M. S.; Nowoswiat, E. F.; Blount, J. F. *J. Med. Chem.* **1976**, *19*, 1017.
- [4] Navidpour, L.; Shadnia, H.; Shafaroodi, H. *Bioorg. Med. Chem.* **2007**, *15*, 1976.
- [5] Singh, V. H.; Chawla, A. S.; Kapoor, V. K.; Paul, D.; Malhotra, R. K. *Prog. Med. Chem.* **1980**, *17*, 151.
- [6] (a) Seco, J. M.; de Araújo Farias, M.; Bachs, N. M.; Caballero, A. B.; Salinas-Castillo, A.; Rodríguez Diéguez, A. *Inorg. Chim. Acta* **2010**, *363*, 3194.
(b) Himo, F.; Demko, Z. P.; Noodleman, L.; Sharpless, K. B. *J. Am. Chem. Soc.* **2003**, *125*, 9983.
(c) Fischer, N.; Karaghiosoff, K.; Klapotke, T. M.; Stierstorfer, J. Z. *Anorg. Allg. Chem.* **2010**, *636*, 735.
(d) Klapotke, T. M.; Sabaté, C. M.; Welch, J. M. *Eur. J. Inorg. Chem.* **2009**, *6*, 769.
(e) List, B.; Lerner, R. A.; Barbas, C. F. *J. Am. Chem. Soc.* **2000**, *122*, 2395.
(f) Odedra, A.; Seeberger, P. H. *Angew. Chem., Int. Ed.* **2009**, *48*, 2699.
- [7] (a) Yella, R.; Khatun, N.; Rout, S. K.; Patel, B. K. *Org. Biomol. Chem.* **2011**, *9*, 3235.
(b) Seelam, M.; Kammela, P. R.; Shaikh, B.; Tamminana, R.; Bogiri, S. *Chem. Heterocycl. Compd.* **2018**, *54*, 535.
(c) Mandapati, U.; Mandapati, P.; Pinapati, S.; Tamminana, R.; Rudraraju, R. *Synth. Commun.* **2018**, *48*, 500.
(d) L'abbé, G.; Verhelst, G.; Toppet, S. *J. Org. Chem.* **1977**, *42*, 1159.
(e) Han, S. Y.; Lee, J. W.; Kim, H.; Kim, Y.; Lee, S. W.; Gyoung, Y. S. *Bull. Korean Chem. Soc.* **2012**, *33*, 55.
(f) Majji, G.; Sahoo, S. K.; Khatun, N.; Patel, B. K. *Eur. J. Org. Chem.* **2015**, 7534.
(g) Guin, S.; Rout, S. K.; Gogoi, A.; Nandi, S.; Ghara, K. K.; Patel, B. K. *Adv. Synth. Catal.* **2012**, *354*, 2757.
(h) Batey, R. A.; Powell, D. A. *Org. Lett.* **2000**, *2*, 3237.
(i) Sathishkumar, M.; Shanmugavelan, P.; Nagarajan, S.; Dinesh, M.; Ponnuswamy, A. *New J. Chem.* **2013**, *37*, 488.
- [8] (a) Hantzsch, A.; Vagt, A. *Annalen* **1901**, *314*, 339.
(b) Demko, Z. P.; Sharpless, K. B. *J. Org. Chem.* **2001**, *66*, 7945.
(c) Demko, Z. P.; Sharpless, K. B. *Org. Lett.* **2001**, *3*, 4091.
(d) Demko, Z. P.; Sharpless, K. B. *Org. Lett.* **2002**, *4*, 2525.
(e) Batey, R. A.; Powell, D. A. *Org. Lett.* **2000**, *2*, 3237.
(f) Kadaba, P. K. *J. Org. Chem.* **1976**, *41*, 1073.

- (g) Gyoung, Y. S.; Shim, J. G.; Yamamoto, Y. *Tetrahedron Lett.* **2000**, *41*, 4193.
- (h) Mancheno, O. G.; Bolm, C. *Org. Lett.* **2007**, *9*, 2951.
- (i) Gutmann, B.; Roduit, J. P.; Roberge, D.; Kappe, C. O. *Angew. Chem., Int. Ed.* **2010**, *49*, 7101.
- (j) Razzaq, T.; Kappe, C. O. *Chem.-Asian J.* **2010**, *5*, 1274.
- (k) Bonnamour, J.; Bolm, C. *Chem.-Eur. J.* **2009**, *15*, 4543.
- (l) Lang, L. M.; Li, B. J.; Liu, W.; Jiang, L.; Xu, Z.; Yin, G. *Chem. Commun.* **2010**, *46*, 448.
- (m) Kantam, M. L.; Shiva Kumar, K. B.; Sridhar, C. *Adv. Synth. Catal.* **2005**, *347*, 1212.
- (n) Alterman, M.; Hallberg, A. *J. Org. Chem.* **2000**, *65*, 7984.
- (o) Finnegan, W. G.; Henry, R. A.; Lofquist, R. *J. Am. Chem. Soc.* **1958**, *80*, 3908.
- [9] Miao, J. K.; Zhang, Y. H.; Sang, X. Y.; Hao, W. Y. *Org. Biomol. Chem.* **2019**, *17*, 2336.
- [10] (a) Ley, S. V.; Thomas, A. W. *Angew. Chem., Int. Ed.* **2003**, *42*, 5400.
- (b) Monnier, F.; Taillefer, M. *Angew. Chem., Int. Ed.* **2009**, *48*, 6954.
- (c) Ma, D.; Cai, Q. *Acc. Chem. Res.* **2008**, *41*, 1450.
- (d) Chemler, S. R.; Fuller, P. H. *Chem. Soc. Rev.* **2007**, *36*, 1153.
- (e) Moses, J. E.; Moorhouse, A. D. *Chem. Soc. Rev.* **2007**, *36*, 1249.
- (f) Jerphagnon, T.; Pizzuti, M. G.; Minnaard, A. J.; Feringa, B. L. *Chem. Soc. Rev.* **2009**, *38*, 1039.
- (g) Surry, D. S.; Buchwald, S. L. *Chem. Sci.* **2010**, *1*, 13.
- (h) Hassan, J.; Sévignon, M.; Gozzi, C.; Schulz, E.; Lemaire, M. *Chem. Rev.* **2002**, *102*, 1359.
- (i) Yuan, C. C.; Tu, G. L.; Zhao, Y. S. *Org. Lett.* **2017**, *19*, 356.
- (c) Tu, G. L.; Yuan, C. C.; Li, Y. T.; Zhang, J. Y.; Zhao, Y. S. *Angew. Chem., Int. Ed.* **2018**, *57*, 15597.
- [11] (a) Rao, R. K.; Naidu, A. B.; Sekar, G. *Org. Lett.* **2009**, *11*, 1923.
- (b) Bhadra, S.; Adak, L.; Samanta, S.; Islam, A. K. M. M.; Mukherjee, M.; Ranu, B. C. *J. Org. Chem.* **2010**, *75*, 8533.
- (c) Melkonyan, F.; Topolyan, A.; Karchava, A.; Yurovskaya, V. *Tetrahedron* **2011**, *67*, 6826.
- (d) Jangili, P.; Kashanna, J.; Das, B. *Tetrahedron Lett.* **2013**, *54*, 3453.
- (e) Liu, Z.; Chen, Y. *Tetrahedron Lett.* **2009**, *50*, 3790.
- (f) Feng, E.; Huang, H.; Zhou, Y.; Ye, D.; Jiang, H.; Liu, H. *J. Org. Chem.* **2009**, *74*, 2846.
- (g) Chen, D.; Shen, G.; Bao, W. *Org. Biomol. Chem.* **2009**, *7*, 4067.
- (h) Sequeira, F. C.; Chemler, S. R. *Org. Lett.* **2012**, *14*, 4482.
- (i) Deb, M. L.; Dey, S. S.; Bento, I.; Barros, M. T.; Maycock, C. D. *Angew. Chem., Int. Ed.* **2013**, *52*, 9791.
- (j) Xiong, T.; Li, Y.; Bi, X.; Lv, Y.; Zhang, V. *Angew. Chem., Int. Ed.* **2011**, *50*, 7140.
- (k) Li, Y.; Li, Z.; Xiong, T.; Zhang, Q.; Zhang, X. *Org. Lett.* **2012**, *14*, 3522.
- [12] (a) Basak, A.; Ghosh, S. C.; Bhowmick, T.; Das, A. K.; Bertolasi, V. *Tetrahedron Lett.* **2002**, *43*, 5499.
- (b) Khangarot, R. K.; Kaliappan, K. P. *Eur. J. Org. Chem.* **2011**, 6117.
- (c) Grzeszczyk, B.; Polawska, K.; Shaker, Y. M.; Stecko, S.; Mames, A.; Woznica, M.; Chmielewski, M.; Furman, B. *Tetrahedron* **2012**, *68*, 10633.
- (d) Saito, T.; Kikuchi, T.; Tanabe, H.; Yahiro, J.; Otani, T. *Tetrahedron Lett.* **2009**, *50*, 4969.
- (e) Mames, A.; Stecko, S.; Mikolajczyk, P.; Soluch, M.; Furman, B.; Chmielewski, M. *J. Org. Chem.* **2010**, *75*, 7580.
- (f) Ye, M. C.; Zhou, J.; Tang, Y. *J. Org. Chem.* **2006**, *71*, 3576.
- (g) Stecko, S.; Mames, A.; Furman, B.; Chmielewski, M. *J. Org. Chem.* **2009**, *74*, 3094.
- (h) Zhang, X.; Hsung, R. P.; Li, H.; Zhang, Y.; Johnson, W. L.; Figueroa, R. *Org. Lett.* **2008**, *10*, 3477.
- (i) Lo, M. M. C.; Fu, G. C. *J. Am. Chem. Soc.* **2002**, *124*, 4572.
- (j) Shintani, R.; Fu, G. C. *Angew. Chem., Int. Ed.* **2003**, *42*, 4082.
- (k) Ye, M. C.; Zhou, J.; Huang, Z. Z.; Tang, Y. *Chem. Commun.* **2003**, 2554.
- (l) Zhao, L.; Li, C. *J. Chem.-Asian J.* **2006**, *1*, 203.
- (m) Liang, J.; Chen, J.; Du, F.; Zeng, X.; Li, L.; Zhang, H. *Org. Lett.* **2009**, *11*, 2820.
- [13] (a) Ma, D.; Geng, Q.; Zhang, H.; Jiang, Y. *Angew. Chem., Int. Ed.* **2010**, *49*, 1291.
- (b) Dai, C.; Sun, X.; Tu, X.; Wu, L.; Zhan, D.; Zeng, Q. *Chem. Commun.* **2012**, *48*, 5367.
- (c) Yang, D.; Liu, H.; Yang, H.; Fu, H.; Hu, L.; Jiang, Y.; Zhao, Y. *Adv. Synth. Catal.* **2009**, *351*, 1999.
- (d) Chen, D.; Wu, J.; Yang, J.; Huang, L.; Xiang, Y.; Bao, W. *Tetrahedron Lett.* **2012**, *53*, 7104.
- (e) Huang, W. S.; Xu, R.; Dodd, R.; Shakespeare, W. C. *Tetrahedron Lett.* **2013**, *54*, 5214.
- (f) Chen, D.; Wang, Z. J.; Bao, W. *J. Org. Chem.* **2010**, *75*, 5768.
- (g) Kaneko, K.; Yoshino, T.; Matsunaga, S.; Kanai, M. *Org. Lett.* **2013**, *15*, 2502.
- (h) Shen, C.; Zhang, P. F.; Sun, Q.; Bai, S. Q.; Andy Hor, T. S.; Liu, X. G. *Chem. Soc. Rev.* **2015**, *44*, 291.
- (i) Kaiser, D.; Klodr, I.; Oost, R.; Neuhaus, J.; Maulide, N. *Chem. Rev.* **2019**, *119*, 8701.
- (j) Sangeetha, S.; Muthupandi, P.; Sekar, G. *Org. Lett.* **2015**, *17*, 6006.
- (k) Soria-Castro, S.; Andrada, D. M.; Caminos, D. A.; Argüello, J. E.; Robert, M.; Peñéñory, A. B. *J. Org. Chem.* **2017**, *82*, 11464.
- [14] (a) Duran, F.; Leman, L.; Ghini, A.; Burton, G.; Dauban, P.; Dodd, R. H. *Org. Lett.* **2002**, *4*, 2481.
- (b) Allen, S. E.; Walvoord, R. R.; Padillasalinas, R.; Kozlowski, M. C. *Chem. Rev.* **2013**, *113*, 6234.
- [15] (a) Hao, W. Y.; Zeng, J. B.; Cai, M. Z. *Chem. Commun.* **2014**, *50*, 11686.
- (b) Hao, W. Y.; J. Huang, J.; Jie, S. S.; Cai, M. Z. *Eur. J. Org. Chem.* **2015**, 6655.
- (c) Hao, W. Y.; Sang, X. Y.; J. Jiang, J.; Cai, M. Z. *Tetrahedron Lett.* **2016**, *57*, 1511.
- (d) Hao, W. Y.; Sang, X. Y.; Jiang, J.; Cai, M. Z. *Tetrahedron Lett.* **2016**, *57*, 4207.
- (e) Hao, W. Y.; Jiang, Y. Y.; Cai, M. Z. *J. Org. Chem.* **2014**, *79*, 3634.
- [16] (a) Miao, J. K.; Sang, X. Y.; Wang, Y.; Deng, S. F.; Hao, W. Y. *Org. Biomol. Chem.* **2019**, *17*, 6994.
- (b) Hao, W. Y.; Zhang, T. L.; Cai, M. Z. *Tetrahedron* **2013**, *69*, 9219.
- (c) Zou, F. H.; Chen, X. W.; Hao, W. Y. *Tetrahedron* **2017**, *73*, 758.
- (d) Hao, W. Y.; Tian, J.; Li, W.; Shi, R. Y.; Huang, Z. L.; Lei, A. W. *Chem.-Asian J.* **2016**, *11*, 1664.
- (e) Hao, W. Y.; Sha, Y. C.; Deng, Y.; Luo, Y.; Zeng, L.; Tang, S.; Weng, Y.; Chiang, C.-W.; Lei, A. W. *Chem.-Eur. J.* **2019**, *25*, 4931.
- [17] (a) Mizoroki, T.; Mori, K.; Ozaki, A. *Bull. Chem. Soc. Jpn.* **1971**, *44*, 581.
- (b) Heck, R. F.; Nolley, J. P. *J. Org. Chem.* **1972**, *37*, 2320.
- [18] (a) Benati, L.; Calestani, G.; Leadini, R.; Minozzi, M.; Nanni, D.; Spagnolo, P.; Stazzari, S.; Zanardi, G. *J. Org. Chem.* **2003**, *68*, 3454.
- (b) Saito, T.; Nihei, H.; Otani, T.; Suyama, T.; Furukawa, N.; Saito, M. *Chem. Commun.* **2008**, 172.

(Zhao, C.)