银催化 1-溴代炔烃的功能化反应: 高区域和立体选择性合成 (Z)- β -溴-1-芳基乙烯基芳基酯

孙明利^{†,a,b} 张家俊^{†,a,d} 张义成^{*,a,c} 李品华^{a,c} 王磊^{*,a,b,c}

(^a 淮北师范大学化学与材料科学学院 安徽淮北 235000)

(^b 台州学院医药化工与材料工程学院 浙江台州 318000)

(^c 中国科学院上海有机化学研究所金属有机国家重点实验室 上海 200032)

(^d 浙江金华康恩贝生物药业有限公司 浙江金华 321016)

摘要 发展一种银催化 1-溴代炔烃的官能团反应, 用于高度区域和立体选择性合成(Z)- β -溴-1-芳基乙烯基芳基酯。以市售的芳香羧酸为原料, Ag₂O 为催化剂, 以 Et₃N 为碱, 1-溴代炔与芳香羧酸反应, 高收率得到相应的(Z)- β -溴-1-芳基乙烯基芳基酯产物。研究表明, Ag₂O 在反应中起着重要的催化作用。

关键词 1-溴代炔; (Z)- β -溴-1-芳基乙烯基芳基酯; 银催化; 区域选择性; 立体选择性

A Silver-Catalyzed Functionalization of 1-Bromoalkynes: Highly Regio- and Stereo-selective Synthesis of (Z)- β -Bromo-1-arylvinyl Aryl Esters

Sun, Mingli^{†,a,b} Zhang, Jiajun^{†,a,d} Zhang, Yicheng^{*,a,c} Li, Pinhua^{a,c} Wang, Lei^{*,a,b,c}

(^a Key Laboratory of Green and Precise Synthetic Chemistry, Ministry of Education, Department of Chemistry, Huaibei Normal University, Huaibei, Anhui 235000)

(^b Advanced Research Institute and Department of Chemistry, Taizhou University, Taizhou, Zhejiang 318000)

(^c State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Shanghai 200032)

(^d Zhejiang Jinhua CONBA Bio-Pharm. Co., Ltd., Jinhua, Zhejiang 321016)

Abstract A silver-catalyzed functionalization of 1-bromoalkynes for the highly regio- and stereo-selective synthesis of (Z)- β -bromo-1-arylvinyl aryl esters was developed. In the presence of Ag₂O as a catalyst, and Et₃N as a base, the reactions of 1-bromoalkynes with commercially available aromatic carboxylic acids underwent smoothly to afford the corresponding (Z)- β -bromo-1-arylvinyl aryl esters in good yields. The investigation indicates that Ag₂O plays an important role in the reaction.

Keywords 1-bromoalkynes; (Z)- β -bromo-1-arylvinyl aryl esters; silver catalysis; regio-selectivity; stereo-selectivity

1 Introduction

To develop general and effective methods for the synthesis of complex molecular skeletons is one of the focuses of modern organic chemistry. The haloethylene carboxylates are important intermediates in organic synthesis because the vinyl halide moiety is often employed as the partners of transition-metal-catalyzed

cross-coupling reactions and halogen-metal exchange reactions.^[1] Particularly, haloethylene carboxylates are frequently used as the important intermediates in the synthetic chemistry and pharmaceutical chemistry.^[2-3] For the preparation of haloethylene carboxylates, Barluenga *et al.*^[4] described a reaction of terminal alkynes with bis(pyridine)iodonium(I) tetrafluoroborate and acetic acid to (E)- β -iodoenol acetates in 1990. Twenty years later,

* Corresponding authors. E-mail: lbqzhych@163.com; leiwang@chnu.edu.cn

Received March 15, 2020; revised May 18, 2020; published online May 27, 2020.

Project supported by the National Natural Science Foundation of China (No. 21772062).

国家自然科学基金(No. 21772062)资助项目.

[†] 共同第一作者(These authors contributed equally to this work).

Jiang *et al.*^[5] developed an elegant Ag-catalyzed difunctionalization of terminal alkynes with acetic anhydride and *N*-halosuccinimide (NXS, X=Cl, Br and I) to (*Z*)- β -haloenol acetates with high regio- and stereo-selectivity in 2010 (Scheme 1a). Recently, Jiang *et al.*^[6] further reported a Pd-catalyzed sequential nucleophilic addition/oxidative annulation of bromoalkynes with benzoic acids to isocoumarins in one-pot, suggesting via a haloethylene carboxylate intermediate. Although other methods for the preparation of haloethylene carboxylates from terminal alkynes have been reported,^[7] there are some drawbacks of poor yields, harsh reaction conditions, complicated procedure or low stereoselectivity to limit the applications. Therefore, to develop efficient method from the commercially available starting materials for the synthesis of halogenated vinyl carboxylic acid esters is highly desired.

Previous works:

Scheme 1 Transformations of alkynes and 1-bromoalkynes

During the past decades, haloalkynes have become powerful and highly versatile building blocks in organic chemistry, which exhibit abundant and tunable reactivities, particularly in the presence of transition-metal catalyst.^[8] So, the remarkable efforts have been devoted to the synthesis of a valuable scaffold from haloalkynes, and a variety of achievements have been obtained.^[9] For examples, a series of nucleophilic additions of fluoride,^[10] iodide,^[11] acetate,^[5,12] isocyanide,^[13] sulfide,^[14] and phenol^[15] to haloalkynes, have been investigated, providing the corresponding trisubstituted alkenes in exclusive *Z*-type with high yields. Moreover, several organic transformations using haloalkynes as starting materials under visible light irradiation were established, including an alkynylation of α -keto acids with bromoacetylenes to

ynones, an oxidative amidation of bromoalkynes with anilines to α -ketoamides, and a three-component cyclization of 2*H*-azirines, alkynyl bromides and molecular oxygen to oxazoles (Scheme 1b and 1c).^[16] However, most of the above reactions are involved in the cleavage of C—X bonds to complete the further transformations. As part of continuing our study on the chemistry of 1-bromoalkynes, we wish to report a silver-catalyzed functionalization 1-bromoalkynes with aryl carboxylic acids to afford (*Z*)- β -bromo-1-arylvinyl aryl esters in high regio- and stereo-selectivity with good yields under mild reaction conditions (Scheme 1d).

2 Results and discussion

Initially, a model reaction of phenylethynyl bromide (**1a**) with benzoic acid (**2a**) was chosen to optimize reaction conditions. When the model reaction was conducted in the presence of Ag_2O (10 mol%) and KOAc (1.5 equiv.) in *N,N*-dimethylformamide (DMF) under air atmosphere at 80 °C for 12 h, a nucleophilic addition of **1a** proceeded selectively and delivered the desired product (*Z*)-2-bromo-1-phenylvinyl benzoate (**3a**) in 80% yield, and no (*E*)-isomer of **3a** was detected (Table 1, Entry 1). It was found that the base played an important role in the reaction. Screening of the bases indicated that Et_3N was the best one for the reaction among the tested inorganic and organic bases, providing **3a** in 94% yield (Table 1, Entry 2). Other bases exhibited the less reactivity and showed the following order, $Et_3N > KOAc > NaOAc > Cs_2CO_3 > K_2CO_3 > Na_2CO_3 > DABCO > K_3PO_4 > DBU > LiO^tBu = KO^tBu$ to the model reaction (Table 1, Entries 3~11). It was noteworthy that the reaction could not occur when LiO^tBu and KO^tBu were used as bases (Table 1, Entries 10, 11). Then the effect of solvents on the reaction was explored, and the results were also shown in Table 1. When *N,N*-dimethylacetamide (DMA), dimethyl sulfoxide (DMSO), *N*-methyl pyrrolidone (NMP), CH_3CN , CH_3NO_2 and 1,2-dichloroethane (DCE) were used as reaction media, the reactions provided **3a** in 27%~78% yields (Table 1, Entries 12~17), indicating the inferior reactivity to the functionalization of 1-bromoalkyne. It should be noted that Ag-catalyst played an important role in the reaction and it was found that Ag was essential to the reaction (Table 1, Entry 18). The effect of a series of Ag-catalysts on the yield of **3a** was explored, and Ag_2O showed highest reactivity. Meanwhile, Ag_2CO_3 and $AgOAc$ used as catalysts, **3a** was obtained in 72% and 68% yield, respectively (Table 1, Entries 19, 20). In addition, Ag_2SO_4 and $AgCl$ used as catalysts gave trace amounts of **3a** (Table 1, Entries 21, 22). Finally, the optimal amount of Ag_2O and Et_3N were found to be 7.5 mol% and 1.2 equiv., respectively for 0.50 mmol scale reaction (Table 1, Entries 23~26).

Based on the optimized conditions in our hand, the generality of (*Z*)- β -bromo-1-phenylvinyl aryl esters preparation from phenylethynyl bromide (**1a**) was investigated,

Table 1 Optimization of the reaction conditions^a

Entry	Solvent	Base	Ag source	Yield ^b /%
1	DMF	KOAc	Ag ₂ O	80
2	DMF	Et₃N	Ag₂O	94
3	DMF	NaOAc	Ag ₂ O	78
4	DMF	Cs ₂ CO ₃	Ag ₂ O	60
5	DMF	K ₂ CO ₃	Ag ₂ O	47
6	DMF	Na ₂ CO ₃	Ag ₂ O	39
7	DMF	DABCO	Ag ₂ O	33
8	DMF	K ₃ PO ₄	Ag ₂ O	23
9	DMF	DBU	Ag ₂ O	21
10	DMF	LiO ^t Bu	Ag ₂ O	NR
11	DMF	KO ^t Bu	Ag ₂ O	NR
12	DMA	Et ₃ N	Ag ₂ O	78
13	DMSO	Et ₃ N	Ag ₂ O	56
14	NMP	Et ₃ N	Ag ₂ O	31
15	CH ₃ CN	Et ₃ N	Ag ₂ O	28
16	CH ₃ NO ₂	Et ₃ N	Ag ₂ O	27
17	DCE	Et ₃ N	Ag ₂ O	44
18	DMF	Et ₃ N	—	NR
19	DMF	Et ₃ N	Ag ₂ CO ₃	72
20	DMF	Et ₃ N	AgOAc	68
21	DMF	Et ₃ N	Ag ₂ SO ₄	Trace
22	DMF	Et ₃ N	AgCl	Trace
23	DMF	Et ₃ N	Ag ₂ O	94 ^c
24	DMF	Et ₃ N	Ag ₂ O	93 ^d
25	DMF	Et ₃ N	Ag ₂ O	94 ^e
26	DMF	Et ₃ N	Ag ₂ O	92 ^f

^a Reaction conditions: phenylethynyl bromide (**1a**, 0.50 mmol), benzoic acid (**2a**, 1.0 mmol), Ag-catalyst (10 mol%), base (0.75 mmol, 1.5 equiv.), solvent (2.0 mL), 80 °C, sealed tube, air, 12 h. ^b Isolated yield. ^c Ag₂O (7.5 mol%) was used. ^d Ag₂O (5 mol%) was added. ^e Et₃N (0.60 mmol, 1.2 equiv.) was used. ^f Et₃N (0.50 mmol, 1.0 equiv.) was added. NR = No reaction.

as shown in Table 2. When aromatic carboxylic acids with an electron-donating group, such as methyl group on the *ortho*- or *meta*-position of the benzene rings, the corresponding products (**3b** and **3c**) were obtained in 88% and 84% yields, respectively. It is obvious that the reactions of **1a** and substituted benzoic acids with a moderate electron-withdrawing group including Cl, Br, and I on the phenyl rings underwent the nucleophilic addition smoothly to generate the desired products (**3d**~**3i**) in 83%~90% yields, neglecting steric effect. It should be noted that halogens such as Cl, Br, and I on the benzoic acid moiety were well tolerated and the obtained products provided further opportunity for organic transformations via C(sp²)X (X = Cl, Br and I) bonds activation and functionalization. When one or two NO₂ (nitro group), a

strong electron-withdrawing group, was introduced into the benzene rings of aromatic acids, they also reacted with **1a** to afford the anticipated products (**3j**~**3l**) in 67%~81% yields. The structure of **3j** was unambiguously confirmed by X-ray crystallography.^[17] The electronic effect and steric effect of the substituted benzoic acids was found to be neglected in the reactions. Moreover, furan-2-carboxylic acid and acrylic acid could also be transformed into the corresponding products (**3m** and **3n**) in 91% and 72% yields, respectively.

Table 2 Scope of aromatic carboxylic acids^a

^a Reaction conditions: phenylethynyl bromide (**1a**, 0.50 mmol), aryl carboxylic acid (**2**, 1.0 mmol), Ag₂O (7.5 mol%), Et₃N (0.60 mmol, 1.2 equiv.), DMF (2.0 mL), 80 °C, sealed tube, air, 12 h; isolated yields of the products.

Subsequently, we turned our attention to extend the

scope of acetylene bromides, as shown in Table 3. A number of arylethynyl bromides with an electron-withdrawing or an electron-donating substituent at the *para*-position of the benzene rings, including F, Cl, Br, Me and *n*-C₃H₇, were used to react with benzoic acid (**2a**) under the standard conditions, affording the according products (**3o**~**3p** and **3r**~**3t**) in 88%~91% yields, indicating little electronic effect. On the other hand, *meta*-substituted phenylethynyl bromide also reacted with **2a** smoothly to generate the anticipated product (**3q**) in 87% yield. In addition, aliphatic terminal acetylene bromide, such as 1-bromoheptyne was used as substrate, and the expected product (**3u**) was obtained in 69% yield.

Table 3 Scope of acetylene bromides^a

^a Reaction conditions: acetylene bromide (**1**, 0.50 mmol), benzoic acid (**2a**, 1.0 mmol), Ag₂O (7.5 mol%), Et₃N (0.60 mmol, 1.2 equiv.), DMF (2.0 mL), 80 °C, sealed tube, air, 12 h; isolated yields of the products.

On the basis of our result and the previous report,^[6] a possible mechanism is proposed as shown in Scheme 2. Firstly, Ag₂O reacts with Et₃N to generate an Ag(I)-complex along with the formation of OH⁻. Then, the obtained Ag(NEt₃)₄-complex undergoes the reaction with phenylethynyl bromide (**1a**) to afford π-complex **A**, which is subsequently converted to the corresponding σ-complex **C** by nucleophilic attack of the benzoic anion (**B**) from the reaction of benzoic acid (**2a**) with generated OH⁻. Finally, the silver complex **C** takes a reaction with Et₃N and **2a** to provide the desired product (*Z*)-β-bromo-1-phenylvinyl benzoate (**3a**) in high regio- and

stereo-selectivity, and regeneration of Ag/NEt₃-complex and benzoic anion (**B**) for the next run.

Scheme 2 Proposed mechanism

3 Conclusions

In summary, a silver-catalyzed method for the nucleophilic addition of 1-bromoalkyne with aromatic carboxylic acids to afford (*Z*)-β-bromo-1-arylvinyl aryl esters has been developed. This reaction uses commercially available aryl acids as nucleophiles and shows a wide range of substrates, which can provide corresponding *Z*-form addition products in high yields under mild reaction conditions. The reaction has a variety of advantages, including a wide range of substrates, simple operation, easy raw materials, good yields, excellent regio- and stereo-selectivity. A detail mechanistic study is currently underway in our laboratory.

4 Experimental section

4.1 General experimental

All reactions were carried out under air. ¹H NMR and ¹³C NMR spectra were measured on a Bruker Avance NMR spectrometer (400 MHz or 100 MHz, respectively) in CDCl₃ or DMSO-*d*₆ as solvent with internal tetramethylsilane standard. Ethyl acetate and petroleum ether were used for column chromatography without further purification. All solvents and chemicals were obtained from commercial sources and used as received unless otherwise noted.

4.2 General procedure

A 10 mL of reaction tube was charged with phenylethynyl bromide (**1a**, 0.50 mmol), benzoic acid (**2a**, 1.0 mmol), Ag₂O (0.0375 mmol, 7.5 mol%), Et₃N (0.60 mmol, 1.2 equiv.), and DMF (2.0 mL). The reaction vessel was placed in an oil bath. After the reaction was carried out at 80 °C for 12 h, it was cooled to room temperature, extracted with EtOAc (5.0 mL × 3). The organic layers were combined, dried over MgSO₄, and concentrated to yield the crude product, which was further purified by flash chromatography (silica gel, petroleum ether/ethyl acetate, *V* : *V* = 7 : 1 to 12 : 1) to give the desired product **3a**.

4.3 Characterization data

(*Z*)-2-Bromo-1-phenylvinyl benzoate (**3a**):^[6] White solid. m.p. 46.1~47.4 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.25 (d, *J*=8.0 Hz, 2H), 7.70~7.66 (m, 1H), 7.57~7.53 (m, 2H), 7.50~7.49 (m, 2H), 7.37~7.32 (m, 3H), 6.68 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.0, 150.6, 133.8, 133.3, 130.3, 129.3, 128.7, 128.7, 128.6, 124.9, 96.6. HRMS (ESI) calcd for C₁₅H₁₂BrO₂ [M+H]⁺ 303.0015, found 303.0018.

(*Z*)-2-Bromo-1-phenylvinyl 3-methylbenzoate (**3b**): White solid. m.p. 54.6~55.9 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.08~8.07 (m, 2H), 7.53~7.49 (m, 3H), 7.46~7.42 (m, 1H), 7.38~7.37 (m, 3H), 6.70 (s, 1H), 2.48 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.2, 150.6, 138.5, 134.6, 133.3, 130.7, 129.3, 128.7, 128.6, 128.5, 127.5, 124.9, 96.6, 21.2. HRMS (ESI) calcd for C₁₆H₁₄BrO₂ [M+H]⁺ 317.0172, found 317.0169.

(*Z*)-2-Bromo-1-phenylvinyl 2-methylbenzoate (**3c**): White solid. m.p. 52.8~53.7 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.30 (d, *J*=7.6 Hz, 2H), 7.53~7.51 (m, 3H), 7.40~7.38 (m, 3H), 7.36~7.35 (m, 2H), 6.69 (s, 1H), 2.70 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.5, 150.7, 141.6, 133.5, 132.9, 131.9, 131.3, 129.3, 128.8, 127.7, 125.9, 124.9, 96.6, 21.8. HRMS (ESI) calcd for C₁₆H₁₄BrO₂ [M+H]⁺ 317.0172, found 317.0170.

(*Z*)-2-Bromo-1-phenylvinyl 2-chlorobenzoate (**3d**): White solid. m.p. 62.1~64.2 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.20 (d, *J*=8.0 Hz, 2H), 7.56~7.54 (m, 2H), 7.54~7.52 (m, 2H), 7.45~7.42 (m, 1H), 7.40~7.39 (m, 3H), 6.69 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 161.5, 150.5, 134.6, 133.4, 133.0, 132.1, 131.4, 129.4, 128.8, 128.3, 126.7, 125.0, 96.8. HRMS (ESI) calcd for C₁₅H₁₁BrClO₂ [M+H]⁺ 336.9625, found 336.9622.

(*Z*)-2-Bromo-1-phenylvinyl 4-bromobenzoate (**3e**): White solid. m.p. 68.9~70.2 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.11 (d, *J*=8.4 Hz, 2H), 7.69 (d, *J*=8.0 Hz, 2H), 7.50~7.48 (m, 2H), 7.38~7.37 (m, 3H), 6.69 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 162.4, 150.5, 133.1, 132.0, 131.7, 129.4, 129.2, 128.8, 127.6, 124.9, 96.8. HRMS (ESI) calcd for C₁₅H₁₁Br₂O₂ [M+H]⁺ 382.9100, found 382.9104.

(*Z*)-2-Bromo-1-phenylvinyl 2-bromobenzoate (**3f**): White solid. m.p. 65.6~67.9 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.21~7.19 (m, 2H), 7.77~7.75 (m, 1H), 7.55~7.53 (m, 2H), 7.50~7.43 (m, 2H), 7.40~7.39 (m, 3H), 6.69 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 161.9, 150.5, 134.8, 133.4, 133.0, 132.1, 130.3, 129.4, 128.8, 127.3, 125.0, 122.5, 96.9. HRMS (ESI) calcd for C₁₅H₁₁Br₂O₂ [M+H]⁺ 382.9100, found 382.9101.

(*Z*)-2-Bromo-1-phenylvinyl 4-iodobenzoate (**3g**): White solid. m.p. 66.8~68.2 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.11 (d, *J*=8.8 Hz, 2H), 7.69 (d, *J*=8.4 Hz, 2H), 7.50~7.48 (m, 2H), 7.38~7.37 (m, 3H), 6.69 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 162.6, 150.4, 138.0, 133.0, 131.6, 129.4, 128.8, 128.1, 124.8, 101.9, 96.8. HRMS (ESI) calcd for C₁₅H₁₁BrIO₂ [M+H]⁺ 428.8982, found 428.8981.

(*Z*)-2-Bromo-1-phenylvinyl 3-iodobenzoate (**3h**): White

solid. m.p. 56.6~58.3 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.60 (s, 1H), 8.22 (d, *J*=7.6 Hz, 1H), 8.00 (d, *J*=8.0 Hz, 1H), 7.50~7.48 (m, 2H), 7.39~7.38 (m, 3H), 7.31~7.27 (m, 1H), 6.70 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 161.6, 150.4, 142.7, 139.0, 133.0, 130.6, 130.3, 129.5, 129.4, 128.8, 124.9, 96.9, 94.0. HRMS (ESI) calcd for C₁₅H₁₁BrIO₂ [M+H]⁺ 428.8982, found 428.8983.

(*Z*)-2-Bromo-1-phenylvinyl 2-iodobenzoate (**3i**): White solid. m.p. 61.4~62.8 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.24~8.23 (m, 1H), 8.10 (d, *J*=8.0 Hz, 1H), 7.53~7.50 (m, 3H), 7.40~7.38 (m, 3H), 7.28~7.23 (m, 1H), 6.69 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 162.3, 150.5, 141.9, 133.5, 133.0, 133.0, 131.8, 129.4, 128.8, 128.0, 96.9, 94.9. HRMS (ESI) calcd for C₁₅H₁₁BrIO₂ [M+H]⁺ 428.8982, found 428.8980.

(*Z*)-2-Bromo-1-phenylvinyl 4-nitrobenzoate (**3j**): Pale yellow solid. m.p. 149.1~149.8 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.42 (d, *J*=8.8 Hz, 2H), 8.37 (d, *J*=8.8 Hz, 2H), 7.49~7.47 (m, 2H), 7.40~7.38 (m, 3H), 6.71 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 161.2, 150.0, 150.4, 134.0, 132.7, 131.4, 129.6, 128.8, 124.8, 123.7, 97.0. HRMS (ESI) calcd for C₁₅H₁₁BrNO₄ [M+H]⁺ 347.9866, found 347.9866.

(*Z*)-2-Bromo-1-phenylvinyl 2-nitrobenzoate (**3k**): Pale yellow solid. m.p. 146.1~147.4 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.11~8.09 (m, 1H), 7.98~7.96 (m, 1H), 7.78~7.69 (m, 2H), 7.5~7.56 (m, 2H), 7.43~7.41 (m, 3H), 6.67 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 161.3, 150.5, 148.2, 132.9, 132.5, 132.5, 130.4, 129.7, 128.8, 126.1, 125.3, 124.1, 97.1. HRMS (ESI) calcd for C₁₅H₁₁BrNO₄ [M+H]⁺ 347.9866, found 347.9869.

(*Z*)-2-Bromo-1-phenylvinyl 3,5-dinitrobenzoate (**3l**): Yellow solid. m.p. 157.9~158.3 °C; ¹H NMR (400 MHz, CDCl₃) δ: 9.34~9.33 (m, 3H), 7.48~7.47 (m, 2H), 7.42~7.40 (m, 3H), 6.67 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 159.2, 150.2, 148.8, 132.4, 132.2, 129.9, 129.8, 128.9, 124.9, 123.1, 97.4. HRMS (ESI) calcd for C₁₅H₁₀BrN₂O₆ [M+H]⁺ 392.9717, found 392.9718.

(*Z*)-2-Bromo-1-phenylvinyl furan-2-carboxylate (**3m**): White solid. m.p. 86.1~87.2 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.70 (s, 1H), 7.49~7.46 (m, 3H), 7.37~7.36 (m, 3H), 6.68 (s, 1H), 6.62~6.61 (m, 1H). ¹³C NMR (100 MHz, CDCl₃) δ: 154.6, 149.8, 147.5, 143.2, 133.0, 129.4, 128.7, 124.9, 120.0, 112.2, 97.1. HRMS (ESI) calcd for C₁₃H₁₀BrO₃ [M+H]⁺ 292.9808, found 292.9811.

(*Z*)-2-Bromo-1-phenylvinyl acrylate (**3n**): Pale yellow oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.44~7.43 (m, 2H), 7.38~7.36 (m, 3H), 6.70~6.66 (m, 1H), 6.61 (s, 1H), 6.41~6.34 (m, 1H), 6.10~6.07 (m, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 162.3, 150.2, 133.3, 133.1, 129.3, 128.7, 126.9, 124.8, 96.5. HRMS (ESI) calcd for C₁₁H₁₀BrO₂ [M+H]⁺ 252.9859, found 252.9862.

(*Z*)-2-Bromo-1-(*p*-tolyl)vinyl benzoate (**3o**): White solid. m.p. 76.2~77.2 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.27 (d, *J*=8.0 Hz, 2H), 7.70~7.66 (m, 1H), 7.57~7.53 (m, 2H), 7.40 (d, *J*=8.0 Hz, 2H), 7.18 (d, *J*=8.0 Hz, 2H), 6.63 (s, 1H), 2.36 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ:

163.1, 150.7, 139.5, 133.7, 130.5, 130.3, 129.4, 128.8, 128.6, 124.8, 95.6, 21.2. HRMS (ESI) calcd for $C_{16}H_{14}BrO_2$ $[M+H]^+$ 317.0172, found 317.0174.

(Z)-2-Bromo-1-(4-fluorophenyl)vinyl benzoate (**3p**): White solid. m.p. 79.4~80.9 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.26 (d, $J=8.0$ Hz, 2H), 7.70~7.67 (m, 1H), 7.57~7.53 (m, 2H), 7.50~7.47 (m, 2H), 7.08~7.04 (m, 2H), 6.62 (s, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 163.2 (d, $J_{CF}=250.8$ Hz), 163.0, 149.7, 133.9, 130.3, 129.6 (d, $J_{CF}=3.2$ Hz), 128.6, 128.5, 126.9 (d, $J_{CF}=8.4$ Hz), 115.8 (d, $J_{CF}=21.9$ Hz), 96.4 (d, $J_{CF}=1.6$ Hz). HRMS (ESI) calcd for $C_{15}H_{11}BrFO_2$ $[M+H]^+$ 320.9921, found 320.9925.

(Z)-2-Bromo-1-(3-fluorophenyl)vinyl benzoate (**3q**): White solid. m.p. 75.6~76.2 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.06 (d, $J=7.6$ Hz, 2H), 7.94 (d, $J=8.8$ Hz, 2H), 7.56~7.49 (m, 3H), 7.42~7.36 (m, 4H), 6.70 (s, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 162.6 (d, $J_{CF}=248.2$ Hz), 161.9 (d, $J_{CF}=3.2$ Hz), 150.5, 133.0, 130.8 (d, $J_{CF}=7.7$ Hz), 130.3 (d, $J_{CF}=7.9$ Hz), 129.4, 128.8, 126.0 (d, $J_{CF}=3.2$ Hz), 124.9, 120.1 (d, $J_{CF}=21.2$ Hz), 117.1 (d, $J_{CF}=23.1$ Hz), 96.8. HRMS (ESI) calcd for $C_{15}H_{11}BrFO_2$ $[M+H]^+$ 320.9921, found 320.9922.

(Z)-2-Bromo-1-(4-chlorophenyl)vinyl benzoate (**3r**): White solid. m.p. 86.3~88.2 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.25 (d, $J=8.0$ Hz, 2H), 7.70~7.66 (m, 1H), 7.57~7.53 (m, 2H), 7.43 (d, $J=8.4$ Hz, 2H), 7.34 (d, $J=8.0$ Hz, 2H), 6.69 (s, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 163.0, 149.7, 135.3, 133.9, 131.8, 130.3, 129.0, 128.7, 128.5, 126.2, 97.3. HRMS (ESI) calcd for $C_{15}H_{11}BrClO_2$ $[M+H]^+$ 336.9625, found 336.9628.

(Z)-2-Bromo-1-(4-bromophenyl)vinyl benzoate (**3s**): White solid. m.p. 83.3~84.8 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.24 (d, $J=7.6$ Hz, 2H), 7.70~7.67 (m, 1H), 7.57~7.53 (m, 2H), 7.49 (d, $J=8.0$ Hz, 2H), 7.36 (d, $J=8.4$ Hz, 2H), 6.69 (s, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 163.0, 149.7, 133.9, 132.3, 131.9, 130.3, 128.7, 128.4, 126.4, 123.5, 97.4. HRMS (ESI) calcd for $C_{15}H_{11}Br_2O_2$ $[M+H]^+$ 382.9100, found 382.9101.

(Z)-2-Bromo-1-(4-propylphenyl)vinyl benzoate (**3t**): White solid. m.p. 98.7~100.3 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.25 (d, $J=7.6$ Hz, 2H), 7.69~7.66 (m, 1H), 7.56~7.53 (m, 2H), 7.40 (d, $J=8.0$ Hz, 2H), 7.17 (d, $J=8.0$ Hz, 2H), 6.63 (s, 1H), 2.59 (t, $J=7.6$ Hz, 2H), 1.65 (q, $J=7.2$ Hz, 2H), 0.95 (t, $J=7.6$ Hz, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 163.1, 150.6, 144.2, 133.7, 130.7, 130.3, 128.8, 128.8, 28.6, 124.8, 95.6, 37.7, 24.2, 13.6. HRMS (ESI) calcd for $C_{18}H_{18}BrO_2$ $[M+H]^+$ 345.0485, found 345.0487.

(Z)-1-Bromohept-1-en-2-yl benzoate (**3u**):^[6] pale yellow oil. 1H NMR (400 MHz, $CDCl_3$) δ : 8.17~8.15 (m, 2H), 7.65~7.61 (m, 1H), 7.52~7.48 (m, 2H), 5.94 (s, 1H), 2.44 (t, $J=7.6$ Hz, 2H), 1.57~1.54 (m, 2H), 1.38~1.32 (m, 4H), 0.92~0.89 (m, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 163.0, 153.2, 133.5, 130.0, 129.0, 128.5, 93.7, 33.5, 31.0, 25.8, 22.2, 13.8. HRMS (ESI) calcd for $C_{14}H_{18}BrO_2$ $[M+H]^+$ 297.0485, found 297.0486.

Supporting Information 1H NMR and ^{13}C NMR spectra of compounds **3a**~**3u**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Corbet, J.-P.; Mignani, G. *Chem. Rev.* **2006**, *106*, 2651.
(b) Cahiez, G.; Moyeux, A. *Chem. Rev.* **2003**, *103*, 4283.
(c) Amatore, C.; Jutand, A. *Acc. Chem. Res.* **2000**, *33*, 314.
(d) Bettinger, H. F.; Filthaus, M. J. *Org. Chem.* **2007**, *72*, 9750.
(e) Uchiyama, M.; Furuyama, T.; Kobayashi, M.; Matsumoto, Y.; Tanaka, K. *J. Am. Chem. Soc.* **2006**, *128*, 8404.
(f) Boukouvalas, J.; Loach, R. P. *J. Org. Chem.* **2008**, *73*, 8109.
- [2] (a) Goossen, L. J.; Paetzold, J. *Angew. Chem., Int. Ed.* **2004**, *43*, 1095.
(b) Zhang, D.; Ready, J. M. *Org. Lett.* **2005**, *7*, 5681.
(c) DeBergh, J. R.; Spivey, K. M.; Ready, J. M. *J. Am. Chem. Soc.* **2008**, *130*, 7828.
(d) Tang, W.; Liu, D.; Zhang, X. *Org. Lett.* **2003**, *5*, 205.
- [3] (a) Kowalski, C. J.; Haque, M. S. *J. Org. Chem.* **1985**, *50*, 5140.
(b) Kowalski, C. J.; O'Dowd, M. L.; Burke, M. C.; Fields, K. W. *J. Am. Chem. Soc.* **1980**, *102*, 5411.
(c) Kowalski, C. J.; Haque, M. S.; Fields, K. W. *J. Am. Chem. Soc.* **1985**, *107*, 1429.
- [4] Barluenga, J.; Rodríguez, M. A.; Campos, P. J. *J. Org. Chem.* **1990**, *55*, 3104.
- [5] Chen, Z.; Li, J.; Jiang, H.; Zhu, S.; Li, Y.; Qi, C. *Org. Lett.* **2010**, *12*, 3262.
- [6] Jiang, G.; Li, J. X.; Zhu, C.; Wu, W.; Jiang, H. *Org. Lett.* **2017**, *19*, 4440.
- [7] (a) Kowalski, C. J.; Creary, X.; Rollin, A. J.; Burke, M. C. *J. Org. Chem.* **1978**, *43*, 2601.
(b) Pincock, J. A.; Yates, K. *Can. J. Chem.* **1970**, *48*, 3332.
(c) Chawla, R.; Singh, A. K.; Yadav, L. D. S. *Synlett* **2013**, 1558.
(d) Xia, X.-F.; Gu, Z.; Liu, W.; Wang, N.; Wang, H.; Xia, Y.; Gao, H.; Liu, X. *Org. Biomol. Chem.* **2014**, *12*, 9909.
(e) Wang, Y.; Lu, B.; Zhang, L. *Chem. Commun.* **2010**, *46*, 9179.
- [8] (a) Wu, W.; Jiang, H. *Acc. Chem. Res.* **2014**, *47*, 2483.
(b) Trofimov, A.; Chernyak, N.; Gevorgyan, V. *J. Am. Chem. Soc.* **2008**, *130*, 13538.
(c) Yoo, W.-J.; Allen, A.; Villeneuve, K.; Tam, W. *Org. Lett.* **2005**, *7*, 5853.
(d) Hein, J. E.; Tripp, J. C.; Krasnova, L. B.; Sharpless, K. B.; Fokin, V. V. *Angew. Chem., Int. Ed.* **2009**, *48*, 8018.
(e) Crossley, J. A.; Browne, D. L. *J. Org. Chem.* **2010**, *75*, 5414.
(f) Xie, L.; Wu, Y.; Yi, W.; Zhu, L.; Xiang, J.; He, W. *J. Org. Chem.* **2013**, *78*, 9190.
(g) Wezeman, T.; Zhong, S.; Nieger, M.; Bräse, S. *Angew. Chem., Int. Ed.* **2016**, *55*, 3823.
(h) Alcaide, B.; Almendros, P.; Milla, C. L. *Adv. Synth. Catal.* **2017**, *359*, 2630.
- [9] (a) Rasolofonjatovo, E.; Theeramunkong, S.; Bouriaud, A.; Kolodych, S.; Chaumontet, M.; Taran, F. *Org. Lett.* **2013**, *15*, 4698.
(b) Ye, Q.; Cheng, T.; Zhao, Y.; Zhao, J.; Jin, R.; Liu, G. *Chem-CatChem* **2015**, *7*, 1801.
(c) Zhu, G.; Chen, D.; Wang, Y.; Zheng, R. *Chem. Commun.* **2012**, *48*, 5796.
(d) Saueremann, N.; Gonzalez, M. J.; Ackermann, L. *Org. Lett.* **2015**, *17*, 5316.
(e) Huang, H.-Y.; Cheng, L.; Liu, J.-J.; Wang, D.; Liu, L.; Li, C.-J. *J. Org. Chem.* **2017**, *82*, 2656.
(f) Ji, F.; Li, X.; Wu, W.; Jiang, H. *J. Org. Chem.* **2014**, *79*, 11246.
(g) Zeng, X.; Liu, S.; Shi, Z.; Xu, B. *Org. Lett.* **2016**, *18*, 4770.
(h) Kreuzahler, M.; Daniels, A.; Wölper, C.; Haberhauer, G. *J. Am. Chem. Soc.* **2019**, *141*, 1337.
(i) Lyu, X.-L.; Huang, S.-S.; Song, H.-J.; Liu, Y.-X.; Wang, Q.-M.

- RSC Adv.* **2019**, *9*, 36213.
- (j) Pan, R.; Shi, C.; Zhang, D.; Tian, Y.; Guo, S.; Yao, H.; Lin, A. *Org. Lett.* **2019**, *21*, 8915.
- (k) Zeng, X.; Liu, S.; Hammond, G. B.; Xu, B. *ACS Catal.* **2018**, *8*, 904.
- [10] (a) Li, Y.; Liu, X.; Ma, D.; Liu, B.; Jiang, H. *Adv. Synth. Catal.* **2012**, *354*, 2683.
(b) Zhao, M.; Ming, L.; Tang, J.; Zhao, X. *Org. Lett.* **2016**, *18*, 416.
(c) Herrera, A. G.; Nahra, F.; Brill, M.; Nolan, S. P.; Cazin, C. S. J. *ChemCatChem.* **2016**, *8*, 3381.
- [11] (a) Chen, Z.; Jiang, H.; Li, Y.; Qi, C. *Chem. Commun.* **2010**, *46*, 8049.
(b) Kawaguchi, S.-I.; Gonda, Y.; Masuno, H.; Thivū, H.; Yamaguchi, K.; Shinohara, H.; Sonoda, M.; Ogawa, A. *Tetrahedron Lett.* **2014**, *55*, 6779.
- [12] Liste, P. J. G.; Francos, J.; Garrido, S. E. G.; Cadierno, V. J. *Org. Chem.* **2017**, *82*, 1507.
- [13] Li, Y.; Zhao, J.; Chen, H.; Liu, B.; Jiang, H. *Chem. Commun.* **2012**, *48*, 3545.
- [14] (a) Jiang, G.; Zhu, C.; Li, J.; Wu, W.; Jiang, H. *Adv. Synth. Catal.* **2017**, *359*, 1208.
(b) Zeng, X.; Chen, B.; Lu, Z.; Hammond, G. B.; Xu, B. *Org. Lett.* **2019**, *21*, 2772.
- [15] Wang, S.; Li, P.; Yu, L.; Wang, L. *Org. Lett.* **2011**, *13*, 5968.
- [16] (a) Chen, L.; Li, H.; Li, P.; Wang, L. *Org. Lett.* **2016**, *18*, 3646.
(b) Tan, H.; Li, H.; Ji, W.; Wang, L. *Angew. Chem., Int. Ed.* **2015**, *54*, 8374.
(c) Ni, K.; Meng, L.-G.; Wang, K.; Wang, L. *Org. Lett.* **2018**, *20*, 2245.
(d) Ni, K.; Meng, L.-G.; Ruan, H.; Wang, L. *Chem. Commun.* **2019**, *55*, 8438.
- [17] The X-ray single crystal structure of (Z)-2-bromo-1-phenylvinyl 4-nitrobenzoate (**3j**, CCDC: 1990480).

(Li, L.; Fan, Y.)