基于化学和对映选择性转移氢化的 β,γ -炔基 α -氨基酸酯的催化不对称合成研究

张璐^{†,a} 刘爱芹^{†,c} 刘华铮^a 万仁忠^{*,a} 孙书涛^{*,b} 刘磊^{*,b}

(^a 山东农业大学动物科技学院 山东泰安 271018)

(^b 山东大学化学与化工学院 济南 250100)

(^c 山东第一医科大学(山东省医学科学院)药物研究所 济南 250062)

摘要 报道了一个通过 β,γ -炔基 α -亚胺酸酯的化学和对映选择性转移氢化反应来合成光学纯 β,γ -炔基 α -氨基酸酯的方法。该不对称还原反应所展示出的优秀的化学选择性是由手性磷酸作为催化剂以及苯并噻唑啉作为氢负供体实现的。反应展示出了良好的官能团兼容性, 高对映选择性地合成了一系列光学活性的非天然氨基酸酯化合物。

关键词 β,γ -炔基 α -氨基酸酯; 化学选择性; 对映选择性; 转移氢化; 苯并噻唑啉

Catalytic Asymmetric Synthesis of β,γ -Alkynyl α -Amino Esters via Chemo- and Enantio-selective Transfer Hydrogenation

Zhang, Lu^{†,a} Liu, Aiqin^{†,c} Liu, Huazheng^a Wan, Renzhong^{*,a}
Sun, Shutao^{*,b} Liu, Lei^{*,b}

(^a College of Animal Science and Veterinary Medicine, Shandong Agricultural University, Taian, Shandong 271018)

(^b School of Chemistry and Chemical Engineering, Shandong University, Jinan 250100)

(^c Institute of Materia Medica, Shandong First Medical University & Shandong Academy of Medical Sciences, Jinan 250062)

Abstract An effective chemo- and enantioselective transfer hydrogenation of β,γ -alkynyl α -imino esters to prepare optically pure β,γ -alkynyl α -amino esters has been described. The excellent chemoselectivity was achieved by using chiral phosphoric acid catalyzed asymmetric reduction with benzothiazoline as hydride donor. The reaction exhibited good functional group tolerance, providing a range of optically active non-natural amino esters with excellent enantioselectivity.

Keywords β,γ -alkynyl α -amino ester; chemoselectivity; enantioselectivity; transfer hydrogenation; benzothiazoline

1 Introduction

Chemoselectivity, which refers to the ability of a given reagent to react with one particular group in a molecule in preference to another group, is of vital importance for a given reaction.^[1] Reactions that proceed in highly chemoselective manner require minimal reliance on protecting groups^[2] and improve both atom^[3] and step economy.^[4] Optically active non-natural amino acids are basic subunits for a broad range of important molecules in medicine and biology.^[5-6] In particular, optically pure β,γ -alkynyl α -amino acids and their derivatives, which allow for further synthetic elaboration, represent a special

class of these compounds.^[7] Although continuous attention has been paid to explore the efficient methodologies for their synthesis, asymmetric synthesis of β,γ -alkynyl α -amino acids and their derivatives has remained challenging.^[8-9] The enantioselective addition of terminal alkynes to α -imino esters represents a straightforward strategy.^[8] However, the methods suffered from narrow scope of the alkyne component. Asymmetric reduction of imines has emerged as a powerful tool for amines synthesis,^[10] and therefore selective reduction of β,γ -alkynyl α -imino esters represents the other ideal strategy.^[9] However, chemo- and enantioselective reduction of the imine moiety with the alkyne intact proved to be difficult (Scheme 1, A).

* Corresponding authors. E-mail: wrzh63@163.com; sunshutao@sdu.edu.cn; leiliu@sdu.edu.cn

Received May 15, 2020; revised June 3, 2020; published online June 11, 2020.

Project supported by the National Natural Science Foundation of China (Nos. 21722204, 21971148).

国家自然科学基金(Nos. 21722204, 21971148)资助项目.

[†] 共同第一作者(These authors contributed equally to this work).

You group^[9a] reported a chiral phosphoric acid (CPA) catalyzed asymmetric transfer hydrogenation process using Hantzsch ester as reducing agent, giving β,γ -alkynyl α -amino esters through concurrent reduction of both $\text{C}\equiv\text{C}$ bond and $\text{C}=\text{N}$ bond. Zhou group^[9b] demonstrated that the chemoselective reduction of the $\text{C}=\text{N}$ bond over $\text{C}\equiv\text{C}$ bond could be achieved by using 5,6-dihydrophenanthridine as the hydrogen source. However, the expected β,γ -alkynyl α -amino ester was isolated in only 26% yield with 36% *ee*. Recently, we reported a non-enzymatic redox deracemization of β,γ -alkynyl α -amino esters consisting of a copper mediated aerobic oxidation and CPA catalyzed asymmetric transfer hydrogenation (Scheme 1, B).^[11] β,γ -Alkynyl α -imino esters were identified as oxidized intermediates for subsequent asymmetric reduction using

(A) Asymmetric reduction of β,γ -alkynyl α -imino esters (You and Zhou's works)

(B) Aerobic deracemization of β,γ -alkynyl α -amino esters (our previous work)

(C) Chemo- and enantioselective reduction of β,γ -alkynyl α -imino esters (this work)

Scheme 1 Overview of chemo- and enantioselective reduction of β,γ -alkynyl α -imino esters

benzothiazoline as hydride donor with exclusive chemoselectivity at the imine moiety over alkyne.^[12] Given the ready availability of β,γ -alkynyl α -imino esters, we herein report the direct asymmetric transfer hydrogenation method using benzothiazoline as hydride donor to prepare optically pure β,γ -alkynyl α -amino esters with excellent chemo- and enantioselectivity (Scheme 1, C).

2 Results and discussion

The synthetic route of compound **1** is shown in Scheme 2. Initially, asymmetric transfer hydrogenation of aliphatic alkynyl substituted α -imino ester **1a** was selected as a model reaction using CPA **3** as the catalyst and 1.2 equiv. of benzothiazoline **4** as reducing agent in the presence of 3 Å molecular sieves (Table 1). CPA **3a** catalyzed reaction with **4a** proceeded smoothly affording the expected (*S*)-**2a** in 71% yield with 70% *ee* (Entry 1, Table 1). Notably, no unexpected alkyne reduction product was observed. An extensive investigation of CPA catalysts suggested **3e** was the best choice,^[13] and (*S*)-**2a** was obtained in 72% yield with 76% *ee* (Entries 2~8, Table 1). Further optimization of benzothiazolines identified **4d** to be optimal (Entries 9~13, Table 1). The solvents were found to be crucial to the yields and enantioselectivity, and when asymmetric reduction was performed in a 1 : 1 mixture of decane and mesitylene, the asymmetric transfer hydrogen provided (*S*)-**2a** in 89% yield with 90% *ee* (Entries 14~24, Table 1). The reaction was found to be sensitive to the moisture, and 3 Å molecular sieves was identified as the best choice (Entries 25~27, Table 1).

Reagents and conditions: (a) Ethyl glyoxalate (1.2 equiv.), Na_2SO_4 (2.0 equiv.), toluene; (b) alkyne (2.0 eq), AgOTf (10 mol%), CH_2Cl_2 ; (c) $\text{Cu}(\text{OAc})_2$ (10 mol%), O_2 (101 kPa), CH_2Cl_2 .

Scheme 2 Synthetic route of compound **1**

With the optimized conditions in hand, the scope of asymmetric transfer hydrogenation of β,γ -alkynyl α -imino esters was investigated (Table 2). The reaction was found to be fairly general for substrates bearing a wide range of aliphatic alkynes with varied chain lengths, affording corresponding optically pure **2a**~**2c** in good yields with excellent *ee* (90%~98%). The method had good functional group compatibility and good tolerance to aryl (**2d**), benzyl ether (**2e**), acetal (**2f**), alkenyl (**2g**) and silyl ether (**2h**). Substrates containing electronically varied heteroaryl (**2i**) and aryl acetylenes (**2j**~**2l**) were also suitable candidates. In addition, the method was found to be insensitive to the variant of the ester moiety, as demonstrated by effective reduction of **2m** and **2n**.

Table 1 Reaction condition optimization^a

Entry	Solvent	R (3)	Ar (4)	Yield ^b /%	ee ^c /%
1	Hexane	4-CF ₃ C ₆ H ₄ (3a)	Ph (4a)	71	70
2	Hexane	3,5-(CF ₃) ₂ C ₆ H ₃ (3b)	Ph (4a)	74	53
3	Hexane	1-Naphthyl (3c)	Ph (4a)	65	33
4	Hexane	2-Naphthyl (3d)	Ph (4a)	68	50
5	Hexane	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	Ph (4a)	72	76
6	Hexane	2,4,6-(cyclo-C ₆ H ₁₁) ₃ C ₆ H ₂ (3f)	Ph (4a)	62	71
7	Hexane	SiPh ₃ (3g)	Ph (4a)	<5	n.d.
8	Hexane	9-Anthryl (3h)	Ph (4a)	53	50
9	Hexane	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	4-ClC ₆ H ₄ (4b)	60	70
10	Hexane	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	4-MeOC ₆ H ₄ (4c)	64	47
11	Hexane	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	1-Naphthyl (4d)	67	79
12	Hexane	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	2-Naphthyl (4e)	47	65
13	Hexane	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	9-Anthryl (4f)	<5	n.d.
14	Cyclohexane	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	1-Naphthyl (4d)	73	69
15	CH ₂ Cl ₂	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	1-Naphthyl (4d)	89	7
16	THF	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	1-Naphthyl (4d)	<5	n.d.
17	Ethyl acetate	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	1-Naphthyl (4d)	76	55
18	Toluene	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	1-Naphthyl (4d)	85	71
19	Mesitylene	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	1-Naphthyl (4d)	92	85
20	Pentane	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	1-Naphthyl (4d)	52	68
21	Decane	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	1-Naphthyl (4d)	78	86
22	Decane/mesitylene (V : V = 8 : 2)	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	1-Naphthyl (4d)	80	86
23	Decane/mesitylene (V : V = 7 : 3)	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	1-Naphthyl (4d)	84	87
24	Decane/mesitylene (V : V = 1 : 1)	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	1-Naphthyl (4d)	89	90
25 ^d	Decane/mesitylene (V : V = 1 : 1)	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	1-Naphthyl (4d)	75	68
26 ^e	Decane/mesitylene (V : V = 1 : 1)	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	1-Naphthyl (4d)	80	79
27 ^f	Decane/mesitylene (V : V = 1 : 1)	2,4,6- <i>i</i> -Pr ₃ C ₆ H ₂ (3e)	1-Naphthyl (4d)	82	81

^a Reaction conditions: **1a** (0.1 mmol), **3** (5 mol%), **4** (0.12 mmol), and 3Å molecular sieves (40 mg) in solvent (2.0 mL) at room temperature for 4 h. ^b Yield of the isolated product. ^c Determined by chiral HPLC analysis. ^d Reaction without molecular sieves. ^e 4Å molecular sieves was used. ^f 5Å molecular sieves. n.d. = not determined.

Table 2 Scope of β,γ-alkynyl α-imino esters

 2a 89%, 90% ee	 2b 86%, 98% ee	 2c 85%, 94% ee	 2d 87%, 95% ee	 2e 82%, 93% ee
 2f 80%, 92% ee	 2g 89%, 95% ee	 2h 82%, 91% ee	 2i 90%, 80% ee	 2j 88%, 89% ee
 2k 84%, 86% ee	 2l 87%, 88% ee	 2m 90%, 93% ee	 2n 83%, 92% ee	

3 Conclusions

In summary, an effective chemo- and enantioselective transfer hydrogenation of β,γ -alkynyl α -imino esters to afford optically pure β,γ -alkynyl α -amino esters has been described. The excellent chemoselectivity was achieved by chiral phosphoric acid catalyzed asymmetric reduction with benzothiazoline as hydride donor. The reaction exhibited good functional group tolerance, providing a range of optically active non-natural α -amino esters with excellent enantioselectivity.

4 Experimental section

4.1 General information

All reactions were carried out with dry solvents under anhydrous conditions. Standard syringe techniques were applied for transfer of dry solvents and some air-sensitive reagents and were introduced into reaction vessels through a rubber septum. ^1H NMR and ^{13}C NMR spectra were recorded on a Bruker AVANCE III 500 NMR Spectrometer at 500 MHz and 125 MHz, respectively. CDCl_3 was used as solvent and reference ($\delta_{\text{H}}=7.26$, $\delta_{\text{C}}=77.23$). Analytical thin layer chromatography (TLC) was performed on pre-coated silica gel GF254 plates. Column chromatography was carried out on silica gel (200~300 mesh). HRMS were carried out on an Orbitrap analyzer. UV spectra were obtained with an Agilent 8453E UV-Visible spectroscopy system. Optical rotations were measured using a 2.5 mL cell with a 10 cm path length on a Hanon P850 Automatic Polarimeter. Enantiomeric excesses were determined by HPLC using a Daicel Chiralpak AD-H or IB column with hexane/*i*-PrOH as the eluent.

4.2 Substrate preparation

p-Anisidine (5 mmol) was dissolved in toluene (10 mL), and anhydrous sodium sulfate (10 mmol) was added with stirring. Ethyl glyoxalate (6 mmol) was added slowly and the mixture was allowed to stir for 1 h. After completion of the reaction, sodium sulfate was removed by filtration and toluene was removed under vacuum to yield the crude **S1** which was used for the next step without purification.^[14-15]

To a stirred mixture of the corresponding alkyne (0.4 mmol, 2.0 equiv.) and AgOTf (0.02 mmol, 10 mol%) in CH_2Cl_2 (2 mL) was the added **S1** (0.2 mmol, 1.0 equiv.), the mixture was stirred overnight at room temperature. After all the **S1** disappeared monitored by TLC, $\text{Cu}(\text{OAc})_2$ (0.02 mmol, 10 mol%) was added. The mixture was stirred at room temperature under dioxygen atmosphere (dioxygen balloon, 101 kPa) for 3 h and then was purified by flash chromatography to give the desired product **1** by using solution of EtOAc in petroleum ether as eluent.

(*Z*)-Ethyl 2-((4-methoxyphenyl)imino)dec-3-ynoate (**1a**): Using 10% solution of EtOAc in petroleum ether as eluent (54.8 mg, 87% yield). ^1H NMR (500 MHz, CDCl_3) δ : 7.44~7.40 (m, 2H), 6.91~6.87 (m, 2H), 4.41 (q, $J=7.1$ Hz, 2H), 3.82 (s, 3H), 2.39 (t, $J=7.0$ Hz, 2H), 1.55~1.50 (m, 2H), 1.40 (t, $J=7.1$ Hz, 3H), 1.32~1.24 (m, 6H),

0.88~0.86 (m, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 163.27, 159.19, 141.79, 140.14, 124.45, 113.81, 102.12, 75.35, 62.81, 55.58, 31.46, 28.67, 27.86, 22.69, 19.89, 14.37, 14.22; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{26}\text{NO}_3$ [$\text{M}+\text{H}$] $^+$ 316.1907, found 316.1925.

(*Z*)-Ethyl 2-((4-methoxyphenyl)imino)dodec-3-ynoate (**1b**): Using 10% solution of EtOAc in petroleum ether as eluent (61.1 mg, 89% yield). ^1H NMR (500 MHz, CDCl_3) δ : 7.44~7.40 (m, 2H), 6.91~6.87 (m, 2H), 4.40 (q, $J=7.1$ Hz, 2H), 3.82 (s, 3H), 2.39 (t, $J=7.0$ Hz, 2H), 1.55~1.50 (m, 2H), 1.40 (t, $J=7.1$ Hz, 3H), 1.30~1.23 (m, 10H), 0.88~0.85 (m, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 163.26, 159.18, 141.77, 140.08, 124.46, 113.79, 102.07, 75.34, 62.78, 55.55, 31.99, 29.33, 29.23, 28.99, 27.88, 22.82, 19.87, 14.36, 14.26; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{30}\text{NO}_3$ [$\text{M}+\text{H}$] $^+$ 344.2220, found 344.2244.

(*Z*)-Ethyl 2-((4-methoxyphenyl)imino)oct-3-ynoate (**1c**): Using 10% solution of EtOAc in petroleum ether as eluent (49.4 mg, 86% yield). ^1H NMR (500 MHz, CDCl_3) δ : 7.44~7.38 (m, 2H), 6.91~6.87 (m, 2H), 4.40 (q, $J=7.1$ Hz, 2H), 3.81 (s, 3H), 2.39 (t, $J=7.0$ Hz, 2H), 1.51 (dt, $J=14.9$, 7.0 Hz, 2H), 1.41~1.33 (m, 5H), 0.88 (t, $J=7.3$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 163.23, 159.15, 141.77, 140.08, 124.41, 113.78, 102.01, 75.32, 62.77, 55.56, 29.87, 22.01, 19.51, 14.33, 13.67; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{22}\text{NO}_3$ [$\text{M}+\text{H}$] $^+$ 288.1594, found 288.1588.

(*Z*)-Ethyl 2-((4-methoxyphenyl)imino)-5-phenylpent-3-ynoate (**1d**): Using 10% solution of EtOAc in petroleum ether as eluent (57.1 mg, 89% yield). ^1H NMR (500 MHz, CDCl_3) δ : 7.38~7.34 (m, 2H), 7.31~7.26 (m, 1H), 7.24~7.21 (m, 1H), 7.19~7.15 (m, 2H), 6.86~6.82 (m, 2H), 4.41 (q, $J=7.1$ Hz, 2H), 3.83 (s, 3H), 2.87 (t, $J=7.3$ Hz, 2H), 2.71 (t, $J=7.3$ Hz, 2H), 1.41 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 163.27, 159.30, 141.64, 140.02, 128.70, 128.59, 126.74, 124.58, 113.87, 100.67, 75.87, 62.85, 55.63, 34.10, 21.94, 14.41; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{20}\text{NO}_3$ [$\text{M}+\text{H}$] $^+$ 322.1438, found 322.1441.

(*Z*)-Ethyl 6-(benzyloxy)-2-((4-methoxyphenyl)imino)hex-3-ynoate (**1e**): Using 10% solution of EtOAc in petroleum ether as eluent (69.2 mg, 85% yield). ^1H NMR (500 MHz, CDCl_3) δ : 7.45~7.39 (m, 2H), 7.36~7.28 (m, 5H), 6.91~6.88 (m, 2H), 4.49 (s, 2H), 4.40 (q, $J=7.1$ Hz, 2H), 3.81 (s, 3H), 3.44 (t, $J=6.5$ Hz, 2H), 2.41 (t, $J=7.0$ Hz, 2H), 1.64~1.54 (m, 5H), 1.46~1.38 (m, 5H); ^{13}C NMR (125 MHz, CDCl_3) δ : 163.26, 159.20, 141.80, 140.10, 138.73, 128.57, 127.81, 127.74, 124.45, 113.84, 101.75, 75.45, 73.13, 70.29, 62.83, 55.60, 29.42, 27.77, 25.67, 19.85, 14.39; HRMS (ESI) calcd for $\text{C}_{25}\text{H}_{30}\text{NO}_4$ [$\text{M}+\text{H}$] $^+$ 408.2169, found 408.2185.

(*Z*)-Ethyl 8-(1,3-dioxolan-2-yl)-2-((4-methoxyphenyl)imino)oct-3-ynoate (**1f**): Using 20% solution of EtOAc in petroleum ether as eluent (60.3 mg, 84% yield). ^1H NMR (500 MHz, CDCl_3) δ : 7.45~7.39 (m, 2H), 6.93~6.88 (m, 2H), 4.81 (t, $J=4.7$ Hz, 1H), 4.41 (q, $J=7.1$ Hz, 2H), 3.97~3.93 (m, 2H), 3.85~3.82 (m, 5H), 2.41 (t, $J=6.9$ Hz, 2H), 1.68~1.57 (m, 5H), 1.52~1.46 (m, 2H), 1.40 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 163.25,

159.24, 141.80, 140.04, 124.47, 113.87, 104.49, 101.53, 75.51, 65.05, 62.80, 55.60, 33.47, 27.84, 23.49, 19.85, 14.37; HRMS (ESI) calcd for $C_{20}H_{26}NO_5$ $[M+H]^+$ 360.1805, found 360.1811.

(Z)-Ethyl 4-(cyclohex-1-en-1-yl)-2-((4-methoxyphenyl)imino)but-3-ynoate (**1g**): Using 10% solution of EtOAc in petroleum ether as eluent (53.5 mg, 86% yield). 1H NMR (500 MHz, $CDCl_3$) δ : 7.50~7.45 (m, 2H), 6.92~6.87 (m, 2H), 6.30 (d, $J=1.1$ Hz, 1H), 4.40 (q, $J=7.1$ Hz, 2H), 3.81 (s, 3H), 2.13 (t, $J=5.1$ Hz, 4H), 1.64~1.55 (m, 4H), 1.39 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (125 MHz, $CDCl_3$) δ : 163.15, 159.32, 141.78, 141.12, 139.53, 124.88, 119.92, 113.73, 101.00, 81.58, 62.71, 55.58, 28.20, 26.20, 22.08, 21.30, 14.32; HRMS (ESI) calcd for $C_{19}H_{22}NO_3$ $[M+H]^+$ 312.1594, found 312.1596.

(Z)-Ethyl 2-((4-methoxyphenyl)imino)-4-(trimethylsilyl)but-3-ynoate (**1h**): Using 5% solution of EtOAc in petroleum ether as eluent (48.5 mg, 80% yield). 1H NMR (500 MHz, $CDCl_3$) δ : 7.57~7.51 (m, 2H), 6.92~6.87 (m, 2H), 4.41 (q, $J=7.1$ Hz, 2H), 3.84 (s, 3H), 1.41 (t, $J=7.1$ Hz, 3H), 0.21 (s, 9H); ^{13}C NMR (125 MHz, $CDCl_3$) δ : 162.94, 159.83, 141.30, 138.60, 125.31, 113.76, 106.89, 97.51, 62.85, 55.67, 14.34, -0.58; HRMS (ESI) calcd for $C_{16}H_{22}NO_3Si$ $[M+H]^+$ 304.1363, found 304.1378.

(Z)-Ethyl 2-((4-methoxyphenyl)imino)-4-(thiophen-3-yl)but-3-ynoate (**1i**): Using 15% solution of EtOAc in petroleum ether as eluent (52.6 mg, 84% yield). 1H NMR (500 MHz, $CDCl_3$) δ : 7.60 (dd, $J=3.0$, 1.1 Hz, 1H), 7.55~7.51 (m, 2H), 7.31 (dd, $J=5.0$, 3.0 Hz, 1H), 7.14 (dd, $J=5.0$, 1.1 Hz, 1H), 6.97~6.93 (m, 2H), 4.46 (q, $J=7.1$ Hz, 2H), 3.85 (s, 3H), 1.44 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (125 MHz, $CDCl_3$) δ : 163.08, 159.62, 141.85, 139.33, 132.54, 130.06, 126.20, 124.93, 120.49, 113.95, 93.98, 83.58, 62.95, 55.69, 14.43; HRMS (ESI) calcd for $C_{17}H_{16}NO_3S$ $[M+H]^+$ 314.0845, found 314.0844.

(Z)-Ethyl 2-((4-methoxyphenyl)imino)-4-phenylbut-3-ynoate (**1j**): Using 15% solution of EtOAc in petroleum ether as eluent (55.9 mg, 91% yield). 1H NMR (500 MHz, $CDCl_3$) δ : 7.59~7.53 (m, 2H), 7.50~7.45 (m, 2H), 7.43~7.39 (m, 1H), 7.38~7.33 (m, 2H), 6.98~6.94 (m, 2H), 4.47 (q, $J=7.1$ Hz, 2H), 3.86 (s, 3H), 1.45 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (125 MHz, $CDCl_3$) δ : 163.07, 159.66, 141.84, 139.31, 132.72, 130.35, 128.76, 125.02, 121.28, 113.96, 98.51, 83.57, 62.93, 55.70, 14.43; HRMS (ESI) calcd for $C_{19}H_{18}NO_3$ $[M+H]^+$ 308.1281, found 308.1288.

(Z)-Ethyl 2-((4-methoxyphenyl)imino)-4-(*p*-tolyl)but-3-ynoate (**1k**): Using 15% solution of EtOAc in petroleum ether as eluent (55.2 mg, 86% yield). 1H NMR (500 MHz, $CDCl_3$) δ : 7.57~7.54 (m, 2H), 7.38~7.34 (m, 2H), 7.17~7.13 (m, 2H), 6.98~6.93 (m, 2H), 4.46 (q, $J=7.1$ Hz, 2H), 3.85 (s, 3H), 2.36 (s, 3H), 1.44 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (125 MHz, $CDCl_3$) δ : 163.10, 159.53, 141.85, 140.94, 139.42, 132.64, 129.51, 124.96, 118.14, 113.89, 99.11, 83.29, 62.84, 55.63, 21.86, 14.38; HRMS (ESI) calcd for $C_{20}H_{20}NO_3$ $[M+H]^+$ 322.1438, found 322.1443.

(Z)-Ethyl 4-(4-fluorophenyl)-2-((4-methoxyphenyl)imino)but-3-ynoate (**1l**): Using 15% solution of EtOAc in petroleum ether as eluent (55.3 mg, 85% yield). 1H NMR (500 MHz, $CDCl_3$) δ : 7.55~7.50 (m, 2H), 7.49~7.43 (m, 2H), 7.08~7.04 (m, 2H), 6.98~6.94 (m, 2H), 4.46 (q, $J=7.1$ Hz, 2H), 3.86 (s, 3H), 1.45 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (125 MHz, $CDCl_3$) δ : 164.82, 163.02, 162.85, 159.67, 141.82, 139.25, 134.89 (d, $J=8.8$ Hz), 124.91, 116.28 (d, $J=22.3$ Hz), 113.97, 97.38, 83.42, 62.99, 55.71, 14.44; HRMS (ESI) calcd for $C_{19}H_{17}FNO_3$ $[M+H]^+$ 328.1187, found 328.1196.

(Z)-Methyl 2-((4-methoxyphenyl)imino)dec-3-ynoate (**1m**): Using 10% solution of EtOAc in petroleum ether as eluent (52.4 mg, 87% yield). 1H NMR (500 MHz, $CDCl_3$) δ : 7.52~7.38 (m, 2H), 6.94~6.87 (m, 2H), 3.95 (s, 3H), 3.83 (s, 3H), 2.40 (t, $J=7.0$ Hz, 2H), 1.53 (dd, $J=14.9$, 7.2 Hz, 2H), 1.37~1.27 (m, 6H), 0.88 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (125 MHz, $CDCl_3$) δ : 163.79, 159.37, 141.63, 139.55, 124.61, 113.88, 102.28, 75.34, 55.62, 53.65, 31.47, 28.72, 27.88, 22.71, 19.93, 14.23; HRMS (ESI) calcd for $C_{18}H_{24}NO_3$ $[M+H]^+$ 302.1751, found 302.1755.

(Z)-Propyl 2-((4-methoxyphenyl)imino)dec-3-ynoate (**1n**): Using 10% solution of EtOAc in petroleum ether as eluent (55.9 mg, 85% yield). 1H NMR (500 MHz, $CDCl_3$) δ : 7.46~7.40 (m, 2H), 6.92~6.88 (m, 2H), 4.30 (t, $J=6.8$ Hz, 2H), 3.83 (s, 3H), 2.39 (t, $J=7.0$ Hz, 2H), 1.81 (dt, $J=14.4$, 7.2 Hz, 2H), 1.56~1.50 (m, 2H), 1.35~1.24 (m, 6H), 1.01 (t, $J=7.4$ Hz, 3H), 0.87 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (125 MHz, $CDCl_3$) δ : 163.28, 159.19, 141.86, 140.10, 124.48, 113.81, 101.96, 75.41, 68.20, 55.58, 31.47, 28.69, 27.89, 22.69, 22.12, 19.87, 14.22, 10.50; HRMS (ESI) calcd for $C_{22}H_{28}NO_3$ $[M+H]^+$ 330.2064, found 330.2081.

4.3 General procedure for the enantioselective transfer hydrogenation of 1

A mixture of **1** (0.1 mmol, 1.0 equiv.), **3e** (0.01 mmol, 5 mol%), 3 Å MS (40 mg), and **4d** (0.12 mmol, 1.2 equiv.) in the mixture of decane (1.0 mL) and mesitylene (1.0 mL) was stirred at room temperature under N_2 for 4 h, and then directly purified by flash chromatography to give the desired product **2** by using solution of EtOAc in petroleum ether as eluent.

(S)-Ethyl 2-((4-methoxyphenyl)amino)dec-3-ynoate (**2a**): Using 5% solution of EtOAc in petroleum ether as eluent (28.4 mg, 89% yield). $[\alpha]_D^{25} + 36.71$ (c 1.28, $CHCl_3$); 1H NMR (500 MHz, $CDCl_3$) δ : 6.81~6.76 (m, 2H), 6.69~6.65 (m, 2H), 4.70 (t, $J=2.1$ Hz, 1H), 4.26 (q, $J=7.1$ Hz, 2H), 3.74 (s, 3H), 2.17 (td, $J=7.0$, 2.1 Hz, 2H), 1.49~1.43 (m, 2H), 1.33~1.23 (m, 8H), 0.87 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (125 MHz, $CDCl_3$) δ : 169.68, 153.42, 139.77, 116.12, 114.87, 85.59, 75.25, 62.29, 55.80, 50.34, 31.46, 28.57, 28.49, 22.71, 18.87, 14.24, 14.21; HRMS (ESI) calcd for $C_{19}H_{28}NO_3$ $[M+H]^+$ 318.2064, found 318.2072. HPLC: the *ee* value was determined by HPLC analysis (Chiralcel AD-H, *i*-PrOH/hexane, $V:V=25/75$, 1.0 mL/min, 249 nm), retention time: $t_{minor}=6.477$ min, $t_{major}=9.070$ min, *ee*=90%.

(*S*)-Ethyl 2-((4-methoxyphenyl)amino)dodec-3-ynoate (**2b**): Using 5% solution of EtOAc in petroleum ether as eluent (29.7 mg, 86% yield). $[\alpha]_{\text{D}}^{25} + 32.51$ (*c* 1.18, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ : 6.81~6.75 (m, 2H), 6.69~6.65 (m, 2H), 4.70 (t, *J*=2.2 Hz, 1H), 4.26 (q, *J*=7.1 Hz, 2H), 3.74 (s, 3H), 2.17 (td, *J*=7.1, 2.2 Hz, 2H), 1.49~1.43 (m, 2H), 1.33~1.23 (m, 13H), 0.88 (t, *J*=7.0 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ : 169.75, 153.43, 139.87, 116.11, 114.92, 85.61, 75.30, 62.31, 55.85, 50.37, 32.04, 29.39, 29.27, 28.95, 28.57, 22.86, 18.91, 14.30, 14.28; HRMS (ESI) calcd for C₂₁H₃₂NO₃ [M+H]⁺ 346.2377, found 346.2388. HPLC: the *ee* value was determined by HPLC analysis (Chiralcel AD-H, *i*-PrOH/hexane, *V*: *V*=25/75, 1.0 mL/min, 256 nm), retention time: *t*_{minor}=5.617 min, *t*_{major}=7.633 min, *ee*=98%.

(*S*)-Ethyl 2-((4-methoxyphenyl)amino)oct-3-ynoate (**2c**): Using 5% solution of EtOAc in petroleum ether as eluent (24.5 mg, 85% yield). $[\alpha]_{\text{D}}^{25} + 49.62$ (*c* 1.12, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ : 6.81~6.76 (m, 2H), 6.70~6.65 (m, 2H), 4.70 (t, *J*=2.2 Hz, 1H), 4.26 (qd, *J*=7.1, 1.2 Hz, 2H), 3.74 (s, 3H), 2.18 (td, *J*=7.0, 2.2 Hz, 2H), 1.49~1.41 (m, 2H), 1.39~1.32 (m, 2H), 1.29 (t, *J*=7.1 Hz, 3H), 0.87 (t, *J*=7.3 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ : 169.70, 153.42, 139.79, 116.14, 114.88, 85.55, 75.25, 62.29, 55.82, 50.37, 30.58, 21.97, 18.55, 14.23, 13.71; HRMS (ESI) calcd for C₁₇H₂₄NO₃ [M+H]⁺ 290.1751, found 290.1764. HPLC: the *ee* value was determined by HPLC analysis (Chiralcel AD-H, *i*-PrOH/hexane, *V*: *V*=25/75, 1.0 mL/min, 241 nm), retention time: *t*_{minor}=7.163 min, *t*_{major}=10.507 min, *ee*=94%.

(*S*)-Ethyl 2-((4-methoxyphenyl)amino)-6-phenylhex-3-ynoate (**2d**): Using 5% solution of EtOAc in petroleum ether as eluent (29.3 mg, 87% yield). $[\alpha]_{\text{D}}^{25} + 36.7$ (*c* 1.01, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ : 7.39~7.32 (m, 2H), 7.31~7.26 (m, 2H), 7.25~7.20 (m, 1H), 7.20~7.14 (m, 2H), 6.87~6.82 (m, 2H), 4.41 (q, *J*=7.1 Hz, 2H), 3.82 (s, 3H), 2.87 (t, *J*=7.3 Hz, 2H), 2.71 (t, *J*=7.3 Hz, 2H), 1.41 (t, *J*=7.1 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ : 163.26, 159.29, 141.61, 140.01, 139.68, 128.69, 128.59, 126.73, 124.58, 113.86, 100.67, 75.85, 62.85, 55.62, 34.09, 21.93, 14.40; HRMS (ESI) calcd for C₂₀H₂₂NO₃ [M+H]⁺ 338.1751, found 338.1767. HPLC: the *ee* value was determined by HPLC analysis (Chiralcel AD-H, *i*-PrOH/hexane, *V*: *V*=25/75, 1.0 mL/min, 308 nm), retention time: *t*_{minor}=11.233 min, *t*_{major}=15.010 min, *ee*=95%.

(*S*)-Ethyl 9-(benzyloxy)-2-((4-methoxyphenyl)amino)-non-3-ynoate (**2e**): Using 5% solution of EtOAc in petroleum ether as eluent (33.6 mg, 82% yield). $[\alpha]_{\text{D}}^{25} + 32.71$ (*c* 0.61, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ : 7.28~7.19 (m, 5H), 6.73~6.68 (m, 2H), 6.61~6.56 (m, 2H), 4.62 (t, *J*=2.1 Hz, 1H), 4.42 (s, 2H), 4.18 (q, *J*=7.1 Hz, 2H), 3.66 (s, 3H), 3.36 (t, *J*=6.6 Hz, 2H), 2.11 (td, *J*=7.0, 2.2 Hz, 2H), 1.55~1.49 (m, 2H), 1.45~1.39 (m, 2H), 1.37~1.30 (m, 2H), 1.21 (t, *J*=7.1 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ : 169.69, 153.43, 139.84, 138.84, 128.56, 127.81, 127.71, 116.11, 114.91, 85.33, 75.47, 73.10, 70.44, 62.32, 55.84, 50.35, 29.44, 28.39, 25.57,

18.87, 14.27; HRMS (ESI) calcd for C₂₅H₃₂NO₄ [M+H]⁺ 410.2326, found 410.2329. HPLC: the *ee* value was determined by HPLC analysis (Chiralcel AD-H, *i*-PrOH/hexane, *V*: *V*=25/75, 1.0 mL/min, 205 nm), retention time: *t*_{minor}=12.743 min, *t*_{major}=18.110 min, *ee*=93%.

(*S*)-Ethyl 8-(1,3-dioxolan-2-yl)-2-((4-methoxyphenyl)amino)oct-3-ynoate (**2f**): Using 20% solution of EtOAc in petroleum ether as eluent (28.9 mg, 80% yield). $[\alpha]_{\text{D}}^{25} + 49.71$ (*c* 1.6, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ : 6.80~6.76 (m, 2H), 6.69~6.65 (m, 2H), 4.81 (t, *J*=4.8 Hz, 1H), 4.69 (s, 1H), 4.26 (q, *J*=7.1 Hz, 2H), 4.18 (s, 1H), 3.97~3.92 (m, 2H), 3.87~3.81 (m, 2H), 3.74 (s, 3H), 2.19 (td, *J*=6.9, 2.1 Hz, 2H), 1.65~1.61 (m, 2H), 1.55~1.45 (m, 4H), 1.30 (t, *J*=7.1 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ : 169.69, 153.42, 139.86, 116.11, 114.93, 104.64, 85.18, 75.55, 65.05, 62.34, 55.85, 50.36, 33.51, 28.45, 23.43, 18.86, 14.27; HRMS (ESI) calcd for C₂₀H₂₈NO₅ [M+H]⁺ 362.1962, found 362.1981; HPLC: the *ee* value was determined by HPLC analysis (Chiralcel AD-H, *i*-PrOH/hexane, *V*: *V*=25/75, 1.0 mL/min, 241 nm), retention time: *t*_{minor}=14.260 min, *t*_{major}=20.310 min, *ee*=92%.

(*S*)-Ethyl 4-(cyclohex-1-en-1-yl)-2-((4-methoxyphenyl)amino)but-3-ynoate (**2g**): Using 5% solution of EtOAc in petroleum ether as eluent (28.5 mg, 91% yield). $[\alpha]_{\text{D}}^{25} + 98.74$ (*c* 0.66, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ : 6.81~6.77 (m, 2H), 6.71~6.65 (m, 2H), 6.10 (dt, *J*=3.7, 2.0 Hz, 1H), 4.82 (s, 1H), 4.27 (q, *J*=7.1 Hz, 2H), 3.75 (s, 3H), 2.10~2.00 (m, 4H), 1.63~1.53 (m, 4H), 1.30 (t, *J*=7.1 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ : 169.46, 153.44, 139.80, 136.32, 120.00, 116.14, 114.92, 86.39, 81.51, 62.39, 55.85, 50.76, 29.09, 25.79, 22.36, 21.58, 14.27; HRMS (ESI) calcd for C₁₉H₂₄NO₃ [M+H]⁺ 314.1751, found 314.1761; HPLC: the *ee* value was determined by HPLC analysis (Chiralcel AD-H, *i*-PrOH/hexane, *V*: *V*=25/75, 1.0 mL/min, 248 nm), retention time: *t*_{minor}=8.183 min, *t*_{major}=10.597 min, *ee*=95%.

(*S*)-Ethyl 2-((4-methoxyphenyl)amino)-4-(trimethylsilyl)but-3-ynoate (**2h**): Using 3% solution of EtOAc in petroleum ether as eluent (25.0 mg, 82% yield). $[\alpha]_{\text{D}}^{25} + 110.62$ (*c* 1.16, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ : 6.80~6.77 (m, 2H), 6.70~6.66 (m, 2H), 4.72 (s, 1H), 4.33~4.24 (m, 2H), 4.19 (s, 1H), 3.75 (s, 3H), 1.30 (t, *J*=7.1 Hz, 3H), 0.14 (s, 9H); ¹³C NMR (125 MHz, CDCl₃) δ : 169.08, 153.57, 139.72, 116.26, 114.91, 100.19, 90.00, 62.47, 55.87, 51.17, 14.21, -0.09; HRMS (ESI) calcd for C₁₆H₂₄NO₃Si [M+H]⁺ 306.1520, found 306.1511; HPLC: the *ee* value was determined by HPLC analysis (Chiralcel AD-H, *i*-PrOH/hexane, *V*: *V*=25/75, 1.0 mL/min, 248 nm), retention time: *t*_{minor}=4.683 min, *t*_{major}=5.693 min, *ee*=91%.

(*S*)-Ethyl 2-((4-methoxyphenyl)amino)-4-(thiophen-3-yl)but-3-ynoate (**2i**): Using 10% solution of EtOAc in petroleum ether as eluent (28.3 mg, 90% yield). $[\alpha]_{\text{D}}^{25} + 108.71$ (*c* 0.76, CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ : 7.44 (dd, *J*=3.0, 1.1 Hz, 1H), 7.23 (dd, *J*=5.0, 3.0 Hz, 1H), 7.07 (dd, *J*=5.0, 1.1 Hz, 1H), 6.83~6.79 (m, 2H),

6.75~6.71 (m, 2H), 4.93 (s, 1H), 4.36~4.20 (m, 3H), 3.75 (s, 3H), 1.32 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 169.13, 153.59, 139.66, 130.15, 129.81, 125.48, 121.38, 116.22, 115.00, 84.04, 79.75, 62.61, 55.86, 50.85, 14.29; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{18}\text{NO}_3\text{S}$ $[\text{M} + \text{H}]^+$ 316.1002, found 316.1020. HPLC: the *ee* value was determined by HPLC analysis (Chiralcel AD-H, *i*-PrOH/hexane, $V: V=25/75$, 1.0 mL/min, 246 nm), retention time: $t_{\text{minor}}=15.287$ min, $t_{\text{major}}=21.230$ min, *ee*=80%.

(*S*)-Ethyl 2-((4-methoxyphenyl)amino)-4-phenylbut-3-ynoate (**2j**): Using 15% solution of EtOAc in petroleum ether as eluent (27.2 mg, 88% yield). $[\alpha]_{\text{D}}^{25} + 107.82$ (*c* 1.22, CHCl_3); ^1H NMR (500 MHz, CDCl_3) δ : 7.47~7.40 (m, 2H), 7.35~7.28 (m, 3H), 6.86~6.81 (m, 2H), 6.79~6.74 (m, 2H), 4.98 (s, 1H), 4.33 (q, $J=7.1$ Hz, 2H), 3.77 (s, 3H), 1.35 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 169.14, 153.54, 139.61, 132.07, 128.83, 128.41, 122.32, 116.22, 114.95, 84.50, 84.36, 62.57, 55.81, 50.80, 14.26; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{20}\text{NO}_3$ $[\text{M} + \text{H}]^+$ 310.1438, found 310.1442. HPLC: the *ee* value was determined by HPLC analysis (Chiralcel AD-H, *i*-PrOH/hexane, $V: V=25/75$, 1.0 mL/min, 306 nm), retention time: $t_{\text{minor}}=11.110$ min, $t_{\text{major}}=13.573$ min, *ee*=89%.

(*S*)-Ethyl 2-((4-methoxyphenyl)amino)-4-(*p*-tolyl)but-3-ynoate (**2k**): Using 10% solution of EtOAc in petroleum ether as eluent (27.1 mg, 84% yield). $[\alpha]_{\text{D}}^{25} + 104.8$ (*c* 0.9, CHCl_3); ^1H NMR (500 MHz, CDCl_3) δ : 7.32 (d, $J=8.0$ Hz, 2H), 7.10 (d, $J=7.9$ Hz, 2H), 6.87~6.80 (m, 2H), 6.80~6.73 (m, 2H), 4.97 (s, 1H), 4.32 (q, $J=7.1$ Hz, 2H), 3.75 (s, 3H), 2.34 (s, 3H), 1.33 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 169.19, 153.44, 139.64, 138.93, 131.90, 129.12, 119.19, 116.13, 114.88, 84.61, 83.63, 62.45, 55.73, 50.76, 21.59, 14.20; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{22}\text{NO}_3$ $[\text{M} + \text{H}]^+$ 324.1594, found 324.1599. HPLC: the *ee* value was determined by HPLC analysis (Chiralcel IB, *i*-PrOH/hexane, $V: V=20/80$, 1.0 mL/min, 255 nm), retention time: $t_{\text{minor}}=7.047$ min, $t_{\text{major}}=8.553$ min, *ee*=86%.

(*S*)-Ethyl 4-(4-fluorophenyl)-2-((4-methoxyphenyl)amino)but-3-ynoate (**2l**): Using 15% solution of EtOAc in petroleum ether as eluent (28.4 mg, 87% yield). $[\alpha]_{\text{D}}^{25} + 96.85$ (*c* 0.96, CHCl_3); ^1H NMR (500 MHz, CDCl_3) δ : 7.41~7.32 (m, 2H), 7.01~6.96 (m, 2H), 6.83~6.78 (m, 2H), 6.77~6.72 (m, 2H), 4.94 (s, 1H), 4.31 (q, $J=7.1$ Hz, 2H), 3.75 (s, 3H), 1.33 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 169.08, 162.91 (d, $J=250.1$ Hz), 153.63, 139.56, 134.09, 134.02, 118.43, 116.27, 115.85, 115.67, 115.00, 84.13, 83.50, 62.65, 55.86, 50.80, 14.30; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{19}\text{FNO}_3$ $[\text{M} + \text{H}]^+$ 328.1343, found 328.1359. HPLC: the *ee* value was determined by HPLC analysis (Chiralcel AD-H, *i*-PrOH/hexane, $V: V=25/75$, 1.0 mL/min, 220 nm), retention time: $t_{\text{minor}}=11.970$ min, $t_{\text{major}}=14.250$ min, *ee*=88%.

(*S*)-Methyl 2-((4-methoxyphenyl)amino)dec-3-ynoate (**2m**): Using 5% solution of EtOAc in petroleum ether as eluent (27.2 mg, 90% yield). $[\alpha]_{\text{D}}^{25} + 45.78$ (*c* 1.08, CHCl_3); ^1H NMR (500 MHz, CDCl_3) δ : 6.81~6.77 (m,

2H), 6.69~6.65 (m, 2H), 4.73 (t, $J=2.1$ Hz, 1H), 4.17 (s, 1H), 3.81 (s, 3H), 3.74 (s, 3H), 2.17 (td, $J=7.1$, 2.1 Hz, 2H), 1.50~1.43 (m, 2H), 1.34~1.23 (m, 6H), 0.88 (t, $J=7.0$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 170.22, 153.47, 139.77, 116.10, 114.92, 85.77, 75.13, 55.81, 53.21, 50.21, 31.45, 28.59, 28.48, 22.70, 18.88, 14.20; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{26}\text{NO}_3$ $[\text{M} + \text{H}]^+$ 304.1907, found 304.1914. HPLC: the *ee* value was determined by HPLC analysis (Chiralcel AD-H, *i*-PrOH/hexane, $V: V=25/75$, 1.0 mL/min, 206 nm), retention time: $t_{\text{minor}}=10.213$ min, $t_{\text{major}}=14.787$ min, *ee*=93%.

(*S*)-Propyl 2-((4-methoxyphenyl)amino)dec-3-ynoate (**2n**): Using 5% solution of EtOAc in petroleum ether as eluent (27.5 mg, 83% yield). $[\alpha]_{\text{D}}^{25} + 21.88$ (*c* 1.1, CHCl_3); ^1H NMR (500 MHz, CDCl_3) δ : 6.80~6.75 (m, 2H), 6.69~6.64 (m, 2H), 4.71 (t, $J=2.2$ Hz, 1H), 4.20~4.13 (m, 2H), 3.73 (s, 3H), 2.17 (td, $J=7.1$, 2.2 Hz, 2H), 1.72~1.65 (m, 2H), 1.49~1.43 (m, 2H), 1.34~1.22 (m, 6H), 0.94 (t, $J=7.4$ Hz, 3H), 0.87 (t, $J=7.1$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 169.77, 153.34, 139.84, 115.98, 114.84, 85.42, 75.34, 67.71, 55.74, 50.24, 31.44, 28.55, 28.47, 22.66, 22.06, 18.82, 14.17, 10.36; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{30}\text{NO}_3$ $[\text{M} + \text{H}]^+$ 332.2220, found 332.2218. HPLC: the *ee* value was determined by HPLC analysis (Chiralcel AD-H, *i*-PrOH/hexane, $V: V=25:75$, 1.0 mL/min, 255 nm), retention time: $t_{\text{minor}}=7.093$ min, $t_{\text{major}}=11.487$ min, *ee*=92%.

Supporting Information ^1H NMR and ^{13}C NMR spectra and HPLC spectra of **1a**~**1n** and **2a**~**2n**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- (a) Afagh, N. A.; Yudin, A. K. *Angew. Chem., Int. Ed.* **2010**, *49*, 262.
(b) Mahatthananchai, J.; Dumas, A.; Bode, J. W. *Angew. Chem., Int. Ed.* **2012**, *51*, 10954.
(c) Szostak, M.; Spain, M.; Procter, D. J. *Chem. Soc. Rev.* **2013**, *42*, 9155.
- (a) Hoffmann, R. W. *Synthesis* **2006**, 3531.
(b) Young, I. S.; Baran, P. S. *Nat. Chem.* **2009**, *1*, 193.
- Trost, B. M. *Science* **1991**, *254*, 1471.
- Wender, P. A.; Verma, V. A.; Paxton, T. J.; Pillow, T. H. *Acc. Chem. Res.* **2008**, *41*, 40.
- (a) Barrett, G. C. *Chemistry and Biochemistry of the Amino Acids*, Chapman & Hall, London, **1985**.
(b) Williams, R. M. *Synthesis of Optically Active α -Amino Acids*, Pergamon, Oxford, **1989**.
(c) Duthaler, R. O. *Tetrahedron* **1994**, *50*, 1539.
(d) Hegedus, L. S. *Acc. Chem. Res.* **1995**, *28*, 299.
(e) Chin, J. W.; Cropp, T. A.; Anderson, J. C.; Mukherji, M.; Zhang, Z.; Schultz, P. G. *Science* **2002**, *301*, 964.
(f) Wang, L.; Schultz, P. G. *Angew. Chem., Int. Ed.* **2004**, *44*, 34.
(g) Cropp, T. A.; Schultz, P. G. *Trends Genet.* **2004**, *20*, 625.
- (a) Maruoka, K.; Ooi, T. *Chem. Rev.* **2003**, *103*, 3013.
(b) Taggi, A. E.; Hafez, A. M.; Lectka, T. *Acc. Chem. Res.* **2003**, *36*, 10.
(c) Lygo, B.; Angrews, B. I. *Acc. Chem. Res.* **2004**, *37*, 518.
(d) Kobayashi, S.; Mori, Y.; Fossey, J. S.; Salter, M. M. *Chem. Rev.*

- 2011, 111, 2626.
- [7] (a) Abdulganeeva, S. A.; Erzhanov, K. B. *Russ. Chem. Rev.* **1991**, 60, 676.
(b) Williams, R. M.; Aldous, D. J.; Aldous, S. C. *J. Org. Chem.* **1990**, 55, 4657.
(c) Castelano, A. L.; Horne, S.; Taylor, G. J.; Billedeau, R.; Krantz, A. *Tetrahedron* **1988**, 44, 5451.
- [8] (a) Ji, J.-X.; Wu, J.; Chan, A. S. C. *Proc. Natl. Acad. Sci. U. S. A.* **2005**, 102, 11196.
(b) Rueping, M.; Antonchick, A. P.; Brinkmann, C. *Angew. Chem., Int. Ed.* **2007**, 46, 6903.
- [9] (a) Kang, Q.; Zhao, Z.-A.; You, S.-L. *Org. Lett.* **2008**, 10, 2031.
(b) Chen, M.-W.; Wu, B.; Chen, Z.-P.; Shi, L.; Zhou, Y.-G. *Org. Lett.* **2016**, 18, 4650.
- [10] (a) Ge, X.; Chen, X.; Qian, C. *Chin. J. Org. Chem.* **2016**, 36, 1208 (in Chinese).
(葛新, 陈新志, 钱超, 有机化学, **2016**, 36, 1208.)
(b) Hu, S.-B.; Chen, M.-W.; Zhai, X.-Y.; Zhou, Y.-G. *Acta Chim. Sinica* **2018**, 76, 103 (in Chinese).
(胡书博, 陈木旺, 翟小勇, 周永贵, 化学学报, **2018**, 76, 103.)
(c) Liu, Y.; Dong, X.-Q.; Zhang, X. *Chin. J. Org. Chem.* **2020**, 40, 1096 (in Chinese).
(刘元华, 董秀琴, 张绪穆, 有机化学, **2020**, 40, 1096.)
- (d) Dou, X.; Liu, D.; Zhang, W. *Chin. J. Pharm.* **2020**, 51, 551 (in Chinese).
(窦骁勇, 刘德龙, 张万斌, 中国医药工业杂志, **2020**, 51, 551.)
- [11] Zhang, L.; Zhu, R.; Feng, A.; Zhao, C.; Chen, L.; Feng, G.; Liu, L. *Chem. Sci.* **2020**, 11, 4444.
- [12] (a) Zhu, C.; Saito, K.; Yamanaka, M.; Akiyama, T. *Acc. Chem. Res.* **2015**, 48, 388.
(b) Zhu, C.; Falck, J. R. *ChemCatChem* **2011**, 3, 1850.
(c) Zhu, C.; Akiyama, T. *Org. Lett.* **2009**, 11, 4180.
(d) Henseler, A.; Kato, M.; Mori, K.; Akiyama, T. *Angew. Chem., Int. Ed.* **2011**, 50, 8180.
(e) Horiguchi, K.; Yamamoto, E.; Saito, K.; Yamanaka, M.; Akiyama, T. *Chem.-Eur. J.* **2016**, 22, 8078.
(f) Kim, K.-H.; Akiyama, T.; Cheon, C.-H. *Chem.-Asian J.* **2016**, 11, 274.
(g) Wen, W.; Zeng, Y.; Peng, L.-Y.; Fu, L.-N.; Guo, Q.-X. *Org. Lett.* **2015**, 17, 3922.
- [13] Hoffmann, S.; Seayad, A. M.; List, B. *Angew. Chem., Int. Ed.* **2011**, 50, 8180.
- [14] Chen, X. M.; Li, X. S.; Chan, A. S. C. *Chin. Chem. Lett.* **2009**, 20, 407.
- [15] Ji, J.-X.; Au-Yeung, T. T.-L.; Wu, J.; Yip, C. W.; Chan, A. S. C. *Adv. Synth. Catal.* **2004**, 346, 42.

(Zhao, C.)