一种基于罗丹明 B 酰肼的具有高灵敏度和选择性的 “Off-On”型 Cu^{2+} 荧光探针

雷萌萌^{a,b} 周起航^{a,b} 杨莉^{a,b} 许志红^{*,a,b} 杨凤岭^{*,a}

(° 郑州大学化学学院 郑州 450001)

(° 许昌学院化工与材料学院 河南许昌 461000)

摘要 以 5-甲酰基-3,4-二甲基吡咯-2-羧酸乙酯和罗丹明 B 酰肼为原料,合成了一种“Off-On”型罗丹明衍生物 Cu^{2+} 荧光探针 **R1**。探针 **R1** 对 Cu^{2+} 识别具有高选择性和高灵敏度,其检出限(LOD)为 $0.201 \mu\text{mol}\cdot\text{L}^{-1}$,此外探针 **R1** 与 Cu^{2+} 作用后颜色从无色变为粉红色,可以实现 Cu^{2+} 的“裸眼检测”。探针 **R1** 成功用于实际水样中 Cu^{2+} 的检测。

关键词 荧光探针; 罗丹明; 吡咯; Cu^{2+}

A Highly Sensitive and Selective “Off-On” Fluorescent Probe for Cu^{2+} Based on Rhodamine B Hydrazone

Lei, Mengmeng^{a,b} Zhou, Qihang^{a,b} Yang, Li^{a,b} Xu, Zhihong^{*,a,b} Yang, Fengling^{*,a}

(° College of Chemistry, Zhengzhou University, Zhengzhou 450001)

(° College of Chemical and Materials Engineering, Xuchang University, Xuchang, Henan 461000)

Abstract Ethyl 5-formyl-3,4-dimethyl-pyrrole-2-carboxylate and rhodamine B hydrazide were used to synthesize a rhodamine derivative **R1**. Notably, the as-synthesized **R1** displayed great sensitivity and selectivity, which was utilized to be an off-on fluorescence probe for Cu^{2+} detection in mediums that contained water. Besides, the Cu^{2+} ions limit of detection (LOD) was $0.201 \mu\text{mol}\cdot\text{L}^{-1}$, and **R1** was able to sense Cu^{2+} by “naked-eye” according to the change in color to pink from colorless. In addition, Cu^{2+} detection was assessed within the actual water samples for testing the probe actual application.

Keywords fluorescent probe; rhodamine; pyrrole; Cu^{2+}

1 Introduction

More and more attention has been paid to chemosensor which can be applied to detect heavy metal ions in trace amounts,^[1-2] since the heavy metal ions play a crucial role during numerous environmental and biological processes.^[3-6] In comparison with electrical approaches, the fluorescence chemosensors are advantageous in terms of their chemical analyses in a real-time manner, excellent selectivity and great sensitivity.^[7-8] Particularly, copper (Cu) plays a vital part in numerous biological processes, like gene expression, enzymatic function, and redox process, thus, it is an important trace element within human

body.^[9-11] Nonetheless, the excessively great copper content will induce oxidative stress (OS) as well as neurodegenerative disorders, including Parkinson’s disease (PD), Wilson’s disease and Alzheimer’s disease (AD).^[12-14] Therefore, it is of crucial importance to detect Cu^{2+} ion contents within the environment together with numerous other scientific fields, for the sake of protecting human health.

Rhodamine, together with its derivatives, serves as the typical fluorescent dye, and it is highly applicable as the fluorescent chromophore, which is ascribed to the excellent photophysical performances, like great visible-light emission wavelengths, high quantum yield, and great extinction coefficients. Great efforts have been made to develop the

* Corresponding authors. E-mail: xuzhihong1980@yahoo.com; yangfengling2005@yahoo.com.cn

Received May 29, 2020; revised July 2, 2020; published online July 8, 2020.

Project supported by the Joint Program for Fostering Talents of National Natural Science Foundation of China and Henan Province (No. U1304202), the Program for Science and Technology Innovation Talents in Universities of Henan Province (No. 15HASTIT002), the Science and Technology Project of Henan Province (No. 192102310295), the Key Program in Universities of Henan Province (No. 19A150046), and the Key Program of Xuchang City (No. 2019-JC-02).

国家自然科学基金-河南联合基金(No. U1304202)、河南省高校科技创新人才计划(No. 15HASTIT002)、河南省科技厅(No. 192102310295)、河南省高校重点研究计划(No. 19A150046)和许昌市重点(No. 2019-JC-02)资助项目。

rhodamine skeleton-containing selective Cu^{2+} fluorescence probes.^[15-19] Among them, a large amount of fluorescent and colorimetric sensors of rhodamine hydrazones towards Cu^{2+} ions have been developed using substituted salicylaldehyde,^[20] cinnamaldehyde,^[21] 1,8-naphthyridine and quinoline-2-aldehyde,^[22] coumarin-3-aldehyde,^[23] indole-3-carboxaldehyde,^[24] furfural,^[25] formaldehyde.^[26] In addition, the pyrrole derivatives have strong physiological activity and good stability.^[27-28] Therefore, Rhodamine B hydrazone and ethyl 5-formyl-3,4-dimethyl-pyrrole-2-carboxylate were used to synthesize ethyl 5-formyl-3,4-dimethyl-pyrrole-2-carboxylate rhodamine B hydrazone (**R1**) which could exhibit better performance.

On the other hand, it has been demonstrated that Schiff bases bearing multi-substituted pyrrole unit can form the stable complexes with Cu^{2+} ions.^[29-30] As a matter of fact, according to our earlier findings, ethyl 5-formyl-3,4-dimethyl-pyrrole-2-carboxylate-derived rhodamine B hydrazone might be used to be the Cu^{2+} selective sensor in dual-mode via ratiometric displacement as well as rhodamine ring opening approach.^[31] To continue our previous research, ethyl 5-formyl-3,4-dimethyl-pyrrole-2-carboxylate rhodamine B hydrazone (**R1**), together with the possible performances as the fluorescent and colorimetric sensor for Cu^{2+} detection, was reported in this study (Eq. 1). The results suggested that the probe **R1** exhibited its function by forming the complex $[\text{R1-H} + \text{Cu} + \text{NO}_3]$. Firstly, the probe **R1** followed by hydrolyzing to a fluorescent rhodamine B in water-containing mediums, which is quite different from the coordination mechanism in our previous work.^[31] In addition, the application of the probe in detecting Cu^{2+} in real water samples has also been demonstrated.

2 Results and discussion

2.1 R1 selectivity towards Cu^{2+} as well as additional metal cations

The probe **R1** selectivity to a variety of metal cations was examined. According to Figure 1, the sensor **R1** displayed nearly no absorption at the wavelength of 450~650 nm in the 10 mmol·L⁻¹ CH₃CN/HEPES (pH=7.40, $V/V=5/5$) buffer solution, which suggested that **R1** existed as the

spirocycle-closed form, and such observation is in good accordance with previous report.^[32] When Cu^{2+} was added into 2.5 $\mu\text{mol}\cdot\text{L}^{-1}$ **R1** buffer solution, the mixed solution showed marked absorbance within UV-vis spectra at the wavelength of 560 nm (Figure 1), and fluorescence emission of 585 nm (Figure 2, $\lambda_{\text{ex}}=350$ nm). This might be ascribed to the spirolactam ring opening induced by Cu^{2+} .^[33-34] Nonetheless, for additional metal cations, including Zn^{2+} , Pb^{2+} , Na^+ , Mn^{2+} , Mg^{2+} , K^+ , Hg^{2+} , Fe^{3+} , Cr^{3+} , Co^{2+} , Cd^{2+} , Ca^{2+} , Al^{3+} and Ag^+ , except for Ni^{2+} (absorbance intensity at 560 nm was half of that addition of Cu^{2+}), neither fluorescence emission of 585 nm nor absorbance enhancement at 560 nm was detected. It should be noted that Ni^{2+} might form a complex with **R1** in spirolactam ring opening form, which is responsible for the UV-vis spectral change, while is non-emissive due to the paramagnetism effect of Ni^{2+} .^[35] However, in the case of Cu^{2+} , the intermediate complex between Cu^{2+} and **R1** is not stable enough, which occurred a hydrolysis process under the tested condition to produce highly emissive rhodamine B (side infra), according to the fluorescence enhancement.

Figure 1 UV-vis spectra of 2.5 $\mu\text{mol}\cdot\text{L}^{-1}$ sensor **R1** in buffered CH₃CN/HEPES ($V:V=5:5$) solution at pH 7.40 with 2 equiv. of metal ions

Metal ions: Ag^+ , Al^{3+} , Ca^{2+} , Cd^{2+} , Co^{2+} , Cr^{3+} , Cu^{2+} , Fe^{3+} , Hg^{2+} , K^+ , Mg^{2+} , Mn^{2+} , Na^+ , Ni^{2+} , Pb^{2+} , Zn^{2+} and blank. The inset shows the change of the color with the addition of $\text{Cu}^{2+}/\text{Ni}^{2+}$ to **R1** in buffered solution

Similarly, when Cu^{2+} was added, the 10 mmol·L⁻¹ CH₃CN/HEPES (pH=7.40, $V/V=5/5$) buffer solution of 2.5 $\mu\text{mol}\cdot\text{L}^{-1}$ **R1** displayed a marked pink color (Figure 1, inset), besides, the red fluorescence was detected under the 365 nm UV lamp (Figure 2, inset), and the remaining cations (except for Ni^{2+}) displayed almost no interference. As expected, **R1** was a “naked-eye” chemosensor that showed selectivity to Cu^{2+} ions in the neutral buffer solution.

2.2 Absorption and fluorescence spectrum titrations of R1 with Cu^{2+}

For obtaining the details regarding the **R1** sensor and Cu^{2+} binding manner, fluorescence and absorption spec-

Figure 2 Fluorescence emission spectra of $2.5 \mu\text{mol}\cdot\text{L}^{-1}$ sensor **R1** in buffered $\text{CH}_3\text{CN}/\text{HEPES}$ ($V/V=5/5$) solution at pH 7.40 with 2 equiv. of metal ions

Metal ions: Ag^+ , Al^{3+} , Ca^{2+} , Cd^{2+} , Co^{2+} , Cr^{3+} , Cu^{2+} , Fe^{3+} , Hg^{2+} , K^+ , Mg^{2+} , Mn^{2+} , Na^+ , Ni^{2+} , Pb^{2+} and Zn^{2+} and blank. The inset shows the fluorescence color of **R1** with Cu^{2+}

trum titration experiments were conducted respectively. As shown in Figure 3, upon the gradual addition of Cu^{2+} to the $\text{CH}_3\text{CN}/\text{HEPES}$ ($10 \text{ mmol}\cdot\text{L}^{-1}$, $\text{pH}=7.40$, $V/V=5/5$) solution of sensor **R1** ($2.5 \mu\text{mol}\cdot\text{L}^{-1}$), the absorbance peaked at 560 nm markedly increased, which obviously indicated spirolactam ring opening induced by Cu^{2+} addition. At the same time, fluorescence intensity at the wavelength of 585 nm showed gradual increasing when Cu^{2+} was added (Figure 4). The quantum yield of $\text{CH}_3\text{CN}/\text{HEPES}$ solution of **R1** with 2 equiv. Cu^{2+} was tested to be 0.18 using rhodamine B as standard ($\Phi_F=0.69$ in ethanol). Furthermore, the absorbance at 560 nm and the fluorescence intensity at 585 nm for **R1** increased linearly over the Cu^{2+} concentration range of $1.25\sim 6.0$ and $1.25\sim 7.5 \mu\text{mol}\cdot\text{L}^{-1}$ (Figures 3 and 4, inset), respectively, indicating that the sensor **R1** may be used as a candidate for detecting Cu^{2+} in live cell and/or water samples. For **R1**, the LOD for Cu^{2+} ions was $0.201 \mu\text{mol}\cdot\text{L}^{-1}$ based on the equation $DL=3S_{bi}/S$ (in which S_{bi} represents blank measurement standard deviation, while S stands for the intensity/sample content slope), which was much lower than the standard value for Cu^{2+} (about $20 \mu\text{mol}\cdot\text{L}^{-1}$) in drinking water recommended by the United States Environmental Protection Agency.^[36] In addition, Job's plot experiments were also carried out based on those absorption spectra, which revealed the **R1** : Cu^{2+} binding of 1 : 1 stoichiometry. Notably, the optical density reached a peak at the Cu^{2+} molecular fraction of about 0.5. Besides, the **R1** and Cu^{2+} binding constant was calculated as $2.79 \times 10^4 \text{ mol}\cdot\text{L}^{-1}$, which displayed good linearity ($R=0.99275$), when data were fit based on the Benesi-Hildebrand expression.^[37]

2.3 Reaction mechanism between R1 and Cu^{2+}

ESI-MS was also carried out for exploring the reaction mechanism between **R1** and Cu^{2+} . According to our observation, when Cu^{2+} was added to the acetonitrile solution of **R1**, a peak was detected at m/z 757.73, which was as-

Figure 3 Absorption spectra of $2.5 \mu\text{mol}\cdot\text{L}^{-1}$ sensor **R1** upon the addition of Cu^{2+} (0~25 equiv.) in $\text{CH}_3\text{CN}/\text{HEPES}$ ($V/V=5/5$) solution at pH 7.40

The inset shows the absorbance at 560 nm as a function of Cu^{2+} concentration ($1.25\sim 6.0 \mu\text{mol}\cdot\text{L}^{-1}$)

Figure 4 Fluorescence emission spectra of $2.5 \mu\text{mol}\cdot\text{L}^{-1}$ sensor **R1** upon the addition of Cu^{2+} (0~25 equiv.) in $\text{CH}_3\text{CN}/\text{HEPES}$ ($V/V=5/5$) solution at pH 7.40

The inset shows the fluorescence intensity at 585 nm as a function of Cu^{2+} concentration ($1.25\sim 7.50 \mu\text{mol}\cdot\text{L}^{-1}$)

signed to intermediate $[\text{R1-H}+\text{Cu}+\text{NO}_3]$ complex. Simultaneously, a peak at m/z 442.84 was assigned to rhodamine B, which may be formed through intermediate hydrolysis, resulting in color and fluorescence changes. In fact, Czarnik *et al.*^[38] had examined the hydrolysis mechanism. In addition, the emission of fluorescence of sensor **R1** ($2.5 \mu\text{mol}\cdot\text{L}^{-1}$) remained almost unchanged at the excess addition of Na_2EDTA (10 equiv.) into $10 \text{ mmol}\cdot\text{L}^{-1}$ $\text{CH}_3\text{CN}/\text{HEPES}$ ($\text{pH}=7.40$, $V/V=5/5$) buffer solution, while 2 equiv. of Cu^{2+} was added. Such facts also verified that the mechanism of hydrolysis reaction between **R1** and Cu^{2+} based on a hydrolysis process rather than a coordination reaction.

To further clarify the coordination behavior of the **R1**- Cu^{2+} complex, energy optimized structures of **R1** and **R1**- Cu^{2+} complex were obtained with the Becke-3-Lee-

Yang-Parr (B3LYP) exchange function using the Gaussian 09 package. Figure 5 presents the optimization of molecular geometry at the binding stoichiometry between **R1** and Cu^{2+} ions of 1 : 1. In the free sensor, **R1** existed in the form of closed spirocycle, while in the **R1**- Cu^{2+} complex, it exists in the form of spirocycle opening. Besides, pyrrole N, imine N as well as carbonyl O atoms were also utilized for Cu^{2+} coordination, and the bond length was 0.1990~0.2055 nm. By contrast, rhodamine B hydrazone that bears the identical pyrrole unit (**R2**) sensors for Cu^{2+} is synthesized according to the coordination mechanism in our previous work.^[30] In the energy-minimized structure for **R2**- Cu^{2+} complex, Cu^{2+} ions are surrounded by two independent probes, with NO donor set from the carbonyl O and pyrrole N atoms. For Cu—O and Cu—N, their bond lengths are 0.192 and 0.205 nm, respectively, which are slightly shorter than those in the **R1**- Cu^{2+} complex. Notably, a potent intramolecular N—H···O hydrogen bond exists in the **R2**- Cu^{2+} complex between the pyrrole ring N atom and the hydrazone O atom (with the D—H···A distance of 0.276 nm), which may enhance the complex stability and thus prevent the hydrolysis process. In this regard, it can be roughly concluded that the rhodamine backbone structure is responsible for the sensor functional mechanism.

Figure 5 Optimized structures and HOMO/LUMO of **R1** and **R1**- Cu^{2+} complex by DFT calculation
H atoms are omitted for clarity

For **R1** and **R1**- Cu^{2+} complex, their LUMO and HOMO orbital energies and spatial distributions were also obtained (Figure 5). The results suggested that, for **R1**, both LUMO and HOMO were related to pyrrole and hydrazone moieties, at the same time, LUMO was on the rhodamine benzene ring. With regard to the **R1**- Cu^{2+} complex, HOMO was mainly localized near Cu^{2+} ions. Clearly, C—N bond on rhodamine moiety spirocycle was broken for the sake of

Cu^{2+} ion binding. In comparison, in **R1**- Cu^{2+} complex, those electrons within LUMO were localized within the entire xanthene backbone. Typically, for the **R1**- Cu^{2+} complex, the energy gap of LUMO with HOMO was determined as 1.58 eV, and this figure was remarkably reduced relative to the 3.78 eV of **R1**. Such findings better verified the **R1**- Cu^{2+} complex binding pattern, as displayed in Scheme 1.

Scheme 1 Proposed reaction mechanism of **R1** and Cu^{2+}

2.4 Environmental factor impacts and kinetic assay

The pH has been recognized as a leading factor affecting the rhodamine-based chemosensor response. The spectrum response of **R1** ($2.5 \mu\text{mol}\cdot\text{L}^{-1}$) with or without Cu^{2+} (2 equiv.) in the buffer solutions at different pH values was evaluated at ambient temperature. The fluorescence intensity in **R1** as well as **R1**+ Cu^{2+} complex was slightly dependent on pH 6.40~8.00. Specifically, the great difference in the fluorescence intensity of **R1** compared with the **R1**- Cu^{2+} complex suggested that **R1** was able to detect Cu^{2+} within such a range of pH. Therefore, the physiological pH 7.40 was chosen as the optimum experimental condition. In addition, the influence of those co-existing metal cations was also investigated. As suggested by our observations, those co-existing metal cations, such as Zn^{2+} , Pb^{2+} , Ni^{2+} , Na^+ , Mn^{2+} , Mg^{2+} , K^+ , Hg^{2+} , Fe^{3+} , Cr^{3+} , Co^{2+} , Cd^{2+} , Ca^{2+} , Al^{3+} and Ag^+ ions had almost no interference with Cu^{2+} detection.

For a chemosensor, the short response time plays a vital role similar to the great selectivity and sensitivity. For real-time monitoring of those target metal ions, we examined the time-fluorescence change response relationship of $2.5 \mu\text{mol}\cdot\text{L}^{-1}$ **R1** in $10 \text{ mmol}\cdot\text{L}^{-1}$ $\text{CH}_3\text{CN}/\text{HEPES}$ (pH=7.40, $V/V=5/5$) buffer solution when 2 equiv. Cu^{2+} was added. The **R1** fluorescence response towards Cu^{2+} ion occurred quite soon. Such findings suggested the completion of **R1**- Cu^{2+} interaction with no time delay.

2.5 Practical applications

For verifying the sensor practical applicability, Cu^{2+} ions were determined within distilled water as well as the commercial Kangshifu Drinking Water samples according to standard addition approach.^[37] In brief, 1 mL of Cu^{2+} spiked water sample was added into 1 mL of CH_3CN solution containing $1 \text{ mmol} \cdot \text{L}^{-1}$ probe, and then the fluorescence spectra was recorded. The equation to calculate Cu^{2+} contents is shown in Figure 4. As suggested by our results, the recovery obtained was 91.2%~96.8%, which indicated that our sensor had appreciable practicality (Table 1).

Table 1 Determination of Cu^{2+} concentrations in drinking water samples

Probe	Sample	$c/(\mu\text{mol} \cdot \text{L}^{-1})$		Recovery/%
		Spiked	Found ^a	
R1	Distilled water	0.0	0	—
		2.5	2.39 ± 0.04	95.6
		5.0	4.84 ± 0.06	96.8
	Drinking water ^b	0.0	0	—
		2.5	2.28 ± 0.05	91.2
		5.0	4.69 ± 0.08	93.8

^a Average value of three determinations. ^b Kangshifu Drinking water (obtained from the local supermarket): $[\text{K}^+] = 25.64 \sim 700 \mu\text{mol} \cdot \text{L}^{-1}$; $[\text{Mg}^{2+}] = 4.17 \sim 203.12 \mu\text{mol} \cdot \text{L}^{-1}$; $[\text{Cl}^-] = 281.69 \sim 769 \mu\text{mol} \cdot \text{L}^{-1}$; $[\text{SO}_4^{2-}] = 4.17 \sim 71 \mu\text{mol} \cdot \text{L}^{-1}$.

3 Conclusions

To sum up, the efficient “turn-on” fluorescent probe based on rhodamine is developed to detect the Cu^{2+} ion. According to our results, the fluorescent probe exhibits great sensitivity and selectivity to Cu^{2+} , the LOD within the $10 \text{ mmol} \cdot \text{L}^{-1}$ $\text{CH}_3\text{CN}/\text{HEPES}$ (pH 7.40, $V/V=5/5$) buffer solution is $0.201 \mu\text{mol} \cdot \text{L}^{-1}$, and those co-existing metal cations almost have no interference on Cu^{2+} detection. The probe functions via coordination reaction by forming complex $[\text{R1-H} + \text{Cu} + \text{NO}_3]$ and followed by hydrolyzing to a fluorescent rhodamine B. Additionally, the practical application of sensor was also performed in Cu^{2+} spiked water samples.

4 Experimental section

4.1 Materials and methods

The starting materials and solvents in synthesis were bought from the market, which were utilized directly without further purification. In addition, rhodamine B hydrazide,^[39] as well as ethyl 5-formyl-3,4-dimethyl-pyrrole-2-carboxylate,^[40] was prepared in accordance with the methods reported in literature. The XT4-100X microscopic melting point machine (made in Beijing, China) was used to determine the sensor melting point. Meanwhile, the Elemental Vario EL analyzer was adopted for elemental analysis. The sensor ^1H NMR spectra were obtained using the Bruker AV400 NMR spectrometer within the $\text{DMSO}-d_6$ solution. Besides, the Bruker V70 FT-IR spectrophotometer

was used to determine IR spectra ($\nu = 4000 \sim 400 \text{ cm}^{-1}$) according to a KBr pressed disc approach. In the meantime, the Purkinje General TU-1800 spectrophotometer was employed to record UV spectra. The Varian CARY Eclipse spectrophotometer was applied in determining fluorescent spectra, and 5 nm pass width was applied in measuring both excitation and emission spectra. The Bruker Daltonics Esquire 6000 mass spectrometer was utilized to obtain ESI-MS spectra.

4.2 Preparation of ethyl 5-formyl-3,4-dimethyl-pyrrole-2-carboxylate rhodamine B hydrazine (R1)

460 mg (1 mmol) of rhodamine B hydrazide was mixed with 20 mL of ethanol solution supplemented with 195 mg (1 mmol) of ethyl 5-formyl-3,4-dimethyl-pyrrole-2-carboxylate. Then, the mixed solution was refluxed for 3 h, followed by cooling until ambient temperature. Later, the isolated solid was subjected to filtering and ethanol washing. Yield 71%. m.p. $124 \sim 128 \text{ }^\circ\text{C}$; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 11.51 (s, 1H, NH), 9.13 (s, 1H, CH=N), 7.81~7.83 (m, 1H, Aryl-H), 7.52~7.57 (m, 2H, Aryl-H), 7.08~7.09 (m, 1H, Aryl-H), 6.26~6.33 (m, 6H, Aryl-H), 4.11~4.16 (q, $J=7.5 \text{ Hz}$, 2H, CH_2), 3.24~3.26 (q, $J=8.0 \text{ Hz}$, 8H, 4CH_2), 2.03 (s, 3H, CH_3), 1.71 (s, 3H, CH_3), 1.18~1.22 (t, $J=8.0 \text{ Hz}$, 3H, CH_3), 0.99~1.03 (t, $J=8.0 \text{ Hz}$, 12H, 4CH_3); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 170.4, 161.6, 151.9, 151.9, 149.4, 149.4, 139.5, 139.2, 132.6, 131.3, 129.2, 128.8, 128.8, 128.3, 128.0, 126.9, 123.3, 122.7, 122.7, 114.3, 114.3, 106.5, 97.7, 97.7, 76.3, 60.9, 47.1, 47.1, 47.1, 47.1, 14.1, 12.9, 12.9, 12.9, 12.9, 10.4, 10.00. ESI-MS m/z : 633.84. Anal. calcd for $\text{C}_{39}\text{H}_{43}\text{N}_5\text{O}_4$: C 72.01, H 6.84, N 11.05; found C 72.11, H 6.98, N 10.89.

4.3 Measurements of general fluorescence and UV-Vis spectra

The spectral analyses were accomplished in $\text{CH}_3\text{CN}/\text{HEPES}$ buffer ($10 \text{ mmol} \cdot \text{L}^{-1}$, pH=7.40, $V/V=5/5$) solution at room temperature. The concentration of the sensor **R1** for UV-vis and fluorescence measurement was $2.5 \mu\text{mol} \cdot \text{L}^{-1}$. The metal ion solutions were prepared using chloride salts or nitrate under CH_3CN . Besides, the fluorescence and UV-vis spectrophotometric titration was carried out within the cuvette (2 mL) through successively adding relevant chemical reagents by the microliter syringe. The solution was sufficiently mixed after adding any aliquot, then the spectra were determined.

Supporting Information Job plots of **R1** and Cu^{2+} in $\text{CH}_3\text{CN}/\text{HEPES}$ solution; the Benesi-Hildebrand plot of **R1**- Cu^{2+} complex; ESI-MS spectrum; reversible experiment; the effect of pH; the interference experiment; response time experiment; NMR spectrum of **R1**. The Supporting Information associated with this article can be found in the online version at <http://sioc-journal.cn/>.

References

- [1] Wei, J. H.; Yi, J. W.; Han, M. L.; Li, B.; Liu, S.; Wu, Y. P.; Ma, L. F.; Li, D. S. A. *Chem. Asian J.* **2019**, *14*, 3694.

- [2] Wang, H.; Qin, J.; Huang, C.; Han, Y.; Xu, W.; Hou, H. *Dalton Trans.* **2016**, 45, 12710.
- [3] Wu, X.; Guo, Z.; Wu, Y.; Zhu, S.; James, T. D.; Zhu, W. *ACS Appl. Mater. Interfaces* **2013**, 5, 12215.
- [4] Guo, Z.; Kim, G. H.; Yoon, J.; Shin, I. *Nat. Protoc.* **2014**, 9, 1245.
- [5] Ma, D. L.; Wong, S. Y.; Kang, T. S.; Ng, H. P.; Han, Q. B.; Leung, C. H. *Methods* **2019**, 168, 3.
- [6] Jung, J. H.; Lee, J. H.; Shinkai, S. *Chem. Soc. Rev.* **2011**, 40, 4464.
- [7] Patil, P.; Ajey, K. V.; Bhat, M. P.; Sriram, G.; Yu, J.; Jung, H. Y.; Altalhi, T.; Kigga, M.; Kurkuri, M. D. *ChemistrySelect* **2018**, 3, 11593.
- [8] Suo, F.; Chen, X.; Fang, H. Gong, Q.; Yu, C.; Yang, N. D.; Li, S.; Wu, Q.; Li, L.; Huang, W. *Dyes Pigm.* **2019**, 170, 107639.
- [9] Ji, Y.; Dai, F.; Zhou, B. *Free Radical Biol. Med.* **2018**, 129, 215.
- [10] Singh, N.; Paknikar, K. M.; Rajwade, J. *Environ. Res.* **2019**, 175, 367.
- [11] Ogunkunle, C. O.; Jimoh, M. A.; Asogwa, N. T.; Viswanathan, K.; Vishwakarma, V.; Fatoba, P. O. *Ecotoxicol. Environ. Saf.* **2018**, 155, 86.
- [12] Donnelly, P. S.; Xiao, Z.; Wedd, A. G. *Curr. Opin. Chem. Biol.* **2007**, 11, 128.
- [13] Squitti, R.; Ghidoni, R.; Simonelli, I.; Ivanova, I. D.; Colabufo, N. A.; Zuin, M.; Benussi, L.; Binetti, G.; Cassetta, E.; Rongioletti, M.; Siotto, M. *J. Trace Elem. Med. Biol.* **2018**, 45, 181.
- [14] Gardner, B.; Dieriks, B. V.; Cameron, S.; Mendis, L. H. S.; Turner, C.; Faull, R. L. M.; Curtis, M. A. *Sci. Rep.* **2017**, 7, 10454.
- [15] Ren, D.; Liu, Y.; Liu, X.; Li, Z.; Li, H.; Yang, X. F. *Sens. Actuators, B* **2018**, 255, 2321.
- [16] Yoon, J. W.; Chang, M. J.; Hong, S.; Lee, M. H. *Tetrahedron Lett.* **2017**, 58, 3887.
- [17] Jiao, Y.; Zhou, L.; He, H.; Yin, J.; Gao, Q.; Wei, J.; Duan, C.; Peng, X. *Talanta* **2018**, 184, 143.
- [18] Jia, X.; Xiao, Z.; Hui, P.; Liu, C.; Wang, Q.; Qiu, X. He, S.; Zeng, X.; Zhao, L. *Dyes Pigm.* **2019**, 160, 633.
- [19] Kang, H.; Fan, C.; Xu, H.; Pu, G.; Liu, S. *Tetrahedron* **2018**, 74, 4390.
- [20] Zhang, J. F.; Zhou, Y.; Yoon, J.; Kim, Y.; Kim, S. J.; Kim, J. S. *Org. Lett.* **2010**, 12, 3852.
- [21] Yang, Z.; She, M.; Zhang, J.; Chen, X.; Huang, Y.; Zhu, H.; Liu, P.; Li, J.; Shi Z. *Sens. Actuators, B* **2013**, 176, 482.
- [22] Yu, M.; Yuan, R.; Shi, C.; Zhou, W.; Wei, L.; Li, Z. *Dyes Pigm.* **2013**, 99, 887.
- [23] Goswami, S.; Sen, D.; Das, A. K.; Das, N. K.; Aich, K.; Fun, H. K.; Quah, C. K.; Maity, A. K.; Saha, P. *Sensors. Actuators, B* **2013**, 183, 518.
- [24] Kar, C.; Adhikari, M. D.; Ramesh, A.; Das, G. *Inorg. Chem.* **2013**, 52, 743.
- [25] Xiong, K.; Chen, J. G. *Catalysis Today* **2018**, 339, 289.
- [26] Xu, Q.; Guo, Z. *J. East China Univ. Sci. Technol.* **2019**, 45, 357.
- [27] Yi, X.; Li, G.; Huang, L.; Chu, Y.; Liu, Z. Q.; Xia, H.; Zheng, A.; Deng, F. *J. Phys. Chem. C* **2017**, 121, 3887.
- [28] Jr, P. A.; Liu, Y.; Palacios, M. A.; Minami, T. Wang, Z.; Nishiyabu, R. *Chem.-Eur. J.* **2013**, 19, 8497.
- [29] Wang, Y.; Wu, H.; Wu, W. N.; Li, S. J.; Xu, Z. H.; Xu, Z. Q. Fan, Y. C.; Zhao, X. L.; Liu, B. Z. *Sens. Actuators, B* **2018**, 260, 106.
- [30] Moubaraki, R.; Li, B.; Murray, K. S.; Brooker, S. *Eur. J. Inorg. Chem.* **2009**, 19, 2851.
- [31] Wang, Y.; Chang, H. Q.; Wu, W. N.; Peng, W. B.; Yan, Y. F. He, C. M.; Chen, T. T.; Zhao, X. L.; Xu, Z. Q. *Sensors. Actuators B: Chem.* **2016**, 228, 395.
- [32] Li, M.; Lv, H.; Luo, J. Z.; Miao, J. Y.; Zhao, B. X. *Sens. Actuators, B* **2013**, 188, 1235.
- [33] Ge, F.; Ye, H.; Luo, J. Z.; Wang, S.; Sun, Y. J.; Zhao, B. X.; Miao, J. Y. *Sens. Actuators, B* **2013**, 181, 215.
- [34] Huang, K.; Jiao, X.; Liu, C.; Wang, Q.; Qiu, X.; He, S. Zhao, L.; Zeng, X. *Dyes Pigm.* **2017**, 145, 561.
- [35] Artesani, A.; Binet, L.; Tana, F.; Comelli, D.; Nardo, L. D.; Nevin, A.; Touati, N.; Valentini, G.; Gourier, D. *Microchem. J.* **2019**, 151, 104256.
- [36] Pressman, J. G.; Richardson, S. D.; Speth, T. F.; Miltner, R. J. *Environ. Sci. Technol.* **2010**, 44, 7184.
- [37] Xu, Z.; Wang, H.; Hou, X.; Xu, W.; Xiang, T.; Wu, C. *Sens. Actuators, B* **2014**, 201, 469.
- [38] Dujols, V.; Ford, F.; Czarnik, A. W. *J. Am. Chem. Soc.* **1997**, 119, 7386.
- [39] Kim, H.; Rao, B. A.; Jeon, J. W.; Mallick, S.; Kang, S. M.; Choi, J. S.; Lee, C. S.; Son, Y. A. *Sens. Actuators, B* **2015**, 210, 173.
- [40] Ye, X. P.; Zhu, T. F.; Wu, W. N.; Ma, T. L.; Xu, J.; Zhang, Z. P.; Wang, Y.; Jia, L. *Inorg. Chem. Commun.* **2014**, 47, 60.

(Fan, Y.)