新型手性钳形 PNN 类配体在钯催化不对称烯丙基烷基化反应中的应用

吴敦奇 成 轩 刘炎开 成 果 管笑宇 邓清海*

(上海师范大学 教育部资源化学重点实验室和上海市稀土功能材料重点实验室 上海 200234)

摘要 从便宜、易得的手性氨基酸出发通过 5 至 6 步反应制备了一种新型手性钳形 PNN 类配体 oxpma。该类手性配体含有 2 至 3 个手性中心。将其用于钯催化的烯丙基酯的不对称烯丙基烷基化反应, 能够以高的对映选择性(最高可达 96%)获得相应的手性目标产物。

关键词 PNN 配体; 钳形配体; 不对称催化; 不对称烯丙基烷基化反应

A New Class of Chiral Pincer-Type PNN Ligands for Pd-Catalyzed Asymmetric Allylic Alkylation

Wu, Dunqi Cheng, Xuan Liu, Yankai Cheng, Guo Guan, Xiaoyu Deng, Qinghai*

(The Education Ministry Key Lab of Resource Chemistry and Shanghai Key Laboratory of Rare Earth Functional Materials, Shanghai Normal University, Shanghai 200234)

Abstract A new class of chiral tridentate P,N,N-donor pincer ligands bearing two or three stereocenters, 1-(4,5-dihydro-oxazol-2-yl)-N-(2-(diphenylphosphanyl)benzyl)methanamines (oxpma), were synthesized starting from readily available amino acids in five or six steps. They were applied in palladium-catalyzed asymmetric allylic alkylation of allylic acetates to afford the desired products with high enantioselectivities (up to 96% ee).

Keywords PNN ligand; pincer ligand; asymmetric catalysis; asymmetric allylic alkylation

1 Introduction

Chiral tridentate ligands, also called as chiral pincer-type ligands, have achieved widespread applications in asymmetric catalysis since they allow exquisite control of electronic/steric features around the metal center as well as enhance the metal complex stability.^[1] Among these, various chiral PNN-type pincer ligands with different chiral structural units have been designed (Figure 1). According to the source of chirality, these ligands can be majorly divided to chiral spiro PNN ligands (A),^[2-3] the ligands bearing phosphine groups with chiral substituents (B),^[4] chiral binaphthyl ligands (C),^[5] chiral ferrocenyl ligands (D),^[6-7] chiral amine-based ligands (E)^[8] and chiral oxazoline-containing ligands (F).^[9] Surprisingly, these ligands as well as the metal complexes have only been applied to asymmetric hydrogenation and asymmetric transfer hydrogenation. Therefore, the development of new kind of chiral PNN ligands to catalyze other kind of asymmetric

transformations is an attractive target.

We have been interested in the synthesis and applications of chiral oxazoline-based ligands^[10] due to their ready accessibility from cheap and diverse chiral amino alcohols as well as versatile utilities in a wide range of asymmetric transformations.^[11] In contrast, only a few chiral pincer PNN ligands containing oxazoline ring have been reported. Recently, Shi and co-workers^[9] developed oxazoline-based PNN ligands (F) for metal catalyzed polymerization of norbornene. In addition, Zhou and Zhang developed the chiral PNN ligands that associated chiral oxazoline ring with chiral spiro- (A2)^[3] and ferrocenyl (D2)^[7] backbones, respectively, which were both applied in iridium-catalyzed asymmetric hydrogenation. In this context, we report the synthesis of a new class of chiral PNN ligands, 1-(4,5-dihydrooxazol-2-yl)-N-(2-(diphenylphosphanyl)benzyl)-methanamines (oxpma, 1) (Figure 1), which are derived from readily available amino alcohols and amino acids. In an evaluation of their potential in asymmetric catalysis, it

* Corresponding author. E-mail: qinghaideng@shnu.edu.cn

Received May 30, 2020; revised June 16, 2020; published online July 8, 2020.

Dedicated to Professor Henry N. C. Wong on the occasion of his 70th birthday.

Project supported by the National Natural Science Foundation of China (Nos. 21402119, 21772122) and the Shanghai Engineering Research Center of Green Energy Chemical Engineering (No. 18DZ2254200).

国家自然科学基金(Nos. 21402119, 21772122)和上海绿色能源化工工程技术研究中心(No. 18DZ2254200)资助项目。

was found that they induced high enantioselectivity in the asymmetric allyl alkylation reaction catalyzed by palladium medium,^[12] which is a powerful strategy for stereoselective construction of C—C bond.

2 Results and discussion

Chiral oxazoline ligand **1** was readily prepared starting from amino acid **2** in 5 or 6 steps (Scheme 1). First, amino acid **2** were transformed easily to Fmoc-protected amino acid **3**,^[13] which react with amino alcohol **4** to construct the oxazoline ring directly for providing Fmoc-oxazoline amines (for **6a~6d**) in moderate yields (62%~72%).^[14] When R¹ is *tert*-butyl group, **6e** could be obtained through the intermediate **5** in 2 steps.^[15] After deprotection of **6**,^[14]

oxazoline amines **7** were condensed with 2-(diphenylphosphanyl)benzaldehyde in the presence of 2 equiv. of tetraethoxytitanium to get imines **8**, which are unstable upon purification with silica gel chromatography. Thus, the crude compounds **8** were converted directly to ligands **1** through reduction by NaBH₄ or addition of Grignard reagents with sole diastereoselectivity.^[16]

With the ligand library in hand, ligand **1b** and [Pd(η³-C₃H₅)Cl]₂ were chosen to study the asymmetric allylic alkylation of **9a** with **10a** (Table 1). Several commonly used bases were first examined (Table 1, Entries 1~3), and *N,O*-bis(trimethylsilyl)acetamide (BSA) proved to be the optimal one (98% yield, 95% *ee*). Solvent screening showed that both 1,2-dichloroethane and acetonitrile are

Figure 1 Representative chiral pincer-type PNN ligands

Scheme 1 Synthesis of chiral pincer-type PNN ligand oxpma (**1**)

the suitable solvents, and dichloromethane was still chosen for further investigations due to the convenience of operation (Table 1, Entries 3~7). A series of ligands were subsequently screened (Table 1, Entries 3 and 8~14). It was found that the conversion and enantioselectivity were dependent upon the combination of substituents at three stereocenters and ligand **1b** gave the most promising result. Then, decreasing the amount of BSA to 2 equiv. (Table 1, Entry 15) and further reducing the catalyst loading (Table 1, Entry 17) achieved comparative results. Finally, changing the reaction temperature did not give better results (Table 1, Entries 19 and 20).

Table 1 Optimization of the reaction conditions^a

Entry	Base (equiv.)	L	Solvent	t/h	Yield ^b /%	ee ^c /%
1	K ₂ CO ₃ (3)	1b	CH ₂ Cl ₂	13	99	92
2	CS ₂ CO ₃ (3)	1b	CH ₂ Cl ₂	13	99	87
3	BSA (3)	1b	CH ₂ Cl ₂	13	98	95
4	BSA (3)	1b	ClCH ₂ CH ₂ Cl	13	99	95
5	BSA (3)	1b	MeCN	13	99	94
6	BSA (3)	1b	Toluene	13	11	89
7	BSA (3)	1b	THF	13	19	94
8	BSA (3)	1a	CH ₂ Cl ₂	12	22	55
9	BSA (3)	1c	CH ₂ Cl ₂	12	65	78
10	BSA (3)	1d	CH ₂ Cl ₂	12	72	86
11	BSA (3)	1e	CH ₂ Cl ₂	12	60	91
12	BSA (3)	1f	CH ₂ Cl ₂	12	63	89
13	BSA (3)	1g	CH ₂ Cl ₂	12	56	86
14	BSA (3)	1h	CH ₂ Cl ₂	12	58	97
15	BSA (2)	1b	CH ₂ Cl ₂	11	98	95
16	BSA (1.5)	1b	CH ₂ Cl ₂	27	80	95
17 ^d	BSA (2)	1b	CH ₂ Cl ₂	12	97	95
18 ^e	BSA (2)	1b	CH ₂ Cl ₂	12	88	95
19 ^{d,f}	BSA (2)	1b	CH ₂ Cl ₂	12	68	95
20 ^{d,g}	BSA (2)	1b	CH ₂ Cl ₂	12	86	83

^a Reaction conditions: **9a** (0.10 mmol), **10a** (1.2 equiv.), base (x equiv.), [Pd(η³-C₃H₅)Cl]₂ (2 mol%), ligand (5 mol%), solvent (1.0 mL), room temperature, nitrogen. ^b The yields of isolated products. ^c Determined by HPLC analysis. ^d 1 mol% [Pd(η³-C₃H₅)Cl]₂ and 3 mol% **1b** were used. ^e 0.5 mol% [Pd(η³-C₃H₅)Cl]₂ and 1.5 mol% **1b** were used. ^f The reaction was carried out under 0 °C. ^g The reaction was carried out under 40 °C.

The substrate scope of this reaction was subsequently investigated under the optimal reaction conditions (Table 1, Entry 17) which was shown in Table 2. A series of allylic acetates (**9a**~**9e**) are well tolerated to provide the corresponding products with high enantioselectivities (**11aa**~**11ea**, 91%~96% *ee*) regardless of the electronic nature of the substituents. The position of substituent affected the reaction outcome. The reaction of 1,3-bis(3-

methylphenyl)allyl acetate (**9f**) conducted well to afford the desired product **11fa** in 79% yield with 90% *ee*, while the substrate with *ortho*-chlorine (**9g**) only gave product **11ga** with 61% *ee*. In particular, the substrate with 2-naphthyl group (**9h**) is suitable for the reaction condition to get the product **11ga** in 97% yield with 92% *ee*. Various nucleophiles **10** were then examined. The malonates with different substituents gave the corresponding products (**11ab**~**11ae**) with good enantioselectivities (85%~94% *ee*) in moderate to high yields (48%~97%). The reaction of **9a** with acetylacetone (**10f**) was also carried out to get the desired product **11af** in 75% yield with 90% *ee*. However, malononitrile (**10g**) is not tolerated and gave the product **11ag** in low yield with moderate enantioselectivity.

Table 2 Substrate scope of Pd-catalyzed asymmetric allylic alkylation^a

Entry	Ar (9)	R ¹ , R ² (10)	Product	Yield ^b /%	ee ^c /%
1	C ₆ H ₅ (9a)	H, CO ₂ Me (10a)	11aa	97	95
2	4-FC ₆ H ₄ (9b)	H, CO ₂ Me (10a)	11ba	86	96
3	4-ClC ₆ H ₄ (9c)	H, CO ₂ Me (10a)	11ca	68	92
4	4-BrC ₆ H ₄ (9d)	H, CO ₂ Me (10a)	11da	89	92
5	4-MeC ₆ H ₄ (9e)	H, CO ₂ Me (10a)	11ea	75	91
6	3-ClC ₆ H ₄ (9f)	H, CO ₂ Me (10a)	11fa	79	90
7	2-MeC ₆ H ₄ (9g)	H, CO ₂ Me (10a)	11ga	84	61
8	2-Np (9h)	H, CO ₂ Me (10a)	11ha	97	92
9	C ₆ H ₅ (9a)	H, CO ₂ ⁱ Pr (10b)	11ab	51	88
10	C ₆ H ₅ (9a)	H, CO ₂ Bn (10c)	11ac	97	94
11	C ₆ H ₅ (9a)	Ph, CO ₂ Et (10d)	11ad	86	92
12	C ₆ H ₅ (9a)	Me, CO ₂ Et (10e)	11ae	48	85
13	C ₆ H ₅ (9a)	H, COMe (10f)	11af	75	90
14	C ₆ H ₅ (9a)	H, CN (10g)	11ag	21	74

^a Reaction conditions: **9** (0.10 mmol), **10** (1.2 equiv.), BSA (2 equiv.), [Pd(η³-C₃H₅)Cl]₂ (1 mol%), ligand (3 mol%), CH₂Cl₂ (1.0 mL), room temperature, nitrogen. ^b The yields of isolated products. ^c Determined by HPLC analysis.

3 Conclusions

A new class of chiral tridentate PNN ligands, oxpma, were developed. These ligands are derived from readily available amino acids. Their brief utility as stereodirecting ligands in Pd-catalyzed asymmetric allylic alkylation has been first evaluated, which achieved good enantioselectivities (up to 96% *ee*). Further applications on asymmetric transformations are currently being studied in our laboratory.

4 Experimental section

4.1 General information

¹H NMR spectra were recorded on a 400 or 600 MHz

NMR spectrometer and ^{13}C NMR spectra were recorded on a 101 or 150 MHz NMR spectrometer. Infrared spectra were prepared as KBr pellets and recorded on a Varian Excalibur 3100 series FT-IR spectrometer. Mass spectra were recorded by the mass spectrometry service of Shanghai Institute of Organic Chemistry. HPLC analyses on a Waters 1596 or Shimadzu SPD-15C. All manipulations were maintained under an atmosphere of nitrogen unless otherwise stated. Commercially available reagents were used without further purification. Solvents were pre-dried over activated 4 Å molecular sieves and refluxed over sodium-benzophenone (toluene, tetrahydrofuran), phosphorus pentoxide (chloroform) or calcium hydride (dichloromethane, dichloroethane, acetonitrile) under an argon atmosphere and collected by distillation. Column chromatography was performed on silica gel (200~300 mesh).

4.2 Synthesis of ligand 1

4.2.1 General procedure for the synthesis of 3

To a solution of amino acid **2** (40.0 mmol) in dioxane (40 mL) was added Na_2CO_3 (aq., $\omega=10\%$, 100 mL) at room temperature, then the mixture was cooled to 0 °C with an ice bath. Fmoc-Cl (1.1 equiv.) in dioxane (100 mL) was added by dropping funnel slowly, and the mixture was stirred at 0 °C for 1 h, then the mixture was stirred at room temperature overnight. 100 mL of water was added to the solution, and the aqueous layer was washed with diethyl ether. The organic layer was extracted with saturated NaHCO_3 and the combined aqueous layers were cooled to 0 °C and acidified to pH 1 with 1 mol/L HCl. The aqueous layer was then extracted with EtOAc (40 mL $\times$ 3). The organic layers were combined and dried over MgSO_4 . After filtration and concentration, the resulting Fmoc-protected amino acid **3** was obtained.^[13]

((9H-Fluoren-9-yl)methoxycarbonyl)-L-valine (**3a**): White solid, 97% yield (13.17 g, 38.8 mmol). m.p. 74.3~75.7 °C (lit.^[18] 144~145 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.77 (d, $J=7.2$ Hz, 2H), 7.60 (d, $J=6.0$ Hz, 2H), 7.40 (t, $J=7.2$ Hz, 2H), 7.31 (t, $J=7.2$ Hz, 2H), 5.28 (d, $J=8.8$ Hz, 1H), 4.42 (d, $J=6.8$ Hz, 2H), 4.36~4.33 (m, 1H), 4.24 (t, $J=6.8$ Hz, 1H), 2.27~2.21 (m, 1H), 1.01 (d, $J=6.4$ Hz, 3H), 0.95 (d, $J=6.8$ Hz, 3H).

(S)-2-(((9H-Fluoren-9-yl)methoxycarbonyl)amino)-3,3-dimethylbutanoic acid (**3b**):^[18] White solid, 97% yield (13.71 g, 38.8 mmol). m.p. 72.3~73.4 °C (lit.^[18] 71 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.76 (d, $J=7.6$ Hz, 2H), 7.59 (d, $J=7.2$ Hz, 2H), 7.40 (t, $J=7.2$ Hz, 2H), 7.31 (t, $J=7.6$ Hz, 2H), 5.34 (d, $J=5.6$ Hz, 1H), 4.48~4.38 (m, 2H), 4.24~4.22 (m, 2H), 1.04 (s, 9H).

4.2.2 General procedure for the synthesis of 6a~6d

To a flame-dried 500 mL three-neck bottle equipped with a constant dropping funnel, *N*-Fmoc-amino acid **3a** (12 mmol), amino alcohol **4** (12 mmol), PPh_3 (36 mmol) and CH_2Cl_2 (60 mL) were added. After further addition of diisopropylethylamine (DIPEA) (36 mmol), the mixture was cooled down to 0 °C with an ice bath. Then, CCl_4 (60 mmol) in another 60 mL of CH_2Cl_2 was added through a

funnel over 3 h. The mixture was stirred at room temperature overnight. After removal of solvent, the residue was dissolved in EtOAc (100 mL). After filtration and subsequent concentration in vacuo, the residue was purified by flash column chromatography (petroleum ether/EtOAc, $V:V=5:1$) to get products **6a**~**6d**.^[14]

(9H-Fluoren-9-yl)methyl-((S)-2-methyl-1-((R)-4-phenyl-4,5-dihydrooxazol-2-yl)propyl)carbamate (**6a**):^[15] White solid, 68% yield (3.59 g, 8.2 mmol). m.p. 119.2~121.3 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.76 (d, $J=7.6$ Hz, 2H), 7.61 (t, $J=6.8$ Hz, 2H), 7.41~7.25 (m, 9H), 5.54 (d, $J=8.8$ Hz, 1H), 5.23 (t, $J=9.6$ Hz, 1H), 4.72 (t, $J=9.4$ Hz, 1H), 4.50~4.11 (m, 5H), 2.28~2.20 (m, 1H), 1.05 (d, $J=6.8$ Hz, 3H), 1.02 (d, $J=6.8$ Hz, 3H).

(9H-Fluoren-9-yl)methyl-((S)-1-((R)-4-isopropyl-4,5-dihydrooxazol-2-yl)-2-methylpropyl)carbamate (**6b**):^[14] White solid, 72% yield (3.51 g, 8.6 mmol). m.p. 117.1~117.3 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.77 (d, $J=7.6$ Hz, 2H), 7.62 (t, $J=7.6$ Hz, 2H), 7.40 (t, $J=7.6$ Hz, 2H), 7.31 (t, $J=7.6$ Hz, 2H), 5.41 (d, $J=8.8$ Hz, 1H), 4.42~4.23 (m, 5H), 4.01~3.91 (m, 2H), 2.17~2.09 (m, 1H), 1.80~1.73 (m, 1H), 0.98~0.89 (m, 12H).

(9H-Fluoren-9-yl)methyl-((S)-1-((R)-4-benzyl-4,5-dihydrooxazol-2-yl)-2-methylpropyl)carbamate (**6c**):^[19] White solid, 64% yield (3.49 g, 7.7 mmol). m.p. 89.2~91.3 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.77 (d, $J=7.6$ Hz, 2H), 7.61 (dd, $J=7.6$, 7.2 Hz, 2H), 7.41 (t, $J=7.6$ Hz, 2H), 7.33~7.20 (m, 7H), 5.40 (d, $J=9.2$ Hz, 1H), 4.47~4.23 (m, 6H), 4.02 (t, $J=8.0$ Hz, 1H), 3.12 (dd, $J=13.6$, 4.8 Hz, 1H), 2.65 (dd, $J=13.6$, 8.8 Hz, 1H), 2.14~2.08 (m, 1H), 0.94 (dd, $J=12.4$, 6.8 Hz, 6H).

(9H-Fluoren-9-yl)methyl-(S)-(1-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)-2-methylpropyl)carbamate (**6d**): White solid, 62% yield (2.92 g, 7.4 mmol). m.p. 104~106 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.76 (d, $J=7.6$ Hz, 2H), 7.62 (t, $J=7.8$ Hz, 2H), 7.39 (t, $J=7.4$ Hz, 2H), 7.31 (t, $J=7.2$ Hz, 2H), 5.49 (d, $J=8.8$ Hz, 1H), 4.46~4.33 (m, 3H), 4.24 (t, $J=7.0$ Hz, 1H), 3.97 (s, 2H), 2.17~2.09 (m, 1H), 1.29 (s, 6H), 0.98~0.93 (m, 6H); ^{13}C NMR (100 MHz, CDCl_3) δ : 164.8, 156.2, 144.1, 143.9, 141.4, 127.7, 127.1, 125.2, 120.0, 79.5, 67.2, 66.9, 54.4, 47.3, 31.7, 28.5, 28.3, 18.8, 17.7; IR (KBr) ν_{max} : 2965, 1716, 1663, 1624, 1394, 1366, 1237, 740 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{29}\text{N}_2\text{O}_3$ ($\text{M}+\text{H}^+$) 393.2173, found 393.2168.

4.2.3 Procedures for the synthesis of 6e

Fmoc-protected amino acid **3b** (20 mmol) was dissolved in dichloromethane (150 mL) at 0 °C. Then, *D*-phenylglycinol (24 mmol, 1.2 equiv.), HOBt (30 mmol, 1.5 equiv.) and *N,N'*-carbonyldiimidazole (EDCI) (30 mmol, 1.5 equiv.) were added in sequence. After stirring at 0 °C for 15 min, *N*-methyl morpholine (34 mmol, 1.7 equiv.) was added. The reaction mixture was stirred at ambient temperature until the disappearance of **3b** as monitored by thin layer chromatography (TLC). The resulting solution was washed successively with cold KHSO_4 aqueous solution ($\omega=5\%$, 50 mL $\times$ 3), saturated NaHCO_3 solution (50

mL×3) and brine (75 mL). The combined organic layer was dried over MgSO_4 and filtered. The solvent was removed under vacuum and the resulting residue was crystallized using CH_2Cl_2 -hexanes ($V:V=1:4$) to afford (9*H*-fluoren-9-yl)methyl-((*S*)-1-(((*R*)-2-hydroxy-1-phenylethyl)amino)-3,3-dimethyl-1-oxobutan-2-yl)-carbamate (**5**), 72% yield (6.81 g, 14.4 mmol). m.p. 175~177 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.76 (d, $J=7.6$ Hz, 2H), 7.58 (d, $J=7.6$ Hz, 2H), 7.42~7.27 (m, 9H), 6.51 (d, $J=7.6$ Hz, 1H), 5.60 (d, $J=9.2$ Hz, 1H), 5.11~5.07 (m, 1H), 4.44~4.30 (m, 2H), 4.21 (t, $J=6.8$ Hz, 1H), 3.96 (d, $J=9.2$ Hz, 1H), 3.90~3.81 (m, 2H), 2.70 (br s, 1H), 0.98 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ : 171.4, 157.2, 144.0, 143.6, 141.4, 139.3, 128.6, 127.84, 127.82, 127.7, 127.22, 127.16, 127.0, 125.2, 125.1, 120.1, 120.0, 67.5, 65.7, 63.0, 55.6, 47.1, 34.8, 26.7; IR (KBr) ν_{max} : 3421, 1627, 1398, 1323, 1230, 682, 669, 650 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{29}\text{H}_{33}\text{N}_2\text{O}_4$ ($\text{M}+\text{H}^+$) 473.2435, found 473.2432.

Compound **5** (5.0 mmol) was dissolved in dry CH_2Cl_2 (100 mL) and then the solution was cooled to 0 °C in an ice bath. 4-Dimethylaminopyridine (1.0 mmol, 0.2 equiv.) and freshly distilled triethylamine (17.5 mmol, 3.5 equiv.) were added in sequence. After stirring at 0 °C for 5 min, a solution of *p*-toluenesulfonyl chloride (15 mmol, 1.5 equiv.) in dry CH_2Cl_2 (50 mL) was added dropwise. Then, the reaction was stirred at ambient temperature until the disappearance of **5** as monitored by TLC. The reaction was quenched with saturated NH_4Cl solution. The aqueous phase was extracted with CH_2Cl_2 (50 mL×3) and the combined organic layers were washed with brine, then dried over anhydrous Na_2SO_4 . After removal of the solvent, the residue was purified by flash column chromatography (300 mL of hexane-EtOAc ($V:V=6:4$), then CH_2Cl_2 -MeOH ($V:V=98:2$)) to afford (9*H*-fluoren-9-yl)methyl-((*S*)-2,2-dimethyl-1-((*R*)-4-phenyl-4,5-dihydrooxazol-2-yl)propyl)carbamate (**6e**),^[15] white solid, 86% yield (1.95 g, 4.3 mmol). m.p. 126~129 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.77 (d, $J=7.6$ Hz, 2H), 7.63 (d, $J=7.6$ Hz, 2H), 7.43~7.28 (m, 9H), 5.77 (br s, 1H), 5.24 (t, $J=9.6$ Hz, 1H), 4.68 (t, $J=9.4$ Hz, 1H), 4.48~4.40 (m, 3H), 4.24 (t, $J=7.0$ Hz, 1H), 4.08 (t, $J=8.8$ Hz, 1H), 1.08 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ : 167.8, 156.3, 144.1, 143.9, 141.8, 141.41, 141.39, 128.9, 127.81, 127.75, 127.1, 126.8, 125.2, 120.0, 74.9, 69.7, 67.0, 57.7, 47.4, 35.4, 26.7; IR (KBr) ν_{max} : 3406, 1716, 1656, 1622, 1398, 1369, 1322, 1239 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{29}\text{H}_{31}\text{N}_2\text{O}_3$ ($\text{M}+\text{H}^+$) 455.2329, found 454.2322.

4.2.4 General procedure for the synthesis of **7**

Fmoc-protected oxazoline **6** (6 mmol) was dissolved in MeOH (30 mL) at room temperature. Upon cooling to 0 °C and addition of Et_2NH (30 mL), the reaction mixture was stirred for 1 h at 0~5 °C. After completion of the reaction, the solvent was removed under reduced pressure and the residue was purified by silica gel column chromatography (EtOAc/petroleum ether, $V:V=1:1$ (100 mL) then EtOAc/EtOH, $V:V=10:1$) to furnish (*S*)-2-methyl-

1-((*R*)-4-phenyl-4,5-dihydrooxazol-2-yl)-propan-1-amine (**7a**),^[15] yellow oil, 96% yield (1.26 g, 5.8 mmol). ^1H NMR (400 MHz, CDCl_3) δ : 7.37~7.24 (m, 5H), 5.21 (t, $J=9.6$ Hz, 1H), 4.66 (dd, $J=10.0, 8.8$ Hz, 1H), 4.12 (t, $J=8.4$ Hz, 1H), 3.45 (d, $J=5.6$ Hz, 1H), 2.12~2.04 (m, 1H), 1.04 (d, $J=6.8$ Hz, 3H), 1.01 (d, $J=6.8$ Hz, 3H).

(*S*)-1-((*R*)-4-Isopropyl-4,5-dihydrooxazol-2-yl)-2-methylpropan-1-amine (**7b**):^[14] Yellow oil, 64% yield (710 mg, 3.8 mmol). ^1H NMR (400 MHz, CDCl_3) δ : 4.24 (dd, $J=9.2, 8.0$ Hz, 1H), 3.98~3.87 (m, 2H), 3.31 (d, $J=5.6$ Hz, 1H), 1.99~1.91 (m, 1H), 1.79~1.71 (m, 1H), 0.97~0.92 (m, 9H), 0.88 (d, $J=6.8$ Hz, 3H).

(*S*)-1-((*R*)-4-Benzyl-4,5-dihydrooxazol-2-yl)-2-methylpropan-1-amine (**7c**):^[19] Yellow oil, 82% yield (1.14 g, 4.9 mmol). ^1H NMR (400 MHz, CDCl_3) δ : 7.31~7.19 (m, 5H), 4.44~4.36 (m, 1H), 4.20 (t, $J=9.0$ Hz, 1H), 3.99 (dd, $J=9.2, 8.4$ Hz, 1H), 3.31 (d, $J=5.6$ Hz, 1H), 3.11 (dd, $J=14.0, 5.2$ Hz, 1H), 2.64 (dd, $J=14.0, 8.8$ Hz, 1H), 1.99~1.90 (m, 1H), 0.94 (dd, $J=10.8, 6.8$ Hz, 6H).

(*S*)-1-(4,4-Dimethyl-4,5-dihydrooxazol-2-yl)-2-methylpropan-1-amine (**7d**): Yellow oil, 76% yield (780 mg, 4.56 mmol). ^1H NMR (400 MHz, CDCl_3) δ : 3.87 (d, $J=5.6$ Hz, 2H), 3.23~3.21 (m, 1H), 1.92~1.85 (m, 1H), 1.53 (s, 2H), 1.21 (d, $J=4.4$ Hz, 6H), 0.91~0.85 (m, 6H); ^{13}C NMR (100 MHz, CDCl_3) δ : 168.1, 79.2, 66.7, 55.4, 32.2, 28.4, 28.3, 19.3, 17.5; IR (KBr) ν_{max} : 3386, 2964, 1662, 1464, 1392, 1366, 992, 669 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_9\text{H}_{19}\text{N}_2\text{O}$ ($\text{M}+\text{H}^+$) 171.1492, found 171.1490.

(*S*)-2,2-Dimethyl-1-((*R*)-4-phenyl-4,5-dihydrooxazol-2-yl)propan-1-amine (**7e**):^[20] Yellow oil, 92% yield (1.28 g, 5.5 mmol). ^1H NMR (400 MHz, CDCl_3) δ : 7.37~7.25 (m, 5H), 5.20 (t, $J=9.6$ Hz, 1H), 4.65 (dd, $J=10.4, 8.8$ Hz, 1H), 4.06 (t, $J=8.8$ Hz, 1H), 3.37 (s, 1H), 1.05 (s, 9H).

4.2.5 General procedure for the synthesis of **8**

To a solution of 2-diphenylphosphine benzaldehyde (4 mmol) in anhydrous tetrahydrofuran (THF) (50 mL) was added tetraethyl titanate (8 mmol). The resulting solution was stirred at 0 °C. Compound **7** (4 mmol) in anhydrous THF (15 mL) was added by syringe. After stirring at 0 °C for 2 h, the mixture was diluted with EtOAc, then washed with saturated sodium chloride solution and filtered with celite. The aqueous layer was extracted with EtOAc (40 mL×3). The organic layers were combined and dried over MgSO_4 . After filtration and concentration, the resulting crude compound **8** was obtained without further purification.

4.2.6 Reduction by NaBH_4 for the synthesis of **1a** and **1g**

To a solution of crude compound **8** (4 mmol) in MeOH (30 mL) was added NaBH_4 (14 mmol) at 0 °C. The reaction mixture was stirred at 0 °C until the disappearance of **8** (monitored by TLC). Then, the reaction was quenched with a small amount of saturated ammonium chloride solution, and methanol was removed under reduced pressure. The residue was purified by flash column chromatography (petroleum ether/ Et_2O , $V:V=10:1$) to afford product **1a**

or **1g**.

(*S*)-*N*-(2-(Diphenylphosphanyl)benzyl)-2-methyl-1-((*R*)-4-phenyl-4,5-dihydrooxazol-2-yl)propan-1-amine (**1a**): Colorless oil, 77% yield (1.52 g, 3.1 mmol). ^1H NMR (400 MHz, CDCl_3) δ : 7.65~7.62 (m, 1H), 7.40~7.26 (m, 17H), 7.18 (t, $J=7.6$ Hz, 1H), 6.92~6.89 (m, 1H), 5.19 (t, $J=9.6$ Hz, 1H), 4.62 (dd, $J=8.8$, 8.4 Hz, 1H), 4.18 (dd, $J=1.6$, 1.6 Hz, 1H), 4.05 (t, $J=9.2$ Hz, 1H), 4.02 (dd, $J=1.6$, 1.6 Hz, 1H), 3.25 (d, $J=6.8$ Hz, 1H), 1.96~1.88 (m, 1H), 1.01 (d, $J=6.8$ Hz, 3H), 0.98 (d, $J=6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 169.4, 144.5 ($J_{\text{c,p}}=23.6$ Hz), 142.2, 137.0 ($J_{\text{c,p}}=4.4$ Hz), 136.9 ($J_{\text{c,p}}=4.6$ Hz), 135.7 ($J_{\text{c,p}}=14.0$ Hz), 134.0, 133.9 ($J_{\text{c,p}}=12.9$ Hz), 133.7, 133.6, 129.0, 128.9 ($J_{\text{c,p}}=5.4$ Hz), 128.8, 128.63 ($J_{\text{c,p}}=5.7$ Hz), 128.57 ($J_{\text{c,p}}=1.7$ Hz), 128.50 ($J_{\text{c,p}}=1.0$ Hz), 127.5, 127.2, 126.9, 74.6, 69.7, 62.0, 50.5 ($J_{\text{c,p}}=22.3$ Hz), 31.7, 19.4, 19.3; ^{31}P NMR (162 MHz, CDCl_3) δ : -15.67; IR (KBr) ν_{max} : 3428, 2961, 1656, 1434, 983, 745, 698, 504 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{32}\text{H}_{34}\text{N}_2\text{OP}$ ($\text{M}+\text{H}^+$) 493.2403, found 493.2424.

(*S*)-*N*-(2-(Diphenylphosphanyl)benzyl)-2,2-dimethyl-1-((*R*)-4-phenyl-4,5-dihydrooxazol-2-yl)propan-1-amine (**1g**): Colorless oil, 79% yield (1.60 g, 3.2 mmol). ^1H NMR (400 MHz, CDCl_3) δ : 7.67~7.64 (m, 1H), 7.40~7.27 (m, 17H), 7.20~7.16 (m, 1H), 6.92~6.89 (m, 1H), 5.18 (t, $J=10.0$ Hz, 1H), 4.62 (t, $J=9.4$ Hz, 1H), 4.17~3.97 (m, 3H), 3.21 (s, 1H), 1.02 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ : 169.4, 144.7 ($J_{\text{c,p}}=23.7$ Hz), 142.2, 137.2, 137.1, 135.9 ($J_{\text{c,p}}=14.1$ Hz), 134.1, 134.0 ($J_{\text{c,p}}=12.3$ Hz), 133.9, 133.8, 129.0, 128.9 ($J_{\text{c,p}}=5.3$ Hz), 128.8, 128.7, 128.6 ($J_{\text{c,p}}=2.5$ Hz), 128.5 ($J_{\text{c,p}}=2.5$ Hz), 127.5, 127.2, 127.0, 74.5, 69.8, 65.4, 51.0 ($J_{\text{c,p}}=3.0$ Hz), 34.3, 27.0; ^{31}P NMR (162 MHz, CDCl_3) δ : -15.60; IR (KBr) ν_{max} : 3406, 2957, 1652, 1434, 1395, 1365, 744, 698 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{33}\text{H}_{36}\text{N}_2\text{OP}$ ($\text{M}+\text{H}^+$) 507.2560, found 507.2565.

4.2.7 Addition of Grignard reagents for the synthesis of **1b**~**1f** and **1h**

Crude product **8** (ca. 4 mmol) and anhydrous dichloromethane (20 mL) were added to a 50 mL oven-dried reaction Schlenk tube. The mixture was cooled to -70 $^{\circ}\text{C}$, and Grignard reagent (16 mmol, 3 mol/L in THF) was added. After stirring at -70 $^{\circ}\text{C}$ for 1 h, the reaction mixture was allowed to warm up to ambient temperature. The reaction was quenched with a small amount of saturated ammonium chloride solution, and dichloromethane was removed under reduced pressure. The residue was purified by flash column chromatography (petroleum ether/ Et_2O , $V:V=10:1$) to afford product **1b**~**1f** or **1h**.

(*S*)-*N*-((*R*)-2-(Diphenylphosphanyl)phenyl)(phenyl)methyl-2-methyl-1-((*R*)-4-phenyl-4,5-dihydrooxazol-2-yl)propan-1-amine (**1b**): White solid, 70% yield (1.59 g, 2.8 mmol). m.p. 53~55 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 8.01~7.98 (m, 1H), 7.41~7.12 (m, 18H), 7.04~6.92 (m, 6H), 6.10 (d, $J=8.8$ Hz, 1H), 5.24 (t, $J=9.6$ Hz, 1H), 4.55 (t, $J=9.2$ Hz, 1H), 4.01 (t, $J=8.8$ Hz, 1H), 3.14 (d,

$J=6.0$ Hz, 1H), 2.02~1.94 (m, 1H), 1.07 (d, $J=6.4$ Hz, 3H), 1.02 (d, $J=6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.1, 148.0 ($J_{\text{c,p}}=23.1$ Hz), 143.9, 142.5, 138.3 ($J_{\text{c,p}}=11.8$ Hz), 136.3 ($J_{\text{c,p}}=2.3$ Hz), 136.1 ($J_{\text{c,p}}=13.2$ Hz), 134.7 ($J_{\text{c,p}}=1.4$ Hz), 134.1 ($J_{\text{c,p}}=20.1$ Hz), 133.6 ($J_{\text{c,p}}=18.9$ Hz), 129.4, 128.8, 128.5 ($J_{\text{c,p}}=6.1$ Hz), 128.4, 128.3, 128.2 ($J_{\text{c,p}}=1.3$ Hz), 128.2 ($J_{\text{c,p}}=1.1$ Hz), 128.1, 127.9 ($J_{\text{c,p}}=5.4$ Hz), 127.5, 127.2, 127.0 ($J_{\text{c,p}}=1.3$ Hz), 126.7, 74.8, 70.1, 61.3 ($J_{\text{c,p}}=27.1$ Hz), 59.8, 32.3, 27.0, 19.8, 19.0; ^{31}P NMR (162 MHz, CDCl_3) δ : -18.50; IR (KBr) ν_{max} : 3427, 2964, 1655, 1454, 1434, 1089, 744, 697 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{38}\text{H}_{38}\text{N}_2\text{OP}$ ($\text{M}+\text{H}^+$) 569.2716, found 569.2693.

(*R*)-1-(2-(Diphenylphosphanyl)phenyl)-2-methyl-*N*-((*S*)-2-methyl-1-((*R*)-4-phenyl-4,5-dihydrooxazol-2-yl)propyl)propan-1-amine (**1c**): Colorless oil, 66% yield (1.41 g, 2.6 mmol). ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 7.69 (s, 1H), 7.40~7.11 (m, 17H), 6.97~6.94 (m, 1H), 5.15 (t, $J=9.6$ Hz, 1H), 4.74 (s, 1H), 4.47 (t, $J=9.2$ Hz, 1H), 3.81 (t, $J=8.5$ Hz, 1H), 2.69 (d, $J=4.8$ Hz, 1H), 1.82~1.73 (m, 2H), 0.91 (d, $J=6.4$ Hz, 3H), 0.83 (d, $J=6.4$ Hz, 6H), 0.59 (d, $J=6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.5, 149.4 ($J_{\text{c,p}}=23.4$ Hz), 142.8, 138.9 ($J_{\text{c,p}}=12.5$ Hz), 137.3 ($J_{\text{c,p}}=11.7$ Hz), 136.5 ($J_{\text{c,p}}=13.1$ Hz), 135.0, 134.3 ($J_{\text{c,p}}=20.2$ Hz), 133.3 ($J_{\text{c,p}}=18.7$ Hz), 129.0, 128.7, 128.6, 128.5, 128.45 ($J_{\text{c,p}}=2.0$ Hz), 128.38, 128.0, 127.4, 127.0 ($J_{\text{c,p}}=1.3$ Hz), 126.7, 74.5, 70.0, 59.3, 34.8, 32.5, 27.1, 20.4, 19.8, 19.0, 17.8; ^{31}P NMR (162 MHz, CDCl_3) δ : -19.96; IR (KBr) ν_{max} : 2959, 1655, 1625, 1434, 1387, 743, 697, 669 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{35}\text{H}_{40}\text{N}_2\text{OP}$ ($\text{M}+\text{H}^+$) 535.2873, found 535.2882.

(*S*)-*N*-((*R*)-2-(Diphenylphosphanyl)phenyl)(phenyl)methyl-1-((*R*)-4-isopropyl-4,5-dihydrooxazol-2-yl)-2-methylpropan-1-amine (**1d**): Colorless oil, 75% yield (1.60 g, 3.0 mmol). ^1H NMR (400 MHz, CDCl_3) δ : 8.03~8.00 (m, 1H), 7.42~7.38 (m, 1H), 7.32~7.11 (m, 12H), 7.06~6.92 (m, 6H), 6.00 (d, $J=8.8$ Hz, 1H), 4.20 (dd, $J=13.6$, 12.0 Hz, 1H), 3.97~3.90 (m, 2H), 3.06 (d, $J=6.4$ Hz, 1H), 1.94~1.87 (m, 1H), 1.86~1.77 (m, 1H), 1.03 (t, $J=6.0$ Hz, 6H), 0.97 (d, $J=6.8$ Hz, 3H), 0.92 (d, $J=6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 168.3, 148.1 ($J_{\text{c,p}}=23.2$ Hz), 144.1, 138.4 ($J_{\text{c,p}}=12.2$ Hz), 136.4 ($J_{\text{c,p}}=10.8$ Hz), 136.2 ($J_{\text{c,p}}=14.7$ Hz), 134.8 ($J_{\text{c,p}}=1.5$ Hz), 134.0 ($J_{\text{c,p}}=20.0$ Hz), 133.5 ($J_{\text{c,p}}=19.0$ Hz), 129.4, 128.5 ($J_{\text{c,p}}=6.2$ Hz), 128.33 ($J_{\text{c,p}}=1.6$ Hz), 128.31, 128.24, 128.18, 128.07, 127.9 ($J_{\text{c,p}}=5.7$ Hz), 127.2, 126.6, 72.6, 70.0, 61.3 ($J_{\text{c,p}}=26.7$ Hz), 59.9, 32.8, 32.2, 19.7, 19.6, 19.0, 18.6; ^{31}P NMR (162 MHz, CDCl_3) δ : -18.52; IR (KBr) ν_{max} : 2989, 1662, 1625, 1434, 1386, 1365, 743, 696 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{35}\text{H}_{40}\text{N}_2\text{OP}$ ($\text{M}+\text{H}^+$) 535.2873, found 535.2881.

(*S*)-1-((*R*)-4-Benzyl-4,5-dihydrooxazol-2-yl)-*N*-((*R*)-2-(diphenylphosphanyl)phenyl)(phenyl)methyl-2-methylpropan-1-amine (**1e**): White solid, 69% yield (1.61 g, 2.8 mmol). m.p. 50~52 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.98~7.96 (m, 1H), 7.41~7.38 (m, 1H), 7.32~7.01 (m, 22H), 6.95~6.92 (m, 1H), 6.04 (d, $J=8.8$ Hz, 1H), 4.48~

4.40 (m, 1H), 4.14 (t, $J=9.0$ Hz, 1H), 3.94 (t, $J=8.0$ Hz, 1H), 3.19 (dd, $J=5.2, 4.8$ Hz, 1H), 3.07 (d, $J=6.8$ Hz, 1H), 2.65 (dd, $J=10.0, 9.6$ Hz, 1H), 1.93~1.85 (m, 1H), 1.03 (d, $J=6.8$ Hz, 3H), 0.94 (d, $J=6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 169.3, 148.0 ($J_{\text{c,p}}=23.2$ Hz), 144.0, 138.4, 138.2 ($J_{\text{c,p}}=11.8$ Hz), 136.3 ($J_{\text{c,p}}=10.4$ Hz), 136.1 ($J_{\text{c,p}}=14.2$ Hz), 134.8 ($J_{\text{c,p}}=1.4$ Hz), 134.0 ($J_{\text{c,p}}=19.9$ Hz), 133.5 ($J_{\text{c,p}}=18.9$ Hz), 129.5 ($J_{\text{c,p}}=5.5$ Hz), 129.3, 128.6, 128.5 ($J_{\text{c,p}}=6.2$ Hz), 128.4, 128.3, 128.23, 128.19 ($J_{\text{c,p}}=1.4$ Hz), 128.1, 127.8 ($J_{\text{c,p}}=5.4$ Hz), 127.2, 126.7, 126.5, 71.9, 67.4, 61.3 ($J_{\text{c,p}}=27.2$ Hz), 59.9, 42.1 ($J_{\text{c,p}}=1.3$ Hz), 32.1, 19.6, 19.1; ^{31}P NMR (162 MHz, CDCl_3) δ : -18.51; IR (KBr) ν_{max} : 2960, 1656, 1454, 1434, 1396, 1231, 744, 696 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{39}\text{H}_{40}\text{N}_2\text{OP}$ ($\text{M}+\text{H}^+$) 583.2873, found 593.2847.

(*S*)-1-(4,4-Dimethyl-4,5-dihydrooxazol-2-yl)-*N*-((*R*)-(2-(diphenylphosphanyl)phenyl)(phenyl)methyl)-2-methylpropan-1-amine (**1f**): Colorless oil, 80% yield (1.67 g, 3.2 mmol); ^1H NMR (400 MHz, CDCl_3) δ : 8.04~8.01 (m, 1H), 7.42~7.38 (m, 1H), 7.32~7.08 (m, 12H), 7.05~6.91 (m, 6H), 5.98 (d, $J=9.2$ Hz, 1H), 3.94 (d, $J=8.0$ Hz, 1H), 3.90 (d, $J=7.6$ Hz, 1H), 3.01 (d, $J=6.8$ Hz, 1H), 1.91~1.93 (m, 1H), 1.37 (s, 3H), 1.33 (s, 3H), 1.06 (d, $J=6.8$ Hz, 3H), 0.96 (d, $J=6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 167.4, 147.9 ($J_{\text{c,p}}=23.2$ Hz), 143.9, 138.4 ($J_{\text{c,p}}=12.1$ Hz), 136.3 ($J_{\text{c,p}}=10.6$ Hz), 136.1 ($J_{\text{c,p}}=14.5$ Hz), 134.9 ($J_{\text{c,p}}=1.6$ Hz), 134.0 ($J_{\text{c,p}}=19.9$ Hz), 133.5 ($J_{\text{c,p}}=18.8$ Hz), 129.4, 128.5 ($J_{\text{c,p}}=6.1$ Hz), 128.35 ($J_{\text{c,p}}=1.3$ Hz), 128.30, 128.23, 128.16, 128.07, 127.7 ($J_{\text{c,p}}=5.7$ Hz), 127.2, 126.7, 79.1, 67.2, 61.5 ($J_{\text{c,p}}=27.4$ Hz), 59.9, 32.3, 28.81, 28.79, 19.6, 19.2; ^{31}P NMR (162 MHz, CDCl_3) δ : -18.87; IR (KBr) ν_{max} : 3406, 1625, 1434, 1397, 1365, 1230, 743, 695 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{34}\text{H}_{38}\text{N}_2\text{OP}$ ($\text{M}+\text{H}^+$) 521.2716, found 521.2718.

(*S*)-*N*-((*R*)-(2-(Diphenylphosphanyl)phenyl)(phenyl)methyl)-2,2-dimethyl-1-((*R*)-4-phenyl-4,5-dihydrooxazol-2-yl)propan-1-amine (**1h**): White solid, 73% yield (1.70 g, 2.9 mmol). m.p. 58~61 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 8.07~8.04 (m, 1H), 7.44~7.13 (m, 18H), 7.07~6.97 (m, 6H), 6.11 (d, $J=9.6$ Hz, 1H), 5.39 (t, $J=9.8$ Hz, 1H), 4.63 (dd, $J=10.4, 8.4$ Hz, 1H), 4.03 (t, $J=9.0$ Hz, 1H), 3.12 (s, 1H), 1.11 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ : 169.9, 147.9 ($J_{\text{c,p}}=23.2$ Hz), 144.0, 142.5, 138.5 ($J_{\text{c,p}}=12.0$ Hz), 136.3 ($J_{\text{c,p}}=14.2$ Hz), 136.1 ($J_{\text{c,p}}=9.8$ Hz), 134.8, 134.1 ($J_{\text{c,p}}=20.1$ Hz), 133.5 ($J_{\text{c,p}}=18.6$ Hz), 129.4, 128.8, 128.5 ($J_{\text{c,p}}=5.9$ Hz), 128.4, 128.3, 128.23, 128.20, 128.1, 128.0, 127.5, 127.2, 127.1 ($J_{\text{c,p}}=1.3$ Hz), 126.6, 74.6, 70.4, 62.9, 61.3 ($J_{\text{c,p}}=28.0$ Hz), 34.7, 27.3; ^{31}P NMR (162 MHz, CDCl_3) δ : -18.92; IR (KBr) ν_{max} : 3409, 3027, 2958, 1625, 1434, 1396, 1232, 696 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{39}\text{H}_{40}\text{N}_2\text{OP}$ ($\text{M}+\text{H}^+$) 583.2873, found 583.2847.

4.3 Catalytic asymmetric allylic alkylation

In a dried schlenk tube, ligand **1b** (0.003 mmol, 3 mol%) and $[\text{Pd}(\eta^3\text{-C}_3\text{H}_5)\text{Cl}]_2$ (0.001 mmol, 1 mol%) were dissolved in anhydrous dichloromethane under N_2

atmosphere, and the mixture was stirred for 30 min. Then, **9** (0.1 mmol, 1.0 equiv.), **10** (0.12 mmol, 1.2 equiv.) and *N,O*-bis(trimethylsilyl)acetamide (BSA) (0.2 mmol, 2.0 equiv.) were added sequentially. The reaction was stirred at room temperature. After completion (monitored by TLC plate), the residue was purified by column chromatography to get the desired product **11**.

Dimethyl (*R,E*)-2-(1,3-diphenylallyl)malonate (**11aa**):^[21] Yellow oil, 97% yield (31.5 mg, 0.097 mmol). ^1H NMR (400 MHz, CDCl_3) δ : 7.32~7.26 (m, 8H), 7.23~7.18 (m, 2H), 6.48 (d, $J=15.6$ Hz, 1H), 6.34 (dd, $J=16.2, 9.0$ Hz, 1H), 4.27 (t, $J=9.6$ Hz, 1H), 3.95 (d, $J=11.4$ Hz, 1H), 3.70 (s, 3H), 3.52 (s, 3H). HPLC: AD-H, hexane/isopropanol ($V:V=95:5$), 1.0 mL/min, $\lambda=254$ nm, 18 $^{\circ}\text{C}$, t_{R1} (major)=18.0 min, t_{R2} (minor)=27.1 min, 95% ee.

Dimethyl (*R,E*)-2-(1,3-bis(4-fluorophenyl)allyl)malonate (**11ba**):^[21] Yellow oil, 86% yield (31.0 mg, 0.086 mmol). ^1H NMR (400 MHz, CDCl_3) δ : 7.29~7.25 (m, 4H), 7.03~6.95 (m, 4H), 6.43 (d, $J=15.6$ Hz, 1H), 6.25 (dd, $J=16.0, 8.8$ Hz, 1H), 4.27 (t, $J=9.4$ Hz, 1H), 3.90 (d, $J=10.8$ Hz, 1H), 3.71 (s, 3H), 3.54 (s, 3H). HPLC: AD-H, hexane/isopropanol ($V:V=95:5$), 1.0 mL/min, $\lambda=254$ nm, 22 $^{\circ}\text{C}$, t_{R1} (major)=20.8 min, t_{R2} (minor)=35.0 min, 96% ee.

Dimethyl (*R,E*)-2-(1,3-bis(4-chlorophenyl)allyl)malonate (**11ca**):^[21] Yellow oil, 68% yield (26.7 mg, 0.068 mmol). ^1H NMR (400 MHz, CDCl_3) δ : 7.33~7.27 (m, 5H), 7.25 (t, $J=7.6$ Hz, 3H), 6.44 (d, $J=16.0$ Hz, 1H), 6.31 (dd, $J=16.0, 8.4$ Hz, 1H), 4.28~4.23 (m, 1H), 3.92 (d, $J=10.8$ Hz, 1H), 3.72 (s, 3H), 3.57 (s, 3H). HPLC: AD-H, hexane/isopropanol ($V:V=80:20$), 1.0 mL/min, $\lambda=254$ nm, 20 $^{\circ}\text{C}$, t_{R1} (major)=13.7 min, t_{R2} (minor)=21.1 min, 92% ee.

Dimethyl (*R,E*)-2-(1,3-bis(4-bromophenyl)allyl)malonate (**11da**):^[21] Yellow oil, 89% yield (42.9 mg, 0.089 mmol). ^1H NMR (400 MHz, CDCl_3) δ : 7.46 (d, $J=8.4$ Hz, 2H), 7.41 (d, $J=8.8$ Hz, 2H), 7.18 (d, $J=8.4$ Hz, 4H), 6.40 (d, $J=16.0$ Hz, 1H), 6.30 (dd, $J=16.0, 8.4$ Hz, 1H), 4.24~4.19 (m, 1H), 3.90 (d, $J=10.8$ Hz, 1H), 3.70 (s, 3H), 3.55 (s, 3H). HPLC: AD-H, hexane/isopropanol ($V:V=80:20$), 1.0 mL/min, $\lambda=254$ nm, 23 $^{\circ}\text{C}$, t_{R1} (major)=16.8 min, t_{R2} (minor)=25.2 min, 92% ee.

Dimethyl (*R,E*)-2-(1,3-di-*p*-tolylallyl)malonate (**11ea**):^[21] Yellow oil, 75% yield (26.4 mg, 0.075 mmol). ^1H NMR (400 MHz, CDCl_3) δ : 7.21~7.16 (m, 4H), 7.12~7.06 (m, 4H), 6.44 (d, $J=15.6$ Hz, 1H), 6.28~6.22 (m, 1H), 4.24~4.19 (m, 1H), 3.93 (d, $J=10.8$ Hz, 1H), 3.69 (s, 3H), 3.53 (s, 3H), 2.30 (s, 6H). HPLC: AD-H, hexane/isopropanol ($V:V=80:20$), 1.0 mL/min, $\lambda=254$ nm, 28 $^{\circ}\text{C}$, t_{R1} (major)=7.7 min, t_{R2} (minor)=9.9 min, 91% ee.

Dimethyl (*R,E*)-2-(1,3-bis(3-chlorophenyl)allyl)malonate (**11fa**):^[22] Yellow oil, 79% yield (31.1 mg, 0.079 mmol). ^1H NMR (400 MHz, CDCl_3) δ : 7.31~7.27 (m, 2H), 7.24~7.16 (m, 6H), 6.44 (d, $J=15.6$ Hz, 1H), 6.31 (dd, $J=15.6, 8.4$ Hz, 1H), 4.26~4.21 (m, 1H), 3.92 (d, $J=10.8$ Hz, 1H), 3.71 (s, 3H), 3.57 (s, 3H). HPLC: AD-H,

hexane/isopropanol ($V: V=80:20$), 1.0 mL/min, $\lambda=254$ nm, 24 °C, $t_{R1}(\text{major})=6.8$ min, $t_{R2}(\text{minor})=8.9$ min, 90% *ee*.

Dimethyl (*R,E*)-2-(1,3-di-*o*-tolylallyl)malonate (**11ga**):^[23] Yellow oil, 84% yield (29.6 mg, 0.084 mmol). ¹H NMR (400 MHz, CDCl₃) δ : 7.37~7.28 (m, 1H), 7.23~7.09 (m, 7H), 6.67 (d, $J=15.6$ Hz, 1H), 6.04 (dd, $J=15.2, 8.4$ Hz, 1H), 4.58~4.53 (m, 1H), 4.08 (d, $J=11.2$ Hz, 1H), 3.73 (s, 3H), 3.52 (s, 3H), 2.47 (s, 3H), 2.27 (s, 3H). HPLC: AD-H, hexane/isopropanol ($V: V=98:2$), 1.0 mL/min, $\lambda=254$ nm, 26 °C, $t_{R1}(\text{major})=19.2$ min, $t_{R2}(\text{minor})=20.8$ min, 61% *ee*.

Dimethyl (*R,E*)-2-(1,3-di(naphthalen-2-yl)allyl)malonate (**11ha**):^[22] White solid, 97% yield (41.2 mg, 0.097 mmol). ¹H NMR (400 MHz, CDCl₃) δ : 7.84~7.68 (m, 8H), 7.55~7.40 (m, 6H), 6.70 (d, $J=16.0$ Hz, 1H), 6.57 (dd, $J=15.6, 8.4$ Hz, 1H), 4.54~4.49 (m, 1H), 4.15 (d, $J=10.8$ Hz, 1H), 3.73 (s, 3H), 3.50 (s, 3H). HPLC: AD-H, hexane/isopropanol ($V: V=80:20$), 1.0 mL/min, $\lambda=254$ nm, 28 °C, $t_{R1}(\text{major})=31.7$ min, $t_{R2}(\text{minor})=42.0$ min, 92% *ee*.

Diisopropyl (*R,E*)-2-(1,3-diphenylallyl)malonate (**11ab**):^[21] Yellow oil, 51% yield (19.4 mg, 0.051 mmol). ¹H NMR (400 MHz, CDCl₃) δ : 7.30~7.17 (m, 10H), 6.48 (d, $J=16.0$ Hz, 1H), 6.36 (dd, $J=15.6, 8.4$ Hz, 1H), 5.08~5.00 (m, 1H), 4.85~4.79 (m, 1H), 4.26~4.22 (m, 1H), 3.87 (d, $J=11.2$ Hz, 1H), 1.22 (d, $J=6.0$ Hz, 3H), 1.17 (d, $J=6.4$ Hz, 3H), 1.06 (d, $J=6.4$ Hz, 3H), 0.97 (d, $J=6.4$ Hz, 3H). HPLC: AD-H, hexane/isopropanol ($V: V=95:5$), 1.0 mL/min, $\lambda=254$ nm, 25 °C, $t_{R1}(\text{major})=11.1$ min, $t_{R2}(\text{minor})=14.7$ min, 88% *ee*.

Dibenzyl (*R,E*)-2-(1,3-diphenylallyl)malonate (**11ac**):^[24] Yellow oil, 97% yield (46.2 mg, 0.097 mmol). ¹H NMR (400 MHz, CDCl₃) δ : 7.34~7.33 (m, 2H), 7.29~7.18 (m, 16H), 7.05~7.03 (m, 2H), 6.43 (d, $J=15.6$ Hz, 1H), 6.33 (dd, $J=15.6, 8.4$ Hz, 1H), 5.14~5.07 (m, 2H), 4.96~4.89 (m, 2H), 4.32~4.27 (m, 1H), 4.05 (d, $J=10.8$ Hz, 1H). HPLC: AD-H, hexane/isopropanol ($V: V=95:5$), 1.0 mL/min, $\lambda=254$ nm, 28 °C, $t_{R1}(\text{major})=38.5$ min, $t_{R2}(\text{minor})=48.0$ min, 94% *ee*.

Diethyl (*S,E*)-2-(1,3-diphenylallyl)-2-phenylmalonate (**11ad**):^[21] Yellow oil, 86% yield (36.9 mg, 0.086 mmol). ¹H NMR (400 MHz, CDCl₃) δ : 7.39~7.12 (m, 10H), 7.15~7.11 (m, 3H), 6.70~6.98 (m, 2H), 6.51~6.39 (m, 2H), 4.65 (t, $J=7.6$ Hz, 1H), 4.25~4.13 (m, 4H), 1.20 (m, 6H). HPLC: AD-H, hexane/isopropanol ($V: V=99:1$), 0.5 mL/min, $\lambda=254$ nm, 23 °C, $t_{R1}(\text{major})=29.5$ min, $t_{R2}(\text{minor})=32.0$ min, 92% *ee*.

Diethyl (*S,E*)-2-(1,3-diphenylallyl)-2-methylmalonate (**11ae**):^[21] Yellow oil, 48% yield (17.6 mg, 0.048 mmol). ¹H NMR (600 MHz, CDCl₃) δ : 7.34~7.32 (m, 4H), 7.30~7.27 (m, 3H), 7.24~7.18 (m, 2H), 6.72 (dd, $J=16.2, 9.0$ Hz, 1H), 6.45 (d, $J=15.6$ Hz, 1H), 4.30 (d, $J=9.0$ Hz, 1H), 4.22~4.19 (m, 2H), 4.10~4.05 (m, 2H), 3.44 (dd, $J=14.4, 7.2$ Hz, 1H), 1.47 (s, 3H), 1.24 (t, $J=7.2$ Hz, 3H), 1.18 (t, $J=7.2$ Hz, 3H). HPLC: AD-H, hexane/isopropanol ($V: V=98:2$), 1.0 mL/min, $\lambda=254$ nm,

25 °C, $t_{R1}(\text{minor})=17.8$ min, $t_{R2}(\text{major})=18.9$ min, 85% *ee*.

(*R,E*)-3-(1,3-Diphenylallyl) pentane-2,4-dione (**11af**):^[21] Yellow oil, 75% yield (21.9 mg, 0.075 mmol). ¹H NMR (400 MHz, CDCl₃) δ : 7.32~7.25 (m, 6H), 7.24~7.20 (m, 4H), 6.45 (d, $J=16.0$ Hz, 1H), 6.22~6.16 (m, 1H), 4.37~4.30 (m, 2H), 2.54 (s, 3H), 1.93 (s, 3H). HPLC: AD-H, hexane/isopropanol ($V: V=95:5$), 1.0 mL/min, $\lambda=254$ nm, 23 °C, $t_{R1}(\text{major})=11.0$ min, $t_{R2}(\text{minor})=11.9$ min, 90% *ee*.

(*R,E*)-2-(1,3-Diphenylallyl) malononitrile (**11ag**):^[25] Yellow oil, 21% yield (5.4 mg, 0.021 mmol). ¹H NMR (400 MHz, CDCl₃) δ : 7.51~7.32 (m, 10H), 6.69~6.59 (m, 1H), 6.50 (d, $J=15.6$ Hz, 1H), 4.05~3.91 (m, 1H), 3.79 (d, $J=9.2$ Hz, 1H). HPLC: OJ-H, hexane/isopropanol ($V: V=80:20$), 1.0 mL/min, $\lambda=254$ nm, 26 °C, $t_{R1}(\text{major})=27.2$ min, $t_{R2}(\text{minor})=31.7$ min, 74% *ee*.

Supporting Information HPLC chromatographic data for chiral products and NMR spectra of new compounds. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] For selected books and reviews on chiral pincer ligands, see: (a) Liu, J.-K.; Gong, J.-F.; Song, M.-P. *Org. Biomol. Chem.* **2019**, *17*, 6069. (b) Morales-Morales, D. *Pincer Compounds: Chemistry and Applications*, Elsevier, Amsterdam, **2018**. (c) Asay, M.; Morales-Morales, D. *Dalton Trans.* **2015**, *44*, 17432. (d) Deng, Q.-H.; Melen, R. L.; Gade, L. H. *Acc. Chem. Res.* **2014**, *47*, 3162. (e) Szabó, K. L.; Wendt, O. F. *Pincer and Pincer-Type Complexes*, Wiley-VCH, Weinheim, **2014**.
- [2] (a) Hua, Y.-Y.; Bin, H.-Y.; Wei, T.; Cheng, H.-A.; Lin, Z.-P.; Fu, X.-F.; Li, Y.-Q.; Xie, J.-H.; Yan, P.-C.; Zhou, Q.-L. *Org. Lett.* **2020**, *22*, 818. (b) Gu, X.-S.; Yu, N.; Yang, X.-H.; Zhu, A.-T.; Xie, J.-H.; Zhou, Q.-L. *Org. Lett.* **2019**, *21*, 4111. (c) Chen, G.-Q.; Lin, B.-J.; Huang, J.-M.; Zhao, L.-Y.; Chen, Q.-S.; Jia, S.-P.; Yin, Q.; Zhang, X. J. *Am. Chem. Soc.* **2018**, *140*, 8064. (d) Zhang, Y.-M.; Yuan, M.-L.; Liu, W.-P.; Xie, J.-H.; Zhou, Q.-L. *Org. Lett.* **2018**, *20*, 4486. (e) Liu, Y.-T.; Chen, J.-Q.; Li, L.-P.; Shao, X.-Y.; Xie, J.-H.; Zhou, Q.-L. *Org. Lett.* **2017**, *19*, 3231. (f) Zhu, G.-L.; Zhang, X.-D.; Yang, L.-J.; Xie, J.-H.; Che, D.-Q.; Zhou, Q.-L.; Yan, P.-C.; Li, Y.-Q. *Org. Process Res. Dev.* **2016**, *20*, 81. (g) Yan, P.-C.; Xie, J.-H.; Zhang, X.-D.; Chen, K.; Li, Y.-Q.; Zhou, Q.-L.; Che, D.-Q. *Chem. Commun.* **2014**, *50*, 15987. (h) Yang, X.-H.; Wang, K.; Zhu, S.-F.; Xie, J.-H.; Zhou, Q.-L. *J. Am. Chem. Soc.* **2014**, *136*, 17426. (i) Yang, X.-H.; Xie, J.-H.; Zhou, Q.-L. *Org. Chem. Front.* **2014**, *1*, 190. (j) Yan, P.-C.; Zhu, G.-L.; Xie, J.-H.; Zhang, X.-D.; Zhou, Q.-L.; Li, Y.-Q.; Shen, W.-H.; Che, D.-Q. *Org. Process Res. Dev.* **2013**, *17*, 307. (k) Zhang, Q.-Q.; Xie, J.-H.; Yang, X.-H.; Xie, J.-B.; Zhou, Q.-L. *Org. Lett.* **2012**, *14*, 6158. (l) Xie, J.-H.; Liu, X.-Y.; Xie, J.-B.; Wang, L.-X.; Zhou, Q.-L. *Angew. Chem., Int. Ed.* **2011**, *50*, 7329.
- [3] Zhang, F.-H.; Wang, C.; Xie, J.-H.; Zhou, Q.-L. *Adv. Synth. Catal.*

- 2019, 361, 2832.
- [4] Zhang, L.; Tang, Y.; Han, Z.; Ding, K. *Angew. Chem., Int. Ed.* **2019**, 58, 4973.
- [5] (a) Yamamura, T.; Nakane, S.; Nomura, Y.; Tanaka, S.; Kitamura, M. *Tetrahedron* **2016**, 72, 3781.
(b) Yamamura, T.; Nakatsuka, H.; Tanaka, S.; Kitamura, M. *Angew. Chem., Int. Ed.* **2013**, 52, 9313.
- [6] (a) Ling, F.; Chen, J.; Nian, S.; Hou, H.; Yi, X.; Wu, F.; Xu, M.; Zhong, W. *Synlett* **2020**, 31, 285.
(b) Qin, C.; Hou, C.-J.; Liu, H.; Liu, Y.-J.; Huang, D.-Z.; Hu, X.-P. *Tetrahedron Lett.* **2018**, 59, 719.
(c) Widegren, M. B.; Harkness, G. J.; Slawin, A. M. Z.; Cordes, D. B.; Clarke, M. L. *Angew. Chem., Int. Ed.* **2017**, 56, 5825.
(d) Chen, X.-S.; Hou, C.-J.; Qin, C.; Liu, H.; Liu, Y.-J.; Huang, D.-Z.; Hu, X.-P. *RSC Adv.* **2017**, 7, 12871.
(e) Hou, C.-J.; Hu, X.-P. *Org. Lett.* **2016**, 18, 5592.
(f) Nie, H.; Zhou, G.; Wang, Q.; Chen, W.; Zhang, S. *Tetrahedron: Asymmetry* **2013**, 24, 1567.
(g) Yamamura, T.; Nakatsuka, H.; Tanaka, S.; Kitamura, M. *Angew. Chem., Int. Ed.* **2013**, 52, 9313.
- [7] (a) Gu, G.; Hu, Y.; Liu, S.; Dong, X.-Q.; Zhang, X. *Chin. J. Org. Chem.* **2020**, 40, 997 (in Chinese).
(古国贤, 胡杨, 刘绍东, 董秀琴, 张绪穆, 有机化学, **2020**, 40, 997.)
(b) Gu, G.; Yang, T.; Yu, O.; Qian, H.; Wang, J.; Wen, J.; Dang, L.; Zhang, X. *Org. Lett.* **2017**, 19, 5920.
(c) Wu, W.; You, C.; Yin, C.; Liu, Y.; Dong, X.-Q.; Zhang, X. *Org. Lett.* **2017**, 19, 2548.
(d) Wu, W.; Liu, S.; Duan, M.; Tan, X.; Chen, C.; Xie, Y.; Lan, Y.; Dong, X.-Q.; Zhang, X. *Org. Lett.* **2016**, 18, 2938.
- [8] (a) Demmans, K. Z.; Olson, M. E.; Morris, R. H. *Organometallics* **2018**, 37, 4608.
(b) Diaz-Valenzuela, M. B.; Phillips, S. D.; France, M. B.; Gunn, M. E.; Clarke, M. L. *Chem.-Eur. J.* **2009**, 15, 1227.
- [9] Liu, H.; Yuan, H.; Shi, X. *Dalton Trans.* **2019**, 48, 609.
- [10] (a) Blasius, C. K.; Ren, B.-T.; Bürgy, D.; Liu, Y.-K.; Li, B.; Michalsky, I.; Wadepohl, H.; Deng, Q.-H.; Gade, L. H. *J. Org. Chem.* **2020**, 85, 6719.
(b) Wang, C.-J.; Sun, J.; Zhou, W.; Xue, J.; Ren, B.-T.; Zhang, G.-Y.; Mei, Y.-L.; Deng, Q.-H. *Org. Lett.* **2019**, 21, 7315.
(c) Bleith, T.; Deng, Q.-H.; Wadepohl, H.; Gade, L. H. *Angew. Chem., Int. Ed.* **2016**, 55, 7852.
(d) Deng, Q.-H.; Rettenmeier, C.; Wadepohl, H.; Gade, L. H. *Chem.-Eur. J.* **2014**, 20, 93.
(e) Deng, Q.-H.; Bleith, T.; Wadepohl, H.; Gade, L. H. *J. Am. Chem. Soc.* **2013**, 135, 5356.
(f) Deng, Q.-H.; Wadepohl, H.; Gade, L. H. *J. Am. Chem. Soc.* **2012**, 134, 2946.
(g) Deng, Q.-H.; Wadepohl, H.; Gade, L. H. *J. Am. Chem. Soc.* **2012**, 134, 10769.
(h) Deng, Q.-H.; Wadepohl, H.; Gade, L. H. *Chem.-Eur. J.* **2011**, 17, 14922.
- [11] For selected reviews on chiral oxazoline-containing ligands, see:
(a) Liu, X.; Chen, P.; Wu, F. *Chin. J. Org. Chem.* **2016**, 36, 1797 (in Chinese).
(刘小建, 陈品红, 吴范宏, 有机化学, **2016**, 36, 1797.)
(b) Liu, L.; Ma, H.; Wu, Y.; Yuan, D.; Liu, J.; Fu, B.; Ma, X. *Chin. J. Org. Chem.* **2013**, 33, 2283 (in Chinese).
(刘磊, 麻红利, 吴燕华, 袁德凯, 刘吉平, 傅滨, 马晓东, 有机化学, **2013**, 33, 2283.)
(c) Desimoni, G.; Faita, G.; Jørgensen, K. A. *Chem. Rev.* **2011**, 111, PR284.
(d) Rasappan, R.; Laventure, D.; Reiser, O. *Coord. Chem. Rev.* **2008**, 252, 702.
(e) McManus, H. A.; Guiry, P. J. *Chem. Rev.* **2004**, 104, 4151.
- [12] For selected reviews and examples on metal-catalyzed asymmetric allylic alkylation, see: (a) Cheng, Q.; Tu, H.-F.; Zheng, C.; Qu, J.-P.; Helmchen, G.; You, S.-L. *Chem. Rev.* **2019**, 119, 1855.
(b) James, J.; Jackson, M.; Guiry, P. J. *Adv. Synth. Catal.* **2019**, 361, 3016.
(c) Yu, F.-L.; Bai, D.-C.; Liu, X.-Y.; Jiang, Y.-J.; Ding, C.-H.; Hou, X.-L. *ACS Catal.* **2018**, 8, 3317.
(d) Farkas, G.; Császár, Z.; Stágel, K.; Nemes, E.; Balogh, S.; Tóth, I.; Bényei, A.; Lendvay, G.; Bakos, J. *J. Organomet. Chem.* **2017**, 846, 129.
(e) Jiang, Y.-J.; Zhang, G.-P.; Huang, J.-Q.; Chen, D.; Ding, C.-H.; Hou, X.-L. *Org. Lett.* **2017**, 19, 5932.
(f) Zheng, N.; Song, W. *Chin. J. Org. Chem.* **2017**, 37, 1099 (in Chinese).
(郑楠, 宋汪洋, 有机化学, **2017**, 37, 1099.)
(g) Fu, J.; Huo, X.; Li, B.; Zhang, W. *Org. Biomol. Chem.* **2017**, 15, 9747.
(h) Hethcox, J. C.; Shockley, S. E.; Stoltz, B. M. *ACS Catal.* **2016**, 6, 6207.
(i) Bai, D.-C.; Yu, F.-L.; Wang, W.-Y.; Chen, D.; Li, H.; Liu, Q.-R.; Ding, C.-H.; Chen, B.; Hou, X.-L. *Nat. Commun.* **2016**, 7, 11806.
(j) Bai, D.-C.; Wang, W.-Y.; Ding, C.-H.; Hou, X.-L. *Synlett* **2015**, 26, 1510.
(k) Zhuo, C.-X.; Zheng, C.; You, S.-L. *Acc. Chem. Res.* **2014**, 47, 2558.
(l) Ding, C.-H.; Hou, X.-L. *Top. Organomet. Chem.* **2011**, 36, 247.
(m) You, S.-L.; Dai, L.-X. *Angew. Chem., Int. Ed.* **2006**, 45, 5246.
(n) Trost, B. M.; Van Vranken, D. L. *Chem. Rev.* **1996**, 96, 395.
- [13] Singh, S. P.; Michaelides, A.; Merrill, A. R.; Schwan, A. L. *J. Org. Chem.* **2011**, 76, 6825.
- [14] Qi, M.-H.; Wang, F.-J.; Shi, M. *Tetrahedron: Asymmetry* **2010**, 21, 247.
- [15] Li, X.; Taichalertpaisarn, J.; Xin, D.; Burgess, K. *Org. Lett.* **2015**, 17, 632.
- [16] (a) Miao, C.; Wang, B.; Wang, Y.; Xia, C.; Lee, Y.-M.; Nam, W.; Sun, W. *J. Am. Chem. Soc.* **2016**, 138, 936.
(b) Steinig, A. G.; Spero, D. M. *J. Org. Chem.* **1999**, 64, 2406.
- [17] Wang, Y.; Du, S.; Armstrong, D. W. *Anal. Bioanal. Chem.* **2018**, 410, 4725.
- [18] Gagnon, D.; Lauzon, S.; Godbout, C.; Spino, C. *Org. Lett.* **2005**, 7, 4769.
- [19] Lee, J.-Y.; Miller, J. J.; Hamilton, S. S.; Sigman, M. S. *Org. Lett.* **2005**, 7, 1837.
- [20] Jung, B.; Kang, S. H. *Proc. Natl. Acad. Sci. U. S. A.* **2007**, 104, 1471.
- [21] Cheng, H.-G.; Feng, B.; Chen, L.-Y.; Guo, W.; Yu, X.-Y.; Lu, L.-Q.; Chen, J.-R.; Xiao, W.-J. *Chem. Commun.* **2014**, 50, 2873.
- [22] Jayakumar, S.; Prakash, M.; Balaraman, K.; Kesavan, V. *Eur. J. Org. Chem.* **2014**, 606.
- [23] Hao, X.-Q.; Dong, Y.-N.; Gao, B.; Li, K.; Zhao, X.-M.; Xu, Y.; Song, M.-P. *Tetrahedron: Asymmetry* **2015**, 26, 1360.
- [24] Jin, Y.; Du, D.-M. *Tetrahedron* **2012**, 68, 3633.
- [25] Xu, J.-X.; Ye, F.; Bai, X.-F.; Zhang, J.; Xu, Z.; Zheng, Z.-J.; Xu, L.-W. *RSC Adv.* **2016**, 6, 45495.

(Zhao, C.)