钯催化全取代吡唑导向的 C(sp²)—H 键后期芳基化反应付小盼^a 王杨阳^a 杨金月^a 吴高荣^a 夏成才^{*,b} 冀亚飞^{*,a}^(a) 华东理工大学药学院 制药工程与过程化学教育部研究中心 上海 200237)^(b) 山东第一医科大学(山东省医学科学院)药学院 山东泰安 271016)

摘要 发展了一种钯催化全取代吡唑类化合物的邻位 C(sp²)—H 键后期芳基化的反应体系。通过对反应条件的优化,得到了最高效的芳基化反应条件。该催化体系表现出广泛的底物适用性、良好的官能团耐受性以及优良的分收率。进行了控制实验,如分子间的竞争实验和放大克级实验,氘代动力学实验证明了 C—H 键的断裂是决速步骤。最后,通过制备双核二聚体环钯化中间体,提出了合理的钯催化循环反应机理。

关键词 芳基化; 后期官能化; 全取代的吡唑; C(sp²)—H 键

Fully Substituted Pyrazoles Assisted Palladium-Catalyzed Late-Stage Arylation of C(sp²)—H BondFu, Xiaopan^a Wang, Yangyang^a Yang, Jinyue^a Wu, Gaorong^a
Xia, Chengcai^{*,b} Ji, Yafei^{*,a}^(a) Engineering Research Centre of Pharmaceutical Process Chemistry, Ministry of Education, School of Pharmacy, East China University of Science & Technology, Shanghai 200237)^(b) Pharmacy College, Shandong First Medical University & Shandong Academy of Medical Sciences, Taian, Shandong 271016)

Abstract A successful protocol has been developed for palladium-catalyzed late-stage arylation of fully substituted pyrazoles. Through screening of optimization of reaction parameters, the most efficient reaction conditions for mono-*ortho*-position arylation were obtained. This reaction features a broad substrate scope, good functional group tolerance as well as good to excellent yield. Moreover, the intermolecular competition experiments and gram scale reaction were also performed. The kinetic isotopic effect (KIE) result revealed C—H bond cleavage was involved in the rate-limiting step and a plausible mechanism was proposed based on the dual-core dimeric palladacycle.

Keywords *mono*-arylation; late-stage functionalization; fully substituted pyrazole; C(sp²)—H bond

1 Introduction

Over the past few decades, the field of C—H bond functionalization has gained considerable attention, mainly because it is an efficient and powerful synthetic method for C—C or C—X bond formation.^[1] The biaryl scaffolds play an indispensable role in numerous medicinal chemistry, natural products and organic materials.^[2] Therefore, searching simple and efficient methods for construction of these structures is becoming more attractive. The oxidative coupling reaction with fewer reaction step has emerged as the most elegant method to construct biaryl scaffolds.^[3] However, poor regioselectivity hinders its development.^[4]

Increasingly, directing group (DG) assisted C—H arylation has been extensively investigated with highly-site- selectivity in atom- and step-economic mode.^[5] In these reactions, diverse DGs have exhibited a superior ability to assist C(sp²)—H arylation, such as pyridine,^[6] 8-amino-quinoline,^[7] oxime-ester,^[8] cyano,^[9] amide^[10] and PIP-amine.^[11]

Recently, late-stage functionalization has become a valuable strategy, which can directly transform complicated structures into new bioactive compounds.^[12] Pyrazoles are widely distributed in drug molecules.^[13] In previous works, transition metal-catalyzed diverse C—H bond functionalizations have been realized using pyrazole as DG through

* Corresponding authors. E-mail: xiachc@163.com; jyf@ecust.edu.cn

Received May 28, 2020; revised June 28, 2020; published online July 22, 2020.

Project supported by the National Natural Science Foundation of China (No. 21676088).

国家自然科学基金(No. 21676088)资助项目.

reactions such as arylation,^[14a-14b,14h-14i,14k,14m-14n] alkenylation,^[14c-14d] amidation,^[14e] arylation^[14g,14j,14l] and others.^[14f]

We recently reported Pd-catalyzed C(sp²)-H late-stage *mono*-acetoxylation,^[15a] iodination^[15a] and arylation^[15b] of pyrazoles (Scheme 1, A). Despite so many advances, efforts are still in high demand to develop late-stage functionalization containing a pharmacophore. Herein, we disclose the first C(sp²)-H bond late-stage *mono*-arylation using fully substituted pyrazole as auxiliary. This protocol shows a broad substrate scope and wide functional group tolerance, which provides a potential tool for the screening of potential natural products and medicines.

(A) Our previous work^[15]

(B) This work

Scheme 1 Previous related works and our present work

2 Results and discussion

As an initial experiment, the reaction of the model substrate **1a** and **2a** (2.0 equiv.) was performed in the presence of Pd(OAc)₂ (10 mol%), AgTFA (2.0 equiv.) in AcOH at 120 °C (Table 1), and **3a** was isolated in the yield of 45%. Firstly, different solvents including 1,1,1,3,3,3-hexafluoro-2-propanol (HFIP), toluene, *N,N*-dimethylformamide (DMF) and the mixture solvents of HFIP/AcOH were screened, HFIP/AcOH (*V* : *V* = 3 : 1) was suitable for the reaction (Entries 1~7). Subsequently, it was found that Pd(OAc)₂ was the best catalyst compared with other Pd salts (Entry 6 vs. 8~12). Furthermore, AgTFA was considered as the ideal additive (Entry 6 vs. 13~15). On the other hand, the yield was obviously improved when in-

creased the loading of **2a** from 2.0 equiv. to 3.0 equiv. (Entry 16). However, when the amount of AgTFA or the reaction temperatures were changed, the results were unsatisfactory (Entries 16~17). Therefore, the standard conditions were as followed: **1a** (0.3 mmol), **2a** (0.9 mmol), AgTFA (2.0 equiv.), HFIP/AcOH (*V* : *V* = 3 : 1, 3.0 mL), 120 °C, 20 h.

Table 1 Optimization of the reaction conditions^a

Entry	Catalyst	Solvent (<i>V</i> : <i>V</i>)	Yield ^b /%
1	Pd(OAc) ₂	AcOH	45
2	Pd(OAc) ₂	HFIP	43
3	Pd(OAc) ₂	Toluene	0
4	Pd(OAc) ₂	DMF	0
5	Pd(OAc) ₂	HFIP/AcOH (1 : 1)	42
6	Pd(OAc) ₂	HFIP/AcOH (3 : 1)	70
7	Pd(OAc) ₂	HFIP/AcOH (5 : 1)	43
8	Pd(CH ₃ CN) ₂ Cl ₂	HFIP/AcOH (3 : 1)	47
9	PdCl ₂	HFIP/AcOH (3 : 1)	45
10	Pd(PPh ₃) ₂ Cl ₂	HFIP/AcOH (3 : 1)	42
11	Pd(TFA) ₂	HFIP/AcOH (3 : 1)	33
12	Pd(PPh ₃) ₃	HFIP/AcOH (3 : 1)	32
13	Pd(OAc) ₂	HFIP/AcOH (3 : 1)	31 ^c (28 ^d)
14	Pd(OAc) ₂	HFIP/AcOH (3 : 1)	30 ^e (0 ^f)
15	Pd(OAc) ₂	HFIP/AcOH (3 : 1)	47 ^g (38 ^h)
16	Pd(OAc) ₂	HFIP/AcOH (3 : 1)	81 ⁱ (70 ^j)
17	Pd(OAc) ₂	HFIP/AcOH (3 : 1)	64 ^k (69 ^l)

^a Reaction conditions: **1a** (0.3 mmol), **2a** (0.6 mmol), catalyst (10 mol%), AgTFA (2.0 equiv.), solvent (3.0 mL), 120 °C, 20 h. ^b Isolated yield. ^c Ag₃PO₄ as the additive. ^d Ag₂CO₃ as the additive. ^e AgOAc as the additive. ^f K₂S₂O₈ as the additive. ^g CsOAc as the additive. ^h Cs₂CO₃ as the additive. ⁱ **2a** (0.9 mmol). ^j AgTFA (3.0 equiv.). ^k 110 °C. ^l 130 °C.

With the established conditions in hand, the scope of functionalized pyrazoles was explored and the representative products are shown in Table 2. Different functional groups including Me, OMe, F, Cl, Br, I, NO₂ and 2-naphthyl groups were tolerated and can be transformed into the desired products (**3a**~**3q**). Electron-donating groups such as methyl and methoxy are substituted at the *para* of 1-phenyl could favorably deliver the desired products in good yields (**3a**, **3b**). In comparison, the substrates containing electron-withdrawing groups were less reactive (**3d**~**3h**). Especially, the strongly electron-withdrawing group NO₂ was also provided the arylation product in 48% yield (**3h**). It was found that the *meta*-substituted of 1-phenyl pyrazoles reacted in the less hindered positions

Table 2 Substrate scope of functionalized pyrazoles^{a,b}

^a Reaction conditions: **1** (0.3 mmol), **2a** (0.9 mmol), Pd(OAc)₂ (10 mol %), AgTFA (2.0 equiv.), HFIP/AcOH (V : V = 3 : 1, 3.0 mL), 120 °C, 20 h. ^b Isolated yield.

with **2a** giving the corresponding products (**3i**~**3m**). To our disappointment, the *ortho*-position substituted substrates were not suitable for this reaction, which possibly due to the fact that steric constraints toward the formation of the complex intermediate (**3n**~**3p**).^[15b] In addition, 1-(2-naphthyl) substrates could also be applied in the transformation to deliver the desired product in good yield (**3q**). Furthermore, the yield was decreased when alkyl group instead of the aryl at the 5-site (**3r**), which means the aryl group was important for the reaction. Moreover, 5-aryl bearing electron-donating group (R²=*p*-Me) and electron-withdrawing group (R²=*p*-Cl) were well tolerated (**3s**, **3t**). Meanwhile, the alkyl substituted substrates could also be applied to the transformation to deliver the desired products in the moderate yield (**3u**, **3v**).

In the next stage, various of aryl iodides were examined. A range of electronically biased aryl iodides such as OMe, Me, COOEt, F, Cl, Br, CF₃, COCH₃ and COOCH₃ were tolerated in this protocol (Table 3). In general, the electron-rich partners gave higher yield than those bearing electron-withdrawing groups in the *para*-positions and *meta*-positions (**4b**~**4l**), but the ethyl 4-iodobenzoate could furnish the corresponding product in 81% yield (**4g**). Importantly, the reactive groups like F, Cl, Br, COOEt and COCH₃ can be easily transformed into another functional molecular. Steric effect plays a critical role in the transformation, no products were obtained for the *ortho*-substituted aryl iodides (**4m**, **4n**), maybe due to the fact that the larger steric could block the formation of the complex palladacyclic intermediate. Then, the electron-rich

6-iodo-1,4-benzodioxane also showed the excellent reactivity and gave the good yield (**4o**). To our surprise, the 3,5-substituted aryl iodides also afforded the moderate yield (**4p**). Unfortunately, 2-substituted 6-iodopyridine and 2-iodothiophene as the coupling partners independently reacted with **1c**, no desired products were obtained (**4q**, **4r**).

To gain insight into the reaction mechanism, a few control experiments were conducted. The intermolecular competition experiments revealed the donating-group was more active than those withdrawing-group (Scheme 2). Meanwhile, the kinetic isotopic effect experiment was performed between **1c** and **1c'** with **2a** under the standard conditions, the result (*k_H*/*k_D*=3.5) implicated the C—H bond cleavage as the rate-limiting step (Scheme 3). Besides, gram scale reaction was also performed (Scheme 4). According to the literature, the key palladacycle intermediate **I** was successfully obtained^[15b,16] (Scheme 5). When the intermediate **II** was exposed to the standard conditions in the absence of substrate **1c** and Pd(OAc)₂, the target product **3c** was obtained in the yield of 67%.

Based on the experiments and the relevant documented reports,^[6-8,15b,16] a plausible reaction pathway is proposed in Scheme 6. First, Pd(OAc)₂ coordinates with the substrate **I** to form the key dual-core dimeric cyclopalladated intermediate **II** through the C—H bond activation, the oxidative addition with aryl iodide and the key intermediate **II** will generate the Pd(IV) intermediate **III**. Finally, the product is obtained by reductive elimination and regenerating the Pd(II) catalyst. The role of the Ag salt may facilitate

Table 3 Substrate scope of aryl iodide^{a,b}

^a Reaction conditions: **1a** (0.3 mmol), **2** (0.9 mmol), Pd(OAc)₂ (10 mol%), AgTFA (2.0 equiv.), HFIP/AcOH (V:V = 3:1, 3.0 mL), 120 °C, 20 h. ^b Isolated yield.

Scheme 4 Gram scale reaction

Scheme 5 Preparation and functionization of the dual-core dimeric cyclopalladated intermediate **II**

oxidative addition of aryl iodides to Pd(II) centers and be used as the iodide scavengers.^[5f]

3 Conclusions

In summary, an easily method for late-stage arylation of fully substituted pyrazoles via palladium-catalyzed have been developed. A variety of pyrazoles and aryl iodides were tolerated in this protocol. Importantly, the arylation reaction of the highly functionalized substrates can be transformed into the complex bioactive molecules in a mono-site-selectivity and simple operation way. This protocol will facilitate the potential development of the medicine and material chemistry.

Scheme 6 Plausible mechanism

4 Experimental section

4.1 General methods

All reactions were carried out in oven-dried glassware and monitored by thin layer chromatography (TLC, pre-coated silica gel plates containing HF254). NMR spectra were determined on Bruker AV400 in CDCl₃ with TMS as internal standard for ¹H NMR (400 MHz) and ¹³C NMR (100 MHz), respectively. HRMS were measured on a QSTAR Pulsar I LC/TOF MS mass spectrometer or Micromass GCTTM gas chromatograph-mass spectrometer.

4.2 General procedure for arylation reaction

A mixture of substrate **1** (0.3 mmol), 4-iodoanisole **2** (0.9 mmol), Pd(OAc)₂ (10 mol%, 6.7 mg), AgTFA (2.0 equiv., 132.6 mg) in HFIP/AcOH (*V* : *V* = 3 : 1, 3.0 mL) was charged in a glass sealed-tube and stirred under air atmosphere at 120 °C for 20 h. Upon completion of the reaction, the solution was concentrated *in vacuo* for 30 min, then saturated sodium bicarbonate (30 mL) and dichloromethane (15 mL) were added into the mixture, then the aqueous layer was extracted with dichloromethane (15 mL × 2). The combined organic layer was dried over anhydrous MgSO₄. Finally, the solution was concentrated *in vacuo* to provide a crude product, which was further purified via a column chromatography on silica gel [eluent: *V*(petroleum ether) : *V*(ethyl acetate) = 5 : 1] to supply the desired product.

Ethyl 1-(4'-methoxy-5-methyl-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**3a**): Yellow solid, 103.6 mg (81% yield). m.p. 131~133 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.55 (d, *J* = 8.0 Hz, 1H), 7.21 (dd, *J* = 8.0, 1.6 Hz, 1H), 7.12 (t, *J* = 7.2 Hz, 1H), 7.03 (t, *J* = 8.0 Hz, 2H), 6.94 (s, 2H), 6.59 (d, *J* = 8.8 Hz, 2H), 6.45 (d, *J* = 7.2 Hz, 2H), 6.41 (d, *J* = 8.8 Hz, 2H), 4.47 (q, *J* = 7.2 Hz, 2H), 3.76 (s, 3H), 2.35 (s, 3H), 2.17 (s, 3H), 1.45 (t,

$J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.5, 158.8, 143.5, 141.6, 139.2, 137.8, 134.9, 130.7, 130.5, 129.23 (2C), 129.17 (2C), 128.8, 128.5, 128.4, 127.69 (2C), 127.6, 118.7, 113.49 (2C), 60.7, 55.3, 21.2, 14.5, 9.7; HRMS (EI) calcd for $\text{C}_{27}\text{H}_{26}\text{N}_2\text{O}_3$ 426.1942, found 426.1944.

Ethyl 1-(4',5'-dimethoxy-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**3b**): Yellow solid, 96.8 mg (73% yield). m.p. 158~160 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.58 (d, $J=8.8$ Hz, 1H), 7.13 (t, $J=7.6$ Hz, 1H), 7.04 (t, $J=8.0$ Hz, 2H), 6.93 (dd, $J=8.8$, 2.8 Hz, 3H), 6.65 (d, $J=2.8$ Hz, 1H), 6.60 (d, $J=8.8$ Hz, 2H), 6.47~6.43 (m, 4H), 4.47 (q, $J=7.2$ Hz, 2H), 3.81 (s, 3H), 3.77 (s, 3H), 2.16 (s, 3H), 1.46 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.5, 159.8, 159.0, 143.6, 141.5, 139.5, 130.5, 130.4, 129.9, 129.21 (2C), 129.20 (2C), 128.8, 127.7 (2C), 127.6, 118.6, 114.9, 113.5 (2C), 112.9, 60.7, 55.5, 55.3, 14.5, 9.7; HRMS (EI) calcd for $\text{C}_{27}\text{H}_{26}\text{N}_2\text{O}_4$ 442.1987, found 442.1990.

Ethyl 1-(4'-methoxy-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**3c**): Yellow solid, 81.6 mg (66% yield). m.p. 130~132 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.68 (d, $J=7.6$ Hz, 1H), 7.44~7.36 (m, 2H), 7.15 (d, $J=6.8$ Hz, 1H), 7.12 (d, $J=7.2$ Hz, 1H), 7.03 (t, $J=7.6$ Hz, 2H), 6.60 (d, $J=8.8$ Hz, 2H), 6.45 (d, $J=8.0$ Hz, 2H), 6.42 (d, $J=8.8$ Hz, 2H), 4.48 (q, $J=7.2$ Hz, 2H), 3.77 (s, 3H), 2.18 (s, 3H), 1.46 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.4, 158.9, 143.5, 141.8, 138.2, 137.3, 130.4, 130.1, 129.3, 129.3 (2C), 129.1 (2C), 128.83, 128.7, 127.8, 127.7 (2C), 127.7, 118.8, 113.6 (2C), 60.8, 55.3, 14.6, 9.7; HRMS (EI) calcd for $\text{C}_{26}\text{H}_{24}\text{N}_2\text{O}_3$ 412.1816, found 412.1817.

Ethyl 1-(5-fluoro-4'-methoxy-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**3d**): Yellow solid, 74.8 mg (58% yield). m.p. 105~107 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.60 (dd, $J=8.8$, 5.6 Hz, 1H), 7.15 (t, $J=7.2$ Hz, 1H), 7.11 (m, 1H), 7.05 (t, $J=7.6$ Hz, 2H), 6.86 (dd, $J=9.6$, 3.2 Hz, 1H), 6.61 (d, $J=8.8$ Hz, 2H), 6.44 (d, $J=6.8$ Hz, 2H), 6.41 (d, $J=8.8$ Hz, 2H), 4.48 (q, $J=7.2$ Hz, 2H), 3.78 (s, 3H), 2.17 (s, 3H), 1.46 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.5 (d, $^1J_{\text{CF}}=247.5$ Hz), 163.3, 159.3, 143.7, 141.9, 140.5 (d, $^3J_{\text{CF}}=8.6$ Hz), 133.4 (d, $^4J_{\text{CF}}=2.9$ Hz), 130.7 (d, $^3J_{\text{CF}}=9.2$ Hz), 129.3, 129.3, 129.2 (2C), 129.1 (2C), 128.5, 127.8 (2C), 118.9, 116.6 (d, $^2J_{\text{CF}}=22.9$ Hz), 114.66 (d, $^3J_{\text{CF}}=26.0$ Hz), 113.7 (2C), 60.8, 55.3, 14.5, 9.7; HRMS (EI) calcd for $\text{C}_{26}\text{H}_{23}\text{N}_2\text{O}_3\text{F}$ 430.1736, found 430.1737.

Ethyl 1-(5-chloro-4'-methoxy-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**3e**): Yellow solid, 97.7 mg (73% yield). m.p. 138~140 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.63 (d, $J=8.4$ Hz, 1H), 7.40 (dd, $J=8.4$, 2.4 Hz, 1H), 7.18~7.14 (m, 2H), 7.06 (t, $J=8.0$ Hz, 2H), 6.61 (d, $J=8.8$ Hz, 2H), 6.44 (d, $J=7.2$ Hz, 2H), 6.40 (d, $J=8.4$ Hz, 2H), 4.48 (q, $J=7.2$ Hz, 2H), 3.78 (s, 3H), 2.17 (s, 3H), 1.46 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.2, 159.3, 143.5, 142.1, 139.8, 135.9, 134.9, 130.2, 130.0, 129.2 (2C), 129.1 (2C), 128.4, 127.9

(2C), 127.9 (2C), 127.7, 119.0, 113.7 (2C), 60.9, 55.3, 14.5, 9.7; HRMS (EI) calcd for $\text{C}_{26}\text{H}_{23}\text{N}_2\text{O}_3\text{Cl}$ 446.1497, found 446.1498.

Ethyl 1-(5-bromo-4'-methoxy-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**3f**): Yellow solid, 81.0 mg (55% yield). m.p. 136~138 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.58~7.53 (m, 2H), 7.31 (s, 1H), 7.15 (t, $J=7.6$ Hz, 1H), 7.05 (t, $J=7.6$ Hz, 2H), 6.60 (d, $J=8.8$ Hz, 2H), 6.43 (d, $J=7.2$ Hz, 2H), 6.39 (d, $J=8.8$ Hz, 2H), 4.48 (q, $J=7.2$ Hz, 2H), 3.77 (s, 3H), 2.16 (s, 3H), 1.45 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.2, 159.3, 143.5, 142.2, 140.0, 136.4, 132.9, 130.7, 130.4, 129.2 (2C), 129.0 (2C), 129.0, 128.4, 127.9, 127.9, 123.0, 119.0, 113.7 (2C), 60.9, 55.3, 55.3, 14.5, 9.7; HRMS (EI) calcd for $\text{C}_{26}\text{H}_{23}\text{N}_2\text{O}_3\text{Br}$ 490.0981, found 490.0979.

Ethyl 1-(5-iodo-4'-methoxy-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**3g**): Yellow solid, 114.6 mg (71% yield). m.p. 157~159 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.74 (dd, $J=8.4$, 2.0 Hz, 1H), 7.50 (d, $J=2.0$ Hz, 1H), 7.41 (d, $J=8.0$ Hz, 1H), 7.15 (t, $J=7.6$ Hz, 1H), 7.05 (t, $J=7.6$ Hz, 2H), 6.59 (t, $J=8.8$ Hz, 2H), 6.43 (d, $J=7.2$ Hz, 2H), 6.37 (d, $J=7.2$ Hz, 2H), 6.37 (d, $J=8.4$ Hz, 2H), 4.47 (q, $J=7.2$ Hz, 2H), 3.77 (s, 3H), 2.16 (s, 3H), 1.45 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.3, 159.3, 143.5, 142.2, 140.1, 138.9, 137.1, 136.7, 130.4, 129.2 (2C), 129.0 (2C), 128.8, 128.4, 127.9, 127.9 (2C), 119.0, 113.7 (2C), 94.8, 60.9, 55.3, 14.5, 9.7; HRMS (EI) calcd for $\text{C}_{26}\text{H}_{23}\text{N}_2\text{O}_3\text{I}$ 538.0836, found 538.0838.

Ethyl 1-(4'-methoxy-5-nitro-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**3h**): Yellow solid, 65.8 mg (48% yield). m.p. 170~172 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.28 (dd, $J=8.8$, 2.8 Hz, 1H), 8.05 (d, $J=2.4$ Hz, 1H), 7.90 (d, $J=8.8$ Hz, 1H), 7.18 (t, $J=7.6$ Hz, 1H), 7.06 (t, $J=7.6$ Hz, 2H), 6.64 (d, $J=8.8$ Hz, 2H), 6.43 (d, $J=8.8$ Hz, 4H), 4.49 (q, $J=7.2$ Hz, 2H), 3.80 (s, 3H), 2.19 (s, 3H), 1.47 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.0, 159.7, 147.8, 143.6, 143.0, 142.2, 139.6, 130.1, 129.2 (2C), 128.9 (2C), 128.2, 128.2, 128.1 (2C), 128.0, 125.4, 122.5, 119.5, 114.0 (2C), 61.1, 55.4, 14.5, 9.6; HRMS (EI) calcd for $\text{C}_{26}\text{H}_{23}\text{N}_3\text{O}_5$ 457.1687, found 457.1688.

Ethyl 1-(4,4'-dimethoxy-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**3i**): Yellow solid, 70.3 mg (53% yield). m.p. 108~110 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.23 (d, $J=2.8$ Hz, 1H), 7.14 (t, $J=7.6$ Hz, 1H), 7.06~7.02 (m, 3H), 6.95 (dd, $J=8.8$, 2.8 Hz, 1H), 6.58 (d, $J=8.8$ Hz, 2H), 6.47 (d, $J=7.2$ Hz, 2H), 6.35 (d, $J=8.8$ Hz, 2H), 4.48 (q, $J=7.2$ Hz, 2H), 3.89 (s, 3H), 3.76 (s, 3H), 2.18 (s, 3H), 1.46 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.4, 159.0, 158.5, 143.5, 141.8, 137.9, 131.0, 130.8, 130.2, 129.1 (2C), 129.1 (2C), 128.7, 127.7 (2C), 127.7, 118.8, 116.0, 113.5 (2C), 113.3, 60.8, 55.7, 55.32, 14.5, 9.7; HRMS (EI) calcd for $\text{C}_{27}\text{H}_{26}\text{N}_2\text{O}_4$ 442.1941, found 442.1937.

Ethyl 1-(4'-methoxy-4-methyl-[1,1'-biphenyl]-2-yl)-4-

methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**3j**): Yellow solid, 93.3 mg (73% yield). m.p. 113~115 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.51 (d, *J*=0.8 Hz, 1H), 7.17 (dd, *J*=7.6, 0.8 Hz, 1H), 7.11 (t, *J*=7.2 Hz, 1H), 7.03~7.00 (m, 3H), 6.58 (d, *J*=8.8 Hz, 2H), 6.44 (d, *J*=7.2 Hz, 2H), 6.36 (d, *J*=8.8 Hz, 2H), 4.48 (q, *J*=7.2 Hz, 2H), 3.75 (s, 3H), 2.42 (s, 3H), 2.17 (s, 3H), 1.45 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.4, 158.7, 143.4, 141.7, 137.8, 137.0, 135.2, 130.4, 130.1, 129.9, 129.1 (2C), 129.1, 129.1 (2C), 128.7, 127.7 (2C), 127.6, 118.7, 113.5 (2C), 60.8, 55.3, 20.9, 14.6, 9.7; HRMS (EI) calcd for C₂₇H₂₆N₂O₃ 426.1936, found 426.1938.

Ethyl 1-(4-fluoro-4'-methoxy-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**3k**): Yellow solid, 68.4 mg (53% yield). m.p. 155~157 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.46 (d, *J*=8.8 Hz, 1H), 7.17~7.11 (m, 3H), 7.05 (t, *J*=7.6 Hz, 2H), 6.60 (d, *J*=8.8 Hz, 2H), 6.46 (d, *J*=7.2 Hz, 2H), 6.36 (d, *J*=8.8 Hz, 2H), 4.49 (q, *J*=7.2 Hz, 2H), 3.77 (s, 3H), 2.17 (s, 3H), 1.47 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.2, 162.6 (d, ¹*J*_{CF}=247.0 Hz), 158.9, 143.5, 142.2, 138.2 (d, ³*J*_{CF}=9.8 Hz), 134.5 (d, ⁴*J*_{CF}=3.5 Hz), 131.5 (d, ³*J*_{CF}=8.5 Hz), 129.4, 129.2 (2C), 129.0 (2C), 128.4, 127.8 (2C), 119.0, 116.5 (d, ²*J*_{CF}=20.8 Hz), 116.0 (d, ²*J*_{CF}=23.5 Hz), 113.6 (2C), 60.9, 55.3, 14.5, 9.6; HRMS (EI) calcd for C₂₆H₂₃N₂O₃F 430.1736, found 430.1735.

Ethyl 1-(4-chloro-4'-methoxy-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**3l**): Yellow solid, 84.3 mg (63% yield). m.p. 98~100 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.74 (d, *J*=2.0 Hz, 2H), 7.37 (dd, *J*=8.4, 2.0 Hz, 1H), 7.15 (t, *J*=7.2 Hz, 2H), 7.08 (d, *J*=8.4 Hz, 1H), 7.05 (t, *J*=8.0 Hz, 2H), 6.60 (d, *J*=8.4 Hz, 2H), 6.45 (d, *J*=7.2 Hz, 2H), 6.36 (d, *J*=8.8 Hz, 2H), 4.49 (q, *J*=7.2 Hz, 2H), 3.77 (s, 3H), 2.17 (s, 3H), 1.47 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.2, 159.1, 143.5, 142.2, 138.0, 136.7, 133.1, 131.2, 129.5, 129.2, 129.1 (2C), 129.0 (2C), 128.9, 128.3, 127.9 (3C), 119.0, 113.7 (2C), 60.9, 55.3, 14.5, 9.6; HRMS (EI) calcd for C₂₆H₂₃N₂O₃Cl 446.1436, found 446.1437.

Ethyl 1-(4-bromo-4'-methoxy-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**3m**): Yellow solid, 95.6 mg (65% yield). m.p. 103~105 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.89 (d, *J*=2.0 Hz, 1H), 7.51 (dd, *J*=8.4, 2.0 Hz, 1H), 7.15 (t, *J*=7.6 Hz, 1H), 7.07~7.00 (m, 3H), 6.59 (d, *J*=8.8 Hz, 2H), 6.44 (d, *J*=7.2 Hz, 2H), 6.36 (d, *J*=8.8 Hz, 2H), 4.49 (q, *J*=7.2 Hz, 2H), 3.77 (s, 3H), 2.17 (s, 3H), 1.47 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.2, 159.1, 143.5, 142.2, 138.2, 137.2, 132.4, 131.7, 131.4, 129.3, 129.1 (2C), 129.0 (2C), 128.3, 127.9 (3C), 120.8, 119.0, 113.7 (2C), 60.9, 55.3, 14.5, 9.7; HRMS (EI) calcd for C₂₆H₂₃N₂O₃Br 490.0936, found 490.0937.

Ethyl 1-(3-(4-methoxyphenyl)naphthalen-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**3q**): Yellow oil, 117.9 mg (85% yield). ¹H NMR (400 MHz, CDCl₃) δ: 8.25 (s, 1H), 7.94~7.91 (m, 1H), 7.81~7.79 (m, 1H), 7.61 (s, 1H), 7.54~7.52 (m, 2H), 7.11 (t, *J*=7.2 Hz, 1H),

6.99 (t, *J*=7.6 Hz, 2H), 6.64 (d, *J*=8.8 Hz, 2H), 6.50 (d, *J*=8.8 Hz, 2H), 6.47 (d, *J*=7.2 Hz, 2H), 4.51 (q, *J*=7.2 Hz, 2H), 3.79 (s, 3H), 2.22 (s, 3H), 1.48 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.4, 159.0, 143.7, 141.9, 136.1, 136.0, 133.4, 132.3, 130.5, 129.4 (2C), 129.2, 129.1 (2C), 128.6, 128.0, 127.9, 127.8 (2C), 127.8, 127.7, 127.2, 126.6, 118.9, 113.6 (2C), 60.9, 55.3, 14.6, 9.8; HRMS (EI) calcd for C₃₀H₂₆N₂O₃ 462.1943, found 462.1942.

Ethyl 5-ethyl-1-(4'-methoxy-[1,1'-biphenyl]-2-yl)-4-methyl-1*H*-pyrazole-3-carboxylate (**3r**): Yellow solid, 68.8 mg (63% yield). m.p. 104~106 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.51~7.47 (m, 2H), 7.42~7.40 (m, 2H), 7.00 (d, *J*=8.8 Hz, 2H), 6.76 (d, *J*=8.8 Hz, 2H), 4.43 (q, *J*=7.2 Hz, 2H), 3.77 (s, 3H), 2.16 (s, 3H), 2.05~1.99 (m, 2H), 1.42 (t, *J*=7.2 Hz, 3H), 0.64 (t, *J*=7.6 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.6, 159.1, 144.7, 141.2, 138.8, 137.0, 130.2, 130.2, 129.8, 129.7 (2C), 129.2, 127.7, 117.3, 113.9 (2C), 60.6, 55.2, 17.3, 14.5, 12.6, 9.1; HRMS (EI) calcd for C₂₂H₂₄N₂O₃ 364.1887, found 364.1888.

Ethyl 1-(4'-methoxy-[1,1'-biphenyl]-2-yl)-4-methyl-5-(*p*-tolyl)-1*H*-pyrazole-3-carboxylate (**3s**): Yellow solid, 113.8 mg (89% yield). m.p. 143~145 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.65 (dd, *J*=7.6, 2.0 Hz, 1H), 7.43~7.35 (m, 2H), 7.16 (dd, *J*=7.2, 1.6 Hz, 1H), 6.83 (d, *J*=8.0 Hz, 2H), 6.61 (d, *J*=8.8 Hz, 2H), 6.45 (d, *J*=8.8 Hz, 2H), 6.34 (d, *J*=8.0 Hz, 2H), 4.47 (q, *J*=7.2 Hz, 2H), 3.77 (s, 3H), 2.24 (s, 3H), 2.16 (s, 3H), 1.45 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.4, 158.9, 143.6, 141.7, 138.3, 137.5, 137.4, 130.4, 130.1, 129.3, 129.3 (2C), 129.0 (2C), 128.8, 128.4 (2C), 127.7, 125.7, 118.6, 113.5 (2C), 60.7, 55.3, 21.2, 14.5, 9.7; HRMS (EI) calcd for C₂₇H₂₆N₂O₃ 426.1987, found 426.1992.

Ethyl 5-(4-chlorophenyl)-1-(4'-methoxy-[1,1'-biphenyl]-2-yl)-4-methyl-1*H*-pyrazole-3-carboxylate (**3t**): White solid, 97.7 mg (73% yield). m.p. 103~105 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.67 (dd, *J*=6.8, 1.2 Hz, 1H), 7.44~7.37 (m, 2H), 7.17 (dd, *J*=7.2, 1.6 Hz, 1H), 6.73 (t, *J*=8.8 Hz, 2H), 6.63 (d, *J*=8.8 Hz, 2H), 6.45 (d, *J*=8.4 Hz, 2H), 6.42~6.38 (m, 2H), 4.48 (q, *J*=7.2 Hz, 2H), 3.77 (s, 3H), 2.15 (s, 3H), 1.45 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.4, 163.3, 160.9, 159.0, 142.5, 141.8, 138.0, 137.0, 131.0, 130.9, 130.3, 130.2, 129.5, 129.2, 128.8, 127.9, 124.8, 118.8, 115.0, 114.7, 113.6 (2C), 60.8, 55.3, 14.5, 9.7; HRMS (EI) calcd for C₂₆H₂₃N₂O₃Cl 446.1436, found 446.1438.

1-(4'-Methoxy-[1,1'-biphenyl]-2-yl)-3,4,5-trimethyl-1*H*-pyrazole (**3u**): Colourless oil, 55.2 mg (63% yield). ¹H NMR (400 MHz, CDCl₃) δ: 7.47~7.37 (m, 4H), 6.98 (d, *J*=8.0 Hz, 2H), 6.77 (d, *J*=8.0 Hz, 2H), 3.77 (s, 3H), 2.24 (s, 3H), 1.81 (s, 3H), 1.52 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 158.9, 147.4, 138.7, 137.5, 137.4, 131.0, 130.2, 129.6 (2C), 128.9, 128.9, 127.7, 113.7 (2C), 111.9, 55.2, 11.9, 9.7, 8.1; HRMS (EI) calcd for C₁₉H₂₀N₂O 292.1618, found 292.1617.

1-(4'-Methoxy-[1,1'-biphenyl]-2-yl)-3-methyl-5-phenyl-

1*H*-pyrazole (**3v**): White solid, 62.2 mg (61% yield). m.p. 146~148 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.69 (dd, *J*=8.0, 4.0 Hz, 1H), 7.47~7.38 (m, 2H), 7.22 (dd, *J*=8.0, 4.0 Hz, 1H), 7.09 (t, *J*=8.0 Hz, 1H), 7.00 (t, *J*=8.0 Hz, 2H), 6.63 (d, *J*=8.0 Hz, 2H), 6.57 (d, *J*=8.0 Hz, 2H), 6.45 (d, *J*=8.0 Hz, 2H), 6.08 (s, 1H), 3.75 (s, 3H), 2.40 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 158.7, 148.9, 145.3, 138.6, 138.0, 130.8, 130.4, 130.3, 129.2 (2C), 128.7 (2C), 128.4 (2C), 127.9, 127.7, 127.5, 127.3, 113.4 (2C), 106.1, 55.3, 13.7; HRMS (EI) calcd for C₂₃H₂₀N₂O 340.1617, found 340.1619.

Ethyl 1-([1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**4a**): Yellow oil, 71.1 mg (62% yield). ¹H NMR (400 MHz, CDCl₃) δ: 7.71 (d, *J*=7.6 Hz, 1H), 7.46 (t, *J*=7.6 Hz, 1H), 7.40 (t, *J*=7.6 Hz, 1H), 7.19~7.10 (m, 3H), 7.06~6.98 (m, 4H), 6.50 (d, *J*=7.2 Hz, 2H), 6.40 (d, *J*=7.2 Hz, 2H), 4.48 (q, *J*=7.2 Hz, 2H), 2.17 (s, 3H), 1.46 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.4, 143.5, 141.8, 138.5, 137.8, 137.4, 130.3, 129.4, 129.2 (2C), 128.8, 128.6, 128.2, 128.1 (2C), 128.1 (2C), 127.7 (2C), 127.7, 127.0, 118.8, 60.8, 14.6, 9.7; HRMS (EI) calcd for C₂₅H₂₂N₂O₂ 382.1793, found 382.1791.

Ethyl 4-methyl-1-(4'-methyl-[1,1'-biphenyl]-2-yl)-5-phenyl-1*H*-pyrazole-3-carboxylate (**4b**): Yellow oil, 83.2 mg (70% yield). ¹H NMR (400 MHz, CDCl₃) δ: 7.68 (d, *J*=7.2 Hz, 1H), 7.45~7.39 (m, 2H), 7.17 (dd, *J*=7.2, 1.2 Hz, 1H), 7.13 (t, *J*=7.6 Hz, 1H), 7.05 (t, *J*=8.0 Hz, 2H), 6.87 (d, *J*=8.0 Hz, 2H), 6.41 (t, *J*=6.8 Hz, 4H), 4.48 (q, *J*=7.2 Hz, 2H), 2.30 (s, 3H), 2.18 (s, 3H), 1.46 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.4, 143.5, 141.7, 138.6, 137.4, 136.8, 134.9, 130.2, 130.1, 129.3, 129.2 (2C), 128.8, 128.8 (2C), 128.7, 128.0, 127.9, 127.7, 127.6, 125.4, 118.8, 60.8, 21.1, 14.6, 9.7; HRMS (EI) calcd for C₂₆H₂₄N₂O₂ 396.1842, found 396.1845.

Ethyl 1-(4'-fluoro-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**4c**): Yellow oil, 72.0 mg (60% yield). ¹H NMR (400 MHz, CDCl₃) δ: 7.70 (d, *J*=8.0 Hz, 1H), 7.46 (td, *J*=7.6, 1.2 Hz, 1H), 7.40 (td, *J*=7.6, 1.2 Hz, 1H), 7.15~7.12 (m, 2H), 7.04 (t, *J*=7.6 Hz, 2H), 6.45 (d, *J*=8.0 Hz, 4H), 4.47 (q, *J*=7.2 Hz, 2H), 2.18 (s, 3H), 1.45 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.3, 162.2 (d, ¹*J*_{CF}=245.4 Hz), 143.4, 142.0, 137.5, 137.4, 133.9 (d, ³*J*_{CF}=3.3 Hz, 2C), 130.2, 129.8, 129.7, 129.4, 129.1 (2C), 128.9, 128.6, 128.4, 127.9 (2C), 127.8, 118.8, 115.1 (d, ²*J*_{CF}=21.3 Hz, 2C), 60.8, 14.5, 9.7; HRMS (EI) calcd for C₂₅H₂₁N₂O₂F 400.1642, found 400.1641.

Ethyl 1-(4'-chloro-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**4d**): Yellow solid, 59.9 mg (48% yield), m.p. 110~112 °C. ¹H NMR (400 MHz, CDCl₃) δ: 7.71 (d, *J*=8.0 Hz, 1H), 7.49 (t, *J*=7.6 Hz, 1H), 7.42 (t, *J*=7.6 Hz, 1H), 7.16~7.14 (m, 2H), 7.07~7.03 (m, 4H), 6.46 (d, *J*=7.2 Hz, 2H), 6.42 (d, *J*=8.4 Hz, 2H), 4.48 (q, *J*=7.2 Hz, 2H), 2.20 (s, 3H), 1.46 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.3, 143.4, 142.1, 137.4, 137.2, 136.3, 133.3, 130.1, 129.5, 129.4 (2C), 129.2

(2C), 129.0, 128.6, 128.5, 128.2 (2C), 127.9 (2C), 127.8, 118.9, 60.9, 14.5, 9.7; HRMS (EI) calcd for C₂₅H₂₁N₂O₂Cl 416.1346, found 416.1349.

Ethyl 4-methyl-5-phenyl-1-(4'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-1*H*-pyrazole-3-carboxylate (**4e**): Yellow solid, 51.3 mg (38% yield). m.p. 103~105 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.75 (d, *J*=7.6 Hz, 1H), 7.53 (t, *J*=7.6 Hz, 1H), 7.45 (t, *J*=7.6 Hz, 1H), 7.31 (d, *J*=8.4 Hz, 2H), 7.19 (d, *J*=7.6 Hz, 1H), 7.15 (t, *J*=7.6 Hz, 1H), 7.02 (t, *J*=7.6 Hz, 2H), 6.60 (d, *J*=8.0 Hz, 2H), 6.38 (d, *J*=7.2 Hz, 2H), 4.48 (q, *J*=7.2 Hz, 2H), 2.18 (s, 3H), 1.46 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.2, 143.3, 142.3, 141.5, 137.5, 136.8, 130.1, 129.5, 129.2, 129.1 (q, ²*J*_{CF}=32.2 Hz, 2C), 129.1, 129.0 (2C), 128.4, 128.4, 128.0, 127.8, 125.49, 125.0 (q, ³*J*_{CF}=3.5 Hz, 2C), 124.1 (q, ¹*J*_{CF}=270.3 Hz), 118.9, 60.9, 14.5, 9.7; HRMS (EI) calcd for C₂₆H₂₁N₂O₂F₃ 450.1693, found 450.1694.

Ethyl 1-(4'-acetyl-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**4f**): Yellow solid, 49.6 mg (39% yield). m.p. 169~171 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.73 (d, *J*=8.0 Hz, 1H), 7.65 (d, *J*=8.4 Hz, 2H), 7.52 (t, *J*=7.6 Hz, 1H), 7.44 (t, *J*=7.6 Hz, 1H), 7.19 (d, *J*=6.8 Hz, 2H), 7.14 (t, *J*=7.6 Hz, 1H), 7.01 (t, *J*=8.0 Hz, 2H), 6.60 (d, *J*=8.4 Hz, 2H), 6.38 (d, *J*=7.2 Hz, 2H), 4.48 (q, *J*=7.2 Hz, 2H), 2.57 (s, 3H), 2.16 (s, 3H), 1.46 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 197.9, 163.2, 143.4, 142.7, 142.2, 137.5, 137.3, 135.5, 130.1, 129.5, 129.2 (2C), 129.1, 129.1, 128.4, 128.3 (2C), 128.1 (2C), 127.9 (2C), 127.9, 118.9, 60.9, 26.7, 14.5, 9.7; HRMS (EI) calcd for C₂₇H₂₄N₂O₃ 424.1898, found 424.1896.

Ethyl 1-(4'-(ethoxycarbonyl)-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**4g**): Yellow solid, 110.4 mg (81% yield). m.p. 105~107 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.72 (d, *J*=8.0 Hz, 3H), 7.49 (t, *J*=7.6 Hz, 1H), 7.41 (t, *J*=7.6 Hz, 1H), 7.17 (d, *J*=7.6 Hz, 1H), 7.12 (t, *J*=7.2 Hz, 1H), 6.99 (t, *J*=7.6 Hz, 2H), 6.55 (d, *J*=8.0 Hz, 2H), 6.38 (d, *J*=8.0 Hz, 2H), 4.46 (q, *J*=7.2 Hz, 2H), 4.35 (q, *J*=7.2 Hz, 2H), 2.15 (s, 3H), 1.44 (t, *J*=7.2 Hz, 3H), 1.38 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 166.4, 163.2, 143.4, 142.4, 142.1, 137.4, 137.4, 130.1, 129.5, 129.3 (2C), 129.2 (2C), 129.0 (2C), 128.9, 128.4, 128.1 (2C), 127.9 (2C), 127.8, 118.9, 61.0, 60.8, 14.5, 14.3, 9.7; HRMS (EI) calcd for C₂₈H₂₆N₂O₄ 454.1903, found 454.1901.

Ethyl 1-(3'-methoxy-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**4h**): Brown oil, 87.8 mg (71% yield). ¹H NMR (400 MHz, CDCl₃) δ: 7.71 (d, *J*=7.6 Hz, 1H), 7.46 (t, *J*=7.2 Hz, 1H), 7.41 (t, *J*=7.2 Hz, 1H), 7.21 (d, *J*=7.2 Hz, 1H), 7.14 (t, *J*=7.6 Hz, 1H), 7.03 (t, *J*=7.6 Hz, 2H), 6.97 (t, *J*=8.0 Hz, 1H), 6.72 (dd, *J*=8.4, 1.8 Hz, 1H), 6.45 (d, *J*=7.2 Hz, 2H), 6.13 (d, *J*=7.6 Hz, 1H), 6.03 (s, 1H), 4.47 (q, *J*=7.2 Hz, 2H), 3.57 (s, 3H), 2.18 (s, 3H), 1.44 (t, *J*=7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 1163.3, 159.4, 143.7, 141.8, 139.0, 138.4, 137.5, 130.2, 129.4, 129.2 (2C), 129.0, 128.8, 128.6, 128.3, 127.7 (2C), 127.7, 120.6, 118.7, 114.2, 112.3, 60.8,

54.9, 14.5, 9.7; HRMS (EI) calcd for $C_{26}H_{24}N_2O_3$ 412.1891, found 412.1890.

Ethyl 1-(3'-bromo-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**4i**): Yellow solid, 85.6 mg (62% yield). m.p. 134~136 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.74 (d, $J=8.0$ Hz, 1H), 7.50 (t, $J=7.6$ Hz, 1H), 7.42 (t, $J=7.6$ Hz, 1H), 7.29 (d, $J=8.0$ Hz, 1H), 7.18 (t, $J=7.6$ Hz, 1H), 7.14 (d, $J=7.6$ Hz, 1H), 7.08 (t, $J=7.6$ Hz, 2H), 6.93 (t, $J=7.6$ Hz, 1H), 6.51 (s, 1H), 6.47 (d, $J=8.0$ Hz, 1H), 6.43 (d, $J=7.2$ Hz, 2H), 4.48 (q, $J=7.2$ Hz, 2H), 2.19 (s, 3H), 1.45 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 163.2, 143.5, 142.1, 139.8, 137.5, 137.0, 131.2, 130.0, 130.0, 129.5, 129.4, 129.0 (2C), 128.9, 128.9, 128.3, 128.0 (2C), 127.9, 126.5, 122.3, 118.8, 60.8, 14.5, 9.7; HRMS (EI) calcd for $C_{25}H_{21}N_2O_2Br$ 460.0806, found 460.0808.

Ethyl 4-methyl-5-phenyl-1-(3'-(trifluoromethyl)-[1,1'-biphenyl]-2-yl)-1*H*-pyrazole-3-carboxylate (**4j**): Yellow solid, 56.7 mg (42% yield). m.p. 103~105 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.77 (d, $J=7.6$ Hz, 1H), 7.53 (td, $J=7.6$, 1.2 Hz, 1H), 7.46 (d, $J=7.6$ Hz, 1H), 7.43 (d, $J=6.4$ Hz, 1H), 7.21~7.12 (m, 3H), 7.02 (t, $J=7.6$ Hz, 2H), 6.71 (d, $J=8.0$ Hz, 1H), 6.68 (s, 1H), 6.38 (d, $J=7.2$ Hz, 2H), 4.48 (q, $J=7.2$ Hz, 2H), 2.17 (s, 3H), 1.45 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 163.2, 143.4, 142.2, 138.5, 137.6, 136.9, 131.3, 130.68 (q, $^2J_{CF}=32.0$ Hz), 130.0, 129.6, 129.1, 129.0, 128.9 (2C), 128.5, 128.3, 128.0 (2C), 127.9, 125.2 (q, $^1J_{CF}=270.8$ Hz), 125.1 (q, $^3J_{CF}=3.6$ Hz), 123.8 (q, $^3J_{CF}=3.7$ Hz), 119.0, 60.9, 14.5, 9.6; HRMS (EI) calcd for $C_{26}H_{21}N_2O_2F_3$ 450.1642, found 450.1641.

Ethyl 1-(3'-acetyl-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**4k**): White solid, 61.1 mg (48% yield). m.p. 119~121 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.79 (d, $J=7.6$ Hz, 1H), 7.75 (d, $J=7.6$ Hz, 1H), 7.52 (t, $J=7.6$ Hz, 1H), 7.45 (t, $J=7.6$ Hz, 1H), 7.22~7.10 (m, 3H), 7.01~6.97 (m, 3H), 6.73 (d, $J=7.6$ Hz, 1H), 6.38 (d, $J=7.6$ Hz, 2H), 4.47 (q, $J=7.2$ Hz, 2H), 2.38 (s, 3H), 2.14 (s, 3H), 1.44 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 197.7, 163.1, 143.5, 142.1, 138.2, 137.5, 137.4, 137.0, 132.7, 130.1, 129.6, 129.0 (2C), 128.9, 128.9, 128.7, 128.4, 128.3, 127.9 (2C), 127.8, 126.6, 118.8, 60.9, 26.4, 14.5, 9.7; HRMS (EI) calcd for $C_{27}H_{24}N_2O_3$ 424.1841, found 424.1837.

Ethyl 1-(3'-(methoxycarbonyl)-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**4l**): Colorless oil, 91.1 mg (69% yield). 1H NMR (400 MHz, $CDCl_3$) δ : 8.43 (d, $J=8.0$ Hz, 1H), 7.73 (d, $J=7.6$ Hz, 1H), 7.49 (td, $J=7.6$, 1.2 Hz, 1H), 7.42 (t, $J=7.6$ Hz, 1H), 7.20 (dd, $J=7.6$, 0.8 Hz, 1H), 7.15~7.08 (m, 2H), 7.05 (s, 1H), 6.97 (t, $J=7.6$ Hz, 2H), 6.71 (d, $J=7.6$ Hz, 1H), 6.36 (d, $J=7.2$ Hz, 2H), 4.47 (q, $J=7.2$ Hz, 2H), 3.84 (s, 3H), 2.14 (s, 3H), 1.44 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 166.6, 163.2, 143.4, 142.1, 138.0, 137.5, 137.4, 132.4, 130.2, 130.1, 129.5 (2C), 129.0 (2C), 128.9, 128.80, 128.3, 128.2, 128.1, 127.8, 127.8 (2C), 118.9, 60.8, 52.0, 14.5, 9.7; HRMS (EI) calcd for $C_{27}H_{24}N_2O_4$ 440.1749,

found 440.1750.

Ethyl 1-(2-(2,3-dihydrobenzo[*b*][1,4]dioxin-6-yl)-phenyl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**4o**): Yellow solid, 108.3 mg (82% yield). m.p. 128~130 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.67 (d, $J=6.8$ Hz, 1H), 7.42 (t, $J=7.2$ Hz, 1H), 7.37 (t, $J=7.2$ Hz, 1H), 7.14 (t, $J=7.3$ Hz, 2H), 7.05 (t, $J=7.6$ Hz, 2H), 6.55 (d, $J=8.4$ Hz, 1H), 6.51 (d, $J=7.6$ Hz, 2H), 5.97 (s, 2H), 4.47 (q, $J=7.2$ Hz, 2H), 4.20 (dd, $J=13.2$, 4.4 Hz, 4H), 2.21 (s, 3H), 1.45 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 163.4, 143.4, 143.3, 142.9, 141.8, 137.8, 137.3, 131.2, 130.1, 129.3, 129.1 (2C), 128.8, 128.6, 127.9, 127.7 (3C), 121.2, 118.7, 117.2, 116.7, 64.5, 64.2, 60.8, 14.5, 9.7; HRMS (EI) calcd for $C_{27}H_{24}N_2O_4$ 440.1787, found 440.1788.

Ethyl 1-(3',5'-dichloro-[1,1'-biphenyl]-2-yl)-4-methyl-5-phenyl-1*H*-pyrazole-3-carboxylate (**4p**): White solid, 59.4 mg (44% yield). m.p. 163~165 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.76 (d, $J=8.0$ Hz, 1H), 7.54 (t, $J=7.2$ Hz, 1H), 7.44 (t, $J=7.6$ Hz, 1H), 7.21 (t, $J=7.2$ Hz, 1H), 7.17 (s, 1H), 7.15~7.09 (m, 3H), 6.50 (d, $J=7.6$ Hz, 2H), 6.33 (s, 2H), 4.49 (q, $J=7.2$ Hz, 2H), 2.24 (s, 3H), 1.46 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 163.1, 143.4, 142.3, 140.6, 137.6, 135.9, 134.6 (2C), 129.8, 129.6, 129.5, 129.0 (3C), 128.2, 128.1 (2C), 128.0, 127.1, 126.6 (2C), 119.0, 60.9, 14.5, 9.7; HRMS (EI) calcd for $C_{25}H_{20}N_2O_2Cl_2$ 450.0910, found 450.0909.

4.3 Procedures for derivatizations of preparation and functionalization of the dual-core dimeric cyclopalladated intermediate II

A mixture of $PdCl_2$ (35.4 mg, 0.2 mmol, 1.0 equiv.), $NaOAc$ (49.2 mg, 0.6 mmol, 3.0 equiv.) and substrate **1c** (61.3 mg, 0.2 mmol, 1.0 equiv.) in $AcOH$ (1.0 mL) was charged in a glass sealed-tube and stirred under N_2 atmosphere at 100 °C for 4 h. Upon completion of the reaction, the reaction was diluted with dichloromethane and sequentially washed with water and brine. The crude mixture was purified by column chromatography to give the title complex **II** as a yellow solid (54.4 mg, 61%). A mixture of intermediate **II** (44.6 mg, 0.05 mmol), 4-iodoanisole (**2a**, 35.1 mg, 3.0 equiv.), $AgTFA$ (22.1 mg, 2.0 equiv.) in $HFIP/AcOH$ ($V:V=3:1$, 0.5 mL) was charged in a glass sealed-tube and stirred under air atmosphere at 120 °C for 20 h. Upon completion of the reaction, saturated brine (15 mL) and dichloromethane (15 mL) were added to the mixture, then the aqueous layer was extracted with dichloromethane (15 mL $\times$ 2). The combined organic layer was dried over anhydrous $MgSO_4$. Finally, the solution was concentrated *in vacuo* to provide a crude product, which was further purified *via* a column chromatography on silica gel [eluent: $V(\text{petroleum ether}):V(\text{ethyl acetate})=5:1$] to supply the product **3c** (20.6 mg, 67% yield).

References

- [1] (a) Giri, R.; Shi, B. F.; Engle, K. M.; Maugel, N.; Yu, J. Q. *Chem. Soc. Rev.* **2009**, 38, 3242.

- (b) Li, B. J.; Shi, Z. J. *Chem. Soc. Rev.* **2012**, *41*, 5588.
 (c) Louillat, M. L.; Patureau, F. W. *Chem. Soc. Rev.* **2014**, *43*, 901.
 (d) Chen, Z. K.; Wang, B. J.; Zhang, J. T.; Yu, W. L.; Liu, Z. X.; Zhang, Y. H. *Org. Chem. Front.* **2015**, *2*, 1107.
 (e) Gensch, T.; James, M. J.; Dalton, T.; Glorius, F. *Angew. Chem., Int. Ed.* **2018**, *57*, 2296.
 (f) Li, J. X.; Yang, S. R.; Wu, W. Q.; Jiang, H. F. *Eur. J. Org. Chem.* **2018**, 1284.
 (g) Li, J. X.; Yang, S. R.; Wu, W. Q.; Jiang, H. F. *Chem.-Asian J.* **2019**, *14*, 4114.
- [2] (a) Alberico, D.; Scott, M. E.; Lautens, M. *Chem. Rev.* **2007**, *107*, 174.
 (b) Seth, K.; Garg, S. K.; Kumar, R.; Purohit, P.; Meena, V. S.; Goyal, R.; Banerjee, U. C.; Chakraborti, A. K. *ACS Med. Chem. Lett.* **2014**, *5*, 512.
 [3] (a) Ashenurst, J. A. *Chem. Soc. Rev.* **2010**, *39*, 540.
 (b) Bugaut, X.; Glorius, F. *Angew. Chem., Int. Ed.* **2011**, *50*, 7479.
 (c) Cho, S. H.; Kim, J. Y.; Kwak, J.; Chang, S. *Chem. Soc. Rev.* **2011**, *40*, 5068.
 (d) Li, B.; Shi, Z. *Chem. Soc. Rev.* **2012**, *41*, 5588.
 [4] (a) Li, R.; Jiang, L.; Lu, W. *Organometallics* **2006**, *25*, 5973.
 (b) Kar, A.; Mangu, N.; Kaiser, H. M.; Tse, M. K. *J. Organomet. Chem.* **2009**, *694*, 524.
 (c) Zhou, L.; Lu, W. *Organometallics* **2012**, *31*, 2124.
 [5] (a) Ye, M. C.; Edmunds, A.; Morris, J.; Sale, D.; Zhang, Y.; Yu, J. Q. *Chem. Sci.* **2013**, *4*, 2374.
 (b) Han, J.; Liu, P.; Wang, C.; Wang, Q.; Zhang, J. Y.; Zhao, Y. W.; Shi, D. Q.; Huang, Z. B.; Zhao, Y. S. *Org. Lett.* **2014**, *16*, 5682.
 (c) Yang, Z.; Qiu, F. C.; Gao, J.; Li, Z. W.; Guan, B. T. *Org. Lett.* **2015**, *17*, 4316.
 (d) Xu, J. C.; Liu, Y.; Wang, Y.; Li, Y. J.; Xu, X. H.; Jin, Z. *Org. Lett.* **2017**, *19*, 1562.
 (e) Hu, Y. H.; Xu, Z.; Shao, L. Y.; Ji, Y. F. *Synlett* **2018**, *29*, 1875.
 (f) Yang, J. Y.; Fu, X. P.; Tang, S. B.; Deng, K. Z.; Zhang, L. L.; Yang, X. J.; Ji, Y. F. *J. Org. Chem.* **2019**, *84*, 10221.
 [6] (a) Deprez, N.; Sanford, M. S. *J. Am. Chem. Soc.* **2009**, *131*, 11234.
 (b) Li, W.; Yin, Z.; Jiang, X.; Sun, P. J. *Org. Chem.* **2011**, *76*, 8543.
 (c) Guo, D. D.; Li, B.; Guo, S. H.; Pan, G. F.; Gao, Y. R.; Wang, Y. Q. *ChemCatChem* **2017**, *9*, 2001.
 [7] (a) Shabasho, D.; Daugulis, O. *J. Am. Chem. Soc.* **2010**, *132*, 3965.
 (b) Yokota, A.; Aihara, Y.; Chatani, N. *J. Org. Chem.* **2014**, *79*, 11922.
 [8] (a) Sun, C. L.; Liu, N.; Li, B. J.; Yu, D. G.; Wang, Y.; Shi, Z. J. *Org. Lett.* **2010**, *12*, 184.
 (b) Thirunavukkarasu, V. S.; Cheng, C. H. *Chem.-Eur. J.* **2011**, *17*, 14723.
 (c) Shao, L. Y.; Xing, L. H.; Guo, Y.; Yu, K. K.; Wang, W.; Liu, H. W.; Liao, D. H.; Ji, Y. F. *Adv. Synth. Catal.* **2018**, *360*, 2925.
 [9] Li, W.; Xu, Z.; Sun, P.; Jiang, X.; Fan, M. *Org. Lett.* **2011**, *13*, 1286.
 [10] (a) Nishikata, T.; Abela, A. R.; Lipshutz, B. H. *Angew. Chem., Int. Ed.* **2010**, *49*, 781.
 (b) Jiang, Z.; Zhang, L.; Dong, C.; Su, X.; Li, H.; Tang, W.; Xu, L.; Fan, Q. *RSC Adv.* **2013**, *3*, 1025.
 (c) Li, D.; Xu, N.; Zhang, Y.; Wang, L. *Chem. Commun.* **2014**, *50*, 14862.
- [11] Zhang, Q.; Yin, X. S.; Zhao, S.; Fang, S. L.; Shi, B. F. *Chem. Commun.* **2014**, *50*, 8353.
 [12] (a) Cernak, T.; Dykstra, K. D.; Tyagarajan, S.; Vachal, P.; Krska, S. W. *Chem. Soc. Rev.* **2016**, *45*, 546.
 (b) Margrey, K. A.; Czaplinski, W. L.; Nicewicz, D. A.; Alexanian, E. J. *J. Am. Chem. Soc.* **2018**, *140*, 4213.
 (c) Brodney, M. A.; Sharma, R.; Lazzaro, J. T.; Walker, G. S.; Obach, R. S. *Bioorg. Med. Chem. Lett.* **2018**, *28*, 2068.
 (d) Graßl, S.; Chen, Y. H.; Hamze, C.; Tüllmann, C. P.; Knochel, P. *Org. Lett.* **2019**, *21*, 494.
 [13] (a) Paulis, T. D.; Hemstapat, K.; Chen, Y. L.; Zhang, Y. Q.; Saleh, S.; Alagille, D.; Baldwin, R. M.; Tamagnan, G. D.; Conn, P. J. *J. Med. Chem.* **2006**, *49*, 3332.
 (b) Lahm, G. P.; Cordova, D.; Barry, J. D. *Bioorg. Med. Chem.* **2009**, *17*, 4127.
 (c) Mowbray, C. E.; Burt, C.; Corbau, R.; Gayton, S.; Hawes, M.; Perros, M.; Tran, I.; Price, D. A.; Quinton, F. J.; Selby, M. D.; Stuppel, P. A.; Webster, R.; Wood, A. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 5857.
 (d) Alvarez, G.; Varela, J.; Cruces, E.; Fernández, M.; Gabay, M.; Leal, S. M.; Escobar, P.; Sanabria, L.; Serna, E.; Torres, S.; Thiel, S. J. F.; Yaluff, G.; Vera de Bilbao, N. I.; Cerecetto, H.; González, M. *Antimicrob. Agents Chemother.* **2015**, *59*, 1398.
 [14] (a) Shabashov, D.; Daugulis, O. *Org. Lett.* **2005**, *7*, 3657.
 (b) Cheng, K.; Zhang, Y.; Zhao, J.; Xie, C. *Synlett* **2008**, 1325.
 (c) Umeda, N.; Hirano, K.; Satoh, T.; Miura, M. *J. Org. Chem.* **2009**, *74*, 7094.
 (d) Arockiam, P. B.; Fischmeister, C.; Bruneau, C.; Dixneuf, P. H. *Green Chem.* **2011**, *13*, 3075.
 (e) Thirunavukkarasu, V. S.; Raghuvanshi, K.; Ackermann, L. *Org. Lett.* **2013**, *15*, 3286.
 (f) Yu, X.; Yu, S.; Xiao, J.; Wan, B.; Li, X. J. *Org. Chem.* **2013**, *78*, 5444.
 (g) Liu, P. M.; Frost, C. G. *Org. Lett.* **2013**, *15*, 5862.
 (h) Fabre, I.; Wolff, N.; Duc, G.; Flegeau, E.; Bruneau, C.; Dixneuf, P.; Jutand, A. *Chem.-Eur. J.* **2013**, *19*, 7595.
 (i) Lu, M. Z.; Lu, P.; Xu, Y. H.; Loh, T.-P. *Org. Lett.* **2014**, *16*, 2614.
 (j) Han, S.; Sharma, S.; Park, J.; Kim, M.; Shin, Y.; Mishra, N. K.; Bae, J. J.; Kwak, J. H.; Jung, Y. H.; Kim, I. S. *J. Org. Chem.* **2014**, *79*, 275.
 (k) Reddy, G. M.; Rao, N. S.; Satyanarayana, P.; Maheswaran, H. *RSC Adv.* **2015**, *5*, 105347.
 (l) Yang, P.; Bao, Y. S. *RSC Adv.* **2017**, *7*, 53878.
 (m) Kwak, S. H.; Gulia, N.; Daugulis, O. *J. Org. Chem.* **2018**, *83*, 5844.
 (n) Abidi, O.; Boubaker, T.; Hierso, J.; Roger, J. *Org. Biomol. Chem.* **2019**, *17*, 5916.
- [15] (a) Fan, X. M.; Guo, Y.; Li, Y. D.; Yu, K. K.; Liu, H. W.; Liao, D. H.; Ji, Y. F. *Asian J. Org. Chem.* **2016**, *5*, 499.
 (b) Chen, M. M.; Shao, L. Y.; Lun, L. J.; Wu, Y. L.; Fu, X. P.; Ji, Y. F. *Chin. Chem. Lett.* **2019**, *30*, 702.
- [16] Ren, Z.; Dong, G. B. *Organometallics* **2016**, *35*, 1057.

(Cheng, F.)