KX/Oxone 体系实现的 *N*-烯丙基芳酰胺的氧化卤环化反应:
5-卤甲基-2-噁唑啉的绿色合成刘新明^a 李 路^a 晋李成^a 赵金灿^{*,b} 华远照^c
王敏灿^c 刘澜涛^{*,a,c}^(a) 商丘师范学院化学化工学院 河南商丘 476000)^(b) 河北大学化学与环境科学学院 河北保定 071002)^(c) 郑州大学化学学院 郑州 450001)

摘要 以廉价的卤化钾为卤源, Oxone 为氧化剂, 一系列烯丙胺在温和的条件下经过卤环化反应以较好的产率得到 5-卤甲基-2-噁唑啉。该方法的优势在于应用了便宜的卤源, 没有有机副产物的产生, 具有成本低廉和环保友好的特点。5-卤甲基-2-噁唑啉还可以通过不同的亲核取代反应得到许多有用的衍生物。

关键词 Oxone; 卤环化; 噁唑啉

Oxidative Halocyclization of *N*-Allylarylamides with KX/Oxone
System: Green Synthesis of 5-Halomethyl-2-OxazolinesLiu, Xinming^a Li, Lu^a Jin, Licheng^a Zhao, Jincan^{*,b} Hua, Yuanzhao^c
Wang, Mincan^c Liu, Lantao^{*,a,c}^(a) College of Chemistry and Chemical Engineering, Shangqiu Normal University, Shangqiu, Henan 476000)^(b) College of Chemistry & Environmental Science, Hebei University, Baoding, Hebei 071002)^(c) College of Chemistry, Zhengzhou University, Zhengzhou 450001)

Abstract With inexpensive potassium halide (KX) as halogen source, and oxone as oxidant, a series of allylamides underwent halocyclization reaction and generated 5-halomethyl-2-oxazolines in good to excellent isolated yields under mild conditions. The protocol showed attractive advanced features including low cost of halogen source, absence of organic byproduct, and resultant environmental-friendly nature. In addition, various useful derivatives could be expected by proper nucleophilic substitution reactions.

Keywords oxone; halocyclization; oxazolines

1 Introduction

2-Oxazolines as a class of prominent compounds, are important building blocks of natural products, biologically active compounds and pharmaceuticals, and have also been widely utilized as key structures, protective groups,^[1] chiral ligands,^[2-3] chiral auxiliaries^[4-8] and catalyst^[9-17] in organic synthesis or materials science. Great efforts have been devoted to their efficient synthesis, and up to now, significant

methods have been developed by using carboxylic acids, carboxylic esters, nitriles, aldehydes, olefins and β -hydroxyamides as starting materials.^[18-20]

5-Halomethyl-2-oxazolines with a halomethyl group are more attractive for their versatile reactivities. Several efficient methods for their synthesis have been developed during the past decades. Particularly, the methods based on the halocyclization reaction of cheap and readily available *N*-allylamides attracted more and more attentions.^[21-28]

* Corresponding authors. E-mail: liult05@iccas.ac.cn.; jczhao@hbu.edu.cn

Received May 29, 2020; revised July 2, 2020; published online July 22, 2020.

Project supported by the National Natural Science Foundation of China (No. 21572126), the Program of Science and Technology Innovation Talents of Henan Province (No. 2018JQ0011), the Natural Science Foundation of Hebei Province (No. B2019201415), and the Fundamental Research Funds for the Midwest Universities Comprehensive Strength Promotion Project (No. 521000981026).

国家自然科学基金(No. 21572126)、河南省科技创新人才(No. 2018JQ0011)、河北省自然科学基金(No. B2019201415)和中西部高校综合提升工程(No. 521000981026)资助项目。

While several elegant asymmetric versions with organic halogenating reagents have been realized under mild reaction conditions (Eq. 1a).^[29-33] Much attention has been paid for the development of different oxidant promoter/halogenating reagent systems, such as PIDA/TMSX (Eq. 1b),^[24] CuBr₂,^[27] and Selectfluor/Pyr•(HF)_x/Et₃N•3HF systems,^[34] for the transformation. Wang *et al.*^[26] reported the iodine-promoted regioselective cyclization of *N*-allylamides with sulfonyl hydrazides for the synthesis of 5-methyl-arylthio-oxazolines *via* sulfonylation reactions (Eq. 1c). Very recently, Waldvogel *et al.*^[35] achieved a hypervalent iodoarene-mediated fluorocyclization of *N*-allylcarboxamide to 5-fluoromethyl-2-oxazoline by using electric current as the only oxidant, and Et₃N•5HF as fluorinating agents (Eq. 1d). Despite of the great progress, some of them suffered from the use of expensive organic halogenating agents or organic terminal oxidants, special equipment, and especially, the generation of hazardous organic byproducts, or heavy metal residues. The transition-metal-free systems using inorganic oxidant promoter/halogenating reagents are highly desirable.

Considering to the multifaceted advantages including low-cost, stability, non-toxicity, and air/moisture-insensitiveness, oxone has been widely used as inorganic oxidant for halides to generate halogenating agents.^[36] Meanwhile KX is a kind of cheap and ideal halogenating reagent source. Thus, KX/oxone system is an ideal environmentally friendly protocol for organic reactions with inorganic K₂SO₄ as the byproduct.^[37-40] Herein, the green synthesis of

5-halomethyl-2-oxazolines *via* oxidative halocyclization of allylamides with KX/oxone is reported (Eq. 1e).

2 Results and discussion

Inspired by recent progress, an intramolecular iodoxy-generation reaction was proposed using *N*-allyl carboxamide **1a** as the model substrate and KI as halogen source. With the success of reaction, the conditions including the amount of oxone and KI, reaction time and temperature were optimized. It is indicated that 5 equiv. of KI and 2 equiv. of oxone to **1a** are enough for the conversion. 4 h is best for this reaction (Table 1, Entries 5~8), and 0 °C promised best conversion (Table 1, Entries 9~11). It was interesting to find that when the ratio of H₂O to CH₃CN increased from 0.5 : 1 to 1 : 1 (Table 1, Entries 11~12), the yield reached nearly 90%, which decreased with further increasing of water ratio of 1 : 0.5. The dependence of the conversion on the water indicated that the solubility of the media toward both inorganic salt of KI and the organic reagents should be well-balanced.^[41-44]

Table 1 Reaction optimization studies

Entry	KI/ equiv.	Oxone/ equiv.	CH ₃ CN/ mL	H ₂ O/ mL	Temp/ °C	Time/ h	Yield ^a / %
1	2.5	1	1	0.5	25	6	21.9
2	5.0	2	1	0.5	25	6	29.6
3	7.5	3	1	0.5	25	6	29.3
4	10.0	4	1	0.5	25	6	30.8
5	5.0	2	1	0.5	25	2	45.8
6	5.0	2	1	0.5	25	4	51.7
7	5.0	2	1	0.5	25	6	29.8
8	5.0	2	1	0.5	25	8	23.1
9	5.0	2	1	0.5	-20	4	36.3
10	5.0	2	1	0.5	0	4	80.9
11	5.0	2	1	0.5	50	4	48.9
12	5.0	2	1	1.0	0	4	88.0
13	5.0	2	0.5	1.0	0	4	56.8

^a Isolated yield.

On the optimized conditions, carboxamide substrates **1b**~**1w** were then subjected to the same reaction to test the scope of the reaction (Eq. 2). Seen from the results, 5-iodomethyl-2-aryloxazolines could be obtained in good to excellent yields. All substrates **2b**~**2t** with either electron-donating or electron-withdrawing groups on aromatic rings could be cyclized in 66.5%~98.8% yields. The yields of reagents with electron-donating substituents including Me, MeO, ^tBu, Ph were higher than those with electron-withdrawing substituents including F and NO₂. Moreover, the reagents with 2- and 4-position occupied gave higher yields than those with 3-position substituted. Especially, substrate **1t** bearing 4-phenyl group worked very

well in the current reaction system, and generated the corresponding 2-([1,1'-biphenyl]-4-yl)-5-(iodomethyl)-4,5-dihydrooxazoline (**2t**) with up to 98.8% yield. Oxazoline structures bearing heterocycles have been used as important functional groups in antibiotic, antidiabetic and antihypertensive agents. To this end, structurally diversified heterocyclic oxazoline compounds were also prepared using the same method. As shown in Eq. 2, the thiophene- and pyridine-containing oxazoline products **2u** and **2v** could also be obtained in 84.6% and 55.8% yields, respectively. Furthermore, the current protocol was also employed to the synthesis of bis(oxazoline) compound **2w** in 58.1% yield.

Reaction conditions: substrate (0.5 mmol), KI (5.0 mmol), oxone (2 mmol), CH₃CN (1 mL), H₂O (1 mL), 0 °C, 4 h. 10 mmol of KI and 4 mmol of oxone were used for the preparation of **2w**. Isolated yield.

To demonstrate the scalability of the protocol, a gram-scale synthesis of **2d** was also carried out. Cyclization of *N*-allyl-4-methyl benzamide **1d** on 1.0 g scale afforded the desired 2,5-disubstituted oxazoline **2d** in 90.6% yield (Eq. 3).

To test the synthetic utility of current reaction, functional group transformations of **2d** were also pursued, and the C—I bond in **2d** provided an easy access to other functional groups (Scheme 1). For example, iodine group in **2d** was replaced by different nucleophiles, such as azide (**3a**) and

benzenethiol (**3b**) under mild conditions.

Scheme 1 Derivatization of oxazoline **2d**

To demonstrate the universality of the protocol, some less active KX reagents such as KCl, KBr were tested with *N*-allyl-4-methyl benzamide **1d** as the substrate, the corresponding 5-bromo- and 5-chloro-methyl-2-oxazolines **4a** and **4b** were obtained in 81.3% and 55.8% isolated yields respectively under the similar experimental conditions (Scheme 2).

Scheme 2 Synthesis of 5-chloro- and 5-bromo-methyl-2-aryl-4,5-dihydrooxazolines

Based on the results and previous literature reports,^[36,45] a preliminary reaction pathway is put forward (Scheme 3). Firstly, the hypoiodous acid is generated *in situ* from the reaction of oxone with KI.^[36] Afterwards, *N*-allylbenzamide forms an intermediate (iodonium ion herein) which is then intramolecular nucleophilic attacked by oxygen atom. A five-membered ring is formed and haloxygenation products are provided.

3 Conclusions

In summary, a practical method for preparation of 5-halomethyloxazolines with oxone as the reaction promoter and KX (X = I, Br and Cl) as the halogen sources was reported. 5-Halomethyl-2-oxazoline products could be achieved in good to excellent yields. The method has the following advanced features: (i) all the starting materials such as acid and acyl chloride are cheap and commercial available; (ii) the reaction is operationally simple, without

Scheme 3 Plausible mechanism for intramolecular haloxygenation reaction

requirement of strict anhydrous or inert reaction conditions only inorganic byproducts K_2SO_4 are generated from the reaction; (iii) the reaction scope is universal and tolerate for both electron withdrawing groups and electron-donating groups. What's more, the 5-chloro and 5-bromo products can also be achieved effectively. It is reasonable to believe that the current method will have widespread application in organic and medicinal chemistry.

4 Experiment

4.1 General information

Reagents were used as received without further purification unless otherwise indicated. Reactions were monitored with thin layer chromatography using silica gel GF254 plates. Organic solutions were concentrated *in vacuo* with a rotavapor. Flash column chromatography was performed using silica gel (200~300 meshes). Petroleum ether used has a boiling point range of 60~90 °C. Melting points were measured on a digital melting point apparatus without correction of the thermometer. Nuclear magnetic resonance spectra were recorded at ambient temperature (unless otherwise stated) at 400 MHz (100 MHz for ^{13}C NMR) in CDCl_3 . Chemical shifts were reported in δ using TMS as internal standard. High resolution mass spectrometry (HRMS) analyses were carried out on an Accurate-Mass Q-TOF. Substrates used were prepared by coupling of carboxylic acids or acyl chloride and allylamine described by Borhan *et al.*^[30]

4.2 General procedure for synthesis of 5-halomethyl-2-oxazolines.

In a 10 mL long neck tube, *N*-allylcarboxamides (0.5 mmol), oxone (1.0 mmol), KX (2.5 mmol), H_2O (1 mL) and CH_3CN (1 mL) were added. The reaction mixture was then stirred at 0 °C for 4 h. Ethyl acetate (10 mL) was then added, and the mixture was washed with aqueous $\text{Na}_2\text{S}_2\text{O}_3$. The combined organic layer was dried (Na_2SO_4) and concentrated to give crude residue, which was purified by flash column chromatography to give the corresponding products.

5-(Iodomethyl)-2-phenyl-4,5-dihydrooxazole (**2a**): Ob-

tained as an oil in 88.0% yield (126.3 mg) after flash chromatography (silica gel, petroleum ether/ethyl acetate, $V:V=5:1$). ^1H NMR (400 MHz, CDCl_3) δ : 7.94~7.91 (m, 2H), 7.51~7.44 (m, 1H), 7.42~7.37 (m, 2H), 4.81~4.75 (m, 1H), 4.19~4.11 (m, 1H), 3.82~3.75 (m, 1H), 3.40~3.26 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.4, 131.4, 128.3, 128.1, 127.4, 78.2, 60.7, 7.7. Spectral data are in good agreement with literature values.^[24-45]

5-(Iodomethyl)-2-(*o*-tolyl)-4,5-dihydrooxazole (**2b**): Obtained as an oil in 93.3% yield (140.5 mg) after flash chromatography (silica gel, petroleum ether/ethyl acetate, $V:V=4:1$). ^1H NMR (400 MHz, CDCl_3) δ : 7.81 (d, $J=8.0$ Hz, 1H), 7.36~7.31 (m, 1H), 7.24~7.20 (m, 2H), 4.73~4.70 (m, 1H), 4.22~4.14 (m, 1H), 3.85~3.80 (m, 1H), 3.35~3.32 (m, 2H), 2.60 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.8, 138.8, 131.2, 130.7, 129.8, 126.7, 125.5, 77.3, 61.0, 21.9, 8.1. Spectral data are in good agreement with literature values.^[24-45]

5-(Iodomethyl)-2-(*m*-tolyl)-4,5-dihydrooxazole (**2c**): Obtained as an oil in 83.3% yield (125.4 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=4:1$). ^1H NMR (400 MHz, CDCl_3) δ : 7.77 (s, 1H), 7.73 (t, $J=3.6$ Hz, 1H), 7.33~7.28 (m, 2H), 4.83~4.77 (m, 1H), 4.18 (dd, $J=15.2, 9.6$ Hz, 1H), 3.80 (dd, $J=15.2, 6.8$ Hz, 1H), 3.39 (dd, $J=10.4, 4.8$ Hz, 1H), 3.32 (dd, $J=10.4, 3.2$ Hz, 1H), 2.39 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.4, 137.9, 132.1, 128.6, 128.1, 127.1, 125.1, 78.0, 60.5, 21.1, 7.7. Spectral data are in good agreement with literature values.^[24-45]

5-(Iodomethyl)-2-(*p*-tolyl)-4,5-dihydrooxazole (**2d**): Obtained as a white solid in 93.4% yield (140.6 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=4:1$); m.p. 94~96 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.82 (d, $J=8.4$ Hz, 2H), 7.22 (d, $J=8.0$ Hz, 2H), 4.82~4.78 (m, 1H), 4.17 (dd, $J=15.2, 9.6$ Hz, 1H), 3.80 (dd, $J=14.8, 6.4$ Hz, 1H), 3.38 (dd, $J=10.4, 5.2$ Hz, 1H), 3.31 (dd, $J=10.0, 7.2$ Hz, 1H), 2.39 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.4, 141.7, 128.9, 128.0, 124.5, 78.0, 60.5, 21.4, 7.8. Spectral data are in good agreement with literature values.^[24-45]

5-(Iodomethyl)-2-(2-methoxyphenyl)-4,5-dihydrooxazole (**2e**): Obtained as an oil in 91.2% yield (144.6 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). ^1H NMR (400 MHz, CDCl_3) δ : 7.80 (dd, $J=8.0, 1.6$ Hz, 1H), 7.46~7.42 (m, 1H), 7.01~6.97 (m, 2H), 4.78~4.73 (m, 1H), 4.24 (dd, $J=15.2, 9.6$ Hz, 1H), 3.93 (s, 3H), 3.88 (dd, $J=14.4, 6.8$ Hz, 1H), 3.38 (dd, $J=10.4, 5.2$ Hz, 1H), 3.31 (dd, $J=10.0, 7.2$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.6, 158.4, 132.4, 131.0, 120.0, 116.2, 111.4, 76.9, 61.2, 55.8, 7.7. HRMS-ESI calcd for $\text{C}_{11}\text{H}_{12}\text{INO}_2$ $[\text{M}+\text{H}]^+$ 317.9991, found 317.9981.

5-(Iodomethyl)-2-(3-methoxyphenyl)-4,5-dihydrooxazole (**2f**): Obtained as an oil in 80.2% yield (127.2 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). ^1H NMR (400 MHz, CDCl_3) δ : 7.44 (d, $J=2.4$ Hz, 1H), 7.38~7.37 (m, 1H), 7.23 (t, $J=8.0$ Hz, 1H), 6.95~6.92 (m, 1H), 4.73~4.66 (m, 1H), 4.07 (dd,

$J=15.2, 9.2$ Hz, 1H), 3.74 (s, 3H), 3.70 (dd, $J=15.2, 6.8$ Hz, 1H), 3.27 (dd, $J=10.4, 5.2$ Hz, 1H), 3.21 (dd, $J=10.4, 7.2$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.2, 159.3, 129.3, 128.4, 120.4, 118.0, 112.4, 78.1, 60.5, 55.2, 7.7. Spectral data are in good agreement with literature values.^[24-45]

5-(Iodomethyl)-2-(4-methoxyphenyl)-4,5-dihydrooxazole (**2g**): Obtained as a white solid in 80.2% yield (151.4 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). m.p. 104~106 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.87 (d, $J=8.8$ Hz, 2H), 6.91 (d, $J=8.8$ Hz, 2H), 4.81~4.74 (m, 1H), 4.15 (dd, $J=15.2, 9.6$ Hz, 1H), 3.83 (s, 3H), 3.77 (dd, $J=15.2, 6.8$ Hz, 1H), 3.36 (dd, $J=10.4, 5.2$ Hz, 1H), 3.30 (dd, $J=10.0, 6.8$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.3, 162.1, 129.8, 119.6, 113.6, 78.1, 60.4, 55.2, 7.7. Spectral data are in good agreement with literature values.^[24-45]

5-(Iodomethyl)-2-(2-fluorophenyl)-4,5-dihydrooxazole (**2h**): Obtained as an oil in 66.5% yield (101.4 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). ^1H NMR (400 MHz, CDCl_3) δ : 7.89~7.84 (m, 1H), 7.49~7.43 (m, 1H), 7.23~7.13 (m, 2H), 4.83~4.77 (m, 1H), 4.22 (dd, $J=15.2, 9.6$ Hz, 1H), 3.86 (dd, $J=15.6, 6.8$ Hz, 1H), 3.41~3.28 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.0 (d, $J=256.8$ Hz), 159.9 (d, $J=5.7$ Hz), 132.9 (d, $J=8.7$ Hz), 130.8 (d, $J=1.4$ Hz), 123.9 (d, $J=3.8$ Hz), 116.5 (d, $J=21.7$ Hz), 115.6 (d, $J=10.3$ Hz), 77.5, 60.9, 7.5; ^{19}F NMR (376 MHz, CDCl_3) δ : -109.4. Spectral data are in good agreement with literature values.^[24-45]

5-(Iodomethyl)-2-(4-fluorophenyl)-4,5-dihydrooxazole (**2i**): Obtained as a white solid in 88.6% yield (135.2 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). m.p. 112~113 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.94~7.90 (m, 2H), 7.08 (t, $J=8.8$ Hz, 2H), 4.81~4.77 (m, 1H), 4.15 (dd, $J=15.2, 9.6$ Hz, 1H), 3.77 (dd, $J=15.2, 6.8$ Hz, 1H), 3.36 (dd, $J=10.4, 5.2$ Hz, 1H), 3.31 (dd, $J=10.0, 7.2$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 164.7 (d, $J=250.5$ Hz), 162.5, 130.4 (d, $J=89.0$ Hz), 123.6 (d, $J=3.1$ Hz), 115.5 (d, $J=21.9$ Hz), 78.4, 60.7, 69.5, 7.6; ^{19}F NMR (376 MHz, CDCl_3) δ : -107.8. Spectral data are in good agreement with literature values.^[24-45]

5-(Iodomethyl)-2-(2-nitrophenyl)-4,5-dihydrooxazole (**2j**): Obtained as an oil in 81.3% yield (135.0 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). ^1H NMR (400 MHz, CDCl_3) δ : 7.86 (dd, $J=8.0, 1.6$ Hz, 1H), 7.83 (dd, $J=8.0, 1.6$ Hz, 1H), 7.67~7.62 (m, 2H), 4.88~4.84 (m, 1H), 4.20 (dd, $J=15.2, 9.6$ Hz, 1H), 3.86 (dd, $J=15.2, 6.4$ Hz, 1H), 3.38 (dd, $J=10.4, 5.2$ Hz, 1H), 3.33 (dd, $J=10.4, 7.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 160.7, 148.8, 132.5, 131.5, 131.0, 123.8, 122.7, 79.2, 60.8, 6.9. HRMS-ESI calcd for $\text{C}_{11}\text{H}_{12}\text{INO}_2$ $[M+H]^+$ 332.9736, found 332.9729.

5-(Iodomethyl)-2-(3-nitrophenyl)-4,5-dihydrooxazole (**2k**): Obtained as a white solid in 73.0% yield (121.2 mg) after flash chromatography (Silica gel, petroleum ether/

ethyl acetate, $V:V=3:1$). m.p. 85~88 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.77 (s, 1H), 8.35 (d, $J=8.4$ Hz, 1H), 8.28 (d, $J=8.0$ Hz, 1H), 7.63 (t, $J=8.0$ Hz, 1H), 4.90~4.83 (m, 1H), 4.24 (dd, $J=15.6, 9.6$ Hz, 1H), 3.86 (dd, $J=15.6, 9.6$ Hz, 1H), 3.43~3.29 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.4, 148.0, 133.8, 129.5, 129.1, 125.9, 123.1, 78.6, 60.8, 7.56. HRMS-ESI calcd for $\text{C}_{11}\text{H}_{12}\text{INO}_2$ $[M+H]^+$ 332.9736, found 332.9724.

5-(Iodomethyl)-2-(4-nitrophenyl)-4,5-dihydrooxazole (**2l**): Obtained as a white solid in 80.4% yield (133.5 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). m.p. 124~125 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.27 (dd, $J=8.8, 1.6$ Hz, 2H), 8.10 (dd, $J=8.8, 2.0$ Hz, 2H), 4.89~4.81 (m, 1H), 4.23 (dd, $J=15.6, 9.6$ Hz, 1H), 3.85 (dd, $J=15.6, 6.8$ Hz, 1H), 3.42~3.34 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 161.7, 149.5, 133.2, 129.2, 123.6, 78.7, 61.1, 7.4. Spectral data are in good agreement with literature values.^[24-45]

5-(Iodomethyl)-2-(3-bromophenyl)-4,5-dihydrooxazole (**2m**): Obtained as an oil in 81.4% yield (149.0 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). ^1H NMR (400 MHz, CDCl_3) δ : 8.07 (t, $J=4.0$ Hz, 1H), 7.86~7.84 (m, 1H), 7.61~7.58 (m, 1H), 7.30~7.26 (m, 1H), 4.83~4.76 (m, 1H), 4.17 (dd, $J=15.2, 9.2$ Hz, 1H), 3.79 (dd, $J=15.2, 6.8$ Hz, 1H), 3.36 (dd, $J=10.4, 4.8$ Hz, 1H), 3.33 (dd, $J=10.0, 7.2$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.1, 134.4, 131.1, 129.9, 129.3, 126.7, 122.3, 78.4, 60.7, 7.5. Spectral data are in good agreement with literature values.^[24-45]

5-(Iodomethyl)-2-(4-bromophenyl)-4,5-dihydrooxazole (**2n**): Obtained as a white solid in 90.9% yield (166.3 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). m.p. 115~117 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.78 (d, $J=8.8$ Hz, 2H), 7.54 (d, $J=8.8$ Hz, 2H), 4.82~4.75 (m, 1H), 4.15 (dd, $J=15.2, 9.6$ Hz, 1H), 3.77 (dd, $J=15.2, 6.8$ Hz, 1H), 3.36 (dd, $J=10.4, 5.2$ Hz, 1H), 3.31 (dd, $J=10.0, 6.8$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 162.6, 131.6, 129.6, 126.3, 126.2, 78.4, 60.8, 7.6. Spectral data are in good agreement with literature values.^[24-45]

5-(Iodomethyl)-2-(4-(*tert*-butyl)phenyl)-4,5-dihydrooxazole (**2o**): Obtained as a white solid in 94.8% yield (162.7 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). m.p. 104~106 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.86 (d, $J=8.5$ Hz, 2H), 7.44 (d, $J=8.6$ Hz, 2H), 4.88~4.74 (m, 1H), 4.17 (dd, $J=15.2, 9.2$ Hz, 1H), 3.80 (dd, $J=15.2, 6.4$ Hz, 1H), 3.38 (dd, $J=10.4, 4.8$ Hz, 1H), 3.31 (dd, $J=10.0, 7.2$ Hz, 1H), 1.33 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.3, 154.8, 127.9, 125.2, 124.5, 78.0, 60.6, 34.8, 31.0, 7.8. Spectral data are in good agreement with literature values.^[24-45]

N-Allyl-4-ethylbenzamide (**1p**): Obtained as an oil, ^1H NMR (400 MHz, CDCl_3) δ : 7.72 (d, $J=8.0$ Hz, 2H), 7.22 (d, $J=8.0$ Hz, 2H), 6.58 (s, 1H), 5.70~5.95 (m, 1H), 5.30~5.10 (m, 2H), 4.13~4.00 (m, 2H), 2.67 (q, $J=7.6$ Hz, 2H), 1.23 (t, $J=7.6$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 167.4, 148.0, 134.2, 131.7, 127.9, 127.0, 116.3,

42.3, 28.6, 15.2. HRMS-ESI calcd for $C_{12}H_{14}INO$ $[M+H]^+$ 190.1232, found 190.1225.

5-(Iodomethyl)-2-(4-ethylphenyl)-4,5-dihydrooxazole (**2p**): Obtained as a white solid in 83.7% yield (131.9 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). m.p. 104~106 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.84 (d, $J=8.0$ Hz, 2H), 7.24 (d, $J=8.4$ Hz, 2H), 4.83~4.73 (m, 1H), 4.15 (dd, $J=15.2, 9.6$ Hz, 1H), 3.78 (dd, $J=15.2, 6.4$ Hz, 1H), 3.37 (dd, $J=10.4, 4.8$ Hz, 1H), 3.30 (dd, $J=10.0, 7.2$ Hz, 1H), 2.68 (q, $J=7.6$ Hz, 2H), 1.24 (t, $J=7.6$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 163.5, 148.1, 128.2, 127.9, 124.8, 78.1, 60.7, 28.8, 15.3, 7.8. HRMS-ESI calcd for $C_{12}H_{14}INO$ $[M+H]^+$ 316.0198, found 316.0190.

5-(Iodomethyl)-2-(4-chlorophenyl)-4,5-dihydrooxazole (**2q**): Obtained as an oil in 86.3% yield (138.7 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). 1H NMR (400 MHz, $CDCl_3$) δ : 7.85 (d, $J=8.4$ Hz, 2H), 7.38 (d, $J=8.8$ Hz, 2H), 4.83~4.76 (m, 1H), 4.16 (dd, $J=15.2, 9.6$ Hz, 1H), 3.78 (dd, $J=15.6, 6.8$ Hz, 1H), 3.36 (dd, $J=10.4, 4.8$ Hz, 1H), 3.31 (dd, $J=10.0, 6.8$ Hz, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 162.6, 137.7, 129.5, 128.7, 125.8, 78.4, 60.7, 7.6. Spectral data are in good agreement with literature values.^[24-45]

5-(Iodomethyl)-2-(3,5-dimethylphenyl)-4,5-dihydrooxazole (**2r**): Obtained as an oil in 84.6% yield (133.3 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). 1H NMR (400 MHz, $CDCl_3$) δ : 7.55 (s, 2H), 7.11 (s, 1H), 4.80~7.5 (m, 1H), 4.15 (dd, $J=15.2, 9.6$ Hz, 1H), 3.78 (dd, $J=15.2, 6.8$ Hz, 1H), 3.37 (dd, $J=10.4, 4.8$ Hz, 1H), 3.31 (dd, $J=10.0, 7.2$ Hz, 1H), 2.34 (s, 6H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 163.8, 138.0, 133.2, 127.1, 125.9, 78.1, 60.6, 21.1, 7.8. Spectral data are in good agreement with literature values.^[24-45]

5-(Iodomethyl)-2-(2,4,6-trimethylphenyl)-4,5-dihydrooxazole (**2s**): Obtained as an oil in 94.5% yield (155.5 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). m.p. 104~106 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 6.86 (s, 2H), 4.81~4.76 (m, 1H), 4.20 (dd, $J=15.2, 9.6$ Hz, 1H), 3.85 (dd, $J=15.2, 7.2$ Hz, 1H), 3.41 (dd, $J=10.4, 5.2$ Hz, 1H), 3.32 (dd, $J=10.0, 7.2$ Hz, 1H), 2.31 (s, 6H), 2.28 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 163.8, 139.3, 136.9, 128.2, 125.4, 77.9, 60.7, 21.1, 19.8, 7.5. HRMS-ESI calcd for $C_{13}H_{16}INO$ $[M+H]^+$ 330.0355, found 330.0350.

5-(Iodomethyl)-2-(1,1'-biphenyl)-4,5-dihydrooxazole (**2t**): Obtained as a white solid in 98.8% yield (179.4 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). m.p. 107~109 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.01 (d, $J=8.4$ Hz, 2H), 7.67~7.62 (m, 4H), 7.46 (t, $J=7.6$ Hz, 2H), 7.39 (t, $J=7.2$ Hz, 1H), 4.88~4.81 (m, 1H), 4.21 (dd, $J=15.2, 9.6$ Hz, 1H), 3.84 (dd, $J=15.2, 6.4$ Hz, 1H), 3.41 (dd, $J=10.4, 4.8$ Hz, 1H), 3.35 (dd, $J=10.0, 7.2$ Hz, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 163.4, 144.2, 140.1, 128.9, 128.6, 127.9, 127.2, 127.0, 126.2, 60.8, 7.7. HRMS-ESI calcd for $C_{16}H_{14}INO$ $[M+H]^+$ 364.0198, found 364.0197.

5-(Iodomethyl)-2-(thiophen-2-yl)-4,5-dihydrooxazole (**2u**): Obtained as a white solid in 84.6% yield (123.6 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). m.p. 100~102 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.60 (d, $J=3.6$ Hz, 1H), 7.47 (dd, $J=5.2, 1.2$ Hz, 1H), 7.09 (dd, $J=4.8, 4.0$ Hz, 1H), 4.84~4.80 (m, 1H), 4.17 (dd, $J=15.2, 9.6$ Hz, 1H), 3.79 (dd, $J=15.2, 6.4$ Hz, 1H), 3.39 (dd, $J=10.4, 4.8$ Hz, 1H), 3.31 (dd, $J=10.0, 7.2$ Hz, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 159.0, 130.3, 129.9, 129.8, 127.4, 78.5, 60.6, 7.4. Spectral data are in good agreement with literature values.^[24-45]

5-(Iodomethyl)-2-(pyridine-4-yl)-4,5-dihydrooxazole (**2v**): Obtained as a white solid in 55.8% yield (80.4 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). m.p. 103~104 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 8.73 (d, $J=6.0$ Hz, 1H), 7.77 (d, $J=6.0$ Hz, 1H), 4.89~4.80 (m, 1H), 4.22 (dd, $J=15.7, 9.6$ Hz, 1H), 3.84 (dd, $J=15.7, 6.8$ Hz, 1H), 3.42~3.32 (m, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 161.7, 150.2, 134.6, 121.7, 78.4, 60.8, 7.3. Spectral data are in good agreement with literature values.^[24-45]

1,4-Bis(5-(iodomethyl)-4,5-dihydrooxazol-2-yl)benzene (**2w**): Obtained as a white solid in 58.1% yield (144.1 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). m.p. 189~191 °C; 1H NMR (400 MHz, $CDCl_3$) δ : 7.98 (s, 4H), 4.86~4.81 (m, 2H), 4.21 (dd, $J=15.2, 9.6$ Hz, 2H), 3.83 (dd, $J=15.7, 6.8$ Hz, 2H), 3.40 (dd, $J=10.4, 4.8$ Hz, 2H), 3.34 (dd, $J=10.0, 7.2$ Hz, 2H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 162.8, 130.1, 128.2, 78.4, 60.8, 7.5. HRMS-ESI calcd for $C_{14}H_{14}I_2N_2O_2$, 496.9223 $[M+H]^+$ found 496.9197.

5-(Azidomethyl)-2-(*p*-tolyl)-4,5-dihydrooxazole (**3a**): Obtained as an oil in 93.2% yield (100.8 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). 1H NMR (400 MHz, $CDCl_3$) δ : 7.83 (d, $J=8.0$ Hz, 2H), 7.21 (d, $J=8.0$ Hz, 2H), 4.88~4.83 (m, 1H), 4.11 (dd, $J=14.8, 10.0$ Hz, 1H), 3.78 (dd, $J=14.8, 10.0$ Hz, 1H), 3.38~3.46 (m, 2H), 2.37 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 163.7, 141.9, 129.0, 128.1, 124.4, 78.0, 57.6, 53.9, 21.5. HRMS-ESI calcd for $C_{11}H_{12}N_4O$ $[M+H]^+$ 217.1089, found 217.1081.

5-(((4-Methoxyphenyl)thio)methyl)-2-(*p*-tolyl)-4,5-dihydrooxazole (**3b**): Obtained as an oil in 84.6% yield (132.6 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). 1H NMR (400 MHz, $CDCl_3$) δ : 7.74 (d, $J=8.2$ Hz, 2H), 7.42 (d, $J=8.8$ Hz, 2H), 7.18 (d, $J=7.7$ Hz, 2H), 6.83 (d, $J=8.8$ Hz, 2H), 4.80~4.73 (m, 1H), 4.12 (dd, $J=15.2, 9.6$ Hz, 1H), 3.84 (dd, $J=15.2, 6.8$ Hz, 1H), 3.79 (s, 3H), 3.20 (dd, $J=13.7, 5.5$ Hz, 1H), 2.96 (dd, $J=13.7, 7.2$ Hz, 1H), 2.38 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 163.8, 159.4, 141.7, 134.1, 129.0, 128.1, 125.0, 124.8, 114.7, 78.3, 59.5, 55.3, 40.3, 21.5. HRMS-ESI calcd for $C_{18}H_{19}NO_2S$ $[M+H]^+$ 314.1215, found 314.1207.

5-(Bromomethyl)-2-(*p*-tolyl)-4,5-dihydrooxazole (**4a**): Obtained as a white solid in 73.1% yield (103.3 mg) after flash chromatography (Silica gel, petroleum ether/ethyl

acetate, $V:V=4:1$). m.p. 86~88 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.82 (d, $J=8.0$ Hz, 2H), 7.22 (d, $J=8.0$ Hz, 2H), 4.97~4.86 (m, 1H), 4.18 (dd, $J=15.2$, 9.6 Hz, 1H), 3.90 (dd, $J=15.2$, 6.8 Hz, 1H), 3.55 (dd, $J=10.8$, 5.2 Hz, 1H), 3.49 (dd, $J=10.8$, 6.4 Hz, 1H), 2.39 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.7, 141.9, 129.1, 128.1, 124.5, 77.8, 59.3, 33.7, 21.6. Spectral data are in good agreement with literature values.^[24-45]

5-(Chloromethyl)-2-(*p*-tolyl)-4,5-dihydrooxazole (**4b**): Obtained as a white solid in 58.1% yield (117.0 mg) after flash chromatography (Silica gel, petroleum ether/ethyl acetate, $V:V=3:1$). m.p. 65~67 °C; ^1H NMR (400 MHz, Chloroform-*d*) δ : 7.95 (d, $J=8.2$ Hz, 2H), 7.26 (d, $J=8.2$ Hz, 2H), 5.64~5.57 (m, 1H), 4.13~4.04 (m, 2H), 3.87 (dd, $J=12.0$, 4.4 Hz, 1H), 3.80 (dd, $J=12.0$, 4.0 Hz, 1H), 2.42 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 165.3, 144.4, 129.9, 129.2, 126.3, 74.6, 70.0, 43.5, 21.7. Spectral data are similar with literature values.^[24-45]

Supporting Information NMR and HRMS spectra of new compounds **1e**, **2e**, **1j**, **2j**, **1k**, **2k**, **1p**, **2p**, **1s**, **2s**, **1t**, **2t**, **1w**, **2w**, **3a** and **3b**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] Saravanan, P.; Corey, E. J. *J. Org. Chem.* **2003**, *68*, 2760.
- [2] Harper, K. C.; Sigman, M. S. *Science* **2011**, *333*, 1875.
- [3] Yeston, J. *Science* **2008**, *321*, 17a.
- [4] Nganga, J. K.; Samanamu, C. R.; Tanski, J. M.; Pacheco, C.; Saucedo, C.; Batista, V. S.; Grice, K. A.; Ertem, M. Z.; Angeles-Boza, A. M. *Inorg. Chem.* **2017**, *56*, 3214.
- [5] Jia, W. G.; Zhang, T.; Xie, D.; Xu, Q. T.; Ling, S.; Zhang, Q. *Dalton Trans.* **2016**, *45*, 14230.
- [6] Deng, H. P.; Wei, Y.; Shi, M. *Adv. Synth. Catal.* **2009**, *351*, 2897.
- [7] Yeston, J. S. *Science* **2005**, *310*, 1249a.
- [8] Sodeoka, M. *Science* **2011**, *334*, 1651.
- [9] Yao, W.; Duan, Z. C.; Zhang, Y.; Sang, X.; Xia, X. F.; Wang, D. *Adv. Synth. Catal.* **2019**, *361*, 5695.
- [10] Sang, X.; Hu, X.; Tao, R.; Zhang, Y.; Zhu, H.; Wang, D. *ChemPlusChem* **2020**, *85*, 123.
- [11] Yao, W.; Zhang, Y.; Zhu, H.; Ge, C.; Wang, D. *Chin. Chem. Lett.* **2020**, *31*, 701.
- [12] Hu, X.; Yang, B.; Yao, W.; Wang, D. *Chin. J. Org. Chem.* **2018**, *38*, 3296 (in Chinese). (胡昕宇, 杨伯斌, 姚玮, 王大伟, 有机化学, **2018**, *38*, 3296.)
- [13] Wang, D.; Yu, X.; Ge, B.; Miao, H.; Ding, Y. *Chin. J. Org. Chem.* **2015**, *35*, 676 (in Chinese). (王大伟, 余晓丽, 葛冰洋, 苗红艳, 丁玉强, 有机化学, **2015**, *35*, 676.)
- [14] Hu, W.; Zhang, Y.; Zhu, H.; Ye, D.; Wang, D. *Green Chem.* **2019**, *21*, 5345.
- [15] Yang, Q.; Zhang, Y.; Zeng, W.; Duan, Z.-C.; Sang, X.; Wang, D. *Green Chem.* **2019**, *21*, 5683.
- [16] Qiu, Y.; Zhang, Y.; Jin, L.; Pan, L.; Du, G.; Ye, D.; Wang, D. *Org. Chem. Front.* **2019**, *6*, 3420.
- [17] Ye, D.; Huang, R.; Zhu, H.; Zou, L.-H.; Wang, D. *Org. Chem. Front.* **2019**, *6*, 62.
- [18] Bellotti, P.; Brocus, J.; El Orfi, F.; Selkti, M.; Koenig, B.; Belmont, P.; Brachet, E. *J. Org. Chem.* **2019**, *84*, 6278.
- [19] Lamberth, C. *J. Heterocycl. Chem.* **2018**, *55*, 2035.
- [20] Seebach, D. *Helv. Chim. Acta* **2019**, *102*, e1900072.
- [21] Rajesh, N.; Manisha, B.; Ranjith, J.; Krishna, P. R. *RSC Adv.* **2016**, *6*, 6058.
- [22] Kawamura, S.; Sekine, D.; Sodeoka, M. *J. Fluorine Chem.* **2017**, *203*, 115.
- [23] Minakata, S.; Morino, Y.; Oderaotoshi, Y.; Komatsu, M. *Org. Lett.* **2006**, *8*, 3335.
- [24] Liu, G. Q.; Yang, C. H.; Li, Y. M. *J. Org. Chem.* **2015**, *80*, 11339.
- [25] Zhou, W.; Xie, C.; Han, J.; Pan, Y. *Org. Lett.* **2012**, *14*, 4766.
- [26] Senadi, G. C.; Guo, B. C.; Hu, W. P.; Wang, J. *J. Chem. Commun. (Camb.)* **2016**, *52*, 11410.
- [27] Yang, C.-H.; Xu, Z.-Q.; Duan, L.; Li, Y.-M. *Tetrahedron* **2017**, *73*, 6747.
- [28] Morino, Y.; Hidaka, I.; Oderaotoshi, Y.; Komatsu, M.; Minakata, S. *Tetrahedron* **2006**, *62*, 12247.
- [29] Kawato, Y.; Kubota, A.; Ono, H.; Egami, H.; Hamashima, Y. *Org. Lett.* **2015**, *17*, 1244.
- [30] Jagannathan, A.; Garzan, A.; Whitehead, D. C.; Staples, R. J.; Borhan, B. *Angew. Chem. Int. Ed.* **2011**, *50*, 2593.
- [31] Egami, H.; Ide, T.; Kawato, Y.; Hamashima, Y. *Chem. Commun.* **2015**, *51*, 16675.
- [32] Egami, H.; Yoneda, T.; Uku, M.; Ide, T.; Kawato, Y.; Hamashima, Y. *J. Org. Chem.* **2016**, *81*, 4020.
- [33] Nagao, Y.; Hisanaga, T.; Egami, H.; Kawato, Y.; Hamashima, Y. *Chem.-Eur. J.* **2017**, *23*, 16758.
- [34] Scheidt, F.; Thiehoff, C.; Yilmaz, G.; Meyer, S.; Daniliuc, C. G.; Kehr, G.; Gilmour, R. *Beilstein J. Org. Chem.* **2018**, *14*, 1021.
- [35] Haupt, J. D.; Berger, M.; Waldvogel, S. R. *Org. Lett.* **2019**, *21*, 242.
- [36] Hussain, H.; Green, I. R.; Ahmed, I. *Chem. Rev.* **2013**, *113*, 3329.
- [37] Xu, J.; Tong, R. *Green Chem.* **2017**, *19*, 2952.
- [38] Moriyama, K.; Sugiue, T.; Nishinohara, C.; Togo, H. *J. Org. Chem.* **2015**, *80*, 9132.
- [39] Li, Y.; Ma, L. F.; Wang, X. J.; Lei, B. W.; Zhao, Y.; Yang, J. Y.; Li, Z. Y. *Chin. J. Org. Chem.* **2017**, *37*, 1213 (in Chinese). (李瑶, 马丽芳, 王晓姣, 雷博文, 赵怡, 杨嘉宇, 李子元, 有机化学, **2017**, *37*, 1213.)
- [40] Tao, X. C.; Cao, X. J.; Yu, W.; Zhang, J. T. *Chin. J. Org. Chem.* **2010**, *30*, 250 (in Chinese). (陶晓春, 曹雄杰, 余伟, 张钧陶, 有机化学, **2010**, *30*, 250.)
- [41] Xie, L.-Y.; Peng, S.; Liu, F.; Liu, Y.-F.; Sun, M.; Tang, Z.-L.; Jiang, S.; Cao, Z.; He, W.-M. *ACS Sustainable Chem. Eng.* **2019**, *7*, 7193.
- [42] Park, S.; Lee, H.; Lee, Y. *Adv. Synth. Catal.* **2019**, *362*, 572.
- [43] Peng, S.; Song, Y.-X.; He, J.-Y.; Tang, S.-S.; Tan, J.-X.; Cao, Z.; Lin, Y.-W.; He, W.-M. *Chin. Chem. Lett.* **2019**, *30*, 2287.
- [44] Yue, H.; Bao, P.; Wang, L.; Lü, X.; Yang, D.; Wang, H.; Wei, W. *Chin. J. Org. Chem.* **2019**, *39*, 463 (in Chinese). (岳会兰, 鲍鹏丽, 王雷雷, 吕晓霞, 杨道山, 王桦, 魏伟, 有机化学, **2019**, *39*, 463.)
- [45] Nagaraju, K.; Rajesh, N.; Krishna, P. R. *Synth. Commun.* **2018**, *48*, 1001.

(Li, L.; Fan, Y.)