铜(I)促进的 3-重氮吲哚-2-亚胺与 AgSCF_3 的三氟甲硫基化反应: 3-三氟甲硫基-2-氨基吲哚的合成

郎 勃 Muhammad Suleman 吕 萍* 王彦广*
(浙江大学化学系 杭州 310027)

摘要 报道了一种通过铜(I)促进的 3-重氮吲哚-2-亚胺与 AgSCF_3 的三氟甲硫基化反应, 合成 3-三氟甲硫基-2-氨基吲哚类化合物的简洁、高效新方法, 其主要特征包括底物范围广, 铜(I)催化剂廉价, 重氮原料和三氟甲硫基化试剂容易获得, 反应条件温和且时间短; 此外, 适量的水能够促进这一反应。

关键词 重氮化合物; 金属卡宾; 三氟甲硫基化; 铜催化; 吲哚; 杂环

Copper(I)-Promoted Trifluoromethylthiolation of 3-Diazoindolin-2-imines with AgSCF_3 : Synthesis of 3-((Trifluoromethyl)thio)-2-aminoindoles

Lang, Bo Suleman, Muhammad Lu, Ping* Wang, Yanguang*
(Department of Chemistry, Zhejiang University, Hangzhou 310027)

Abstract A facile and efficient method for the synthesis of novel 3-((trifluoromethyl)thio)-2-aminoindoles through copper (I)-promoted trifluoromethylthiolation of 3-diazoindolin-2-imines with AgSCF_3 is reported. This methodology features a broad substrate scope of diazo compounds, economical copper(I) catalyst, readily available and stable diazo materials and trifluoromethylthiolating reagent, mild reaction conditions, short reaction time and water assisted higher yields.

Keywords diazo compounds; metal carbenes; trifluoromethylthiolation; copper catalysis; indoles; heterocycles

1 Introduction

Recent decades have witnessed great industrial interests in organic compounds having SCF_3 group incorporated in various aromatics. Pharmaceutical and agrochemical industries are particularly invested in these structural motifs due to enhanced bioavailability, transmembrane permeation and metabolic stability of host molecules by strong electron-withdrawing effect and lipophilicity (Hansch π value = 1.44) of SCF_3 group.^[1] Toltrazuril, Losartan, Vaniliprole and Triflorlex are a few prominent examples of biologically active drugs containing SCF_3 group (Figure 1).^[2] Installation of SCF_3 group into smaller molecules for design and discovery of novel bioactive compounds has always been a hot topic of research.^[3] Numerous methodologies have been developed for the introduction of SCF_3 group in different organic compounds over the course of

several decades.^[4] Some of the most commonly used methods includes nucleophilic,^[2a,5] electrophilic,^[4d,6] and radical trifluoromethylthiolation,^[7] trifluoromethylation of sulfur compounds,^[8] fluorine-halogen exchange reactions.^[9] Rapid growth of transition metal catalyzed protocols has led to the development of more efficient synthetic methods for the direct trifluoromethylthiolation of organic compounds.^[10] Elegant examples include transition metal catalyzed coupling reactions of various SCF_3 sources with different aryl halides,^[11] diazonium salts,^[12] arenes,^[13] and aliphatics.^[14]

Transition metal catalyzed reactions through a metal carbene intermediate has emerged as a powerful tool in organic synthesis.^[15] Among these, the use of diazo compounds as metal carbene precursors for the introduction of SCF_3 group has attracted much attention.^[16-19] Elegant examples include the copper catalyzed coupling reactions of

* Corresponding authors. E-mail: pinglu@zju.edu.cn; orgwyg@zju.edu.cn

Received May 5, 2020; revised June 16, 2020; published online July 30, 2020.

Dedicated to Professor Henry N. C. Wong on the occasion of his 70th birthday.

Project supported by the National Natural Science Foundation of China (Nos. 21772169, 21632003).

国家自然科学基金(Nos. 21772169, 21632003)资助项目。

Figure 1 Examples of SCF₃ containing biologically active compounds

diazo compounds with various nucleophilic SCF₃ sources^[17] and the rhodium^[18] or zinc^[19] catalyzed multicomponent reactions of diazo compounds with SCF₃ sources. Notably, only acyclic diazo compounds like α -diazo esters have been employed in trifluoromethylthiolations giving mostly aliphatic SCF₃ derivatives (Scheme 1a),^[17] while the use of cyclic diazo compounds for these transformations remains unexplored.

Scheme 1 Trifluoromethylthiolation of acyclic diazo compounds and our work

In 2014, we developed practical methods for the preparation of 3-diazoindolin-2-imines, a new class of cyclic α -diazo imidamides,^[20] and used them as metal carbene precursors for the syntheses of structurally divers indole and indoline derivatives.^[20-21] It was envisioned that 3-diazoindolin-2-imines could be used for copper-catalyzed trifluoromethylthiolation reactions and would result in important indole embedded aromatic SCF₃ derivatives (Scheme 1b). The results of our effort are described below.

2 Results and discussion

We began our trial by treating 3-diazoindolin-2-imine (**1p**) with silver trifluoromethylthiolate (AgSCF₃, **2**) in the presence of different copper salts (Table 1). In the first attempt, **1p** was treated with 1 equiv. AgSCF₃ in the presence of CuI catalyst in 1,2-dichloroethane (DCE) solvent at room temperature and in ambient air for 1 h as a model

reaction. Fortunately, trifluoromethylthiolated product **3p** was obtained in 62% yield (Table 1, Entry 1).

Table 1 Optimization of reaction conditions^a

Entry	[Cu]/equiv.	1p /equiv.	Additive (equiv.)	Solvent	Time/h	Yield ^b /%
1	CuI (1.0)	1.0	—	DCE	1	62
2	CuBr (1.0)	1.0	—	DCE	1	Trace
3	CuCl (1.0)	1.0	—	DCE	1	Trace
4 ^c	CuI (1.0)	1.0	—	DCE	1	53
5	CuI (1.0)	1.2	—	DCE	1	77
6	CuI (1.0)	1.4	—	DCE	1	77
7	CuI (1.0)	1.2	H ₂ O (10)	DCE	1	87
8	CuI (1.0)	1.2	H ₂ O (20)	DCE	1	67
9	CuI (1.0)	1.2	H ₂ O (30)	DCE	1	64
10	CuI (1.0)	1.2	H ₂ O (40)	DCE	1	55
11	CuI (1.25)	1.2	H ₂ O (10)	DCE	1	66
12	CuI (0.75)	1.2	H ₂ O (10)	DCE	1	74
13	CuI (0.50)	1.2	H ₂ O (10)	DCE	1	71
14	CuI (0.25)	1.2	H ₂ O (10)	DCE	1	30
15	CuI (1.0)	1.2	H ₂ O (10)	DCM	1	84
16	CuI (1.0)	1.2	H ₂ O (10)	PhMe	1	87
17	CuI (1.0)	1.2	H ₂ O (10)	CH ₃ CN	1	trace
18	CuI (1.0)	1.2	H ₂ O (10)	NMP	1	trace
19	CuI (1.0)	1.2	H ₂ O (10)	DCE	0.5	85
20	CuI (1.0)	1.2	H ₂ O (10)	DCE	2	87

^a Reaction conditions: **1a** (0.12 mmol), **2** (0.1 mmol), CuI (0.1 mmol), solvent (2 mL), room temperature; ^b isolated yield; ^c N₂ atmosphere.

The product **3p** was characterized by HRMS, ¹H NMR and ¹³C NMR analysis, and its structure was confirmed by X-ray single crystal analysis.^[22] These results prompted us to further screen the reaction conditions. When CuBr and CuCl were used as catalysts, only a trace amount of **3p** was detected on thin-layer chromatography (TLC) (Table 1, Entries 2 and 3). When the reaction was performed under nitrogen atmosphere, **3p** was obtained in lower yield (53%) (Table 1, Entry 4). The yield could be increased to 77% when the amount of diazo compound was risen to 1.2 and 1.4 equiv. (Table 1, Entries 5 and 6). When 10 equiv. of water was added to the reaction as additive, the yield of **3p** was increased to 87% (Table 1, Entry 7). A gradual decrease in the yield was observed when the amount of water was increased (Table 1, Entries 8~10). Both increase and decrease in the amount of CuI led to a decrease in the yield of **3p** (Table 1, Entries 11~14). The solvents were also screened. Dichloromethane (DCM) and toluene gave 84% and 87% yields, respectively (Table 1, Entries 15 and 16), while only trace amount of **3p** was observed on TLC when acetonitrile and *N*-methylpyrrolidone (NMP) were used as

solvents (Table 1, Entries 17 and 18). Finally, the reaction time was optimized. Decreasing the reaction time led to a slightly reduction in the yield (85%) (Table 1, Entry 19), while increasing the reaction time did not improve the reaction (Table 1, Entry 20). Thus, the best reaction conditions were determined (Table 1, Entry 7).

After determining the optimal reaction conditions, the substrate scope of 3-diazoindolin-2-imines **1** was investigated (Table 2). Substituent R^1 at 1-position of diazo component **1** could be varied from methyl to ethyl, isopropyl and benzyl, giving the corresponding products **3a**~**3d** in 78%~86% yields. However, the diazo compound **1e** without substituent at 1-position of indole ring ($R^1=H$) did not undergo this reaction to give the desired product **3e**. Substituent at 5-position of 3-diazoindolin-2-imines **1** could be varied from electron-withdrawing groups (F, Cl, Br) to electron-donating groups (Me, OMe). Thus, the corresponding products **3f** (89%), **3g** (81%), **3h** (80%) and **3i** (83%) and **3j** (93%) were obtained in excellent yields, indicating that the presence of electron-donating groups at phenyl ring of diazo component slightly favors the reaction (Scheme 2). The presence of substituents at 4-, 6- or 7-positions of diazo compounds **1** also tolerated the reaction conditions and the corresponding products **3k** (54%), **3l** (80%) and **3m** (82%) were isolated in moderate to excellent yields. 3-diazoindolin-2-imines **1** with different sulfonyl groups attached (e.g. methylsulfonyl, phenylsulfonyl, (4-chlorophenyl)sulfonyl and (4-fluorophenyl)sulfonyl) were also favorable for these transformations. The corresponding products **3n**, **3o**, **3p** and **3q** were isolated in 74%, 75%, 87% and 82% yields, respectively. When the sulfonyl group of diazo compound was changed to 4-nitrophenyl-sulfonyl, the desired product **3r** was not formed. On the other hand, diazo compounds **1** bearing 4-methoxyphenyl sulfonyl or naphthalen-2-yl sulfonyl group generated the corresponding products **3s** and **3t** in 86% and 83% yields, respectively.

A gram scale reaction of **3p** was performed to evaluate the synthetic potential of this approach. Diazo compound **1p** (3.36 mmol) smoothly reacted with **2** (2.8 mmol) under the standard conditions and the corresponding product **3p** was isolated in 85% yield (1.008 g).

Based on our results and previous report,^[17] a plausible mechanism is proposed for the formation of **3** (Scheme 2). First, $AgSCF_3$ interacts with CuI and a more reactive specie $CuSCF_3$ is formed along with the removal of AgI precipitates. This nucleophilic specie then interacts with diazo compound **1** and SCF_3 containing copper carbene intermediate **A** is formed with the release of nitrogen gas. Subsequently, the Cu carbene **A** undergoes migratory insertion and the copper complex **B** is formed. Finally, water assisted protonation results in the formation of SCF_3 containing intermediate **C**, which undergoes tautomerization to give the final product **3** (path a). Aromatization is the driving force of this step. Alternatively, **B** tautomerizes to complex **D**, which gives **3** after hydrolysis.

Table 2 Substrate scope of 3-diazoindolin-2-imines^a

3a , $R^1 = Me$, 78%	3f , $R^2 = F$, 89%	3n , $R^3 = Me$, 74%
3b , $R^1 = Et$, 85%	3g , $R^2 = Cl$, 81%	3o , $R^3 = Ph$, 75%
3c , $R^1 = i-Pr$, 82%	3h , $R^2 = Br$, 80%	3p , $R^3 = p-ClC_6H_4$, 87%
3d , $R^1 = Bn$, 86%	3i , $R^2 = Me$, 83%	3q , $R^3 = p-FC_6H_4$, 82%
3e , $R^1 = H$, 0	3j , $R^2 = OMe$, 93%	3r , $R^3 = p-NO_2C_6H_4$, 0
		3s , $R^3 = p-MeOC_6H_4$, 86%
		3t , $R^3 = 2-Naphthalenyl$, 83%

3k, 56% **3l**, 80% **3m**, 82%

X-ray crystal structure of **3p**

^a Reaction conditions: **1** (0.12 mmol), **2** (0.1 mmol), CuI (0.1 mmol), H_2O (1 mmol), DCE (2 mL), room temperature, 1 h; Isolated yield.

Scheme 2 Mechanism for the formation of 3-((trifluoromethyl)thio)-2-aminoindoles (**3**)

3 Conclusions

We have developed an efficient cross-coupling reaction between 3-diazoindolin-2-imines **1** and nucleophilic trifluoromethylthiolating reagent AgSCF₃ for the synthesis of novel indole embedded SCF₃ derivatives. It is the first example of trifluoromethylthiolation of cyclic diazo compounds. Readily available starting materials, cheap catalysts, water assisted higher yields and mild reaction conditions are the merits of this reaction. As indole ring system is part of numerous natural products and drugs, these indole embedded SCF₃ derivatives may find applications in medicinal chemistry and be valuable in future.

4 Experimental section

4.1 Instruments and reagents

¹H NMR spectra were obtained on 400 MHz in CDCl₃. The chemical shifts were reported referenced to tetramethylsilane (TMS) as an internal standard. ¹³C NMR spectra were recorded on 100 MHz in CDCl₃. The chemical shifts were reported referenced to the internal solvent signals (δ 77.0 for CDCl₃). ¹⁹F NMR chemical shifts were determined relative to CFCl₃ at δ 0.0. Infrared spectra were obtained on an FTIR spectrometer. High-resolution mass spectra (HRMS) data were obtained by using EI ionization on time-of-flight (TOF) mass spectrometer. Melting points were measured with a SGW X-4 micro melting point apparatus. Flash column chromatography was performed employing 300~400 mesh silica gel. Thin layer chromatography (TLC) was performed on silica gel HSGF254. Unless otherwise mentioned, solvents and reagents were purchased from commercial sources and used as received. 3-Diazoindolin-2-imines **1** were prepared according to our published methods.^[20]

4.2 General procedure for the synthesis of compounds 3

To an over-dried reaction tube with a magnetic stirring bar were added sequentially **1** (0.12 mmol), **2** (0.1 mmol, 21 mg), CuI (0.1 mmol, 19 mg), H₂O (1 mmol, 18 mg) and dichloromethane (DCE) (2 mL). The mixture was stirred at room temperature for 1 h. After the solvent was evaporated in vacuum, the residue was purified by column chromatography on silica gel (petroleum ether/ethyl acetate, *V*: *V*=5:1) to give pure products **3**.

4.3 Characterization data for products 3

4-Methyl-*N*-(1-methyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)benzenesulfonamide (**3a**): White solid, yield 78% (31 mg). m.p. 158.2~159.3 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.62 (d, *J*=8.0 Hz, 1H), 7.49 (d, *J*=8.4 Hz, 2H), 7.42 (d, *J*=8.0 Hz, 1H), 7.36 (td, *J*=7.6 Hz, 1.2 Hz, 1H), 7.29~7.23 (m, 3H), 6.74 (s, 1H), 3.94 (s, 3H), 2.44 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 145.0, 136.7, 135.4, 134.2, 129.8, 129.2 (q, *J*=309 Hz), 128.0, 127.8, 123.8, 121.9, 119.4, 110.5, 89.3, 31.3, 21.7; ¹⁹F NMR (376 MHz, CDCl₃) δ : -44.06 (s); IR (film) ν : 3245, 2959, 1597, 1522, 1479,

1378, 1185, 1124, 849, 738, 672 cm⁻¹. HRMS (EI) calcd for C₁₇H₁₅F₃N₂O₂S₂ 400.0527; found 400.0525.

N-(1-Ethyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)-4-methylbenzenesulfonamide (**3b**): White solid, yield 85% (35 mg). m.p. 129.2~130.1 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.62 (d, *J*=8.0 Hz, 1H), 7.51 (d, *J*=8.0 Hz, 2H), 7.46 (d, *J*=8.4 Hz, 1H), 7.35 (td, *J*=7.6, 1.2 Hz, 1H), 7.27~7.24 (m, 3H), 6.71 (s, 1H), 4.52 (d, *J*=7.6 Hz, 2H), 2.44 (s, 3H), 1.49 (t, *J*=3.2 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ : 145.0, 136.0, 134.4, 134.2, 129.8, 129.2 (q, *J*=309 Hz), 128.3, 127.9, 123.8, 121.8, 119.6, 110.9, 90.0, 39.4, 21.7, 14.5; ¹⁹F NMR (376 MHz, CDCl₃) δ : -44.00 (s); IR (film) ν : 3252, 2979, 2929, 1598, 1525, 1383, 1343, 1230, 1186, 1105, 1018, 813, 749, 672 cm⁻¹. HRMS (EI) calcd for C₁₈H₁₇F₃N₂O₂S₂ 414.0684; found 414.0684.

N-(1-Isopropyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)-4-methylbenzenesulfonamide (**3c**): White solid, yield 82% (35 mg). m.p. 158.2~159.1 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.68 (d, *J*=8.4 Hz, 1H), 7.63 (d, *J*=8.0 Hz, 1H), 7.51 (d, *J*=8.4 Hz, 2H), 7.31 (td, *J*=8.0 Hz, 0.8 Hz, 1H), 7.26~7.21 (m, 3H), 6.68 (s, 1H), 5.39~5.32 (m, 1H), 2.44 (s, 3H), 1.72 (s, 3H), 1.70 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 145.0, 135.8, 134.1, 133.1, 129.8, 129.2 (q, *J*=309 Hz), 129.0, 127.9, 123.2, 121.3, 119.8, 113.2, 89.8, 48.8, 21.7, 21.1; ¹⁹F NMR (376 MHz, CDCl₃) δ : -43.99 (s); IR (film) ν : 3253, 2977, 2940, 1598, 1527, 1456, 1377, 1168, 1102, 1019, 909, 813, 672 cm⁻¹. HRMS (EI) calcd for C₁₉H₁₉F₃N₂O₂S₂ 428.0840; found 428.0839.

N-(1-Benzyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)-4-methylbenzenesulfonamide (**3d**): White solid, yield 86% (41 mg). m.p. 134.0~135.0 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.66~7.64 (m, 1H), 7.53 (d, *J*=8.4 Hz, 2H), 7.27~7.25 (m, 8H), 7.00 (d, *J*=6.4 Hz, 2H), 6.72 (s, 1H), 5.69 (s, 2H), 2.45 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 145.0, 136.6, 136.5, 135.0, 134.3, 129.8, 129.2 (q, *J*=309 Hz), 128.7, 128.2, 127.9, 127.5, 126.3, 124.1, 122.0, 119.6, 115.5, 91.2, 47.9, 21.7; ¹⁹F NMR (376 MHz, CDCl₃) δ : -43.92 (s); IR (film) ν : 3255, 3032, 2929, 1597, 1528, 1384, 1167, 1105, 971, 909, 813, 735 cm⁻¹. HRMS (EI) calcd for C₂₃H₁₉F₃N₂O₂S₂ 476.0840; found 476.0841.

N-(5-Fluoro-1-methyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)-4-methylbenzenesulfonamide (**3f**): White solid, yield 89% (37 mg). m.p. 213.0~214.0 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.58 (d, *J*=8.4 Hz, 2H), 7.35 (dd, *J*=8.8 Hz, 4.0 Hz, 2H), 7.28~7.25 (m, 3H), 7.10 (td, *J*=9.2 Hz, 2.4 Hz, 2H), 6.78 (s, 1H), 3.93 (s, 3H), 2.45 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 159.0 (d, *J*=237.2 Hz), 145.2, 137.9, 134.1, 131.9, 129.9, 129.2 (q, *J*=309 Hz), 128.7 (d, *J*=10.4 Hz), 127.7, 112.5 (d, *J*=22.1 Hz), 111.7 (d, *J*=9.5 Hz), 104.6 (d, *J*=10.2 Hz), 89.4, 31.5, 21.7; ¹⁹F NMR (376 MHz, CDCl₃) δ : -44.06 (s, 3F), -120.62~-120.67 (m, 1F); IR (film) ν : 3232, 2959, 2916, 1383, 1489, 1333, 1169, 1104, 986, 863, 674 cm⁻¹. HRMS (EI) calcd for C₁₇H₁₄F₄N₂O₂S₂ 418.0433; found 418.0433.

N-(5-Chloro-1-methyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)-4-methylbenzenesulfonamide (**3g**): White solid, yield 81% (35 mg). m.p. 181.2~182.3 °C; ¹H NMR (400

MHz, CDCl₃) δ : 7.58 (s, 1H), 7.47 (d, J =8.0 Hz, 2H), 7.35~7.29 (m, 2H), 7.27~7.25 (m, 2H), 6.76 (s, 1H), 3.93 (s, 3H), 2.45 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 145.2, 137.8, 134.0, 133.8, 129.9, 129.02, 129.01 (q, J =309 Hz), 128.0, 127.7, 124.3, 118.9, 111.8, 89.1, 31.5, 21.7; ¹⁹F NMR (376 MHz, CDCl₃) δ : -44.09 (s); IR (film) ν : 3224, 3054, 2954, 1598, 1525, 1373, 1332, 1142, 1116, 907, 713 cm⁻¹. HRMS (EI) calcd for C₁₇H₁₄ClF₃N₂O₂S₂ 434.0137; found 434.0138.

N-(5-Bromo-1-methyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)-4-methylbenzenesulfonamide (**3h**): White solid, yield 80% (38 mg). m.p. 234.6~235.6 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.73 (d, J =1.2 Hz, 1H), 7.48~7.43 (m, 3H), 7.29 (d, J =8.8 Hz, 1H), 7.27~7.25 (m, 2H), 6.76 (s, 1H), 3.93 (s, 3H), 2.45 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 145.3, 137.7, 134.1, 134.0, 129.9, 129.6, 129.0 (q, J =309 Hz), 127.7, 126.9, 122.0, 115.6, 112.1, 89.0, 31.5, 21.7; ¹⁹F NMR (376 MHz, CDCl₃) δ : -43.95 (s); IR (film) ν : 3229, 2959, 2912, 1598, 1472, 1372, 1188, 1117, 1052, 809, 675 cm⁻¹; HRMS (ESI) calcd for C₁₇H₁₄BrF₃N₂O₂S₂Na⁺ [M+Na]⁺: 500.9524; found 500.9537.

N-(1,5-Dimethyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)-4-methylbenzenesulfonamide (**3i**): White solid, yield 83% (34 mg). m.p. 165.2~166.3 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.48 (d, J =8.0 Hz, 2H), 7.39 (s, 1H), 7.30 (d, J =8.4 Hz, 1H), 7.25~7.22 (m, 2H), 7.18 (dd, J =8.4 Hz, 1.2 Hz, 1H), 6.74 (s, 1H), 3.91 (s, 3H), 2.47 (s, 3H), 2.44 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 144.9, 136.5, 134.2, 133.8, 131.5, 129.8, 129.2 (q, J =309 Hz), 128.2, 127.8, 125.5, 118.9, 110.3, 88.6, 31.2, 21.7, 21.5; ¹⁹F NMR (376 MHz, CDCl₃) δ : -44.13 (s); IR (film) ν : 3253, 3028, 2923, 1598, 1491, 1343, 1306, 1186, 1122, 909, 795, 674 cm⁻¹. HRMS (EI) calcd for C₁₈H₁₇F₃N₂O₂S₂ 414.0684; found 414.0686.

N-(5-Methoxy-1-methyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)-4-methylbenzenesulfonamide (**3j**): White solid, yield 93% (40 mg). m.p. 190.1~191.2 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.49 (d, J =8.4 Hz, 2H), 7.30 (d, J =8.8 Hz, 1H), 7.25 (d, J =7.2 Hz, 2H), 7.03 (d, J =2.0 Hz, 1H), 7.00 (dd, J =8.8 Hz, 2.4 Hz, 1H), 6.77 (s, 1H), 3.90 (s, 3H), 3.86 (s, 3H), 2.44 (s, 3H), 2.44 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 155.8, 144.9, 136.6, 134.3, 130.3, 129.8, 129.3 (q, J =309 Hz), 128.8, 114.3, 111.5, 100.6, 88.8, 55.7, 31.3, 21.7; ¹⁹F NMR (376 MHz, CDCl₃) δ : -44.16 (s); IR (film) ν : 3218, 3004, 2965, 1623, 1598, 1378, 1291, 1221, 1120, 1029, 952, 794, 697 cm⁻¹. HRMS (EI) calcd for C₁₈H₁₇F₃N₂O₃S₂ 430.0633; found 430.0635.

N-(4-Bromo-1-methyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)-4-methylbenzenesulfonamide **3k**: White solid, yield 56% (27 mg). m.p. 157~159 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.51 (d, J =8.4 Hz, 2H), 7.44~7.38 (m, 2H), 7.27 (d, J =8.0 Hz, 2H), 7.18 (t, J =7.6 Hz, 1H), 6.87 (s, 1H), 3.95 (s, 3H), 2.46 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 145.3, 138.9, 136.6, 134.2, 129.9, 128.9 (q, J =309 Hz), 127.8, 127.1, 124.9, 124.2, 114.2, 110.1, 89.7, 31.8, 21.7; ¹⁹F NMR (376 MHz, CDCl₃) δ : -44.74 (s); IR (film) ν : 3254, 3051, 2948, 1624, 1514, 1343, 1285, 1186,

1122, 909, 795, 674 cm⁻¹. HRMS (EI) calcd for C₁₇H₁₄BrF₃N₂O₂S₂ 477.9632; found 477.9635.

N-(6-Chloro-1-methyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)-4-methylbenzenesulfonamide (**3l**): White solid, yield 80% (35 mg). m.p. 159~161 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.53~7.47 (m, 3H), 7.42 (d, J =1.6 Hz, 1H), 7.26~7.21 (m, 3H), 6.80 (s, 1H), 3.90 (s, 3H), 2.45 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 145.1, 137.3, 135.8, 134.1, 129.89, 129.86, 129.0 (q, J =309 Hz), 127.8, 126.4, 122.8, 120.5, 110.6, 89.9, 31.4, 21.7; ¹⁹F NMR (376 MHz, CDCl₃) δ : -43.92 (s); IR (film) ν : 3236, 3032, 2956, 1598, 1491, 1343, 1306, 1186, 1122, 909, 795, 674 cm⁻¹. HRMS (EI) calcd for C₁₇H₁₄ClF₃N₂O₂S₂ 434.0137; found 434.0138.

N-(1,7-Dimethyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)-4-methylbenzenesulfonamide (**3m**): White solid, yield 82% (34 mg). m.p. 170.1~171.2 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.41 (d, J =8.0 Hz, 2H), 7.37 (d, J =7.6 Hz, 1H), 7.16 (d, J =8.4 Hz, 2H), 7.03 (t, J =3.6 Hz, 1H), 6.97 (d, J =7.2 Hz, 1H), 6.69 (s, 1H), 4.09 (s, 3H), 2.74 (s, 3H), 2.36 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 145.0, 137.1, 134.6, 134.3, 129.8, 129.2 (q, J =309 Hz), 128.9, 127.8, 126.6, 122.5, 121.8, 117.4, 89.5, 34.6, 21.7, 20.0; ¹⁹F NMR (376 MHz, CDCl₃) δ : -44.02 (s); IR (film) ν : 3253, 3028, 2923, 1598, 1491, 1343, 1306, 1186, 1122, 909, 795, 674 cm⁻¹. HRMS (EI) calcd for C₁₈H₁₇F₃N₂O₂S₂ 414.0684; found 414.0686.

N-(1-Methyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)-methanesulfonamide (**3n**): White solid, yield 74% (24 mg). m.p. 158.2~159.3 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.77 (d, J =8.0 Hz, 1H), 7.42~7.36 (m, 2H), 7.34~7.30 (m, 1H), 6.84 (s, 1H), 3.87 (s, 3H), 3.14 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 136.4, 135.4, 129.2 (q, J =309 Hz), 128.1, 124.2, 122.3, 119.4, 89.0, 40.4, 31.2; ¹⁹F NMR (376 MHz, CDCl₃) δ : -43.64 (s); IR (film) ν : 3256, 2075, 2936, 1528, 1477, 1382, 1162, 973, 746 cm⁻¹. HRMS (EI) calcd for C₁₁H₁₁F₃N₂O₂S₂ 324.0214; found 324.0215.

N-(1-Methyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)-benzenesulfonamide (**3o**): White solid, yield 75% (29 mg). m.p. 200.1~201.1 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.67~7.61 (m, 4H), 7.49~7.44 (m, 3H), 7.37 (td, J =8.4, 1.2 Hz, 1H), 7.29~7.26 (m, 1H), 6.78 (s, 1H), 3.95 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 137.1, 136.5, 135.4, 133.9, 129.3, 129.1 (q, J =309 Hz), 128.0, 127.8, 123.9, 122.0, 119.4, 89.5, 31.3; ¹⁹F NMR (376 MHz, CDCl₃) δ : -44.08 (s); IR (film) ν : 3232, 3062, 2929, 1526, 1385, 1238, 1171, 1121, 1012, 847, 751, 686 cm⁻¹. HRMS (EI) calcd for C₁₆H₁₃F₃N₂O₂S₂ 386.0371; found 386.0372.

4-Chloro-*N*-(1-methyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)benzenesulfonamide (**3p**): White solid, yield 87% (37 mg). m.p. 199.8~200.8 °C; ¹H NMR (400 MHz, CDCl₃) δ : 7.64 (d, J =8.0 Hz, 1H), 7.58~7.55 (m, 2H), 7.46~7.43 (m, 3H), 7.38 (td, J =7.2, 1.2 Hz, 1H), 7.31~7.27 (m, 1H), 6.83 (s, 1H), 3.95 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ : 140.7, 136.1, 135.8, 135.5, 129.5, 129.2, 129.1 (q, J =309 Hz), 127.9, 124.1, 122.1, 119.5, 110.6, 89.7, 31.3; ¹⁹F NMR (376 MHz, CDCl₃) δ : -43.94 (s); IR (film) ν : 3250, 3062, 2959, 1574, 1475, 1379, 1172, 1092,

1016, 848, 668 cm^{-1} . HRMS (EI) calcd for $\text{C}_{16}\text{H}_{12}\text{ClF}_3\text{N}_2\text{O}_2\text{S}_2$ 419.9981; found 419.9981.

4-Fluoro-*N*-(1-methyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)benzenesulfonamide (**3q**): White solid, yield 82% (33 mg). m.p. 156.9~157.8 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.66~7.63 (m, 3H), 7.43 (d, $J=8.4$ Hz, 1H), 7.40~7.36 (m, 1H), 7.29~7.26 (m, 1H), 7.27 (t, $J=7.2$ Hz, 1H), 7.15 (t, $J=8.4$ Hz, 1H), 6.78 (s, 1H), 3.95 (s, 3H), 3.87 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 165.9 ($J=255.6$ Hz), 136.2, 135.5, 133.3, 130.6 ($J=9.6$ Hz), 129.1 (q, $J=308$ Hz), 127.9, 124.1, 122.1, 119.5, 116.5 ($J=22.6$ Hz), 110.6, 89.5, 31.3; ^{19}F NMR (376 MHz, CDCl_3) δ : -43.99 (s); IR (film) ν : 3255, 3067, 2959, 1593, 1527, 1386, 1157, 1014, 949, 838, 735 cm^{-1} . HRMS (EI) calcd for $\text{C}_{16}\text{H}_{12}\text{F}_4\text{N}_2\text{O}_3\text{S}_2$ 404.0276; found 404.0277.

4-Methoxy-*N*-(1-methyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)benzenesulfonamide (**3s**): White solid, yield 86% (36 mg). m.p. 143.0~144.1 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.62 (d, $J=8.0$ Hz, 1H), 7.55~7.52 (m, 2H), 7.42 (d, $J=8.0$ Hz, 1H), 7.39~7.37 (m, 1H), 7.29~7.26 (m, 1H), 6.90 (d, $J=8.8$ Hz, 2H), 6.71 (s, 1H), 3.95 (s, 3H), 3.87 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.9, 136.9, 135.5, 130.0, 129.2 (q, $J=309$ Hz), 128.7, 128.0, 123.8, 121.9, 119.4, 114.4, 110.5, 89.2, 55.7, 31.3; ^{19}F NMR (376 MHz, CDCl_3) δ : -43.99 (s, 3F), -102.60~-102.66 (m, 1F); IR (film) ν : 3253, 2946, 1596, 1527, 1386, 1264, 1162, 1121, 1026, 943, 833, 675 cm^{-1} . HRMS (EI) calcd for $\text{C}_{17}\text{H}_{15}\text{F}_3\text{N}_2\text{O}_3\text{S}_2$ 416.0476; found 416.0478.

N-(1-Methyl-3-((trifluoromethyl)thio)-1*H*-indol-2-yl)-naphthalene-2-sulfonamide (**3t**): White solid, yield 83% (36 mg). m.p. 205.8~206.6 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.93~7.89 (m, 2H), 7.83 (d, $J=8.0$ Hz, 1H), 7.67~7.62 (m, 1H), 7.62~7.56 (m, 3H), 7.43 (d, $J=8.4$ Hz, 1H), 7.39~7.35 (m, 1H), 7.25 (td, $J=8.0, 1.2$ Hz, 1H), 6.94 (s, 1H), 3.97 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 136.5, 135.5, 135.2, 134.4, 132.0, 129.5, 129.42, 129.35, 129.34, 129.1 (q, $J=309$ Hz), 128.0, 127.7, 123.9, 122.5, 122.0, 119.5, 89.7, 31.3; ^{19}F NMR (376 MHz, CDCl_3) δ : -44.02 (s); IR (film) ν : 3236, 3057, 2943, 1625, 1525, 1240, 1169, 1103, 1013, 939, 816, 670 cm^{-1} . HRMS (EI) calcd for $\text{C}_{20}\text{H}_{15}\text{F}_3\text{N}_2\text{O}_3\text{S}_2$ 436.0527; found 436.0526.

Supporting Information The copies of NMR of newly synthesized products **3**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Manteau, B.; Pazenok, S.; Vors, J. P.; Leroux, F. R. *J. Fluorine Chem.* **2010**, *131*, 140.
(b) Leroux, F.; Jeschke, P.; Schlosser, M. *Chem. Rev.* **2005**, *105*, 827.
(c) Hansch, C.; Leo, A.; Taft, R. W. *Chem. Rev.* **1991**, *91*, 165.
(d) Arnott, J. A.; Planey, S. L. *Expert Opin. Drug Discovery* **2012**, *7*, 863.
(e) Liu, X.; Testa, B.; Fahr, A. *Pharm. Res.* **2011**, *28*, 962.
- [2] (a) Zheng, H.; Huang, Y.; Weng, Z. *Tetrahedron Lett.* **2016**, *57*, 1397.
(b) Yagupolskii, L. M.; Maletina, I. I.; Petko, K. I.; Fedyuk, D. V.; Handrock, R.; Shavaran, S. S.; Klebanov, B. M.; Herzig, S. *J. Fluorine Chem.* **2001**, *109*, 87.
- [3] For selected reviews on trifluoromethylthiolation, see:
(a) Tlili, A.; Billard, T. *Angew. Chem. Int. Ed.* **2013**, *52*, 6818.
(b) Liu, H.; Jiang, X. *Chem. Asian. J.* **2013**, *8*, 2546.
(c) Boiko, V. N. *Beilstein J. Org. Chem.* **2010**, *6*, 880.
- [4] For selected reviews on the introduction of SCF_3 group, see:
(a) Chu, L.; Qing, F. L. *Acc. Chem. Res.* **2014**, *47*, 1513.
(b) Xu, X. H.; Matsuzaki, K.; Shibata, N. *Chem. Rev.* **2015**, *115*, 731.
(c) Chachignon, H.; Cahard, D. *Chin. J. Chem.* **2016**, *34*, 445.
(d) Shao, X.; Xu, C.; Lu, L.; Shen, Q. *Acc. Chem. Res.* **2015**, *48*, 1227.
(e) Leroux, F.; Jeschke, P.; Schlosser, M. *Chem. Rev.* **2005**, *105*, 827.
(f) Li, S. S.; Wang, J. B. *Acta Chim. Sinica* **2018**, *76*, 913 (in Chinese).
(李树森, 王剑波, 化学学报, **2018**, *76*, 913.)
- [5] (a) Man, E. H.; Coffman, D. D.; Muetterties E. L. *J. Am. Chem. Soc.* **1959**, *81*, 3575.
(b) Munavalli, S.; Wagner, G. W.; Hashemi, A. B.; Rohrbaugh, D. K.; Durst, H. D. *Synth. Commun.* **1997**, *27*, 2847.
(c) Tyrra, W.; Naumann, D.; Hoge, B.; Yagupolskii, Y. L. *J. Fluorine Chem.* **2003**, *119*, 101.
(d) Chen, H.-T.; Huang, Y.; Qing, F. L.; Xu, X. H. *Tetrahedron Lett.* **2020**, *61*, 151628.
(e) Liu, Y. L.; Qing, F. L.; Xu, X. H. *Eur. J. Org. Chem.* **2020**, 1015.
- [6] (a) Wang, Q.; Qi, Z.; Xie, F.; Li, X. *Adv. Synth. Catal.* **2015**, *357*, 355.
(b) Baert, F.; Colomb, J.; Billard, T. *Angew. Chem. Int. Ed.* **2012**, *51*, 10382.
(c) Shao, X.; Wang, X.; Yang, T.; Lu, L.; Shen, Q. *Angew. Chem. Int. Ed.* **2013**, *52*, 3457.
(d) Bootwicha, T.; Liu, X.; Pluta, R.; Atodiresei, I.; Rueping, M. *Angew. Chem. Int. Ed.* **2013**, *52*, 12856.
(e) Zhu, X. Li; Xu, J. H.; Cheng, D. J.; Zhao, L. J.; Liu, X. Y.; Tan, B. *Org. Lett.* **2014**, *16*, 2192.
(f) Franco, F.; Meninno, S.; Bengalia, M.; Lattanzi, A. *Chem. Commun.* **2020**, *56*, 3073.
- [7] For selected examples, see:
(a) Harris, J. F. *J. Org. Chem.* **1966**, *31*, 931.
(b) Zhu, M.; Li, R.; You, Q.; Fu, W.; Guo, W. *Asian J. Org. Chem.* **2019**, *8*, 2002.
(c) Hu, J. J.; Huang, Y. G.; Xu, X. H.; Qing, F. L. *Chin. J. Org. Chem.* **2019**, *39*, 177 (in Chinese).
(解晓娟, 张忠, 赵华欣, 万文, 郝健, 有机化学, **2019**, *39*, 177.)
(d) Chen, D.; Ji, M. S.; Yao, Y. M.; Zhu, C. *Acta Chim. Sinica* **2018**, *76*, 951 (in Chinese).
(陈栋, 吉梅山, 姚英明, 朱晨, 化学学报, **2018**, *76*, 951.)
(e) Xiao, Z.; Liu, Y.; Zheng, L.; Liu, C.; Guo, Y.; Chen, Q. Y. *J. Org. Chem.* **2018**, *83*, 5836.
(f) Pan, S.; Huang, Y.; Xu, X. H.; Qing, F. L. *Org. Lett.* **2017**, *19*, 4624.
(g) Xiao, Z.; Liu, Y.; Zheng, L.; Zhou, X.; Xie, Y.; Liu, C.; Guo, Y.; Chen, Q. Y. *Tetrahedron* **2018**, *74*, 6213.
(h) Wu, Hao.; Xiao, Z.; Wu, J.; Guo, Y.; Xiao, J. C.; Liu, C.; Chen, Q. Y. *Angew. Chem. Int. Ed.* **2015**, *54*, 4070.
(i) Dagousset, G.; Simon, C.; Anselmi, E.; Tuccio, B.; Billard, T.; Magnier, E. *Chem.-Eur. J.* **2017**, *23*, 4282.
(j) Pan, S.; Li, H.; Huang, Y.; Xu, X. H.; Qing, F. L. *Org. Lett.* **2017**, *19*, 3247.
(k) Li, H.; Liu, S.; Huang, Y.; Xu, X. H.; Qing, F. L. *Chem. Commun.* **2017**, *53*, 10136.
- [8] (a) Pooput, C.; Medebielle, M.; Dolbier, W. R. *Org. Lett.* **2004**, *6*, 301.
(b) Pooput, C.; Dolbier, W. R.; Médebille, M. J. *Org. Chem.* **2006**,

- 71, 3564.
(c) Kieltisch, I.; Eisenberger, P.; Togni, A. *Angew. Chem. Int. Ed.* **2007**, *46*, 754.
- [9] (a) Scherer, O. *Angew. Chem.* **1939**, *52*, 457.
(b) Saint-Jalmes, L. *J. Fluorine Chem.* **2006**, *127*, 85.
- [10] For selected reviews on direct trifluoromethylthiolations, see:
(a) Zhang, K.; Xu, X. H.; Qing, F. L. *Chin. J. Org. Chem.* **2015**, *35*, 556 (in Chinese).
(张柯, 徐修华, 卿凤翎, 有机化学, **2015**, *35*, 556.)
(b) Zhang, P. P.; Lu, L.; Shen, Q. L. *Acta Chim. Sinica* **2017**, *75*, 744 (in Chinese).
(张盼盼, 吕龙, 沈其龙, 化学学报, **2017**, *75*, 744.)
- [11] For selected examples, see:
(a) Toulgoat, F.; Alazet, S.; Billard, T. *Eur. J. Org. Chem.* **2014**, *2014*, 2415.
(b) Zhang, M.; Weng, Z. *Org. Lett.* **2019**, *21*, 5838.
(c) Teverovskiy, G.; Surry, D. S.; Buchwald, S. L. *Angew. Chem. Int. Ed.* **2011**, *50*, 7312.
(d) Zhang, C.-P.; Vicić, D. *J. Am. Chem. Soc.* **2012**, *134*, 183.
(e) Chen, C.; Xie, Y.; Chu, L.; Wang, R.-W.; Zhang, X.; Qing, F.-L. *Angew. Chem. Int. Ed.* **2012**, *51*, 2492.
(f) Zhang, C.-P.; Vicić, D. *Chem. Asian J.* **2012**, *7*, 1756.
(g) Chen, C.; Chu, L.; Qing, F.-L. *J. Am. Chem. Soc.* **2012**, *134*, 12454.
(h) Tran, L. D.; Popov, I.; Daugulis, O. *J. Am. Chem. Soc.* **2012**, *134*, 18237.
(i) Weng, Z.; He, W.; Chen, C.; Lee, R.; Tan, D.; Lai, Z.; Kong, D.; Yuan, Y.; Huang, K.-W. *Angew. Chem. Int. Ed.* **2013**, *52*, 1548.
(j) Li, S.-G.; Zard, S. *Org. Lett.* **2013**, *15*, 5898.
(k) Wang, X.; Yang, T.; Cheng, X.; Shen, Q. *Angew. Chem. Int. Ed.* **2013**, *52*, 12860.
(l) Vinogradova, E. V.; Müller, P.; Buchwald, S. L. *Angew. Chem. Int. Ed.* **2014**, *53*, 3125.
- [12] (a) Adams, D. J.; Goddard, A.; Clark, J. H.; Macquarrie, D. J. *Chem. Commun.* **2000**, 987.
(b) Danoun, G.; Bayarmagnai, B.; Grünberg, M.; Gooßen, L. J. *Chem. Sci.* **2014**, *5*, 1312.
(c) Bertoli, G.; Exner, B.; Evers, M. V.; Tschuik, K.; Gooben, L. J. *J. Fluorine Chem.* **2018**, *210*, 132.
(d) Zheng, C.; Liu, Y.; Hong, J.; Huag, S.; Zhang, W.; Yang, Y.; Fang, G. *Tetrahedron Lett.* **2019**, *60*, 1404.
- [13] Liu, S.; Zeng, X.; Xu, B. *Asian J. Org. Chem.* **2019**, *8*, 1372.
- [14] Modak, A.; Pinter, E. N.; Cook, S. P. *J. Am. Chem. Soc.* **2019**, *141*, 18405.
- [15] (a) Sahani, R. L.; Ye, L. W.; Liu, R. S. *Adv. Organomet. Chem.* **2020**, *73*, 195.
(b) Xia, Y.; Qiu, D.; Wang, J. B. *Chem. Rev.* **2017**, *117*, 13810.
(c) Li, L.; Tan, T. D.; Zhang, Y. Q.; Liu, X.; Ye, L. W. *Org. Biomol. Chem.* **2017**, *15*, 8483.
(d) Davies, P. W.; Garzón, M. *Asian J. Org. Chem.* **2015**, *4*, 694.
(e) Davies, H. M. L.; Alford, J. S. *Chem. Soc. Rev.* **2014**, *43*, 5151.
(f) Chattopadhyay, B.; Gevorgyan, V. *Angew. Chem. Int. Ed.* **2012**, *51*, 862.
- [16] For a recent review, see:
Khanal, H. D.; Thombal, R. S.; Maezono, S. M. B.; Lee, Y. R. *Adv. Synth. Catal.* **2018**, *360*, 3185.
- [17] (a) Matheis, C.; Krause, T.; Bragani, V.; Goossen, L. J. *Chem.-Eur. J.* **2016**, *22*, 12270.
(b) Lefebvre, Q.; Fava, E.; Niko-laienko, P.; Rueping, M. *Chem. Commun.* **2014**, *50*, 6617.
(c) Wang, X.; Zhou, Y.; Ji, G.; Wu, G.; Li, M.; Zhang, Y.; Wang, J. *Eur. J. Org. Chem.* **2014**, *2014*, 3093.
(d) Hu, M.; Rong, J.; Miao, W.; Ni, C.; Han, Y.; Hu, J. *Org. Lett.* **2014**, *16*, 2030.
- [18] (a) Lubcke, M.; Yunan, W.; Szabo, K. J. *Org. Lett.* **2017**, *19*, 4548.
(b) Mai, B. K.; Szabo, K. J.; Himo, F. *Org. Lett.* **2018**, *20*, 6646.
- [19] Lübecke, M.; Bezhan, D.; Szabó, K. J. *Chem. Sci.* **2019**, *10*, 5990.
- [20] (a) Xing, Y. P.; Sheng, G. R.; Wang, J.; Lu, P.; Wang, Y. G. *Org. Lett.* **2014**, *16*, 1244.
(b) Sheng, G. R.; Huang, K.; Chi, Z. H.; Ding, H. L.; Xing, Y. P.; Lu, P.; Wang, Y. G. *Org. Lett.* **2014**, *16*, 5096.
- [21] For selected examples, see:
(a) Ma, F. H.; Qian, J.; Lu, P.; Wang, Y. G. *Org. Biomol. Chem.* **2018**, *16*, 439.
(b) Wu, L.; Li, Z. M.; Lu, P.; Wang, Y. G. *J. Org. Chem.* **2018**, *83*, 13956.
(c) Suleman, M.; Li, Z. M.; Lu, P.; Wang, Y. G. *Eur. J. Org. Chem.* **2019**, *27*, 4447.
(d) Sheng, G. R.; Ma, S. C.; Bai, S. L.; Mao, J. M.; Lu, P.; Wang, Y. G. *Chem. Commun.* **2018**, *54*, 1529.
(e) Ding, H. L.; Bai, S. L.; Lu, P.; Wang, Y. G. *Org. Lett.* **2017**, *19*, 4604.
(f) Zhou, Y. X.; Li, Z. M.; Ma, F. H.; Zhao, C.; Lu, P.; Wang, Y. G. *J. Org. Chem.* **2019**, *84*, 6655.
(g) Lang, B.; Zhu, H.; Wang, C.; Lu, P.; Wang, Y. G. *Org. Lett.* **2017**, *19*, 1630.
(h) Sheng, G. R.; Li, Z. M.; Mao, J. M.; Lu, P.; Wang, Y. G. *J. Org. Chem.* **2019**, *84*, 9561.
(i) Sheng, G. R.; Huang, K.; Ma, S. C.; Qian, J.; Lu, P.; Wang, Y. G. *Chem. Commun.* **2015**, *51*, 11056.
- [22] CCDC 1997845 (**3p**) contains supplementary crystallographic data for this paper.

(Li, L.; Fan, Y.)