新型 2,4,6-三取代嘧啶衍生物作为琥珀酸脱氢酶抑制剂的研究

严映坤[†] 程 玮[†] 肖婷婷 张桂兰
张婷婷 陆 童 唐孝荣*
(西华大学理学院 成都 610039)

摘要 设计并合成了 36 个未见文献报道的嘧啶类衍生物,并用 IR、¹H NMR、¹³C NMR 和 HRMS 对其结构进行了表征.以 5 种植物病原真菌即水稻纹枯、小麦赤霉、玉米小斑、油菜菌核和番茄灰霉病菌为靶标,对合成化合物的抑菌活性进行了测定.结果显示,在 20 mg/L 时,其中的大多数都有着显著的抑菌活性.其中,化合物 2-甲基-4-(呋喃-2-基)-6-(对甲苯基)嘧啶(**2c**)和 2-(1-吡唑-1-基)-4-(4-氯苯基)-6-(5-甲基呋喃-2-基)嘧啶(**3d**)对油菜菌核的抑制活性最强,其 EC₅₀ 值分别为 0.072、0.077 mg/L,与市售杀菌剂氟吡菌酰胺(EC₅₀=0.244 mg/L)相比,具有更好的抑菌活性.与此同时,还测定了化合物 **2c** 和 **3d** 对琥珀酸脱氢酶(SDH)的抑制活性.结果表明,它们的半数抑制浓度(IC₅₀)分别为 0.115 和 0.121 mg/L,也比氟吡菌酰胺(IC₅₀=0.356 mg/L)有更好的抑制活性.分子对接研究结果显示,化合物 **2c**、**3d** 和氟吡菌酰胺与 SDH 的结合能分别为-32.2, -31.8 和 -28.9 kJ/mol,这表明它们对 SDH 的亲合力比氟吡菌酰胺更强.化合物 **2c** 和 **3d** 对 SDH 的抑制活性尚属首次报道.

关键词 抑菌活性;嘧啶衍生物;分子对接;琥珀酸脱氢酶抑制剂

Discovery of Novel 2,4,6-Trisubstituted Pyrimidine Derivatives as Succinate Dehydrogenase Inhibitors

Yan, Yingkun[†] Cheng, Wei[†] Xiao, Tingting Zhang, Guilan
Zhang, Tingting Lu, Tong Tang, Xiaorong*
(School of Science, Xihua University, Chengdu 610039)

Abstract Thirty-six unreported pyrimidine analogues were designed, synthesized and characterized by IR, ¹H NMR, ¹³C NMR and HRMS. Their antifungal activities were determined against five plant pathogenic fungi namely *Rhizoctonia solani*, *Fusarium graminearum*, *Helminthosporium maydis*, *Sclerotinia sclerotiorum* and *Botrytis cinerea*. The results indicated that most of them revealed significant antifungal activities at 20 mg/L. Among them, 4-(furan-2-yl)-2-methyl-6-(*p*-tolyl)pyrimidine (**2c**) and 4-(4-chlorophenyl)-6-(5-methylfuran-2-yl)-2-(1*H*-pyrazol-1-yl)pyrimidine (**3d**) showed the strongest activities against *Sclerotinia sclerotiorum* and their median effect concentrations (EC₅₀) were 0.072 and 0.077 mg/L, respectively, which implied that they had better antifungal activities than the commercial fungicide fluopyram (EC₅₀=0.244 mg/L). Meanwhile, the inhibitory activities of compounds **2c** and **3d** were determined against succinate dehydrogenase (SDH). The results exhibited that their half inhibitory concentrations (IC₅₀) were 0.115 and 0.121 mg/L, respectively, indicating that they also had better inhibitory activities than fluopyram (IC₅₀=0.356 mg/L). Molecular docking studies demonstrated that the binding energy of compounds **2c**, **3d** and fluopyram to SDH was -32.2 kJ/mol, -31.8 kJ/mol and -28.9 kJ/mol, respectively, which represented that they had stronger affinities than fluopyram. The inhibitory activities of compounds **2c** and **3d** against SDH have been reported for the first time.

Keywords antifungal activity; pyrimidine derivative; molecular docking; succinate dehydrogenase inhibitor

* Corresponding author. E-mail: tangxr112@126.com

Received May 21, 2020; revised June 30, 2020; published online August 11, 2020.

Project supported by the Chunhui Programs of the Ministry of Education (No. 191653), the Technical Innovation Programs of Chengdu Municipal Bureau of Science and Technology (No. 2018-YF05-00970-SN), the Innovation Fund of Postgraduate of Xihua University (No. YCJJ2019025) and the Undergraduate Innovation and Entrepreneurship Training Programs of Xihua University (No. 201810623009).

教育部春晖计划(No. 191653)、成都市科学技术局技术创新(No. 2018-YF05-00970-SN)、西华大学研究生创新基金(No. YCJJ2019025)、西华大学大学生创新创业训练(No. 201810623009)资助项目.

[†] 共同第一作者(These authors contributed equally to this work.)

1 Introduction

Succinate dehydrogenase (SDH, also known as complex II), which catalyzes the oxidation of succinate to fumarate in the mitochondrial matrix, is the only enzyme complex simultaneously involved in the respiration chain and tricarboxylic acid cycle.^[1-3] Due to its crucial role in life processes, SDH has been particularly considered as a promising target for agrochemical discovery. Succinate dehydrogenase inhibitor (SDHI) fungicides are an important category of fungicides. SDHIs have been widely used to control destructive plant pathogenic fungi. They exert their fungicidal activities by disrupting the mitochondrial tricarboxylic acid cycle and respiration chain.^[4] Moreover, they have no cross-resistance with other commercial fungicides because of their unique mode of action.^[4] Therefore, SDHIs have been extensively investigated in the world.^[5-8] Eighteen SDHIs have been successfully developed and employed since the first SDHI carboxin was used in 1966. Meanwhile, the researches and innovations of SDHIs have been the frontier of the fungicidal field and novel SDHIs are ceaselessly reported.^[9-12] These SDHIs have the following common features in molecular structure: (a) amide scaffolds, (b) halogen atoms, methyl groups, methoxy groups or (c) heterocyclic rings, such as furan rings, thiophene rings, pyrazole rings, pyridine rings (Figure 1).

Figure 1 Representatives of succinate dehydrogenase inhibitor fungicides

Pyrimidine derivatives are an important class of compounds. They have been found in a variety of natural products, such as nucleic acids and thiamine (vitamin B1), which are important to the research and development of

pharmaceutical and agrochemical compounds. In the meantime, pyrimidine derivatives possess diverse and important biological properties, including antibacterial,^[13-15] anticancer,^[16-19] antiviral,^[20-21] anti-inflammatory,^[22-24] analgesic,^[22,24] antimalarial,^[25-26] anti-tuberculosis,^[27-28] anticonvulsant^[29-30] and antioxidant.^[22] In addition, pyrimidine derivatives are important fungicides in the agrochemical field. Dozens of fungicides containing pyrimidine rings have been successfully developed and employed since the first pyrimidine fungicide ethirimol was used in 1968 (Figure 2). Currently, structural modifications on pyrimidine rings are still ongoing to discover more bioactive compounds.

Figure 2 Representatives of commercial pyrimidine derivative fungicides

As mentioned above, in the present study, thirty-six new pyrimidine analogues were designed in accordance with the principle of splicing bioactive substructures and the following modifications of pyrimidine rings: (a) substituting the hydrogen atom at position-2 with the methyl group, pyrazole ring or amide moiety, (b) substituting the hydrogen atom at position-4 with the furan ring/5-methyl furan ring or thiophene ring/5-bromothiophene ring and (c) substituting the hydrogen atom at position-6 with the phenyl ring substituted by the fluorine atom, chlorine atom, bromine atom, methyl group or methoxy group (Scheme 1). Meanwhile, the designed compounds were synthesized and their antifungal activities were evaluated in the laboratory to discover novel SDHIs with simple structures and excellent performances. Molecular docking studies were conducted to detect the probable interactions between the active compounds and SDH.

Scheme 1 Design of target compounds

2 Results and discussion

2.1 Chemistry

The syntheses and chemical structures of the target compounds are shown in Scheme 2. The syntheses of the designed compounds were performed according to the methods reported previously and modified appropriately.^[31-32] Intermediate **1** was obtained by Claisen-Schmidt condensation reaction and the characterizing data of these compounds conformed to the literatures.^[33-36] The target-compounds (**2a~2l**, **3a~3l** and **4a~4l**) were synthesized by Michael addition between the α,β -unsaturated C=C double bond of intermediate **1** and acetimidamide, pyrazole-1-carboximidamide or guanyl urea in the presence of base, which was followed by a cyclization reaction. All the synthesized compounds were purified by crystallization or silica gel chromatography. Their purity was checked by

Compd. 2	X	R ¹	R ²	Compd. 3	X	R ¹	R ²	Compd. 4	X	R ¹	R ²
2a	O	4-F	H	3a	O	4-Br	H	4a	O	4-Cl	H
2b	O	3,4-Cl ₂	H	3b	O	4-CH ₃	H	4b	O	4-OCH ₃	H
2c	O	4-CH ₃	H	3c	O	4-OCH ₃	H	4c	O	4-Cl	CH ₃
2d	O	4-Cl	CH ₃	3d	O	4-Cl	CH ₃	4d	O	4-CH ₃	CH ₃
2e	O	4-CH ₃	CH ₃	3e	O	4-CH ₃	CH ₃	4e	O	4-OCH ₃	CH ₃
2f	O	4-OCH ₃	CH ₃	3f	S	4-Cl	H	4f	S	4-Cl	H
2g	S	4-Cl	H	3g	S	4-CH ₃	H	4g	S	4-CH ₃	H
2h	S	4-CH ₃	H	3h	S	4-OCH ₃	H	4h	S	4-OCH ₃	H
2i	S	4-OCH ₃	H	3i	S	4-Cl	Br	4i	S	4-Cl	Br
2j	S	4-Br	Br	3j	S	4-Br	Br	4j	S	4-Br	Br
2k	S	4-CH ₃	Br	3k	S	4-CH ₃	Br	4k	S	4-CH ₃	Br
2l	S	4-OCH ₃	Br	3l	S	4-OCH ₃	Br	4l	S	4-OCH ₃	Br

Reagents and conditions: (a) 10% NaOH anhydrous ethanol, 0~5 °C, 3~4 h; (b) 10% NaOH anhydrous ethanol, 65~70 °C, 4~5 h; (c) NaOH/MeOH, 70~75 °C, 5~6 h; (d) DMF/K₂CO₃, 55~60 °C, 5~6 h

Scheme 2 Synthesis of target compounds

thin layer chromatography (TLC). The structures of all the synthesized compounds were confirmed by IR, ^1H NMR, ^{13}C NMR and HRMS. The results showed that their spectrum data were in agreement with the proposed structures.

2.2 Antifungal activity

Compared with the efficient fungicide fluopyram, all the synthesized compounds were submitted to laboratorial assay using *R. solani*, *F. graminearum*, *H. maydis*, *S. sclerotiorum* and *B. cinerea* as targets. The results are listed in Table 1. Most of the target compounds displayed remarkable antifungal activities at 20 mg/L. Among them, **2a**, **2c**, **2f**, **3d**, **3j**, **4b** and **4j** possessed excellent antifungal activities and their inhibitory rate reached 100% at 20 mg/L against *S. sclerotiorum*. Moreover, compounds **2c** and **3d** showed better activities than fluopyram against *R. solani*

and *F. graminearum* at 20 mg/L.

As shown in Table 1, all the synthesized compounds showed inhibitory activities against the above five plant pathogenic fungi in the following order: *S. sclerotiorum* > *H. maydis* > *B. cinerea* > *R. solani* > *F. graminearum*. If compounds **2c**, **3b**, **4a** and **4b** were compared with compounds **2e**, **3e**, **4c** and **4e**, respectively, it could be found that when the hydrogen atom at position-5 of the furan ring was substituted by a methyl group, the antifungal activity of the corresponding compounds decreased, which needs to be further studied. Likewise, when compounds **2c**, **3b**, **3c**, **4a** and **4b** were compared with compounds **2h**, **3g**, **3h**, **4f** and **4h**, respectively, it was noticeable that when the furan ring of these compounds was changed into a thiophene ring, the antifungal activities of the corresponding

Table 1 Antifungal activity of target compounds at 20 mg/L

Compound	Inhibition rate ^a /%				
	<i>R. solani</i>	<i>F. graminearum</i>	<i>H. maydis</i>	<i>S. sclerotiorum</i>	<i>B. cinerea</i>
2a	51.8±2.2	30.1±0.9	88.2±3.5	100.0±0.0	76.5±3.5
2b	39.0±1.3	26.6±1.1	50.3±2.0	80.1±3.8	43.8±1.9
2c	75.6±2.9	68.2±3.3	91.6±4.1	100.0±0.0	83.2±2.8
2d	58.3±2.6	40.1±1.9	83.4±3.8	98.6±1.1	73.3±3.2
2e	56.7±2.0	43.4±1.3	81.7±3.4	96.8±2.8	78.2±3.1
2f	61.9±2.4	45.5±2.2	86.7±3.8	100.0±0.0	80.5±3.5
2g	32.8±1.3	26.4±0.9	51.9±2.1	64.3±2.1	40.3±1.9
2h	30.5±0.7	21.3±0.3	55.0±2.3	68.5±2.2	44.7±1.8
2i	44.5±0.5	30.5±0.9	67.5±2.8	73.3±3.3	51.8±2.2
2j	41.2±1.3	38.9±1.0	65.1±2.7	68.0±2.8	55.4±1.9
2k	51.1±1.2	36.5±1.6	84.3±4.0	94.7±2.4	77.9±2.8
2l	50.3±1.7	38.1±1.8	70.3±3.1	81.2±4.0	58.4±2.1
3a	60.3±2.2	48.5±2.1	84.7±3.6	94.9±3.8	74.5±3.3
3b	54.1±1.9	44.2±2.0	82.5±3.6	97.6±2.0	77.0±2.5
3c	40.7±1.7	33.2±1.3	68.2±2.7	82.7±2.6	59.2±2.5
3d	74.1±2.3	67.9±2.5	90.3±3.4	100.0±0.0	81.5±3.2
3e	40.8±2.1	36.4±1.5	67.2±2.6	75.6±3.2	54.3±2.0
3f	32.0±0.8	23.5±1.0	56.4±2.3	66.0±3.1	47.6±1.4
3g	30.9±0.9	20.2±0.2	53.8±2.1	68.0±2.7	47.9±2.5
3h	35.2±1.6	22.1±0.3	56.4±2.4	68.7±3.1	50.5±1.6
3i	60.0±2.2	42.6±1.9	83.5±3.7	97.9±1.7	75.3±2.9
3j	57.0±2.3	49.3±1.5	88.6±4.2	100.0±0.0	76.1±2.4
3k	46.3±1.5	37.6±1.4	61.3±2.0	78.2±3.1	55.7±2.8
3l	45.7±1.9	30.6±1.0	67.0±2.6	84.3±3.9	58.1±2.0
4a	51.4±1.3	42.4±1.7	83.2±3.9	97.2±1.9	75.1±2.1
4b	60.9±2.9	48.2±2.2	87.6±4.0	100.0±0.0	74.3±3.3
4c	32.6±0.7	25.3±1.3	51.9±2.6	65.3±2.3	43.5±2.0
4d	39.1±1.3	28.6±0.4	61.4±3.0	73.8±2.3	58.0±1.8
4e	52.7±1.7	41.1±1.8	81.9±3.5	94.1±3.1	60.2±2.4
4f	32.0±1.0	20.9±1.1	50.6±2.3	61.3±2.8	41.6±1.0
4g	33.2±1.4	23.6±0.6	52.8±2.2	60.2±2.5	43.2±1.3
4h	31.7±1.5	26.9±0.9	45.1±1.8	55.1±1.3	38.4±1.4
4i	50.8±1.7	45.6±1.2	77.8±3.3	95.7±2.1	65.7±1.2
4j	60.6±2.4	45.3±1.6	87.6±4.3	100.0±0.0	76.9±2.2
4k	41.6±1.5	35.4±0.6	60.3±2.7	71.2±1.0	52.3±2.4
4l	51.5±2.3	41.8±1.5	78.9±3.1	92.3±3.3	58.2±2.2
Fluopyram	71.4±1.9	62.9±1.0	100.0±0.0	100.0±0.0	100.0±0.0

^a Data are the mean ± standard deviation (SD) of three replicates.

compounds also decreased. However, if compounds **2h**, **2i**, **3f**, **3g**, **3h**, **4f**, **4g** and **4h** were contrasted with compounds **2k**, **2l**, **3i**, **3k**, **3l**, **4i**, **4k** and **4l**, it could be found that when the hydrogen atom at position-5 of the thiophene ring was replaced by a bromine atom, the antifungal activities of the corresponding compounds increased, whose causes also deserve to be further investigated.

The EC_{50} values of the compounds with excellent effects against *S. sclerotiorum* were further determined to investigate their antifungal activities. The results are displayed in Table 2 and Figure 3. The EC_{50} values of seventeen compounds (**2a**, **2c**~**2f**, **2k**, **3a**, **3b**, **3d**, **3i**, **3j**, **4a**, **4b**, **4e**, **4i**, **4j** and **4l**) ranged from 0.072~1.230 mg/L. Among them, the EC_{50} values of compounds **2c** and **3d** reached 0.072 and 0.077 mg/L, respectively, whereas the EC_{50} value of fluopyram was 0.244 mg/L, indicating that they possessed better inhibitory activities than fluopyram. Similarly, in accordance with the EC_{50} values of compounds **2a**, **2f**, **3j**, **4b** and **4j**, it can be known that their inhibitory activities were close to fluopyram.

Table 2 EC_{50} values of the compounds with excellent effects against *S. sclerotiorum*

Compound	$EC_{50}^a/(mg \cdot L^{-1})$	Compound	$EC_{50}^a/(mg \cdot L^{-1})$
2a	0.258 ± 0.024	3i	0.704 ± 0.081
2c	0.072 ± 0.010	3j	0.299 ± 0.030
2d	0.632 ± 0.081	4a	0.740 ± 0.079
2e	0.768 ± 0.091	4b	0.267 ± 0.026
2f	0.272 ± 0.027	4e	1.106 ± 0.161
2k	1.003 ± 0.102	4i	0.807 ± 0.091
3a	0.984 ± 0.110	4j	0.327 ± 0.032
3b	0.711 ± 0.082	4l	1.230 ± 0.186
3d	0.077 ± 0.010	Fluopyram	0.244 ± 0.019

^a Data are the mean $\pm$ standard deviation (SD) of three replicates.

Figure 3 Comparison of EC_{50} values among compounds with excellent effects against *S. sclerotiorum*

Although definite structure-activity relationships could not be found through the above-mentioned findings, some interesting results obtained can be used for designing and synthesizing more analogues to further study their quantitative structure-activity relationship so that more bioactive

compounds or lead compounds may be discovered. For example, compounds **2c** and **3d** might be further used as lead compounds because they exhibited excellent inhibitory activities and had simple structures.

2.3 Inhibitory activity against enzyme

The assay of the inhibitory activity against SDH was performed to investigate whether SDH is a potential target enzyme of the synthesized compounds. Using fluopyram as the comparative standard, the inhibitory activities of compounds **2c** and **3d** were tested against the SDH from the mitochondria of *S. sclerotiorum*. The results are listed in Table 3. The IC_{50} values of compounds **2c** and **3d** were 0.115, 0.121 mg/L, respectively, which revealed that they possessed better inhibitory activity than fluopyram (IC_{50} = 0.356 mg/L). Therefore, the findings implied that SDH may be probably one action target of these compounds.

Table 3 Inhibitory activities of compounds **2c** and **3d** against SDH

Compound	$IC_{50}^a/(mg \cdot L^{-1})$
2c	0.115 ± 0.011
3d	0.121 ± 0.012
Fluopyram	0.356 ± 0.040

^a Data are the mean $\pm$ standard deviation (SD) of three replicates.

2.4 Homology modelling and molecular docking analysis

The theoretical binding modes of compounds **2c**, **3d** and fluopyram were simulated to investigate whether SDH is a potential target enzyme of the target compounds. Their theoretical binding modes to SDH are displayed in Figure 4. Fluopyram fitted in the gap composed of subunits B, C and D (Figure 4A). The 2-trifluoromethylphenyl group of fluopyram was located in the hydrophobic pocket surrounded by residues B/Pro-226, B/Trp-230 and B/Ile-275, whereas the 3-chloro-5-(trifluoromethyl) pyridine ring of fluopyram stretched into the hydrophobic pocket that consisted of residues C/Trp-81, C/Ile-82 and C/Ile-89, forming strong hydrophobic binding. In addition, the 2-trifluoromethylphenyl group of fluopyram formed a cation- π interaction with residue C/Arg-88.

The binding mode of compound **2c** to SDH is illustrated in Figure 4B. Compound **2c** also fitted in the gap composed of subunits B, C and D. It was located in the hydrophobic pocket surrounded by residues B/Pro-226, B/Trp-230, B/Ile-275, C/Leu-72, C/Trp-81 and C/Ile-82, forming strong hydrophobic binding. Like fluopyram, the furan ring of compound **2c** also formed a cation- π interaction with residue C/Arg-88. Besides, compound **2c** and residue B/Trp-230 formed one hydrogen bond with a length of 3.1 Å.

As mentioned above, compound **2c** and fluopyram shared similar binding modes (Figure 4C). The main difference was that compound **2c** formed an extra hydrogen bond with residue B/Trp-230 when compared with fluopyram, which might make compound **2c** more active than fluopyram against SDH. Meanwhile, the estimated binding

Figure 4 Theoretical binding modes of fluopyram (A), compound **2c** (B), fluopyram and compound **2c** (C), compound **3d** (D), fluopyram and compound **3d** (E) to SDH

energy of compound **2c** and fluopyram was -32.2 and -28.9 kJ/mol, respectively, which implied compound **2c** had stronger affinity to SDH than fluopyram.

The binding mode of compound **3d** to SDH is outlined in Figure 4D. Compound **3d** also fitted in the gap composed of subunits B, C and D. It was located in the hydrophobic pocket surrounded by residues B/Trp-230, B/Ile-275, C/Leu-72, C/Trp-81, C/Ile-82 and C/Ile-89, forming strong hydrophobic binding. The 4-chlorophenyl group of compound **3d** formed a π - π stacking interaction with resi-

due C/Trp-81. Like fluopyram, the pyrazole ring of compound **3d** also formed a cation- π interaction with residue C/Arg-88. In addition, compound **3d** and residue B/Trp-230 formed one hydrogen bond with a length of 3.2 Å.

In a word, compound **3d** and fluopyram also had similar binding modes (Figure 4E). The main difference was also that compound **3d** formed an extra hydrogen bond with residue B/Trp-230 when compared with fluopyram, which might make compound **3d** more active than fluopyram against SDH. Furthermore, the estimated binding energy of

compound **3d** and fluopyram was -31.8 and -28.9 kJ/mol, respectively, which signified compound **3d** had stronger affinity to SDH than fluopyram.

The above molecular docking results displayed that compounds **2c** and **3d** might be potential inhibitors of SDH. Meanwhile, the different binding energy between SDH and compounds **2c**, **3d** or fluopyram revealed that compounds **2c** and **3d** possessed stronger affinity to SDH than fluopyram, which was in agreement with the results of the antifungal activity assay. These results rendered rational explanations of the interactions between SDH and compounds **2c**, **3d** or fluopyram and provided some valuable information for the further research and development of SDHI fungicides.

3 Conclusions

Thirty-six pyrimidine derivatives were designed, synthesized and tested for their antifungal activities against the above five phytopathogenic fungi. The seven compounds (**2a**, **2c**, **2f**, **3d**, **3j**, **4b** and **4j**) with excellent antifungal activity were discovered against *S. sclerotiorum*. They have simple structures and are easily synthesized. In particular, compounds **2c** and **3d** exhibited potent activities against *S. sclerotiorum* and their antifungal activities even surpassed fluopyram. They can be further used as lead compounds to design and synthesize more analogues for the discovery of more active compounds. The results of molecular docking studies may be conducive to further exploring the possible fungicidal mechanism of the above compounds and the interactions between similar fungicidal compounds and SDH.

4 Experimental section

4.1 General

IR spectra were recorded on a Nicolet 380 spectrometer. NMR spectra (^1H NMR and ^{13}C NMR) were taken on an AVANCE NEO 400 spectrometer using deuterated dimethyl sulfoxide ($\text{DMSO}-d_6$) as the solvent. HRMS data were taken on an LTQ Orbitrap Elite mass spectrometer. Melting points were determined using an X-4B micro-melting point apparatus and were not corrected. All chemicals and solvents were purchased from commercial sources unless specified otherwise. *R. solani*, *F. graminearum*, *H. maydis*, *S. sclerotiorum* and *B. cinerea* were obtained from the Chinese Academy of Agricultural Sciences. They were preserved at $4\text{ }^\circ\text{C}$.

4.2 Synthesis of target compounds

4.2.1 Synthesis of intermediate **1**

As shown in Scheme 1, in accordance with the methods reported previously,^[33-36] a mixture of selected aldehyde (0.01 mol), substituted acetophenone (0.01 mol) and anhydrous ethanol (20 mL) was stirred in an ice bath. 10% NaOH anhydrous ethanol (5 mL) was added to the mixture. The reaction was kept at $0\sim 5\text{ }^\circ\text{C}$ for $3\sim 4$ h. The course of the reaction was monitored by thin-layer chro-

matography (TLC). After completion, distilled water was added to the reaction mixture. The pH value of the mixture was adjusted to about 7 by 10% HCl solution. The solid that precipitated was filtered and washed with distilled water. The obtained crude product was recrystallized from anhydrous ethanol to afford intermediate **1**.

4.2.2 Synthesis of target compounds **2a**~**2l**

As shown in Scheme 1, a mixture of intermediate **1** (0.005 mol), anhydrous ethanol (10 mL) and 10% NaOH anhydrous ethanol (5 mL) was stirred at room temperature. The solution of acetimidamide hydrochloride (0.01 mol) and anhydrous ethanol (10 mL) was slowly added to the above mixture. The reaction was kept at $65\sim 70\text{ }^\circ\text{C}$ for $4\sim 5$ h. The process of the reaction was monitored by TLC. After completion, the reaction mixture was neutralized with HCl (10%) till the solution became neutral. The solvent was removed under reduced pressure. The obtained residue was purified by recrystallization from anhydrous ethanol to obtain compounds **2a**~**2l**.

4-(4-Fluorophenyl)-6-(furan-2-yl)-2-methylpyrimidine (**2a**): White powder, yield 81%. m.p. $124\sim 125\text{ }^\circ\text{C}$; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.37~8.32 (m, 2H), 8.11 (s, 1H), 8.01 (dd, $J=1.6, 0.8$ Hz, 1H), 7.50 (dd, $J=3.6, 0.4$ Hz, 1H), 7.40 (t, $J=8.8$ Hz, 2H), 6.77 (dd, $J=3.2, 1.6$ Hz, 1H), 2.69 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 168.2, 164.5 (d, $J=247.0$ Hz), 163.0, 156.1, 151.8, 146.6, 133.2 (d, $J=3.0$ Hz), 130.0 (d, $J=9.0$ Hz), 116.3 (d, $J=21.0$ Hz), 113.5, 113.3, 107.6, 26.5; IR (KBr) ν_{max} : 1603, 1565, 1536, 1509, 1381, 1102, 842 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{12}\text{N}_2\text{OF}$ $[\text{M} + \text{H}]^+$ 255.0928, found 255.0924.

4-(3,4-Dichlorophenyl)-6-(furan-2-yl)-2-methylpyrimidine (**2b**): Yellow powder, yield 89%. m.p. $139\sim 140\text{ }^\circ\text{C}$; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.53 (d, $J=2.0$ Hz, 1H), 8.28 (dd, $J=8.4, 1.6$ Hz, 1H), 8.23 (s, 1H), 7.84 (d, $J=8.4$ Hz, 1H), 7.54 (d, $J=3.2$ Hz, 1H), 6.78 (dd, $J=3.2, 1.6$ Hz, 1H), 2.71 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 168.3, 161.5, 156.5, 151.6, 146.8, 137.3, 134.2, 132.4, 131.6, 129.3, 127.6, 113.9, 113.3, 108.2, 26.4; IR (KBr) ν_{max} : 1605, 1549, 1523, 1379, 1121, 819 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{11}\text{Cl}_2\text{N}_2\text{O}$ $[\text{M} + \text{H}]^+$ 305.0243, found 305.0237.

4-(Furan-2-yl)-2-methyl-6-(*p*-tolyl)pyrimidine (**2c**): Yellow powder, yield 85%. m.p. $86\sim 87\text{ }^\circ\text{C}$; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.18 (d, $J=8.4$ Hz, 2H), 8.07 (s, 1H), 8.00 (d, $J=0.8$ Hz, 1H), 7.48 (d, $J=2.8$ Hz, 1H), 7.37 (d, $J=8.0$ Hz, 2H), 6.77 (dd, $J=3.2, 1.6$ Hz, 1H), 2.69 (s, 3H), 2.40 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 168.1, 164.0, 156.0, 151.9, 146.5, 141.5, 134.0, 130.0, 127.5, 113.3, 113.2, 107.3, 26.5, 21.5; IR (KBr) ν_{max} : 1605, 1522, 1383, 1106, 829 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{15}\text{N}_2\text{O}$ $[\text{M} + \text{H}]^+$ 251.1179, found 251.1179.

4-(4-Chlorophenyl)-2-methyl-6-(5-methylfuran-2-yl)pyrimidine (**2d**): Yellow powder, yield 77%. m.p. $132\sim 133\text{ }^\circ\text{C}$; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.28 (d, $J=8.4$ Hz, 2H), 8.04 (s, 1H), 7.62 (d, $J=8.8$ Hz, 2H), 7.41 (d, $J=$

3.2 Hz, 1H), 6.40 (d, $J=2.8$ Hz, 1H), 2.68 (s, 3H), 2.43 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 168.2, 162.5, 156.2, 156.1, 150.2, 136.3, 135.6, 129.4, 129.3, 115.0, 109.8, 107.2, 26.5, 14.2; IR (KBr) ν_{max} : 1606, 1573, 1517, 1381, 1095, 836 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{14}\text{N}_2\text{OCl} [\text{M}+\text{H}]^+$ 285.0789, found 285.0785.

2-Methyl-4-(5-methylfuran-2-yl)-6-(*p*-tolyl)pyrimidine (**2e**): Brown powder, yield 88%. m.p. 143~144 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 8.16 (d, $J=8.0$ Hz, 2H), 7.98 (s, 1H), 7.39~7.36 (m, 3H), 6.39 (d, $J=2.8$ Hz, 1H), 2.67 (s, 3H), 2.44 (s, 3H), 2.41 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 168.0, 163.7, 156.0, 155.9, 150.4, 141.4, 134.0, 130.0, 127.4, 114.6, 110.0, 106.7, 26.5, 21.5, 14.2; IR (KBr) ν_{max} : 1604, 1515, 1382, 1117, 814 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{17}\text{N}_2\text{O} [\text{M}+\text{H}]^+$ 265.1335, found 265.1338.

4-(4-Methoxyphenyl)-2-methyl-6-(5-methylfuran-2-yl)-pyrimidine (**2f**): Yellow powder, yield 80%. m.p. 107~108 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 8.23 (d, $J=8.8$ Hz, 2H), 7.94 (s, 1H), 7.37 (d, $J=3.2$ Hz, 1H), 7.10 (d, $J=8.8$ Hz, 2H), 6.39 (d, $J=2.8$ Hz, 1H), 3.86 (s, 3H), 2.65 (s, 3H), 2.43 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 167.9, 163.4, 162.1, 155.8, 150.4, 129.1, 127.5, 114.7, 114.5, 109.7, 106.2, 55.9, 26.5, 14.2; IR (KBr) ν_{max} : 2923, 1629, 1573, 1515, 1381, 1108 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{17}\text{N}_2\text{O}_2 [\text{M}+\text{H}]^+$ 281.1285, found 281.1280.

4-(4-Chlorophenyl)-2-methyl-6-(thiophen-2-yl)pyrimidine (**2g**): White powder, yield 83%. m.p. 86~87 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 8.42 (s, 1H), 8.36 (d, $J=8.4$ Hz, 2H), 8.30 (dd, $J=3.6$, 1.2 Hz, 1H), 7.85 (dd, $J=5.2$, 1.2 Hz, 1H), 7.65 (d, $J=8.8$ Hz, 2H), 7.29 (dd, $J=4.8$, 3.6 Hz, 1H), 2.68 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 168.1, 162.7, 160.0, 142.7, 136.4, 135.6, 131.6, 129.6, 129.4, 129.3, 108.2, 26.4; IR (KBr) ν_{max} : 2923, 1629, 1515, 1381, 1097, 817 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{12}\text{N}_2\text{SCl} [\text{M}+\text{H}]^+$ 287.0404, found 287.0410.

2-Methyl-4-(thiophen-2-yl)-6-(*p*-tolyl)pyrimidine (**2h**): White crystal, yield 80%. m.p. 103~104 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 8.34 (s, 1H), 8.28 (d, $J=3.6$ Hz, 1H), 8.23 (d, $J=8.0$ Hz, 2H), 7.83 (d, $J=4.8$ Hz, 1H), 7.38 (d, $J=8.0$ Hz, 2H), 7.29 (t, $J=4.0$ Hz, 1H), 2.67 (s, 3H), 2.41 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 168.0, 163.8, 159.7, 142.9, 141.4, 134.0, 131.3, 129.9, 129.2, 127.6, 125.4, 107.7, 26.4, 21.5; IR (KBr) ν_{max} : 1630, 1523, 1383, 1107 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{15}\text{N}_2\text{S} [\text{M}+\text{H}]^+$ 267.0950, found 267.0945.

4-(4-Methoxyphenyl)-2-methyl-6-(thiophen-2-yl)pyrimidine (**2i**): Yellow powder, yield 86%. m.p. 167~168 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 8.31~8.26 (m, 4H), 7.82 (d, $J=4.8$ Hz, 2H), 7.28 (t, $J=4.0$ Hz, 1H), 7.11 (d, $J=8.8$ Hz, 2H), 3.86 (s, 3H), 2.66 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 167.9, 163.5, 162.2, 159.5, 143.0, 131.1, 129.3, 129.2, 129.1, 114.7, 107.2, 55.9, 26.4; IR (KBr) ν_{max} : 1603, 1565, 1536, 1509, 1381, 1102, 842 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{15}\text{N}_2\text{OS} [\text{M}+\text{H}]^+$ 283.0940, found 283.0970.

4-(4-Bromophenyl)-6-(5-bromothiophen-2-yl)-2-methyl-

pyrimidine (**2j**): White powder, yield 77%. m.p. 161~162 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 8.48 (s, 2H), 8.42 (s, 2H), 8.31 (d, $J=8.0$ Hz, 1H), 8.15 (d, $J=4.0$ Hz, 1H), 7.77 (d, $J=8.0$ Hz, 1H), 7.53 (t, $J=8.0$ Hz, 1H), 7.43 (d, $J=4.0$ Hz, 1H), 2.66 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 168.2, 162.4, 159.1, 144.2, 138.9, 131.5, 130.3, 130.1, 126.6, 122.9, 117.7, 108.0, 26.3; IR (KBr) ν_{max} : 1574, 1525, 1389, 1103, 798 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{11}\text{N}_2\text{SBr}_2 [\text{M}+\text{H}]^+$ 408.9004, found 408.9000.

4-(5-Bromothiophen-2-yl)-2-methyl-6-(*p*-tolyl)pyrimidine (**2k**): Light yellow powder, yield 85%. m.p. 167~168 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 8.35 (s, 1H), 8.22 (d, $J=8.0$ Hz, 2H), 8.13 (d, $J=4.0$ Hz, 1H), 7.42 (d, $J=4.0$ Hz, 1H), 7.38 (d, $J=8.0$ Hz, 2H), 2.66 (s, 3H), 2.40 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 168.0, 164.1, 158.7, 144.5, 141.6, 133.8, 132.7, 130.0, 129.9, 127.6, 117.4, 107.3, 26.3, 21.5; IR (KBr) ν_{max} : 1574, 1523, 1379, 1121, 822, 785 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{14}\text{N}_2\text{SBr} [\text{M}+\text{H}]^+$ 345.0056, found 345.0046.

4-(5-Bromothiophen-2-yl)-6-(4-methoxyphenyl)-2-methylpyrimidine (**2l**): Light yellow powder, yield 82%. m.p. 139~140 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 8.31 (s, 1H), 8.29 (d, $J=8.8$ Hz, 2H), 8.12 (d, $J=4.0$ Hz, 1H), 7.42 (d, $J=4.0$ Hz, 1H), 7.11 (d, $J=8.4$ Hz, 2H), 3.87 (s, 3H), 2.65 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 167.9, 163.7, 162.3, 158.5, 144.6, 132.6, 129.7, 129.3, 128.9, 117.2, 114.7, 106.7, 55.9, 26.3; IR (KBr) ν_{max} : 1605, 1574, 1517, 1390, 1253, 1114, 837, 801 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{14}\text{N}_2\text{OSBr} [\text{M}+\text{H}]^+$ 361.0005, found 361.0004.

4.2.3 Synthesis of target compounds **3a**~**3l**

As shown in Scheme 1, a mixture of intermediate **1** (0.005 mol), anhydrous methanol (10 mL) and NaOH (0.02 mol) was stirred at room temperature. The solution of pyrazole-1-carboximidamide hydrochloride (0.01 mol) and anhydrous methanol (10 mL) was slowly added to the above mixture. The reaction was kept at 70~75 °C for 5~6 h. The process of the reaction was monitored by TLC. After completion, the reaction mixture was poured into ice water and neutralized by HCl solution (10%) till the solution became neutral and extracted with ethyl acetate (10 mL $\times$ 3). The organic phase was dried with anhydrous Na_2SO_4 and concentrated under reduced pressure. The obtained residue was purified by silica gel (200~300 mesh) chromatography using petroleum ether (b.p. 60~90 °C)/ethyl acetate ($V:V=1:2$) as the eluting system to give compounds **3a**~**3l**.

4-(4-Bromophenyl)-6-(furan-2-yl)-2-(1*H*-pyrazol-1-yl)-pyrimidine (**3a**): Brown powder, yield 85%. m.p. 100~101 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 8.89 (s, 1H), 8.37 (d, $J=8.4$ Hz, 2H), 8.26 (s, 1H), 8.10 (s, 1H), 7.91 (s, 1H), 7.82 (d, $J=8.0$ Hz, 2H), 7.70 (d, $J=2.4$ Hz, 1H), 6.85 (s, 1H), 6.66 (s, 1H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 164.8, 157.9, 156.3, 151.2, 147.4, 143.7, 135.2, 132.5, 130.4, 129.9, 126.0, 115.2, 113.6, 109.2, 108.2; IR (KBr)

$\nu_{\max}$: 1601, 1517, 1456, 1397, 1256, 1175, 835, 759 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{12}\text{N}_4\text{OBr}$ $[\text{M}+\text{H}]^+$ 367.0189, found 367.0183.

4-(Furan-2-yl)-2-(1*H*-pyrazol-1-yl)-6-(*p*-tolyl)pyrimidine (**3b**): Black powder, yield 85%. m.p. 71~72 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.88 (s, 1H), 8.36 (d, $J=8.4$ Hz, 2H), 8.26 (s, 1H), 8.09 (s, 1H), 7.91 (s, 1H), 7.81 (d, $J=8.0$ Hz, 2H), 7.69 (d, $J=2.4$ Hz, 1H), 6.84 (s, 1H), 6.65 (s, 1H), 2.42 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 161.9, 154.9, 153.3, 148.2, 144.4, 140.7, 132.2, 129.5, 127.4, 126.9, 123.0, 112.2, 110.6, 106.2, 105.2, 23.0; IR (KBr) $\nu_{\max}$: 1585, 1521, 1455, 1396, 1366, 1077, 824, 769 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{15}\text{N}_4\text{O}$ $[\text{M}+\text{H}]^+$ 303.1246, found 303.1248.

4-(Furan-2-yl)-6-(4-methoxyphenyl)-2-(1*H*-pyrazol-1-yl)pyrimidine (**3c**): Yellow powder, yield 80%. m.p. 114~115 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.88 (s, 1H), 8.39 (d, $J=8.4$ Hz, 2H), 8.16 (s, 1H), 8.08 (s, 1H), 7.90 (s, 1H), 7.65 (d, $J=3.2$ Hz, 1H), 7.15 (d, $J=8.4$ Hz, 2H), 6.83 (s, 1H), 6.65 (s, 1H), 3.89 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 164.2, 157.2, 155.6, 150.5, 146.7, 143.0, 134.5, 131.8, 129.7, 129.2, 125.3, 114.5, 112.9, 108.5, 107.5, 53.4; IR (KBr) $\nu_{\max}$: 1605, 1576, 1516, 1452, 1253, 1177, 1028, 829 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{15}\text{N}_4\text{O}_2$ $[\text{M}+\text{H}]^+$ 319.1195, found 319.1179.

4-(4-Chlorophenyl)-6-(5-methylfuran-2-yl)-2-(1*H*-pyrazol-1-yl)pyrimidine (**3d**): White powder, yield 88%. m.p. 110~111 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.88 (d, $J=2.4$ Hz, 1H), 8.48 (dd, $J=8.8$, 5.6 Hz, 2H), 8.14 (s, 1H), 7.9 (s, 1H), 7.59 (d, $J=3.6$ Hz, 1H), 7.44 (t, $J=8.4$ Hz, 2H), 6.65~6.64 (m, 1H), 6.48 (d, $J=3.2$ Hz, 1H), 2.48 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 159.7, 156.3, 154.5, 147.0, 146.0, 142.6, 135.6, 135.1, 131.5, 129.7, 129.6, 109.4, 107.6, 106.1, 104.6, 14.5; IR (KBr) $\nu_{\max}$: 1582, 1521, 1458, 1395, 1365, 1200, 942, 819, 796 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{14}\text{N}_4\text{OCl}$ $[\text{M}+\text{H}]^+$ 337.0856, found 337.0825.

4-(5-Methylfuran-2-yl)-2-(1*H*-pyrazol-1-yl)-6-(*p*-tolyl)pyrimidine (**3e**): Black powder, yield 83%. m.p. 123~124 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.73 (s, 1H), 8.21 (d, $J=8.4$ Hz, 2H), 8.11 (s, 1H), 7.94 (s, 1H), 7.76 (s, 1H), 7.66 (d, $J=8.0$ Hz, 2H), 7.54 (d, $J=2.4$ Hz, 1H), 6.69 (s, 1H), 2.36 (s, 3H), 2.25 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 159.7, 156.3, 154.5, 147.0, 146.0, 142.6, 135.6, 135.1, 131.5, 129.7, 129.6, 109.4, 107.6, 106.1, 104.6, 24.1, 14.5; IR (KBr) $\nu_{\max}$: 1582, 1522, 1458, 1396, 1365, 1182, 1038, 942, 819, 797 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{17}\text{N}_4\text{O}$ $[\text{M}+\text{H}]^+$ 317.1402, found 317.1475.

4-(4-Chlorophenyl)-2-(1*H*-pyrazol-1-yl)-6-(thiophen-2-yl)pyrimidine (**3f**): Yellow powder, yield 87%. m.p. 217~218 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.89 (d, $J=2.4$ Hz, 1H), 8.48 (dd, $J=8.8$, 5.6 Hz, 2H), 8.16 (s, 1H), 7.91 (s, 1H), 7.60 (d, $J=3.6$ Hz, 1H), 7.45 (t, $J=8.4$ Hz, 2H), 7.27 (s, 1H), 6.65 (s, 1H), 6.48 (s, 1H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 160.4, 153.9, 152.0, 146.2, 136.7, 135.8, 135.3, 132.2, 131.7, 129.9, 129.8, 129.0, 121.5, 109.6, 107.8; IR (KBr) $\nu_{\max}$: 1604, 1576, 1516, 1454, 1390,

1358, 1253, 1178, 1027, 828 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{12}\text{N}_4\text{SCl}$ $[\text{M}+\text{H}]^+$ 339.0471, found 339.0458.

2-(1*H*-pyrazol-1-yl)-4-(thiophen-2-yl)-6-(*p*-tolyl)pyrimidine (**3g**): Black powder, yield 86%. m.p. 235~236 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.88 (s, 1H), 8.36 (d, $J=8.4$ Hz, 2H), 8.26 (s, 1H), 8.09 (s, 1H), 7.91 (s, 1H), 7.81 (d, $J=8.0$ Hz, 2H), 7.69 (s, 1H), 6.84 (s, 1H), 6.65 (s, 1H), 2.42 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 160.7, 154.2, 152.3, 146.5, 142.2, 137.0, 136.8, 132.5, 132.0, 131.0, 129.3, 129.0, 121.8, 110.0, 108.1, 21.7; IR (KBr) $\nu_{\max}$: 1601, 1516, 1455, 1397, 1357, 1255, 1173, 1098, 834, 758 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{15}\text{N}_4\text{S}$ $[\text{M}+\text{H}]^+$ 319.1017, found 319.1054.

4-(4-Methoxyphenyl)-2-(1*H*-pyrazol-1-yl)-6-(thiophen-2-yl)pyrimidine (**3h**): Grey powder, yield 88%. m.p. 125~126 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.83 (s, 1H), 8.34 (d, $J=6.8$ Hz, 2H), 8.11 (s, 1H), 8.03 (s, 1H), 7.85 (s, 1H), 7.60 (s, 1H), 7.10 (d, $J=5.6$ Hz, 2H), 6.78 (s, 1H), 6.60 (s, 1H), 3.84 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 163.4, 161.0, 154.6, 152.7, 146.9, 132.9, 132.4, 132.0, 129.7, 128.9, 122.2, 115.0, 110.3, 108.5, 105.9, 56.9; IR (KBr) $\nu_{\max}$: 2961, 2925, 1600, 1561, 1524, 1487, 1453, 1398, 1261, 1078, 1041, 1010, 950, 820, 757 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{15}\text{N}_4\text{OS}$ $[\text{M}+\text{H}]^+$ 335.0956, found 335.0920.

4-(5-Bromothiophen-2-yl)-6-(4-chlorophenyl)-2-(1*H*-pyrazol-1-yl)pyrimidine (**3i**): Light yellow powder, yield 79%. m.p. 188~189 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.81 (d, $J=2.4$ Hz, 1H), 8.52~8.49 (m, 3H), 8.21 (d, $J=4.0$ Hz, 1H), 7.91 (d, $J=0.8$ Hz, 1H), 7.47~7.42 (m, 3H), 6.64 (dd, $J=2.4$, 1.6 Hz, 1H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 164.9, 160.7, 155.9, 143.8, 143.5, 132.9, 131.2, 130.6, 130.5, 130.2, 118.6, 116.5, 116.3, 109.2, 108.0; IR (KBr) $\nu_{\max}$: 1580, 1522, 1456, 1395, 1367, 1226, 944, 837, 801, 761 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{11}\text{N}_4\text{SClBr}$ $[\text{M}+\text{H}]^+$ 416.9576, found 416.9566.

4-(4-Bromophenyl)-6-(5-bromothiophen-2-yl)-2-(1*H*-pyrazol-1-yl)pyrimidine (**3j**): Light yellow powder, yield 86%. m.p. 179~180 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.87 (d, $J=2.4$ Hz, 1H), 8.47 (dd, $J=8.8$, 5.6 Hz, 2H), 8.14 (s, 1H), 7.90 (s, 1H), 7.58 (d, $J=3.4$ Hz, 1H), 7.44 (t, $J=8.4$ Hz, 2H), 6.65~6.64 (m, 1H), 6.47 (d, $J=3.2$ Hz, 1H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 164.8, 163.5, 160.6, 155.8, 143.7, 143.4, 132.8, 132.3, 131.1, 130.5, 130.4, 130.2, 118.5, 109.1, 107.9; IR (KBr) $\nu_{\max}$: 1585, 1521, 1456, 1446, 1393, 1366, 1011, 943, 824, 768 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{11}\text{Br}_2\text{N}_4\text{S}$ $[\text{M}+\text{H}]^+$ 460.9066, found 460.9072.

4-(5-Bromothiophen-2-yl)-2-(1*H*-pyrazol-1-yl)-6-(*p*-tolyl)pyrimidine (**3k**): White powder, yield 89%. m.p. 182~183 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.79 (d, $J=2.4$ Hz, 1H), 8.44 (s, 1H), 8.33 (d, $J=8.0$ Hz, 2H), 8.22 (d, $J=4.0$ Hz, 1H), 7.91 (s, 1H), 7.46 (d, $J=4.0$ Hz, 1H), 7.40 (d, $J=8.0$ Hz, 2H), 6.64 (t, $J=2.0$ Hz, 1H), 2.42 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 165.9, 160.5, 155.9, 143.7, 142.4, 133.1, 132.9, 131.0, 130.2, 130.0, 127.9, 118.4, 109.2, 107.7, 21.6; IR (KBr) $\nu_{\max}$: 1584,

1523, 1459, 1397, 1365, 1182, 1038, 941, 819, 797, 774, 754 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{14}\text{N}_4\text{SBr}$ $[\text{M}+\text{H}]^+$ 397.0117, found 397.0111.

4-(5-Bromothiophen-2-yl)-6-(4-methoxyphenyl)-2-(1*H*-pyrazol-1-yl)pyrimidine (**3l**): Yellow powder, yield 79%. m.p. 157~158 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 8.78 (d, $J=2.4$ Hz, 1H), 8.49~8.45 (m, 3H), 8.18 (d, $J=4.0$ Hz, 1H), 7.88 (s, 1H), 7.44~7.39 (m, 3H), 6.61 (dd, $J=2.4, 1.6$ Hz, 1H), 3.84 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 165.9, 160.4, 155.8, 143.6, 142.3, 133.0, 132.8, 130.9, 130.1, 129.9, 127.8, 118.3, 109.1, 107.6, 57.1; IR (KBr) ν_{max} : 2916, 1584, 1523, 1459, 1396, 1365, 1200, 1038, 971, 818, 797, 774 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{14}\text{N}_4\text{OSBr}$ $[\text{M}+\text{H}]^+$ 413.0071, found 413.0070.

4.2.4 Synthesis of target compounds **4a**~**4l**

As shown in Scheme 1, a mixture of intermediate **1** (0.005 mol), dimethylformamide (DFM, 10 mL) and K_2CO_3 (0.025 mol) was stirred at room temperature. The solution of guanyl urea sulfate (0.01 mol) and DFM (10 mL) was slowly added to the above mixture. The reaction was kept at 55~60 °C for 5~6 h. The course of the reaction was monitored by TLC. After completion, the reaction mixture was poured into ice water and neutralized with HCl solution (10%) till the solution became neutral and extracted with dichloromethane (DCM, 10 mL $\times$ 3). The organic phase was dried with anhydrous Na_2SO_4 and concentrated under reduced pressure. The obtained residue was purified by silica gel (200~300 mesh) chromatography using petroleum ether (b.p. 60~90 °C)/ethyl acetate ($V:V=1:1$) as the eluting system to give compounds **4a**~**4l**.

1-(4-(4-Chlorophenyl)-6-(furan-2-yl)pyrimidin-2-yl)urea (**4a**): Light yellow powder, yield 80%. m.p. 207~208 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.64 (s, 1H), 8.64 (s, 1H), 8.28 (dd, $J=8.4, 5.6$ Hz, 2H), 8.03 (s, 1H), 7.90 (s, 1H), 7.55 (d, $J=3.2$ Hz, 1H), 7.41 (t, $J=8.8$ Hz, 2H), 7.18 (s, 1H), 6.79 (dd, $J=3.2, 1.6$ Hz, 1H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 164.2, 159.0, 156.1, 155.2, 151.1, 146.9, 130.2, 130.1, 116.5, 116.3, 114.3, 113.4, 104.1; IR (KBr) ν_{max} : 3399, 2928, 1694, 1603, 1573, 1538, 1510, 1394, 1227, 834 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{11}\text{N}_4\text{O}_2\text{ClNa}$ $[\text{M}+\text{Na}]^+$ 337.0464, found 337.0458.

1-(4-(Furan-2-yl)-6-(4-methoxyphenyl)pyrimidin-2-yl)urea (**4b**): Light yellow powder, yield 80%. m.p. 214~215 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.56 (s, 1H), 8.71 (s, 1H), 8.19 (d, $J=8.8$ Hz, 2H), 8.02 (d, $J=0.8$ Hz, 1H), 7.84 (s, 1H), 7.52 (d, $J=3.6$ Hz, 1H), 7.16 (s, 1H), 7.12 (d, $J=8.8$ Hz, 2H), 6.78 (dd, $J=3.6, 1.6$ Hz, 1H), 3.86 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 164.8, 162.5, 159.0, 155.8, 155.3, 151.3, 146.7, 129.3, 128.6, 114.8, 113.9, 113.3, 103.4, 55.9; IR (KBr) ν_{max} : 3380, 1698, 1600, 1538, 1512, 1478, 1399, 1346, 1241, 1176, 835 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{15}\text{N}_4\text{O}_3$ $[\text{M}+\text{Na}]^+$ 333.0959, found 333.0939.

1-(4-(4-Chlorophenyl)-6-(5-methylfuran-2-yl)pyrimidin-2-yl)urea (**4c**): White powder, yield 86%. m.p. 103~

104 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.57 (s, 1H), 8.72 (s, 1H), 8.20 (d, $J=8.8$ Hz, 2H), 8.03 (s, 1H), 7.85 (s, 1H), 7.53 (s, 1H), 7.13 (d, $J=8.8$ Hz, 2H), 6.80 (s, 1H), 2.43 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 164.3, 162.0, 155.3, 150.7, 138.2, 133.4, 132.2, 131.4, 128.9, 127.3, 114.1, 109.4, 100.1, 14.1; IR (KBr) ν_{max} : 3372, 3130, 2922, 2853, 1693, 1585, 1535, 1430, 1396, 1351, 819 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{14}\text{N}_4\text{O}_2\text{Cl}$ $[\text{M}+\text{H}]^+$ 322.0805, found 322.0836.

1-(4-(5-Methylfuran-2-yl)-6-(*p*-tolyl)pyrimidin-2-yl)urea (**4d**): White powder, yield 88%. m.p. 108~109 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.57 (s, 1H), 8.72 (s, 1H), 8.20 (d, $J=8.8$ Hz, 2H), 8.04 (s, 1H), 7.85 (s, 1H), 7.53 (d, $J=3.6$ Hz, 1H), 7.13 (d, $J=8.8$ Hz, 2H), 6.80 (dd, $J=3.6, 1.6$ Hz, 1H), 2.44 (s, 3H), 2.41 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 165.8, 163.5, 160.0, 156.8, 156.3, 152.3, 147.7, 130.3, 129.6, 115.8, 114.9, 114.3, 104.4, 21.2, 14.0; IR (KBr) ν_{max} : 3329, 3133, 2964, 1692, 1584, 1530, 1394, 1367, 1239, 817, 788 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{17}\text{N}_4\text{O}_2$ $[\text{M}+\text{H}]^+$ 309.1351, found 309.1343.

1-(4-(4-Methoxyphenyl)-6-(5-methylfuran-2-yl)pyrimidin-2-yl)urea (**4e**): Grey powder, yield 74%. m.p. 178~179 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.56 (s, 1H), 8.71 (s, 1H), 8.19 (d, $J=8.8$ Hz, 2H), 8.03 (s, 1H), 7.84 (s, 1H), 7.52 (d, $J=3.6$ Hz, 1H), 7.16 (s, 1H), 7.12 (d, $J=8.8$ Hz, 2H), 3.87 (s, 3H), 2.40 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 162.8, 161.3, 155.5, 154.9, 154.8, 153.3, 142.8, 131.4, 128.3, 114.4, 107.9, 106.4, 105.0, 56.3, 14.8; IR (KBr) ν_{max} : 3374, 2922, 2853, 1693, 1585, 1535, 1429, 1396, 1351, 1097, 819 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{17}\text{N}_4\text{O}_3$ $[\text{M}+\text{H}]^+$ 347.1117, found 347.1106.

1-(4-(4-Chlorophenyl)-6-(thiophen-2-yl)pyrimidin-2-yl)urea (**4f**): Green powder, yield 80%. m.p. 134~135 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.64 (s, 1H), 8.49 (s, 1H), 8.33 (dd, $J=8.8, 5.6$ Hz, 2H), 8.29 (dd, $J=3.6, 0.8$ Hz, 1H), 8.15 (s, 1H), 7.88 (dd, $J=5.2, 0.8$ Hz, 1H), 7.43 (t, $J=8.8$ Hz, 2H), 7.31 (dd, $J=5.2, 4.0$ Hz, 2H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 164.2, 160.0, 158.8, 155.2, 142.1, 131.7, 130.2, 130.1, 130.0, 129.5, 116.5, 116.3, 104.4; IR (KBr) ν_{max} : 3327, 1693, 1588, 1536, 1395, 1230, 1096, 828 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{11}\text{N}_4\text{OSClNa}$ $[\text{M}+\text{Na}]^+$ 353.0235, found 353.0232.

1-(4-(Thiophen-2-yl)-6-(*p*-tolyl)pyrimidin-2-yl)urea (**4g**): Light yellow powder, yield 79%. m.p. 201~202 °C; IR (KBr) ν_{max} : 3331, 1692, 1585, 1531, 1394, 1343, 1238, 1099, 818; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.56 (s, 1H), 8.55 (s, 1H), 8.28 (d, $J=3.6$ Hz, 1H), 8.16 (d, $J=8.0$ Hz, 2H), 8.11 (s, 1H), 7.86 (d, $J=4.8$ Hz, 1H), 7.39 (d, $J=8.0$ Hz, 2H), 7.30 (dd, $J=4.8, 4.0$ Hz, 2H), 2.41 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 165.1, 159.9, 158.8, 155.2, 142.2, 141.9, 133.7, 131.5, 130.0, 129.9, 129.5, 127.6, 104.2, 21.5 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{14}\text{N}_4\text{OSNa}$ $[\text{M}+\text{Na}]^+$ 333.0781, found 333.0759.

1-(4-(4-Methoxyphenyl)-6-(thiophen-2-yl)pyrimidin-2-yl)urea (**4h**): White powder, yield 87%. m.p. 98~99 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.62 (s, 1H), 8.47 (s,

1H), 8.31 (dd, $J=8.8$, 5.6 Hz, 2H), 8.26 (dd, $J=3.6$, 0.8 Hz, 1H), 8.13 (s, 1H), 7.85 (dd, $J=5.2$, 0.8 Hz, 1H), 7.41 (t, $J=8.8$ Hz, 2H), 7.29 (dd, $J=5.2$, 4.0 Hz, 2H), 3.88 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 165.3, 163.0, 159.5, 156.3, 155.8, 151.8, 147.3, 129.8, 129.1, 115.3, 114.5, 113.8, 103.9, 56.4; IR (KBr) ν_{max} : 3328, 2964, 1693, 1588, 1536, 1395, 1230, 1097, 828 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{15}\text{N}_4\text{O}_2\text{S}$ $[\text{M} + \text{H}]^+$ 349.0731, found 349.0759.

1-(4-(5-Bromothiophen-2-yl)-6-(4-chlorophenyl)pyrimidin-2-yl)urea (**4i**): Black powder, yield 83%. m.p. 196~197 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 9.67 (s, 1H), 8.35~8.28 (m, 3H), 8.15 (s, 1H), 8.13 (d, $J=4.0$ Hz, 1H), 7.47~7.42 (m, 4H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 164.3, 159.1, 158.8, 155.0, 143.7, 133.0, 130.7, 130.3, 130.2, 117.7, 116.5, 116.3, 104.1; IR (KBr) ν_{max} : 3418, 1605, 1577, 1549, 1510, 1446, 1410, 1364, 1223, 1103, 974, 798 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{11}\text{ON}_4\text{SBrCl}$ $[\text{M} + \text{H}]^+$ 430.9347, found 430.9381.

1-(4-(4-Bromophenyl)-6-(5-bromothiophen-2-yl)pyrimidin-2-yl)urea (**4j**): White powder, yield 84%. m.p. 157~158 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 9.67 (s, 1H), 8.32~8.29 (m, 3H), 8.15 (s, 1H), 8.13 (d, $J=4.0$ Hz, 1H), 7.47~7.42 (m, 4H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 164.7, 159.5, 158.4, 154.8, 141.8, 141.5, 133.2, 131.1, 129.6, 129.4, 129.0, 127.2, 103.8; IR (KBr) ν_{max} : 3419, 1605, 1577, 1548, 1510, 1446, 1410, 1225, 843, 799 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{10}\text{N}_4\text{OSBr}_2\text{Na}$ $[\text{M} + \text{Na}]^+$ 474.8834, found 474.8838.

1-(4-(5-Bromothiophen-2-yl)-6-(*p*-tolyl)pyrimidin-2-yl)urea (**4k**): Grey powder, yield 86%. m.p. 245~246 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 9.60 (s, 1H), 8.41 (s, 1H), 8.14 (d, $J=2.0$ Hz, 2H), 8.11 (d, $J=7.6$ Hz, 2H), 7.44 (d, $J=4.0$ Hz, 1H), 7.39 (d, $J=8.0$ Hz, 2H), 7.22 (s, 1H), 2.41 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 165.2, 159.0, 158.8, 155.1, 143.9, 142.1, 133.5, 132.9, 130.5, 130.0, 127.6, 117.5, 103.8, 21.5; IR (KBr) ν_{max} : 3379, 1693, 1585, 1535, 1396, 1351, 1097, 818 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{13}\text{N}_4\text{OSBrNa}$ $[\text{M} + \text{Na}]^+$ 410.9886, found 410.9888.

1-(4-(5-Bromothiophen-2-yl)-6-(4-methoxyphenyl)pyrimidin-2-yl)urea (**4l**): Yellow powder, yield 81%. m.p. 163~164 $^{\circ}\text{C}$; ^1H NMR (400 MHz, DMSO- d_6) δ : 9.67 (s, 1H), 8.32~8.29 (m, 3H), 8.15 (s, 1H), 8.13 (d, $J=4.0$ Hz, 1H), 7.47~7.42 (m, 4H), 3.83 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 164.2, 159.0, 156.1, 155.2, 151.1, 146.9, 130.1, 130.0, 116.5, 116.3, 114.3, 113.3, 104.1, 55.1; IR (KBr) ν_{max} : 3326, 2963, 1693, 1588, 1536, 1395, 1346, 1262, 1230, 1096, 828 cm^{-1} ; HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{14}\text{N}_4\text{O}_2\text{SBr}$ $[\text{M} + \text{H}]^+$ 426.9836, found 426.9863.

4.3 Antifungal assay

All the synthesized compounds were screened for their antifungal activities against *R. solani*, *F. graminearum*, *H. maydis*, *S. sclerotiorum* and *B. cinerea* by the mycelium growth rate method.^[6] Each tested compound was dissolved in DMSO (1 mL) and diluted with 1% Tween-80

solution into 10 mL to prepare the solution with a concentration of 1000 mg/L. The diluted solution (1 mL) was added to the sterile potato dextrose agar (PDA, 49 mL). Then, mycelion dishes of 5 mm diameter were cut from the culture medium, picked up with a sterilized inoculating needle, and inoculated in the center of the PDA plates (9 cm in diameter). The inoculated plates were incubated at (28 ± 1) $^{\circ}\text{C}$. The compounds were tested at a concentration of 20 mg/L for primary screening. The inhibitory activities of the compounds with excellent effects were further tested at five different concentrations by the two-fold dilution method based on the concentration of 20 mg/L. Fluopyram (purity 99%) was employed as the positive control, whereas the corresponding solution without the tested compound and fluopyram was used as the blank control. Each treatment was performed three times. When the diameter of the colony used as the blank control reached 5~6 cm, the diameter of each colony was measured by the cross method. The corrected inhibitory rates were calculated using the formula $I = [(C - T) / (C - 0.5)] \times 100\%$, where I represents the inhibition rate, C represents the average diameter of mycelia for the blank control and T represents the average diameter of mycelia for the treated PDA. The corresponding EC_{50} values were calculated using the IBM SPSS Statistics 25.0 software.

4.4 Assay of enzyme inhibition activity

4.4.1 Isolation of SDH from *S. sclerotiorum*

Mitochondria from *S. sclerotiorum* were isolated according to a previously reported method.^[37] Cultures were inoculated at 0.05 $\text{OD}_{600 \text{ nm}}$ and grown on a reciprocal shaker (180 r/min, 25 $^{\circ}\text{C}$) for 5 d in Sabouraud maltose broth (SMB) medium. Cells were harvested by vacuum filtration and disrupted in liquid nitrogen using a mortar and pestle. The resultant powder was resuspended to 10% (w/v) in extraction buffer [10 mmol/L KH_2PO_4 , pH 7.2, 10 mmol/L KCl, 10 mmol/L MgCl_2 , 0.5 mol/L sucrose, 0.2 mmol/L ethylene diamine tetraacetic acid (EDTA), 2 mmol/L phenylmethanesulfonyl fluoride (PMSF)]. The extract was clarified by centrifugation (5000 g, 4 $^{\circ}\text{C}$ for 10 min, 2 times), and intact mitochondria were then pelleted at 10000 g for 20 min at 4 $^{\circ}\text{C}$ and resuspended in the same buffer. Mitochondrial suspensions were brought to a concentration of 10 mg/mL and stored at -80 $^{\circ}\text{C}$ until use. SDH activity was found to remain stable for months.

4.4.2 Succinate: ubiquinone/2,6-dichloro-4-[(4-hydroxyphenyl)imino]-2,5-cyclohexadien-1-one (DCPIP) activity inhibition

Mitochondrial suspensions were diluted 1/20 in extraction buffer and preactivated at 30 $^{\circ}\text{C}$ for 30 min in the presence of 10 mmol/L succinate. Succinate: ubiquinone/DCPIP activity inhibition measurements were performed by adding 10 μL of preactivated mitochondria to 200 μL of assay buffer (50 mmol/L phosphate-sodium, pH 7.2, 250 mmol/L sucrose, 3 mmol/L NaN_3 , 10 mmol/L succinate) supplemented with 140 $\mu\text{mol/L}$ dichlorophenolindophenol (DCIP) and 1 mmol/L 2,3-dimethoxy-5-methyl-1,4-

benzoquinone (Q_0). Inhibitor concentrations ranged between 4.4 and 150 $\mu\text{mol/L}$, with uniform $2\times$ dilution factor steps (six inhibitor concentrations + DMSO control). A total of 96 well plates were pre-equilibrated at reaction temperature (30 $^{\circ}\text{C}$) for 10 min before the reactions were started by the addition of 10 μL of preactivated *S. sclerotiorum* mitochondrion suspension. DCPIP reduction was conducted at 30 $^{\circ}\text{C}$ and monitored at 595 nm. Calculated absorbance slopes (OD/h) were used for IC_{50} calculations using the IBM SPSS Statistics 25.0 software.

4.5 Homology modeling and molecular docking

4.5.1 Homology modeling

The NCBI protein database (<https://www.ncbi.nlm.nih.gov/protein>) was used to search the SDH amino acid sequence of *S. sclerotiorum*. The employed hypothetical protein sequences were XP_001591238, XP_001594577, XP_001597467 and XP_001593251, which were reported by Birren. The SDH from porcine heart (PDB ID: 1ZOY) was applied as the template, and the three-dimensional (3D) structure of the SDH was obtained by SWISS-MODEL, a fully automated protein structure homology-modelling server (<http://swissmodel.expasy.org/>).

4.5.2 Molecular docking

Molecular docking studies were performed to investigate the binding modes of compounds **2c**, **3d** and fluopyram to SDH using Autodock vina 1.1.2. The 3D structure of compounds **2c**, **3d** and fluopyram was drawn by ChemBioDraw Ultra 14.0 and ChemBio3D Ultra 14.0 softwares. The AutoDock Tools 1.5.6 package was employed to generate the docking input files. The search grid of SDH was identified as center_x: 86.459, center_y: 65.6, and center_z: 85.537 with dimensions size_x: 15, size_y: 15, and size_z: 15. The value of exhaustiveness was set to 20. For Vina docking, the default parameters were used if it was not mentioned. The best-scoring pose as judged by the Vina docking score was chosen and visually analyzed using PyMol 1.7.6 software (<http://www.pymol.org/>).

Supporting Information ^1H NMR, ^{13}C NMR, IR, HRMS spectra of **2a~2l**, **3a~3l** and **4a~4l**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] Cecchini, G. *Annu. Rev. Biochem.* **2003**, 72, 77.
- [2] Yankovskaya, V.; Horsefield, R.; Törnroth, S.; Luna-Chavez, C.; Miyoshi, H.; Léger, C.; Byrne, B.; Cecchini, G.; Iwata, S. *Science* **2003**, 299, 700.
- [3] Sun, F.; Huo, X.; Zhai, Y.; Wang, A.; Xu, J.; Su, D.; Bartlam, M.; Rao, Z. *Cell* **2005**, 121, 1043.
- [4] Xiong, L.; Shen, Y.; Jiang, L.; Zhu, X.; Yang, W.; Huang, W.; Yang, G. *ACS Symp. Ser.* **2015**, 1204, 175.
- [5] Xiong, L.; Li, H.; Jiang, L.; Ge, J.; Yang, W.; Zhu, X.; Yang, G. *J. Agric. Food Chem.* **2017**, 65, 1021.
- [6] Jin, H.; Zhou, J.; Pu, T.; Zhang, A.; Gao, X.; Tao, K.; Hou, T. *Bioorg. Chem.* **2017**, 73, 76.
- [7] Yao, T.; Xiao, D.; Li, Z.; Cheng, J.; Fang, S.; Du, Y.; Zhao, J.; Dong, X.; Zhu, G. *J. Agric. Food Chem.* **2017**, 65, 5397.
- [8] Yao, T.; Fang, S.; Li, Z.; Xiao, D.; Cheng, J.; Ying, H.; Du, Y.; Zhao, J.; Dong, X. *J. Agric. Food Chem.* **2017**, 65, 3204.
- [9] Zhang, X.; Jin, H.; Deng, Y.; Gao, X.; Li, Y.; Zhao, Y.; Tao, K.; Hou, T. *Chin. Chem. Lett.* **2017**, 28, 1731.
- [10] Wang, H.; Gao, X.; Zhang, X.; Jin, H.; Tao, K.; Hou, T. *Bioorg. Med. Chem. Lett.* **2017**, 27, 90.
- [11] Zhang, A.; Zhou, J.; Tao, K.; Hou, T.; Jin, H. *Bioorg. Med. Chem. Lett.* **2018**, 28, 3042.
- [12] Zhang, A.; Yue, Y.; Yang, Y.; Yang, J.; Tao, K.; Jin, H.; Hou, T. *Pestic. Biochem. Phys.* **2019**, 158, 175.
- [13] Nagarajan, S.; Shanmugavelan, P.; Sathishkumar, M.; Selvi, R.; Ponnuswamy, A.; Harikrishnan, H.; Shanmugaiah, V.; Murugavel, S. *Bioorg. Med. Chem. Lett.* **2014**, 24, 4999.
- [14] Suresh, L.; Kumar, P. S. V.; Chandramouli, G. V. P. *J. Mol. Struct.* **2017**, 1134, 51.
- [15] Misra, A.; Jain, S.; Kishore, D.; Dave, V.; Reddy, K. R.; Sadhu, V.; Dwivedi, J.; Sharma, S. *J. Microbiol. Meth.* **2019**, 163, 105648.
- [16] Hassan, G. S.; El-Messery, S. M.; Abbas, A. *Bioorg. Chem.* **2017**, 74, 41.
- [17] Kumar, B.; Sharma, P.; Gupta, V. P.; Khullar, M.; Singh, S.; Dogra, N.; Kumar, V. *Bioorg. Chem.* **2018**, 78, 130.
- [18] El Sayed, M. T.; Hussein, H. A. R.; Elebiary, N. M.; Hassan, G. S.; Elmessery, S. M.; Elsheakh, A. R.; Nayel, M.; Abdel-Aziz, H. A. *Bioorg. Chem.* **2018**, 78, 312.
- [19] Amin, L. H. T.; Shawer, T. Z.; El-Naggar, A. M.; El-Sehrawi, H. M. A. *Bioorg. Chem.* **2019**, 91, 103159.
- [20] Rashad, A. E.; Hegab, M. I.; Abdel-Megeid, R. E.; Fathalla, N.; Abdel-Megeid, F. M. E. *Eur. J. Med. Chem.* **2009**, 44, 3285.
- [21] Shakya, N.; VEDI, S.; Liang, C.; Yang, F.; Agrawal, B.; Kumar, R. *Bioorg. Med. Chem. Lett.* **2014**, 24, 1407.
- [22] Adhikari, A.; Kalluraya, B.; Sujith, K. V.; Gouthamchandra, K.; Jairam, R.; Mahmood, R.; Sankolli, R. *Eur. J. Med. Chem.* **2012**, 55, 467.
- [23] Somakala, K.; Tariq, S.; Amir, M. *Bioorg. Chem.* **2019**, 87, 550.
- [24] El-Gazzar, A. R. B. A.; Hafez, H. N. *Bioorg. Med. Chem. Lett.* **2009**, 19, 3392.
- [25] Kumar, D.; Khan, S. I.; Tekwani, B. L.; Ponnann, P.; Rawat, D. S. *Eur. J. Med. Chem.* **2015**, 89, 490.
- [26] Boschi, D.; Pippione, A. C.; Sainas, S.; Lolli, M. L. *Eur. J. Med. Chem.* **2019**, 183, 111681.
- [27] Rajesh, S. M.; Kumar, R. S.; Libertsen, L. A.; Perumal, S.; Yogeeswari, P.; Sriram, D. *Bioorg. Med. Chem. Lett.* **2011**, 21, 3012.
- [28] Thanh, N. D.; Hai, D. S.; Ha, N. T. T.; Tung, D. T.; Le, C. T.; Van, H. T. K.; Toan, V. N.; Toan, D. N.; Dang, L. H. *Bioorg. Med. Chem. Lett.* **2019**, 29, 164.
- [29] Sahu, M.; Siddiqui, N.; Iqbal, R.; Sharma, V.; Wakode, S. *Bioorg. Chem.* **2017**, 74, 166.
- [30] Sahu, M.; Siddiqui, N.; Sharma, V.; Wakode, S. *Bioorg. Chem.* **2018**, 77, 56.
- [31] Kumar, N.; Chauhan, A.; Drabu, S. *Biomed. Pharmacother.* **2011**, 65, 375.
- [32] Kahriman, N.; Serdaroglu, V.; Peker, K.; Aydın, A.; Usta, A.; Fandakli, S.; Yayli, N. *Bioorg. Chem.* **2019**, 83, 580.
- [33] Musumma, G.; Ballistreri, F. P. *Org. Magn. Reson.* **1980**, 14, 384.
- [34] Herenciaa, F.; Ferrindiz, M. L.; Ubeda, A.; Dominguez, J. N.; Charris, J. E.; Lobo, G. M.; Alcaraz, M. J. *Bioorg. Med. Chem. Lett.* **1998**, 8, 1169.
- [35] Zheng, C. J.; Jiang, S. M.; Chen, Z. H.; Ye, B. J.; Piao, H. R. *Arch. Pharm. Chem. Life Sci.* **2011**, 344, 689.
- [36] Al-Maqtari, H. M.; Jamalis, J.; Hadda, T. B.; Sankaranarayanan, M.; Chander, S.; Ahmad, N. A.; Sirat, H. M.; Althagafi, I. I.; Mabkhot, Y. N. *Res. Chem. Intermed.* **2017**, 43, 1893.
- [37] Ye, Y.; Ma, L.; Dai, Z.; Xiao, Y.; Zhang, Y.; Li, D.; Wang, J.; Zhu, H. *J. Agric. Food Chem.* **2014**, 62, 4063.

(Cheng, F.)