川芎内生菌 *Fusarium tricinctum* 的四个新吡喃-2-酮类成分曹钰镁^a 旷歧轩^a 巨 凤^a 邓 赞^a 邓 放^a
顾玉诚^b 任 波^{*,a} 郭大乐^{*,a}^(a) 成都中医药大学药学院 成都 611137)^(b) Syngenta Jealott's Hill International Research Centre, Berkshire RG42 6EY, UK)

摘要 采用硅胶、凝胶等柱色谱及反相碳十八制备液相色谱及手性拆分从川芎内生菌 *Fusarium tricinctum* 的糙米培养基中分离得到四个具有相同平面结构的 2-吡喃酮类化合物, 并采用改良的 Snatzke 法对其绝对构型进行鉴定.

关键词 2-吡喃酮; *Fusarium tricinctum*; 川芎; 诱导圆二色谱

Four New 2-Pyrones from *Fusarium tricinctum*, an Endophytic Fungus of *Ligusticum chuanxiong*Cao, Yumei^a Kuang, Qixuan^a Ju, Feng^a Deng, Yun^a Deng, Fang^a
Gu, Yucheng^b Ren, Bo^{*,a} Guo, Dale^{*,a}^(a) School of Pharmacy, Chengdu University of Traditional Chinese Medicine, Chengdu 611137)^(b) Syngenta Jealott's Hill International Research Centre, Berkshire RG42 6EY, UK)

Abstract By silica gel, Sephadex LH-20 column chromatography as well as preparative high performance liquid chromatography (HPLC), four previously undescribed α -pyrones sharing same primary structure were isolated from the brown rice solid medium of *Fusarium tricinctum*. Their structures were elucidated by spectroscopic analyses and their absolute configurations were finally confirmed by a modified Snatzke's method.

Keywords 2-pyrone; *Fusarium tricinctum*; *Ligusticum chuanxiong*; induced circular dichroism

1 Introduction

2-Pyrone, a six-membered cyclic unsaturated ester, is common structural motif in many natural products^[1] belonging to fungi of several genera including *Alternaria*, *Aspergillus*, *Fusarium*, *Penicillium* and *Trichoderma*. Compounds containing 2-pyrone sub-structures such as bufadienolides, fusapyrones, gibepyrone, styrylpyrones and perpyrones also exhibit various activities including cytotoxicity, neurotoxicity as well as phytotoxicity.^[2-6]

In the course of our ongoing search bioactive compounds of endophytes from *Ligusticum chuanxiong*,^[7-8] *Fusarium tricinctum* was obtained from its root. And along with two known compounds, four new 2-pyrone derivatives sharing same planar structure of 6-(2,3-dihydroxybutan-2-yl)-3-methyl-2H-pyran-2-one were firstly isolated as two

pairs of racemes by pre-HPLC and further as four epimers by subsequent chiral separation from the brown rice solid medium of *F. tricinctum*. Because the structures of these compounds contained a diol moiety on the branch (Figure 1) which was confirmed by spectroscopic analyses, induced circular dichroism (ICD) spectra based on a modified Snatzke's method were collected to elucidate their absolute configurations. The details of isolation, structure elucidation as well as bioactivity of these compounds are discussed below.

2 Results and discussion

Compound **1** was obtained as yellow oil. The HRESIMS ion at m/z 199.0948 $[M+H]^+$ indicated its molecular formula should be $C_{10}H_{14}O_4$ (calculated for $C_{10}H_{15}O_4$ $[M+$

* Corresponding authors. E-mail: renbo@cdutcm.edu.cn; guodale@cdutcm.edu.cn

Received June 18, 2020; revised July 19, 2020; published online August 11, 2020.

Dedicated to Professor Youyou Tu, the 2015 Nobel Prize Laureate of Physiology or Medicine, on the Occasion of Her 90th Birthday.

Project supported by the National Natural Science Foundation of China (No. 81973460), the Foundation of Science and Technology Department of Sichuan Province (No. 2018JY0186) and the Xinglin Scholar Research Promotion Project of Chengdu University of Traditional Chinese Medicine (No. BSH2018008). 国家自然科学基金(No. 81973460)、四川省科技厅(No. 2018JY0186)及成都中医药大学杏林学者学科人才科研提升计划(No. BSH2018008)资助项目.

Figure 1 Compounds 1~4 isolated from *F. tricinatum*

H^+], 199.0892). The IR bands indicated the presence of hydroxyl (3414 cm^{-1}) and carbonyl (1701 cm^{-1}) groups. The ^1H NMR, ^{13}C NMR and HSQC spectra displayed signals of three methyl [δ_{H} 2.04 (d, $J=1.2\text{ Hz}$, 3H, H-11), 1.46 (s, 3H, H-10) and 1.10 (d, $J=6.5\text{ Hz}$, 3H, H-9); δ_{C} 22.8 (C-10), 17.6 (C-9), 16.4 (C-11)]; an oxymethine [δ_{H} 3.91 (q, $J=6.5\text{ Hz}$, 1H, H-8); δ_{C} 72.5 (C-8)]; a carbonyl at δ_{C} 165.6 (C-2), two trisubstituted double bonds [δ_{H} 7.34 (dd, $J=6.8$, 1.2 Hz, 1H, H-4) and 6.44 (d, $J=6.8\text{ Hz}$, 1H, H-5); δ_{C} 167.8 (C-6), 142.1 (C-4), 123.6 (C-3), and 103.5 (C-5)] and an oxygen-bearing nonprotonated carbon at δ_{C} 76.3 (C-7).

Detailed analysis of ^1H - ^1H COSY data of **1** revealed the presence of two spin coupling systems (H-4/H-5 and H-8/H-9) as shown in Figure 2. The ^1H detected heteronuclear multiple-bond coherence (HMBC) correlations of H-4/C-2, C-6 and H-5/C-3 confirmed the existence of a 2-pyrone moiety. Furthermore, the HMBC correlations of H-9/C-7, H-10/C-5, C-6, as well as H-11/C-2, C-3, C-4 confirmed the locations of above three methyl groups. The obvious downfield shifts at C-7 and C-8 combined with the molecular formula information indicated that the remaining two hydroxyl groups should be attached to C-7 and C-8 respectively. Thus, the gross structure of **1** was confirmed as shown in Figure 2.

Figure 2 ^1H - ^1H COSY and key HMBC correlations of compounds 1~2

Compound **2** was obtained as yellow oil. It was assigned the same molecular formula $\text{C}_{10}\text{H}_{14}\text{O}_4$ as **1** on the basis of HRESIMS. The NMR spectra of **2** were similar to those of **1**, which indicated that it was an epimer of **1** (Table 1).

HPLC analyses of compounds **1** and **2** with a chiralpak-IC column were carried out to explain their approximate-

Table 1 NMR spectroscopic data of compounds **1** and **2**

Position	1 ^a		2 ^b	
	δ_{C}	δ_{H} (J in Hz)	δ_{C}	δ_{H} (J in Hz)
2	165.6		165.8	
3	123.6		123.5	
4	142.1	7.34, dd (6.8, 1.2)	142.1	7.34, dd (6.8, 1.2)
5	103.5	6.44, d (6.8)	103.5	6.46, d (6.8)
6	167.8		168.2	
7	76.3		76.7	
8	72.5	3.91, q (6.5)	72.3	3.98, q (6.4)
9	17.6	1.10, d (6.5)	17.0	1.18, d (6.4)
10	22.8	1.46, s	22.8	1.37, s
11	16.4	2.04, d (1.2)	16.4	2.04, d (1.2)

^a 400 MHz (^1H NMR), 100 MHz (^{13}C NMR), CD_3OD ; ^b 800 MHz (^1H NMR), 200 MHz (^{13}C NMR), CD_3OD .

to-zero specific rotations and confirmed that there were two pairs of anticipated enantiomers. Subsequent HPLC preparation yielded the enantiomers **1a**, **1b**, **2a** and **2b** successfully.

Due to the presence of a *vic*-diol group in the branch of compounds **1** and **2**, a modified Sneath's method with dimolybdenum tetraacetate [$\text{Mo}_2(\text{OAc})_4$] was applied to determine their absolute configurations. Specifically, the Cotton effects of Mo_2 -complex were determined by the absolute configuration of the diol group and the counter-clockwise O—C—C—O torsional angle (negative sign of torsional angle) in the conformation of chiral Mo_2 -complex displayed negative signs of the bands II and IV (approximately 300 nm) and vice versa.

As Mo_2 -*erythro*-*vic*-diols complex can generate two favored conformations which can lead to opposite optical rotations, Cotton effects produced by Mo_2 -*threo*-*vic*-diols complex should be stronger than that of Mo_2 -*erythro*-*vic*-diols complex when samples were measured at the same concentration.^[9] And the diol moiety of compound **1** was confirmed as *threo*, and of compound **2** should be *erythro* by the means of above accordingly. Furthermore, the absolute configurations of these isomers were assigned by the optical rotation direction indicated by the C-9 and C⁷-O transpositional plane arrangement configuration. According to the favored conformation in the Mo_2 -complexes of **1** and **2** shown in Figure 3, the Mo_2 -complex of **1a** in dimethyl sulfoxide (DMSO) showed a positive Cotton effects, and **1b** showed a negative Cotton effects at approximately 302 nm (band IV), **2a** showed a negative Cotton effects and **2b** showed a positive Cotton effects at approximately 318 nm (band IV), inducing the fact that the absolute configuration of **1a** should be 7*S*,8*S*; of **1b** should be 7*R*,8*R*; of **2a** should be 7*S*,8*R*; and of **2b** should be 7*R*,8*S*.

Compounds **3** and **4** were identified as (–)-4,6'-anhydroxysporidinone^[10] and cyclonerodiol lactone^[11] by spectroscopic analyses and comparing the spectral data with those reported.

A methyl thiazolyl tetrazolium (MTT) method with Taxol as a positive control was also applied to evaluate their cytotoxicity against HTC116, A549 and MV 4-11 cell

Figure 3 ICD spectra from the Mo_2 -complexes of **1** and **2** in DMSO

lines. Unfortunately, none of them showed obvious growth inhibitions at the concentration of 20 $\mu\text{mol/L}$.

3 Conclusions

The rice medium of *F. tricinctum* gave all of four epimers sharing the same plane structure of 6-(2,3-dihydroxybutan-2-yl)-3-methyl-2H-pyran-2-one. Their absolute configurations were elucidated by comprehensive analyses of their spectroscopic spectra. A MTT method was also applied to evaluate their cytotoxicity against HTC116, A549 and MV 4-11 cell lines and none of them showed obvious growth inhibitions at the concentration of 20 $\mu\text{mol/L}$.

4 Experimental section

4.1 General experimental procedures

Optical rotations were determined on a Perkin-Elmer-241 polarimeter (Perkin Elmer, Inc., Waltham, MA, USA) at room temperature. UV spectra were recorded on a Perkin-Elmer Lambda 35 UV-VIS spectrophotometer (Perkin Elmer, Inc., Waltham, MA, USA). IR spectra were measured by a Perkin-Elmer one FT-IR spectrometer (KBr) (Perkin Elmer, Inc., Waltham, MA, USA). The ICD spec-

trum was measured on a Chirascan circular dichroism spectrometer (Applied Photophysics Ltd., Leatherhead, UK). 1D and 2D NMR were carried out on Bruker-Ascend-400 and 800 MHz instruments (Bruker, Bremen, Germany) at 300 K, with TMS as internal standard. The HRESIMS data were obtained using a Q Exactive mass spectrometer (Thermo Fisher Scientific, MA, USA). Preparative HPLC was performed on a NP7000 serials pump (Hanbon Sci. & Tech., Jiangsu, China) equipped with a YMC ODS column (10 mm $\times$ 250 mm, 5 μm , Akzo Nobel Pulp and Performance Chemicals AB, Bohus, Sweden) using a NU3000 serials UV detector (Hanbon Sci. & Tech., Jiangsu, China). A column Chiralpak IC column (4.6 mm $\times$ 250 mm, 5 μm ; Chiral Technologies, West Chester, PA, USA) was applied for chiral resolution. Column chromatographs (CC) were performed with silica gel (200 $\sim$ 300 mesh, Qingdao Haiyang Chemical Co., Qingdao, China) and Sephadex LH-20 (GE-Healthcare Bio-Sciences AB, Uppsala, Sweden).

4.2 Material

L. chuanxiong was collected in Aoping town, suburb of Pengzhou city, Sichuan Province, China, and taxonomically identified by prof. Fei Long of Chengdu University of TCM. A voucher specimen (access number: 2017042411) was deposited at the ministry of education key laboratory of standardization of Chinese herbal medicine, Chengdu university of TCM. *F. tricinctum* was isolated from the root of *L. chuanxiong*, and identified by morphological observation and sequence analyses of the ITS region of rDNA. The strain (GenBank accession No. MN216248) also was deposited at ministry of education key laboratory of standardization of Chinese herbal medicine, Chengdu university of TCM.

4.3 Fermentation and extraction

The fungal strain was inoculated into twenty-four 250 mL Erlenmeyer flasks, each containing 100 mL of liquid medium at 30 $^{\circ}\text{C}$ on an orbital shaker at 120 r/min for 3 d to produce the seed culture. Then large scale fermentation was carried out in 200 tissue culture flasks of 330 mL, each containing 2 g of peptone, 40 g of brown rice, 25 mL of water and 10 mL of seed culture for 40 d at room temperature.

Fermentation broth was ultrasonically extracted with 10 L of methanol for 3 times, and the extract was concentrated to give a crude extract. Then the extract was dispersed with water, and further extracted with EtOAc. The solvent was evaporated under vacuum and finally give an EtOAc extract (65.5 g).

4.4 Isolation of metabolites

The EtOAc extract was then fractionated by silica gel with petroleum ether and acetone mixture from 100 : 0 to 50 : 50 (V/V) to give twenty-seven fractions (Fr.1 $\sim$ Fr.27). The Fr.19 was further subjected to Sephadex LH-20 column chromatography (4 cm $\times$ 180 cm; $\text{CHCl}_3/\text{MeOH}$, V : V = 1 : 1; about 1000 mL), to yield 5 subfractions (Fr.S1 $\sim$

Fr.S5). Fr.S2 (0.38 g) was purified by a preparative HPLC equipped with a YMC ODS-C18 column (10 cm×250 mm, s-5 μ m; 210 nm) to afford **1** (7.9 mg), and the racemic mixture **1** was further purified by HPLC on a chiralpak-IC column (*n*-hexane/isopropanol, *V*: *V*=72:18, 0.8 mL/min) to give **1a** (3.0 mg, *t_R*=5.2 min) and **1b** (3.9 mg, *t_R*=5.6 min). Fr.S3 (0.27 g) was purified by a preparative HPLC equipped with a YMC ODS-C18 column (10 mm×250 mm, s-5 μ m; 210 nm) to afford **2** (7.2mg), and the racemic mixture **2** was further purified by HPLC on a chiralpak-IC column (*n*-hexane/isopropanol, *V*: *V*=72:18, 0.8 mL/min) to give **2a** (2.0 mg, *t_R*=7.2 min) and **2b** (2.8 mg, *t_R*=8.6 min). The Fr.21 (1.21 g) was purified by a preparative HPLC equipped with a YMC ODS-C18 column (10 mm×250 mm, s-5 μ m; 210 nm) to afford **3** (5.8 mg, *t_R*=23 min), and the Fr.22 (1.40 g) was purified by a preparative HPLC equipped with a YMC ODS-C18 column (10 mm×250 mm, s-5 μ m; 210 nm) to afford **4** (3.1 mg, *t_R*=25 min).

6-((2*S*,3*S*)-2,3-Dihydroxybutan-2-yl)-3-methyl-2*H*-pyran-2-one (**1a**): 3.0 mg, light yellow oil. [α]_D²⁰+89.6 (*c* 0.21, MeOH); UV (MeOH) $\lambda_{\max}$ [log ϵ /(L·mol⁻¹·cm⁻¹): 298.7 (4.39), 218.8 (4.05) nm; ¹H NMR (400 MHz, CD₃OD) and ¹³C NMR (100 MHz, CD₃OD) spectroscopic data are listed in Table 1; IR (KBr) ν : 3414.8, 1701.9, 1580.7, 1384.4, 1097.9 cm⁻¹; HRESIMS calcd for C₁₀H₁₅O₄ [M+H⁺] 199.0892, found 199.0948.

6-((2*R*,3*R*)-2,3-Dihydroxybutan-2-yl)-3-methyl-2*H*-pyran-2-one (**1b**): 3.9 mg, Light yellow oil. [α]_D²⁰-89.6 (*c* 0.21, MeOH); UV, IR, NMR, MS and HRESIMS data are the same as those of **1a**.

6-((2*S*,3*R*)-2,3-Dihydroxybutan-2-yl)-3-methyl-2*H*-pyran-2-one (**2a**): 2.0 mg, light yellow oil. [α]_D²⁰-65.8 (*c* 0.28, MeOH); UV (MeOH) $\lambda_{\max}$ [log ϵ /(L·mol⁻¹·cm⁻¹): 298.0 (4.14), 219.1 (3.89) nm; ¹H NMR (800 MHz, CD₃OD) and ¹³C NMR (200 MHz, CD₃OD) spectroscopic data are listed in Table 1; IR (KBr) ν : 3354.9, 1704.9, 1580.5, 1384.1, 1095.3 cm⁻¹; HRESIMS calcd for C₁₀H₁₅O₄ [M+H⁺] 199.0892, found 199.0969.

6-((2*R*,3*S*)-2,3-Dihydroxybutan-2-yl)-3-methyl-2*H*-pyran-2-one (**2b**): 2.8 mg, light yellow oil. [α]_D²⁰+65.8 (*c* 0.28, MeOH); UV, IR, NMR, MS and HRESIMS data are the same as those of **2a**.

4.5 Snatzke's method to determine absolute configurations

Generally, the electrostatic circular dichroism (ECD) spectra of these isomers were first recorded to establish background spectra, and then the ICD spectra of the Mo-complexes of these isomers with same concentration were offered according to the published procedure.^[12]

4.6 Cytotoxic activity assay

HTC116, A549 and MV 4-11 cell lines were cultured in 96-well plates firstly, and then treated with above compounds at specified concentrations for 72 h, followed by stand MTT assays with Taxol as a positive control (which exhibited antiproliferative activity against the HTC116, A549 and MV4-11 cell lines with an IC₅₀ value of (0.013 ± 0.004), (0.014 ± 0.005) and (0.039 ± 0.016) μ mol/L, respectively). The detailed process also were described in our previous work.^[13]

Supporting Information Spectroscopic spectra including HRESIMS, IR, 1D and 2D NMR, and ICD of compounds **1** and **2**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] McGlacken, G. P.; Fairlamb, I. J. *S. Nat. Prod. Rep.* **2005**, *22*, 369.
- [2] Bhat, Z. S.; Rather, M. A.; Maqbool, M.; UL Lah, H.; Yousuf, S. K.; Ahmad, Z. *Biomed. Pharmacother.* **2017**, *91*, 265.
- [3] Lee, J.-S. *Mar. Drugs* **2015**, *13*, 1581.
- [4] Furstner, A. *Angew. Chem., Int. Ed.* **2018**, *57*, 4215.
- [5] Hou, X.-T.; Xu, Y.; Zhu, S.-M.; Zhang, Y.; Guo, L.-D.; Qiu, F.; Che, Y.-S. *RSC Adv.* **2020**, *10*, 15622.
- [6] Guo, D.-L.; Qiu, L.; Feng, D.; He, X.; Li, X.-H.; Cao, Z.-X.; Gu, Y.-C.; Mei, L.; Deng, F.; Deng, Y. *Nat. Prod. Res.* **2020**, *34*, 958.
- [7] Li, X.-H.; Hu, H.-W.; Tan, L.; Wu, W.-L.; Cao, Z.-X.; Gu, Y.-C.; Deng, Y.; Guo, D.-L. *Nat. Prod. Commun.* **2018**, *13*, 1419.
- [8] Cao, Y.-M.; Guo, D.-L.; Jin, M.-Y.; Tan, L.; Yang, T.-L.; Deng, F.; Gu, Y.-C.; Li, X.-H.; Cao, Z.-X.; Deng, Y. *Nat. Prod. Res.* **2020**, DOI: 10.1080/14786419.2020.1712385
- [9] Li, C.-W.; Cui, C.-B. *J. Int. Pharm. Res.* **2015**, *42*, 811 (in Chinese). (李长伟, 崔承彬, 国际药学研究杂志, **2015**, *42*, 811.)
- [10] Zhan, J.-X.; Burns, A. M.; Liu, M.-P. X.; Faeth, S. H.; Gunatilaka, A. A. L. *J. Nat. Prod.* **2007**, *70*, 227.
- [11] Hanson, J. R.; Hitchcock, P. B.; Nyfeler, R. *J. Chem. Soc., Perkin Trans. 1* **1975**, *16*, 1586.
- [12] Liu, Y.-F.; Yue, Y.-F.; Feng, L.-X.; Zhu, H.-J.; Cao, F. *Mar. Drugs* **2019**, *17*, 550.
- [13] Li, S.; Chen, J.-F.; Qin, L.-L.; Li, X.-H.; Cao, Z.-X.; Gu, Y.-C.; Guo, D.-L.; Deng, Y. *J. Asian Nat. Prod. Res.* **2019**, *22*, 138.

(Lu, Y.)