铂催化氧杂二环烯烃与芳基磺酰肼的顺式-立体控制开环反应

王 林^{†,a,c} 杨丽丽^{†,a} 欧云付^a 徐石海^{*,a,b}
林啟福^c 杨定乔^{*,c}

(^a 暨南大学分析测试中心 广州 510632)

(^b 暨南大学化学与材料学院 广州 510632)

(^c 华南师范大学化学学院 广州 510006)

摘要 开发了一种铂催化氧杂二环烯烃与芳基磺酰肼的顺式-立体控制开环反应。该方法显示出高效性和良好的官能团耐受性,在温和条件下能以高达89%的产率得到顺式-2-芳基-1,2-二氢萘-1-醇(**3**)或脱水产物2-芳基萘(**4**)。此外,通过X射线单晶衍射分析确认了产物(1*S**,2*R**)-6,7-二溴-2-(4-甲基苯基)-1,2-二氢萘-1-醇(**3db**)的顺式-1,2-构型。根据实验结果,提出了一种合理的开环反应机理。值得注意的是,在开环反应中,芳基磺酰肼通过释放 N₂ 和 SO₂ 形成碳负离子作为亲核试剂。

关键词 铂催化剂; 开环; 氧杂二环烯烃; 芳基磺酰肼; 亲核试剂

Platinum-Catalyzed *syn*-Stereocontrolled Ring-Opening of Oxabicyclic Alkenes with Arylsulfonyl Hydrazides

Wang, Lin^{†,a,c} Yang, Lili^{†,a} Ou, Yunfu^a Xu, Shihai^{*,a,b}
Lin, Qifu^c Yang, Dingqiao^{*,c}

(^a Analytical and Testing Center, Jinan University, Guangzhou 510632)

(^b College of Chemistry and Materials Science, Jinan University, Guangzhou 510632)

(^c School of Chemistry, South China Normal University, Guangzhou 510006)

Abstract A platinum-catalyzed *syn*-stereocontrolled ring-opening reaction of oxabicyclic alkenes with arylsulfonyl hydrazides was developed. This protocol exhibited high efficiency and good functional group tolerance, affording *cis*-2-aryl-1,2-dihydronaphthalen-1-ols (**3**) or 2-aryl-naphthalenes (**4**) as dehydrated products in good to excellent yields under mild conditions (up to 89%). In addition, the *cis*-1,2-configuration of product (1*S**,2*R**)-6,7-dibromo-2-(*p*-tolyl)-1,2-dihydronaphthalen-1-ol (**3db**) was confirmed by X-ray single crystal diffraction analysis. Based on the results, a plausible mechanism for the ring-opening reaction was proposed. Remarkably, arylsulfonyl hydrazides were used as carboanion nucleophiles in the ring-opening reaction via releasing N₂ and SO₂ *in situ*.

Keywords platinum catalyst; ring-opening; oxabicyclic alkenes; arylsulfonyl hydrazides; nucleophiles

1 Introduction

The transition metal-catalyzed ring-opening reaction of oxa- and aza-bicyclic alkenes is a useful method to construct carbon-carbon and carbon-heteroatom bonds.^[1] It is well known that 1,2-dihydronaphthalene skeleton was found in a wide range of naturally occurring compounds which exhibited a variety of biological activities.^[2] In addition,

this transformation is valuable because multiple stereocenters can be established in a single step. In this regard, numerous metal catalysts, including Ir,^[3] Ni,^[4] Pd,^[5] Cu,^[6] Rh,^[7] Pt,^[8] Ru,^[9] Fe,^[10] and so forth,^[11] have been investigated for the ring-opening reaction of heterobicyclic alkenes with various carbon and heteroatom nucleophiles.

Since the seminal work of Lautens,^[1b,1d] many research

* Corresponding authors. E-mail: txush@jnu.edu.cn; yangdq@snu.edu.cn

Received June 30, 2020; revised July 23, 2020; published online August 18, 2020.

Project supported by the National Natural Science Foundation of China (Nos. 21672084, 21372090) and the Special Fund Project of Department of Natural Resources of Guangdong Province (Guangdong Natural Resources Cooperation) (No. 2020037).

国家自然科学基金(Nos. 21672084, 21372090)和广东省自然资源厅专项资金(粤自然资合) (No. 2020037)资助项目。

[†] 共同第一作者(These authors contributed equally to this work).

groups have been working on the transition metal-catalyzed asymmetric ring-opening (ARO) reactions of oxa(aza)bicyclic alkenes and make some considerable progress. Our group has long-term interest in the ARO reactions of oxa- and aza-bicyclic alkenes with various nucleophiles. The nucleophiles, such as amines,^[3a,3c,3e,12] phenols,^[8a,13] alcohols,^[3f,14] carboxylic acids,^[3b,3d] Grignard reagents^[15] and organic zincs^[16] have been found as feasible nucleophiles used in the ring-opening reaction. In recent years, arylboronic acids and arylsulfonate salts, because of their stability and low toxicity, also have been employed as effective carbanion nucleophiles in ring-opening reactions of oxabicyclic alkenes to afford the corresponding product 2-aryl-1,2-dihydronaphthalen-1-ols in high yields. For instance, our group demonstrated platinum-catalyzed ring-opening of oxabenzonorbornadienes with sodium sulfonates^[17] and arylboronic acids^[18], which generated *cis*-1,2-ring-opening products in high yields.

Arylsulfonyl hydrazides are generally stable in the air. They can be prepared in one step from readily available arylsulfonyl chlorides and hydrazine hydrates,^[19] and increasing attention has recently been paid to the construction of C—C bonds through the extrusion of N₂ and SO₂ *in situ*.^[20] Recently, our group^[21] has reported palladium-catalyzed *syn*-stereocontrolled ring-opening reactions of oxabenzonorbornadienes with arylsulfonyl hydrazides, affording the desired products in good to excellent yields (up to 96%) under an air atmosphere. To the best of our knowledge, there is no literature reported about platinum-catalyzed *syn*-stereocontrolled ring-opening of oxabicyclic alkenes with arylsulfonyl hydrazides till now. Our continuous interest in this type of reactions promotes us to further explore and expand the possibility of this reaction in the presence of platinum catalyst. Herein, we reported platinum-catalyzed ring-opening of oxabicyclic alkenes with arylsulfonyl hydrazides to afford the corresponding products of *cis*-2-aryl-1,2-dihydronaphthalene-1-ols and 2-aryl-naphthalenes in good to excellent yields.

2 Results and discussion

Oxabenzonorbornadiene (**1a**) and phenylsulfonyl hydrazide (**2a**) were initially chosen as model substrates to identify the feasibility of our process. When **2a** reacted with **1a** in 1,2-dichloroethane (DCE) at 80 °C in the presence of PtCl₂ (10 mol%), PPh₃ (20 mol%) and Cu(OAc)₂ (3 equiv.), the desired ring-opening product **3aa** was obtained in 62% yield after 4 h (Table 1, Entry 1). Other two platinum catalysts were then screened, and PtCl₂ was the best one (Entries 1~3). It was worth mentioning that the ring-opening reaction did not take place in the absence of PtCl₂ (Entry 4). When the catalyst loading was 5 mol% and 20 mol%, the ring-opening product **3aa** could be obtained with yields of 39% and 60%, respectively (Entries 5, 6). In order to further improve the yield of **3aa**, series achiral phosphine ligands, including 1,3-bis(diphenylphosphino)propane (DPPP), 1,2-bis(diphenylphosphino)ethane

(DPPE) and 1,1'-bis(diphenylphosphino)ferrocene (DPPF) were examined for the ring-opening reaction (Entries 7~9) (yields up to 62%). Moreover, chiral phosphine ligands, including (*S*)-SEGPHOS, (*R,S*)-PPF-Bu₂, (*R*)-BINAP, (*S*)-DM-SEGPHOS, (4*S*,5*S*)-DIOP and (2*S*,4*S*)-BDPP were also tested. Unfortunately, the product **3aa** was obtained only in moderate yield and low enantioselectivity (up to 53% yield and 43% *ee*) (Entries 10~15). Therefore, among the ligands screened, achiral ligand PPh₃ was the most effective in terms of yield (Entries 1~15). Next, several cupric salts, such as CuO, CuCl₂ and Cu(acac)₂ were also investigated. Unfortunately, all of them gave low yields (Entries 16~18). The effect of solvent was then examined (Entries 1, 19~21). The results indicated that solvents played an important role in the ring-opening reaction. The solvent DCE was the optimal solvent. Furthermore, the impact of reaction temperature on the reaction efficiency was investigated (Entries 22~23). When the reaction temperature was 60 or 100 °C, the product **3aa** was obtained in moderate yield (53% and 48%, respectively). At last, the impact of the amount of Cu(OAc)₂ on the reaction was investigated (Entries 24~26). The results indicated that 1.5 equiv. of Cu(OAc)₂ gave the best yield (82%). It should be noted that in the absence of Cu(OAc)₂, none of the targeted product was observed (Entry 27). Meanwhile, product **3aa** could be obtained in 20% yield in the absence of PPh₃ (Entry 28). Therefore, the optimal reaction conditions were: 10 mol% PtCl₂, 20 mol% PPh₃, 1.5 equiv. of Cu(OAc)₂ in DCE at 80 °C for 4 h.

Next, attention was paid to the ring-opening reaction of **1a** with a variety of arylsulfonyl hydrazides under the optimized reaction conditions (Table 2). In general, all of arylsulfonyl hydrazides **2a**~**2h** as nucleophiles could react smoothly with **1a** and provided the corresponding targeted products **3aa**~**3ah** in moderate to good yields (52%~85%). The substituents attached on arylsulfonyl hydrazides had obvious electronic effect on the ring-opening reaction. Compared with arylsulfonyl hydrazides with strong electron donating groups, the ring opening reaction of arylsulfonylhydrazides with electron withdrawing groups on benzene ring was better. Arylsulfonyl hydrazides **2e**~**2g** bearing electron-withdrawing groups could react smoothly with **1a** and afforded the corresponding products **3ae**~**3ag** in good yields (78%, 85% and 77%, respectively). Arylsulfonyl hydrazides **2b**~**2c** bearing electron-donating groups could react with **1a** and afforded the corresponding products **3ab**~**3ac** in moderate yields (68% and 66%, respectively). 2-Chlorobenzenesulfonyl hydrazide (**2d**) could react with **1a** and afforded ring-opening product **3ad** in 60% yield. In addition to these, when 2,4,6-trimethylbenzenesulfonyl hydrazide (**2h**) as nucleophile was employed in the reaction, the desired product **3ah** was obtained in 52% yield, which probably due to steric hindrance.

The scope of this ring-opening reaction was further expanded to various oxabenzonorbornadienes **1b**~**1f**. The results showed that the reaction of various oxabenzonor-

Table 1 Optimization for the reaction conditions^a

Entry	Pt catalyst (mol%)	Ligand (mol%)	Additive (equiv.)	Solvent	Yield ^b /%	ee ^c /%
1	PtCl ₂ (10)	PPh ₃ (20)	Cu(OAc) ₂ (3)	DCE	62	
2	Pt(COD)Cl ₂ (10)	PPh ₃ (20)	Cu(OAc) ₂ (3)	DCE	58	
3	Pt(acac) ₂ (10)	PPh ₃ (20)	Cu(OAc) ₂ (3)	DCE	36	
4	—	PPh ₃ (20)	Cu(OAc) ₂ (3)	DCE	0	
5	PtCl ₂ (5)	PPh ₃ (10)	Cu(OAc) ₂ (3)	DCE	39	
6	PtCl ₂ (20)	PPh ₃ (40)	Cu(OAc) ₂ (3)	DCE	60	
7	PtCl ₂ (10)	DPPP	Cu(OAc) ₂ (3)	DCE	Trace	
8	PtCl ₂ (10)	DPPE	Cu(OAc) ₂ (3)	DCE	Trace	
9	PtCl ₂ (10)	DPPF	Cu(OAc) ₂ (3)	DCE	41	
10	PtCl ₂ (10)	(S)-SEGPPOS	Cu(OAc) ₂ (3)	DCE	53	36
11	PtCl ₂ (10)	(R,S)-PPF- ^t Bu ₂	Cu(OAc) ₂ (3)	DCE	36	20
12	PtCl ₂ (10)	(R)-BINAP	Cu(OAc) ₂ (3)	DCE	48	10
13	PtCl ₂ (10)	(S)-DM-SEGPPOS	Cu(OAc) ₂ (3)	DCE	50	43
14	PtCl ₂ (10)	(4S,5S)-DIOP	Cu(OAc) ₂ (3)	DCE	10	5
15	PtCl ₂ (10)	(2S,4S)-BDPP	Cu(OAc) ₂ (3)	DCE	8	5
16	PtCl ₂ (10)	PPh ₃	CuO (3)	DCE	Trace	
17	PtCl ₂ (10)	PPh ₃	CuCl ₂ (3)	DCE	23	
18	PtCl ₂ (10)	PPh ₃	Cu(acac) ₂ (3)	DCE	52	
19	PtCl ₂ (10)	PPh ₃	Cu(OAc) ₂ (3)	1,4-Dioxane	46	
20	PtCl ₂ (10)	PPh ₃	Cu(OAc) ₂ (3)	THF	51	
21	PtCl ₂ (10)	PPh ₃	Cu(OAc) ₂ (3)	DMF	37	
22 ^d	PtCl ₂ (10)	PPh ₃	Cu(OAc) ₂ (3)	DCE	53	
23 ^e	PtCl ₂ (10)	PPh ₃	Cu(OAc) ₂ (3)	DCE	48	
24	PtCl ₂ (10)	PPh ₃	Cu(OAc) ₂ (2)	DCE	73	
25	PtCl ₂ (10)	PPh ₃	Cu(OAc) ₂ (1.5)	DCE	82	
26	PtCl ₂ (10)	PPh ₃	Cu(OAc) ₂ (1)	DCE	65	
27	PtCl ₂ (10)	PPh ₃	—	DCE	n.r.	
28	PtCl ₂ (10)	—	Cu(OAc) ₂ (1.5)	DCE	20	

^a The reaction was carried out with Pt catalyst, ligand, additive, **1a** (0.1 mmol, 1.0 equiv.) and **2a** (0.2 mmol, 2.0 equiv.) in the corresponding solvent at 80 °C under an air atmosphere for 4 h. ^b Isolated yield. ^c Determined by HPLC with a Chiralcel OD-H column. ^d 60 °C. ^e 100 °C.

bornadienes with arylsulfonyl hydrazides proceeded smoothly to afford corresponding target product in moderate to excellent yields (Table 3). The ring-opening reactions of oxabenzonorbornadienes **1d**~**1f** with various ar-

ylsulfonyl hydrazides proceeded smoothly to afford the expected products with good to excellent yields (up to 89%). Oxabenzonorbornadienes **1d** and **1f** bearing bromine and fluorine groups showed better reaction efficiency than

Table 2 Platinum-catalyzed ring-opening of oxabenzonorbornadiene **1a** with various arylsulfonyl hydrazides **2a~2h**^a

Entry	Ar	Product	Yield ^b /%
1	C ₆ H ₅	3aa	82
2	4-CH ₃ C ₆ H ₄	3ab	68
3	4-CH ₃ OC ₆ H ₄	3ac	66
4	2-ClC ₆ H ₄	3ad	60
5	4-ClC ₆ H ₄	3ae	78
6	4-FC ₆ H ₄	3af	85
7	4-O ₂ NC ₆ H ₄	3ag	77
8	2,4,6-(CH ₃) ₃ C ₆ H ₂	3ah	52

^a Conditions: **1a** (0.1 mmol, 14.4 mg), **2** (0.2 mmol), Cu(OAc)₂ (1.5 equiv.), PtCl₂ (10 mol%), PPh₃ (20 mol%), DCE (2 mL), 80 °C. ^b Isolated yield.

Table 3 Platinum-catalyzed ring-opening of oxabenzonorbornadienes **1b~1g** with various arylsulfonyl hydrazides^a

Entry	1	Ar	3	Yield ^b /% of 3	4	Yield ^b /% of 4
1	1b	C ₆ H ₅			4ba	68
2	1b	4-CH ₃ C ₆ H ₄			4bb	66
3	1b	4-FC ₆ H ₄			4bf	78
4	1c	C ₆ H ₅			4ca	70
5	1c	4-CH ₃ C ₆ H ₄			4cb	72
6	1d	C ₆ H ₅	3da	83		
7	1d	4-CH ₃ C ₆ H ₄	3db	80		
8	1d	4-FC ₆ H ₄	3df	88		
9	1e	C ₆ H ₅	3ea	71		
10	1e	4-ClC ₆ H ₄	3ee	72		
11	1e	4-O ₂ NC ₆ H ₄	3eg	70		
12	1f	C ₆ H ₅	3fa	87		
13	1f	4-CH ₃ C ₆ H ₄	3fb	80		
14	1f	4-FC ₆ H ₄	3ff	89		
15	1g	C ₆ H ₅	3ga	52		

^a Conditions: **1** (0.1 mmol, 14.4 mg), **2** (0.2 mmol), Cu(OAc)₂ (1.5 equiv.), PtCl₂ (10 mol%), PPh₃ (20 mol%), DCE (2 mL), 80 °C. ^b Isolated yield.

1e bearing methyl group. Besides, bulkier substrate **1g** afforded product **3ga** in 52% yield, indicating that steric hindrance on substrate **1g** had a negative impact on this

ring-opening transformation. Interestingly, the corresponding products 2-arylnaphthalenes **4ba~4cb** could be obtained in good yields (up to 78%) rather than ring-opening products **3** when methoxy substituted oxabenzonorbornadienes **1b~1c** were used to react with various arylsulfonyl hydrazides.

The *cis*-1,2-configuration of compound **3db** was undoubtedly confirmed by X-ray single crystal diffraction analysis (Figure 1). The single crystal was obtained by diffusion method from a mixed solvent of petroleum ether and ethyl acetate.

Figure 1 X-ray single crystal diffraction analysis of compound **3db**

To further evaluate the reaction effect on the scale of preparation, scaled-up reaction was then conducted under the current standard conditions. The desired product **3aa** was obtained in 68% yield (0.302 g), indicating the application of this method in synthetic chemistry (Scheme 1).

Scheme 1 Scaled-up experiment of **3aa**

According to our experimental results and the general mechanism of desulfurization coupling reactions,^[20] a plausible mechanism for this ring-opening reaction is proposed (Scheme 2). Initially, ligand PPh₃ reacts with PtCl₂ to give the active catalyst Pt(PPh₃)₂Cl₂. Then Pt(PPh₃)₂Cl₂ is treated with arylsulfonyl hydrazide **2** to result in complex **A**, which undergoes β-hydride elimination to give sulfonyl diazene **B** and release Pt(PPh₃)₂. Pt(0)(PPh₃)₂ is then oxidized by Cu(OAc)₂ or O₂ from air to regenerate the

Scheme 2 Plausible mechanism

active catalyst $\text{Pt}(\text{PPh}_3)_2\text{Cl}_2[\text{Pt}(\text{PPh}_3)_2\text{X}_2]$. Meanwhile, one of chlorine atom of $\text{Pt}(\text{PPh}_3)_2\text{Cl}_2$ is replaced by sulfonyl diazene **B** to give intermediate **C**. Intermediate **C** releases nitrogen and sulfur dioxide, affording arylplatinum **D**, which then undergoes *exo*-1,2-addition with substrate **1a** to yield intermediate **E**. β -Oxygen atom elimination of intermediate **E** occurs and provides new Pt intermediate **F**. Protonation of the intermediate **F** affords product **3** and a $\text{Pt}(\text{PPh}_3)_2$. Further dehydration of **3** gives 2-arylnaphthalene **4**.

3 Conclusions

In summary, an efficient platinum-catalyzed *syn*-stereocontrolled ring-opening of oxabicyclic alkenes with arylsulfonyl hydrazides via the release of N_2 and SO_2 *in situ* under mild conditions has been developed. This synthetic protocol provides a useful tool for the construction of *cis*-2-aryl-1,2-dihydronaphthalen-1-ols **3** and 2-arylnaphthalenes **4** in good to excellent yields. The different substituents of oxabicyclic alkenes play a crucial role in the control of targeted product. The electron-withdrawing groups in the substrate are advantageous for the ring-opening reaction, and the dehydrated compounds are more readily available when oxabenzonorbornadienes bearing a strong donor group were used. Further studies toward the evaluating of the biological and pharmaceutical activities of these target compounds are going on in our laboratory.

4 Experimental section

4.1 General information

All ^1H NMR and ^{13}C NMR spectra were recorded at 600

and 150 MHz respectively, using $\text{CDCl}_3/\text{CD}_3\text{OD}$ as the solvent. ^{19}F NMR spectra were recorded at 564 MHz at 25 °C in $\text{CDCl}_3/\text{CD}_3\text{OD}$. The chemical shifts of all ^1H NMR and ^{13}C NMR spectra were referenced to the residual signal of CDCl_3 (δ 7.26 for the ^1H NMR spectra and δ 77.00 for the ^{13}C NMR spectra) or CD_3OD (δ 3.31 for the ^1H NMR spectra and δ 49.00 for the ^{13}C NMR spectra). Enantiomeric excesses were determined with a Chiralcel OD-H column eluting with a mixture of hexane and propan-2-ol ($V:V=95:5$, 1.0 mL/min, $\lambda=254$ nm). Infrared spectra (IR) were obtained with thin film samples on a PerkinElmer Spectrum Two spectrometer. HRMS were obtained from mass spectrometer (ESI-TOF). Melting points were recorded on an electrothermal digital melting point apparatus and were uncorrected. The crystal structure determination was carried out by an X-ray diffraction apparatus. Unless otherwise stated, all starting materials, catalysts, and solvents were commercially available and were used as purchased. Anhydrous solvent 1,4-dioxane, tetrahydrofuran (THF) and *N,N*-dimethylformamide (DMF) were used without any pretreatment. Substrates of oxabenzonorbornadienes (**1a**~**1g**) were synthesized according to the known procedures.^[3c] Flash column chromatography was performed using the indicated solvent system on Qingdao-Haiyang silica gel (200~300 mesh).

4.2 General procedure for the synthesis of products **3** and **4**

All experiments were carried out under air. PtCl_2 (2.7 mg, 10 mol%), PPh_3 (5.2 mg, 20 mol%), oxabicyclic alkenes **1** (0.1 mmol), arylsulfonyl hydrazides **2** (2 equiv., 0.2 mmol), $\text{Cu}(\text{OAc})_2$ (27.2 mg, 0.15 mmol) and DCE (2.0

mL) were simultaneously added to a 10 mL round-bottomed flask. The mixture was stirred at 80 °C in oil bath for 4 h. After cooling to room temperature, 10 mL of water was added to the reaction mixture, which then was extracted by ethyl acetate (10 mL × 3). The organic layers were combined and dried with anhydrous MgSO_4 , and then filtered. The filtrate was concentrated under vacuum, and the resulting residue was purified by column chromatography on silica gel (200~300 mesh) using ethyl acetate/petroleum ether or ethyl acetate/hexane to obtain the desired products **3** and **4**.

(1*S**,2*R**)-2-Phenyl-1,2-dihydronaphthalen-1-ol (**3aa**): Colorless oil (18.3 mg, 82%). ^1H NMR (600 MHz, CDCl_3) δ : 7.36~7.27 (m, 6H), 7.28~7.26 (m, 2H), 7.19~7.16 (m, 1H), 6.71 (dd, $J=9.6, 2.0$ Hz, 1H), 6.13 (dd, $J=9.6, 4.0$ Hz, 1H), 4.93 (d, $J=5.9$ Hz, 1H), 3.88 (ddd, $J=6.0, 4.0, 2.1$ Hz, 1H), 1.56 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 137.8, 136.1, 132.7, 129.7, 129.3, 128.7, 128.4, 128.3, 128.0, 127.5, 126.7, 126.4, 71.3, 47.4; IR (film) ν : 3445, 2925, 1495, 1449, 1128, 765, 691 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{16}\text{H}_{11}\text{O}$ [$\text{M}-3\text{H}$] $^-$ 219.0810, found 219.0810.

(1*S**,2*R**)-2-(*p*-Tolyl)-1,2-dihydronaphthalen-1-ol (**3ab**): Colorless oil (16.1 mg, 68%). ^1H NMR (600 MHz, CDCl_3) δ : 7.34 (d, $J=7.2$ Hz, 1H), 7.30~7.22 (m, 2H), 7.17~7.10 (m, 5H), 6.68 (dd, $J=9.6, 1.9$ Hz, 1H), 6.11 (dd, $J=9.6, 4.1$ Hz, 1H), 4.91 (t, $J=6.6$ Hz, 1H), 3.83 (ddd, $J=6.1, 4.2, 2.0$ Hz, 1H), 2.31 (s, 3H), 1.49 (d, $J=7.8$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 137.1, 136.2, 134.3, 132.7, 129.9, 129.4, 129.1, 128.2, 128.1, 127.9, 126.6, 126.3, 71.3, 46.9, 21.0; IR (film) ν : 3437, 2920, 1512, 1451, 1374, 1184, 1071, 786 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{17}\text{H}_{13}\text{O}$ [$\text{M}-3\text{H}$] $^-$ 233.0966, found 233.0970.

(1*S**,2*R**)-2-(4-Methoxyphenyl)-1,2-dihydronaphthalen-1-ol (**3ac**): Colorless oil (16.7 mg, 66%). ^1H NMR (600 MHz, CDCl_3) δ : 7.35 (dd, $J=7.3, 0.5$ Hz, 1H), 7.29~7.23 (m, 2H), 7.18~7.14 (m, 3H), 6.84 (d, $J=8.7$ Hz, 2H), 6.68 (dd, $J=9.6, 1.9$ Hz, 1H), 6.10 (dd, $J=9.6, 4.3$ Hz, 1H), 4.93 (t, $J=5.9$ Hz, 1H), 3.81 (ddd, $J=6.2, 4.3, 2.0$ Hz, 1H), 3.77 (s, 3H), 1.49 (d, $J=6.9$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 159.0, 136.2, 132.7, 130.3, 130.0, 129.1, 128.2, 128.0, 128.0, 126.5, 126.3, 114.1, 71.3, 55.2, 46.4; IR (film) ν : 3448, 2925, 1601, 1512, 1242, 1174, 1025, 796 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{17}\text{H}_{13}\text{O}_2$ [$\text{M}-3\text{H}$] $^-$ 249.0916, found 249.0916.

(1*S**,2*R**)-2-(2-Chlorophenyl)-1,2-dihydronaphthalen-1-ol (**3ad**): Colorless oil (15.4 mg, 60%). ^1H NMR (600 MHz, CDCl_3) δ : 7.45 (dd, $J=7.5, 1.8$ Hz, 1H), 7.42~7.34 (m, 3H), 7.31~7.22 (m, 4H), 6.76 (dd, $J=9.6, 2.7$ Hz, 1H), 6.10~6.04 (m, 1H), 4.91 (d, $J=5.0$ Hz, 1H), 4.51 (dt, $J=5.3, 2.9$ Hz, 1H), 1.64 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 136.9, 135.2, 134.1, 132.1, 131.0, 129.5, 129.0, 128.8, 128.4, 128.3, 128.0, 127.9, 126.9, 126.7, 69.1, 44.0; IR (film) ν : 3414, 2923, 1469, 1456, 1257, 1071, 1031, 808 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{16}\text{H}_{10}\text{ClO}$ [$\text{M}-3\text{H}$] $^-$ 253.0418, found 253.0421.

(1*S**,2*R**)-2-(4-Chlorophenyl)-1,2-dihydronaphthalen-1-ol (**3ae**): White solid (20.0 mg, 78%). m.p. 113.0~114.0 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.33~7.24 (m, 5H), 7.20~7.14 (m, 3H), 6.70 (dd, $J=9.6, 2.0$ Hz, 1H), 6.06 (dd, $J=9.6, 4.1$ Hz, 1H), 4.89 (d, $J=6.0$ Hz, 1H), 3.81 (ddd, $J=6.0, 4.1, 2.0$ Hz, 1H), 1.55 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 136.3, 135.9, 133.1, 132.4, 130.6, 129.2, 128.6, 128.4, 128.4, 128.2, 126.5, 126.4, 71.2, 46.7; IR (film) ν : 3424, 2909, 1484, 1413, 1189, 1089, 1012, 801 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{16}\text{H}_{10}\text{ClO}$ [$\text{M}-3\text{H}$] $^-$ 253.0418, found 253.0422.

(1*S**,2*R**)-2-(4-Fluorophenyl)-1,2-dihydronaphthalen-1-ol (**3af**): White solid (20.4 mg, 85%). m.p. 100.0~101.0 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.34~7.16 (m, 6H), 7.01~6.93 (m, 2H), 6.70 (dd, $J=9.6, 2.0$ Hz, 1H), 6.08 (dd, $J=9.6, 4.1$ Hz, 1H), 4.91 (t, $J=6.8$ Hz, 1H), 3.84 (ddd, $J=6.1, 4.2, 2.0$ Hz, 1H), 1.46 (d, $J=8.1$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 162.2 (d, $J=240.0$ Hz), 136.0, 133.3 (d, $J=3.2$ Hz), 132.5, 130.8 (d, $J=8.0$ Hz), 129.6, 128.4, 128.3, 128.1, 126.5, 126.4, 115.4 (d, $J=21.3$ Hz), 71.2, 46.5; ^{19}F NMR (564 MHz, CDCl_3) δ : -115.4 (s); IR (film) ν : 3420, 2922, 1484, 1456, 1410, 1086, 1012, 832 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{16}\text{H}_{10}\text{FO}$ [$\text{M}-3\text{H}$] $^-$ 237.0713, found 237.0715.

(1*S**,2*R**)-2-(4-Nitrophenyl)-1,2-dihydronaphthalen-1-ol (**3ag**): Pale-yellow solid (20.6 mg, 77%). m.p. 122.3~123.6 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.15 (d, $J=8.7$ Hz, 2H), 7.44 (d, $J=8.6$ Hz, 2H), 7.33 (dd, $J=11.5, 4.1$ Hz, 2H), 7.29~7.27 (m, 1H), 7.21 (d, $J=7.3$ Hz, 1H), 6.77 (dd, $J=9.6, 2.0$ Hz, 1H), 6.08 (dd, $J=9.6, 3.8$ Hz, 1H), 4.93 (s, 1H), 3.95 (t, $J=5.7$ Hz, 1H), 1.60 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 147.2, 146.5, 135.6, 132.1, 130.2, 129.0, 128.8, 128.5, 128.1, 126.8, 126.7, 123.6, 71.2, 47.3; IR (film) ν : 3417, 2922, 1591, 1513, 1339, 1107, 1071, 857 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{16}\text{H}_{10}\text{NO}_3$ [$\text{M}-3\text{H}$] $^-$ 264.0658, found 264.0658.

(1*S**,2*R**)-2-Mesityl-1,2-dihydronaphthalen-1-ol (**3ah**): Colorless oil (13.8 mg, 52%). ^1H NMR (600 MHz, CD_3OD) δ : 7.36~7.31 (m, 2H), 7.26~7.19 (m, 2H), 6.87 (s, 2H), 6.61 (dd, $J=9.6, 3.3$ Hz, 1H), 6.16 (d, $J=9.6$ Hz, 1H), 4.69 (d, $J=4.9$ Hz, 1H), 4.14~4.10 (m, 1H), 2.47 (s, 3H), 2.33 (s, 3H), 2.25 (s, 3H); ^{13}C NMR (150 MHz, CD_3OD) δ : 140.4, 137.7, 137.3, 136.7, 135.1, 134.3, 133.5, 131.8, 130.1, 129.8, 129.6, 128.1, 127.5, 125.2, 71.1, 44.8, 22.0, 21.1, 20.9; IR (film) ν : 3420, 2919, 1520, 1485, 1453, 1079, 812, 761 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{19}\text{H}_{17}\text{O}$ [$\text{M}-3\text{H}$] $^-$ 261.1279, found 261.1277.

1,4-Dimethoxy-6-phenylnaphthalene (**4ba**): Colorless oil (17.9 mg, 68%). ^1H NMR (600 MHz, CDCl_3) δ : 8.44 (d, $J=1.7$ Hz, 1H), 8.27 (d, $J=8.7$ Hz, 1H), 7.78~7.74 (m, 3H), 7.46 (dd, $J=10.6, 4.8$ Hz, 2H), 7.35 (dt, $J=8.4, 1.1$ Hz, 1H), 6.70~6.66 (m, 2H), 3.95 (d, $J=1.9$ Hz, 6H); ^{13}C NMR (150 MHz, CDCl_3) δ : 149.7, 149.4, 141.3, 138.4, 128.7, 127.4, 127.2, 126.6, 125.4, 125.3, 122.4, 119.8, 103.6, 103.3, 55.7; IR (film) ν : 2940, 1599, 1462, 1364, 1272, 1229, 1097, 755 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{18}\text{H}_{17}\text{O}_2$ [$\text{M}+\text{H}$] $^+$ 265.1229, found 265.1225.

1,4-Dimethoxy-6-(*p*-tolyl)naphthalene (**4bb**): White solid (18.4 mg, 66%). m.p. 113.2~114.4 °C; ¹H NMR (600 MHz, CDCl₃) δ: 8.42 (d, *J*=1.6 Hz, 1H), 8.25 (d, *J*=8.7 Hz, 1H), 7.75 (dd, *J*=8.7, 1.9 Hz, 1H), 7.65 (d, *J*=8.1 Hz, 2H), 7.26 (d, *J*=7.8 Hz, 2H), 6.67 (q, *J*=8.3 Hz, 2H), 3.95 (d, *J*=3.0 Hz, 6H), 2.40 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 149.7, 149.5, 138.4, 138.3, 137.0, 129.5, 127.2, 126.6, 125.2, 125.2, 122.4, 119.4, 103.6, 103.1, 55.7, 21.1; IR (film) ν: 2937, 1594, 1520, 1459, 1359, 1267, 1232, 1104, 804 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₉H₁₉O₂ [M+H]⁺ 279.1385, found 279.1380.

6-(4-Fluorophenyl)-1,4-dimethoxynaphthalene (**4bf**): White solid (22.0 mg, 78%). m.p. 110.5~111.6 °C; ¹H NMR (600 MHz, CDCl₃) δ: 8.37 (d, *J*=1.8 Hz, 1H), 8.25 (d, *J*=8.7 Hz, 1H), 7.71~7.68 (m, 3H), 7.14 (t, *J*=8.7 Hz, 2H), 6.70 (dd, *J*=14.6, 7.1 Hz, 2H), 3.96 (d, *J*=1.5 Hz, 6H); ¹³C NMR (150 MHz, CDCl₃) δ: 162.5 (d, *J*=244.9 Hz), 149.2 (d, *J*=27.6 Hz), 137.4, 129.0 (d, *J*=8.06 Hz), 126.4 (d, *J*=33.1 Hz), 125.8, 125.3, 125.1, 122.6, 121.7, 119.7, 115.6 (d, *J*=21.6 Hz), 103.7, 103.2 (d, *J*=28.56 Hz), 55.7; ¹⁹F NMR (564 MHz, CDCl₃) δ: -115.9 (s); IR (film) ν: 2955, 1599, 1510, 1456, 1364, 1268, 1225, 1104, 827 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₈H₁₆FO₂ [M+H]⁺ 283.1134, found 283.1130.

2,3-Dimethoxy-6-phenylnaphthalene (**4ca**): White solid (18.5 mg, 70%). m.p. 121.0~122.0 °C; ¹H NMR (600 MHz, CDCl₃) δ: 7.92 (d, *J*=1.3 Hz, 1H), 7.76 (d, *J*=8.4 Hz, 1H), 7.73~7.71 (m, 2H), 7.62 (dd, *J*=8.4, 1.8 Hz, 1H), 7.48 (t, *J*=7.7 Hz, 2H), 7.37 (t, *J*=7.5 Hz, 1H), 7.19 (s, 1H), 7.15 (s, 1H), 4.03 (d, *J*=1.8 Hz, 6H); ¹³C NMR (150 MHz, CDCl₃) δ: 149.8, 149.5, 141.3, 136.9, 129.4, 128.7, 128.3, 127.2, 127.0, 126.8, 124.3, 123.8, 106.5, 106.0, 55.8; IR (film) ν: 2928, 1601, 1495, 1252, 1235, 1166, 1128, 1003, 854 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₈H₁₇O₂ [M+H]⁺ 265.1229, found 265.1225.

2,3-Dimethoxy-6-(*p*-tolyl)naphthalene (**4cb**): White solid (20.0 mg, 72%). m.p. 136.2~137.3 °C; ¹H NMR (600 MHz, CDCl₃) δ: 7.88 (d, *J*=1.4 Hz, 1H), 7.73 (d, *J*=8.4 Hz, 1H), 7.60~7.58 (m, 3H), 7.27 (d, *J*=7.8 Hz, 2H), 7.15 (d, *J*=22.1 Hz, 2H), 4.01 (d, *J*=2.2 Hz, 6H), 2.41 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 149.7, 149.4, 138.4, 136.9, 136.7, 129.5, 129.4, 128.2, 127.0, 126.7, 124.0, 123.8, 106.5, 106.1, 55.8, 21.1; IR (film) ν: 2948, 1602, 1456, 1360, 1268, 1232, 1097, 812 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₉H₁₉O₂ [M+H]⁺ 279.1385, found 279.1383.

(1*S**,2*R**)-6,7-Dibromo-2-phenyl-1,2-dihydronaphthalen-1-ol (**3da**): Colorless oil (31.5 mg, 83%). ¹H NMR (600 MHz, CD₃OD) δ: 7.59 (s, 1H), 7.50 (s, 1H), 7.26~7.17 (m, 5H), 6.66 (dd, *J*=9.6, 1.7 Hz, 1H), 6.23 (dd, *J*=9.6, 4.6 Hz, 1H), 4.93 (d, *J*=6.4 Hz, 1H), 3.78 (ddd, *J*=6.4, 4.6, 1.8 Hz, 1H); ¹³C NMR (150 MHz, CD₃OD) δ: 139.5, 138.9, 135.6, 134.0, 132.5, 131.8, 130.5, 129.1, 128.0, 127.0, 124.3, 123.9, 71.1, 48.2; IR (film) ν: 3396, 2912, 1655, 1495, 1444, 1257, 1076, 890, 801 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₆H₉Br₂O [M-3H]⁻ 374.9021, found 374.9024.

(1*S**,2*R**)-6,7-Dibromo-2-(*p*-tolyl)-1,2-dihydronaphthalen-1-ol (**3db**) (CCDC 1966855): Colorless oil (31.4 mg, 80%). ¹H NMR (600 MHz, CDCl₃) δ: 7.58 (s, 1H), 7.39 (s, 1H), 7.10 (d, *J*=7.9 Hz, 2H), 7.04 (d, *J*=8.0 Hz, 2H), 6.58 (dd, *J*=9.7, 1.0 Hz, 1H), 6.19 (dd, *J*=9.6, 4.9 Hz, 1H), 4.95 (s, 1H), 3.81~3.77 (m, 1H), 2.31 (s, 3H), 1.52 (s, 1H); ¹³C NMR (150 MHz, CDCl₃) δ: 137.7, 137.3, 133.6, 132.2, 132.0, 131.4, 130.7, 129.6, 129.1, 126.2, 123.9, 123.5, 70.2, 46.2, 21.0; IR (film) ν: 3417, 2915, 1510, 1464, 1260, 1107, 1072, 887 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₇H₁₁Br₂O [M-3H]⁻ 388.9177, found 388.9183.

(1*S**,2*R**)-6,7-Dibromo-2-(4-fluorophenyl)-1,2-dihydronaphthalen-1-ol (**3df**): Colorless oil (35.0 mg, 88%). ¹H NMR (600 MHz, CD₃OD) δ: 7.59 (s, 1H), 7.50 (s, 1H), 7.20~7.14 (m, 2H), 6.98~6.94 (m, 2H), 6.66 (dd, *J*=9.6, 1.5 Hz, 1H), 6.21 (dd, *J*=9.6, 4.7 Hz, 1H), 4.92 (d, *J*=6.4 Hz, 1H), 3.81~3.77 (m, 1H); ¹³C NMR (150 MHz, CD₃OD) δ: 163.5 (d, *J*=241.7 Hz), 139.4, 135.5, 134.6 (d, *J*=2.7 Hz), 133.8, 132.4, 132.2 (d, *J*=8.0 Hz), 131.8, 127.2, 124.3, 124.0, 115.7 (d, *J*=21.6 Hz), 71.0, 47.3; ¹⁹F NMR (564 MHz, CD₃OD) δ: -118.4 (s); IR (film) ν: 3414, 2915, 1596, 1510, 1464, 1219, 1155, 887 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₆H₈Br₂FO [M-3H]⁻ 392.8927, found 392.8932.

(1*S**,2*R**)-6,7-Dimethyl-2-phenyl-1,2-dihydronaphthalen-1-ol (**3ea**): Pale-yellow oil (17.8 mg, 71%). ¹H NMR (600 MHz, CD₃OD) δ: 7.25 (dd, *J*=5.9, 1.2 Hz, 4H), 7.21~7.16 (m, 1H), 7.08 (s, 1H), 6.94 (s, 1H), 6.63 (dd, *J*=9.6, 2.2 Hz, 1H), 6.02 (dd, *J*=9.6, 3.8 Hz, 1H), 4.78 (d, *J*=5.7 Hz, 1H), 3.74 (ddd, *J*=5.9, 3.8, 2.3 Hz, 1H), 2.25 (d, *J*=5.9 Hz, 6H); ¹³C NMR (150 MHz, CD₃OD) δ: 140.9, 137.1, 137.0, 135.4, 132.0, 130.6, 130.2, 129.2, 129.1, 128.9, 128.6, 127.7, 72.3, 49.1, 19.7, 19.5; IR (film) ν: 3399, 2919, 1648, 1487, 1089, 1061, 1018, 887 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₈H₁₅O [M-3H]⁻ 247.1124, found 247.1132.

(1*S**,2*R**)-2-(4-Chlorophenyl)-6,7-dimethyl-1,2-dihydronaphthalen-1-ol (**3ee**): Colorless oil (20.5 mg, 72%). ¹H NMR (600 MHz, CD₃OD) δ: 7.25~7.21 (m, 4H), 7.09 (s, 1H), 6.94 (s, 1H), 6.64 (dd, *J*=9.6, 1.9 Hz, 1H), 5.99 (dd, *J*=9.6, 4.0 Hz, 1H), 4.80 (d, *J*=5.9 Hz, 1H), 3.74 (ddd, *J*=6.0, 4.0, 2.1 Hz, 1H), 2.26 (d, *J*=5.0 Hz, 6H); ¹³C NMR (150 MHz, CD₃OD) δ: 139.6, 137.2, 137.1, 135.2, 133.4, 132.1, 131.8, 129.6, 129.2, 129.1, 129.0, 128.6, 72.0, 48.3, 19.7, 19.5; IR (film) ν: 3414, 2920, 1510, 1489, 1311, 1253, 1094, 887 cm⁻¹; HRMS (ESI-TOF) calcd for C₁₈H₁₄ClO [M-3H]⁻ 281.0734, found 281.0738.

(1*S**,2*R**)-6,7-Dimethyl-2-(4-nitrophenyl)-1,2-dihydronaphthalen-1-ol (**3eg**): Colorless oil (20.7 mg, 70%). ¹H NMR (600 MHz, CDCl₃) δ: 8.17 (d, *J*=8.5 Hz, 2H), 7.46 (d, *J*=8.5 Hz, 2H), 7.10 (s, 1H), 6.99 (s, 1H), 6.71 (d, *J*=8.3 Hz, 1H), 6.00 (dd, *J*=9.5, 3.5 Hz, 1H), 4.86 (d, *J*=5.4 Hz, 1H), 3.93 (s, 1H), 2.27 (d, *J*=9.6 Hz, 6H), 1.59 (s, 1H); ¹³C NMR (150 MHz, CDCl₃) δ: 147.1, 147.0, 137.0, 136.9, 133.0, 130.2, 129.7, 128.9, 128.2, 128.1, 127.0, 123.5, 71.1, 47.6, 19.7, 19.5; IR (film) ν: 3427, 2925,

1591, 1512, 1342, 1306, 1257, 1104, 850 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{18}\text{H}_{14}\text{NO}_3$ $[\text{M}-3\text{H}]^-$ 292.0974, found 292.0982.

(1*S**,2*R**)-6,7-Difluoro-2-phenyl-1,2-dihydronaphthalen-1-ol (**3fa**): Yellow oil (22.4, 87%). ^1H NMR (600 MHz, CDCl_3) δ : 7.33~7.28 (m, 3H), 7.21~7.17 (m, 3H), 6.98 (dd, $J=10.5$, 7.6 Hz, 1H), 6.60 (dd, $J=9.7$, 1.7 Hz, 1H), 6.16 (dd, $J=9.6$, 4.5 Hz, 1H), 4.95 (d, $J=6.4$ Hz, 1H), 3.84 (t, $J=4.8$ Hz, 1H), 1.56 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 150.7 (dd, $J=26.5$, 12.8 Hz), 149.1 (dd, $J=28.5$, 12.9 Hz), 136.1, 133.2 (t, $J=4.1$ Hz), 130.4 (d, $J=2.4$ Hz), 129.5 (dd, $J=6.2$, 4.1 Hz), 129.3, 128.8, 127.8, 126.5, 116.0 (d, $J=18.5$ Hz), 115.0 (d, $J=17.6$ Hz), 70.4, 46.6; ^{19}F NMR (564 MHz, CDCl_3) δ : -138.3 (d, $J=21.5$ Hz), -139.8 (d, $J=20.3$ Hz); IR (film) ν : 3424, 2919, 1591, 1502, 1374, 1306, 1071, 875 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{16}\text{H}_9\text{F}_2\text{O}$ $[\text{M}-3\text{H}]^-$ 255.0622, found 255.0626.

(1*S**,2*R**)-6,7-Difluoro-2-(*p*-tolyl)-1,2-dihydronaphthalen-1-ol (**3fb**): Colorless oil (21.8 mg, 80%). ^1H NMR (600 MHz, CDCl_3) δ : 7.18 (dd, $J=10.5$, 8.0 Hz, 1H), 7.10 (dd, $J=23.9$, 8.0 Hz, 4H), 6.97 (dd, $J=10.5$, 7.7 Hz, 1H), 6.58 (dd, $J=9.7$, 1.3 Hz, 1H), 6.14 (dd, $J=9.7$, 4.6 Hz, 1H), 4.94 (s, 1H), 3.80 (t, $J=5.3$ Hz, 1H), 2.32 (s, 3H), 1.50 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 150.6 (dd, $J=20.2$, 13.2 Hz), 149.0 (dd, $J=22.1$, 12.9 Hz), 137.6, 133.3 (t, $J=4.5$ Hz), 132.7, 130.7, 130.7, 129.6, 129.1, 126.4, 115.9 (d, $J=18.5$ Hz), 114.9 (d, $J=18.0$ Hz), 70.3, 46.1, 21.0; ^{19}F NMR (564 MHz, CDCl_3) δ : -138.4 (d, $J=20.3$ Hz), -140.0 (d, $J=20.3$ Hz); IR (film) ν : 3410, 2923, 1507, 1449, 1378, 1308, 1072, 883 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{17}\text{H}_{11}\text{F}_2\text{O}$ $[\text{M}-3\text{H}]^-$ 269.0779, found 269.0785.

(1*S**,2*R**)-6,7-Difluoro-2-(4-fluorophenyl)-1,2-dihydronaphthalen-1-ol (**3ff**): Colorless oil (24.6 mg, 89%). ^1H NMR (600 MHz, CDCl_3) δ : 7.21~7.12 (m, 3H), 7.04~6.94 (m, 3H), 6.59 (dd, $J=9.6$, 1.3 Hz, 1H), 6.12 (dd, $J=9.6$, 4.6 Hz, 1H), 4.92 (s, 1H), 3.80 (t, $J=5.0$ Hz, 1H), 1.51 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 162.4 (d, $J=244.7$ Hz), 150.7 (dd, $J=19.2$, 12.8 Hz), 149.0 (dd, $J=20.6$, 13.1 Hz), 133.0 (t, $J=4.1$ Hz), 131.8 (d, $J=3.2$ Hz), 130.8 (d, $J=7.8$ Hz), 130.3 (d, $J=2.3$ Hz), 129.4 (dd, $J=6.2$, 4.0 Hz), 126.6, 115.9 (d, $J=18.5$ Hz), 115.6 (d, $J=20.9$ Hz), 115.1 (d, $J=17.9$ Hz), 70.3, 45.8; ^{19}F NMR (564 MHz, CDCl_3) δ : -114.7 (s), -138.0 (d, $J=20.3$ Hz), -139.5 (d, $J=20.3$ Hz); IR (film) ν : 3424, 2922, 1597, 1505, 1313, 1221, 1104, 878 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{16}\text{H}_8\text{F}_3\text{O}$ $[\text{M}-3\text{H}]^-$ 273.0528, found 273.0534.

(1*S**,2*R**)-2-Phenyl-1,2-dihydrotriphenylen-1-ol (**3ga**): White solid (16.8 mg, 52%). m.p. 156~158 $^\circ\text{C}$; ^1H NMR (600 MHz, CDCl_3) δ : 8.80~8.69 (m, 2H), 8.35~8.30 (m, 1H), 8.29~8.24 (m, 1H), 7.73~7.59 (m, 5H), 7.55 (dd, $J=8.0$, 0.9 Hz, 2H), 7.50~7.46 (m, 2H), 7.42~7.37 (m, 1H), 6.49 (ddd, $J=9.8$, 2.3, 1.5 Hz, 1H), 5.44 (s, 1H), 4.07~4.00 (m, 1H), 1.56 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 140.1, 130.8, 130.6, 130.6, 129.8, 129.2, 128.8, 128.7, 128.6, 127.3, 127.2, 126.9, 126.8, 126.5, 126.4,

124.1, 123.9, 123.8, 123.1, 123.1, 67.6, 48.1; IR (film) ν : 3486, 2996, 1525, 1510, 1108, 775, 710 cm^{-1} ; HRMS (ESI-TOF) calcd for $\text{C}_{24}\text{H}_{15}\text{O}$ $[\text{M}-3\text{H}]^-$ 319.1124, found 319.1125.

4.3 Procedure for the scaled-up synthesis of product 3aa

The experiment was carried out under air. PtCl_2 (0.054 g, 10 mol%), PPh_3 (0.105 g, 20 mol%), oxabicyclic alkenes **1a** (0.288 g, 2.0 mmol), arylsulfonyl hydrazides **2a** (0.689 g, 4.0 mmol), $\text{Cu}(\text{OAc})_2$ (0.544 g, 3.0 mmol) and DCE (20.0 mL) were simultaneously added to a 50 mL round-bottomed flask. The mixture was stirred at 80 $^\circ\text{C}$ in oil bath for 4 h. After cooling to room temperature, 30 mL of water was added to the reaction mixture, which then was extracted by ethyl acetate (20 mL $\times$ 3). The organic layers were combined and dried with anhydrous MgSO_4 , and then filtered. The filtrate was concentrated under vacuum, and there sulting residue was purified on silica gel [eluent: $V(\text{ethyl acetate}) : V(\text{petroleum ether}) = 1 : 20$] to obtain product **3aa** as a colorless oil (0.302 g, 68%).

Supporting Information The ^1H NMR, ^{13}C NMR spectra of compounds **3** and **4**, the ^{19}F NMR spectra of compounds **3af**, **3df**, **3fa**, **3fb**, **3ff**, and **4bf**, HPLC chromatograms of compound **3aa**, and the X-ray single crystal diffraction analysis of compound **3db**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Ward, R. S. *Chem. Soc. Rev.* **1990**, 19, 1.
(b) Fagnou, K.; Lautens, M. *Chem. Rev.* **2003**, 103, 169.
(c) Hayashi, T.; Yamasaki, K. *Chem. Rev.* **2003**, 103, 2829.
(d) Lautens, M.; Fagnou, K.; Hiebert, S. *Acc. Chem. Res.* **2003**, 36, 48.
- [2] (a) Lautens, M.; Fagnou, K.; Zunic, V. *Org. Lett.* **2002**, 4, 3465.
(b) McManus, H. A.; Fleming, M. J.; Lautens, M. *Angew. Chem., Int. Ed.* **2007**, 46, 433.
(c) Tsoung, J.; Krämer, K.; Zajdlík, A.; Liébert, C.; Lautens, M. *J. Org. Chem.* **2011**, 76, 9031.
- [3] (a) Yang, D.-Q.; Long, Y.-H.; Wang, H.; Zhang, Z.-M. *Org. Lett.* **2008**, 10, 4723.
(b) Long, Y.-H.; Li, X.-L.; Pan, X.-J.; Ding, D.-D.; Xu, X.; Zuo, X.-J.; Yang, D.-Q.; Wang, S.-Y.; Li, C.-R. *Catal. Lett.* **2014**, 144, 419.
(c) Yang, W.; Luo, R.-S.; Yang, D.-Q. *Molecules* **2015**, 20, 21103.
(d) Zhu, M.-N.; Chen, J.-C.; He, X.-B.; Gu, C.-P.; Xu, J.-B.; Fan, B.-M. *J. Org. Chem.* **2017**, 82, 3167.
(e) Yang, X.; Yang, W.; Yao, Y.-Q.; Deng, Y.-Y.; Zuo, X.-J.; Yang, D.-Q. *J. Org. Chem.* **2018**, 83, 10097.
(f) Yang, X.; Yang, W.; Yao, Y.-Q.; Deng, Y.-Y.; Yang, D.-Q. *Org. Chem. Front.* **2019**, 6, 1151.
- [4] (a) Rayabarapu, D. K.; Chiou, C.-F.; Cheng, C.-H. *Org. Lett.* **2002**, 4, 1679.
(b) Li, L.-P. Rayabarapu, D. K.; Nandi, M.; Cheng, C.-H. *Org. Lett.* **2003**, 5, 1621.
(c) Mannathana, S.; Cheng, C.-H. *Adv. Synth. Catal.* **2014**, 356, 2239.
(d) Shukla, P.; Sharma, A.; Pallavi, B.; Cheng, C.-H. *Tetrahedron* **2015**, 71, 2260.

- [5] (a) Cabrera, S.; Arrayás, R. G.; Carretero, J. C. *Angew. Chem., Int. Ed.* **2004**, *43*, 3944.
 (b) Zhang, T.-K.; Mo, D.-L.; Dai, L.-X.; Hou, X.-L. *Org. Lett.* **2008**, *10*, 3689.
 (c) Liu, S.-S.; Li, S.-F.; Chen, H.-L.; Yang, Q.-J.; Xu, J.-B.; Zhou, Y.-Y.; Yuan, M.-L.; Zeng, W.-M.; Fan, B.-M. *Adv. Synth. Catal.* **2014**, *356*, 2960.
 (d) Li, S.-F.; Xu, J.-B.; Fan, B.-M.; Lu, Z.-W.; Zeng, C.-Y.; Bian, Z.-X.; Zhou, Y.-Y.; Wang, J. *Chem.-Eur. J.* **2015**, *21*, 9003.
 (e) Lu, Z.-W.; Wang, J.; Han, B.-Q.; Li, S.-F.; Zhou, Y.-Y.; Fan, B.-M. *Adv. Synth. Catal.* **2015**, *357*, 3121.
 (f) Zhou, H.; Li, J.-X.; Yang, H.-M.; Xia, C.-G.; Jiang, G.-X. *Org. Lett.* **2015**, *17*, 4628.
 (g) Yang, F.; Chen, J.-C.; Xu, J.-B.; Ma, F.-J.; Zhou, Y.-Y.; Shinde, M. V.; Fan, B.-M. *Org. Lett.* **2016**, *18*, 4832.
- [6] (a) Bertozzi, F.; Pineschi, M.; Macchia, F.; Arnold, L. A.; Minnaard, A. J.; Feringa, B. L. *Org. Lett.* **2002**, *4*, 2703.
 (b) Arrayas, R. G.; Cabrera, S.; Carretero, J. C. *Org. Lett.* **2005**, *7*, 219.
 (c) Zhang, W.; Wang, L.-X.; Shi, W.-J.; Zhou, Q.-L. *J. Org. Chem.* **2005**, *70*, 3734.
 (d) Zhang, W.; Zhu, S.-F.; Qiao, X.-C.; Zhou, Q.-L. *Chem.-Asian J.* **2008**, *3*, 2105.
 (e) Millet, R.; Bernardez, T.; Palais, L.; Alexakis, A. *Tetrahedron Lett.* **2009**, *50*, 3474.
 (f) Millet, R.; Gremaud, L.; Bernardez, T.; Palais, L.; Alexakis, A. *Synthesis* **2009**, *12*, 2101.
 (g) Bos, P. H.; Rudolph, A.; Pérez, M.; Fañanás-Mastral, M.; Harutyunyan, S. R.; Feringa, B. L. *Chem. Commun.* **2012**, *48*, 1748.
- [7] (a) Leong, P.; Lautens, M. J. *Org. Chem.* **2004**, *69*, 2194.
 (b) Cho, Y.-H.; Zunic, V.; Senboku, H.; Olsen, M.; Lautens, M. J. *Am. Chem. Soc.* **2006**, *128*, 6837.
 (c) Nishimura, T.; Tsurumaki, E.; Kawamoto, T.; Guo, X.-X.; Hayashi, T. *Org. Lett.* **2008**, *10*, 4057.
 (d) Tsui, G. C.; Lautens, M. *Angew. Chem., Int. Ed.* **2012**, *51*, 5400.
 (e) Zhu, J.-T.; Tsui, G. C.; Lautens, M. *Angew. Chem., Int. Ed.* **2012**, *51*, 12353.
 (f) Tsui, G. C.; Ninnemann, N. M.; Hosotani, A.; Lautens, M. *Org. Lett.* **2013**, *5*, 1064.
 (g) Zhang, L.; Le, C. M.; Lautens, M. *Angew. Chem., Int. Ed.* **2014**, *53*, 5951.
 (h) Chen, J.-C.; Zou, L.-L.; Zeng, C.-Y.; Zhou, Y.-Y.; Fan, B.-M. *Org. Lett.* **2018**, *20*, 1283.
- [8] (a) Meng, L.; Yang, W.; Pan, X.-J.; Tao, M.; Cheng, G.; Wang, S.-Y.; Zeng, H.-P.; Long, Y.-H.; Yang, D.-Q. *J. Org. Chem.* **2015**, *80*, 2503.
 (b) Pan, X.-J.; Huang, G.-B.; Long, Y.-H.; Zuo, X.-J.; Xu, X.; Gu, F.-L.; Yang, D.-Q. *J. Org. Chem.* **2014**, *79*, 187.
- [9] (a) Villeneuve, K.; Tam, W. J. *Am. Chem. Soc.* **2006**, *128*, 3514.
 (b) Tenaglia, A.; Marc, S.; Giordano, L.; De Raggi, I. *Angew. Chem., Int. Ed.* **2011**, *50*, 9062.
- [10] Ito, S.; Itoh, T.; Nakamura, M. *Angew. Chem., Int. Ed.* **2011**, *50*, 454.
- [11] (a) Sawama, Y.; Kawamoto, K.; Satake, H.; Krause, N.; Kita, Y. *Synlett* **2010**, 2151.
 (b) Huang, Y.; Ma, C.; Lee, Y. X.; Huang, R.-Z.; Zhao, Y. *Angew. Chem., Int. Ed.* **2015**, *54*, 13696.
- [12] (a) Zeng, C.-Y.; Yang, F.; Chen, J.-C.; Wang, J.; Fan, B.-M. *Org. Biomol. Chem.* **2015**, *13*, 8425.
 (b) Xu, X.; Chen, J.-C.; He, Z.-X.; Zhou, Y.-Y.; Fan, B.-M. *Org. Biomol. Chem.* **2016**, *14*, 2480.
 (c) Yang, W.; Cheng, G.; Li, Y.; Zuo, X.-J.; Yang, D.-Q. *Synthesis* **2017**, *49*, 2025.
 (d) Shen, G.-L.; Khan, R.; Lv, H.-P.; Yang, Y.; Zhang, X.; Zhan, Y.; Zhou, Y.-Y.; Fan, B.-M. *Org. Chem. Front.* **2019**, *6*, 1423.
- [13] (a) Cheng, H.-C.; Yang, D.-Q. *J. Org. Chem.* **2012**, *77*, 9756.
 (b) Fang, S.; Liang, X.-L.; Long, Y.-H.; Li, X.-L.; Yang, D.-Q.; Wang, S.-Y.; Li, C.-R. *Organometallics* **2012**, *31*, 3113.
- [14] Yang, D.-Q.; Xia, J.-Y.; Long, Y.-H.; Zeng, Z.-Y.; Zuo, X.-J.; Wang, S.-Y.; Li, C.-R. *Org. Biomol. Chem.* **2013**, *11*, 4871.
- [15] (a) Yang, D.-Q.; Liang, N. *Org. Biomol. Chem.* **2014**, *12*, 2080.
 (b) Chen, G.; Yang, W.; Li, Y.; Yang, D.-Q. *J. Org. Chem.* **2016**, *81*, 7817.
- [16] Deng, Y.-Y.; Yang, W.; Yao, Y.-Q.; Yang, X.; Zuo, X.-J.; Yang, D.-Q. *Org. Biomol. Chem.* **2019**, *17*, 703.
- [17] (a) Li, Y.; Yang, W.; Cheng, G.; Yang, D.-Q. *J. Org. Chem.* **2016**, *81*, 4744.
 (b) Wu, R.-H.; Yang, W.; Chen, W.-K.; Yang, D.-Q. *Org. Chem. Front.* **2017**, *4*, 1921.
- [18] (a) Zeng, Z.-Y.; Yang, D.-Q.; Long, Y.-H.; Pan, X.-J.; Huang, G.-B.; Zuo, X.-J.; Zhou, W. J. *Org. Chem.* **2014**, *79*, 5249.
 (b) Zhang, W.; Chen, J.-C.; Zeng, G.-Z.; Yang, F.; Xu, J.-B.; Sun, W.-Q.; Shinde, M. V.; Fan, B.-M. *J. Org. Chem.* **2017**, *82*, 2641.
- [19] Myers, A. G.; Zheng, B.; Movassaghi, M. *J. Org. Chem.* **1997**, *62*, 7507.
- [20] Aziz, J.; Messaoudi, S.; Alami, M.; Hamze, A. *Org. Biomol. Chem.* **2014**, *12*, 9743.
- [21] Chen, D.-H.; Yao, Y.-Q.; Yang, W.; Lin, Q.-F.; Li, H.-Y.; Wang, L.; Chen, S.-Q.; Tan, Y.; Yang, D.-Q. *J. Org. Chem.* **2019**, *84*, 12481.

(Cheng, F.)