N-氟代双苯磺酰胺作为氧化剂制备二硫化化合物的研究

景丽^{†,a} 金辉^{†,a} 管玫^b 武潇华^a 王乾韬^{*,a} 吴勇^{*,a}

(a 四川大学华西药学院 成都 610041)

(b 四川大学华西医院 成都 610041)

摘要 建立了一种以 N-氟代双苯磺酰胺(NFSI)为氧化剂合成二硫化化合物的新方法。该方法反应温和、操作简便、反应时间短并无过度氧化的副产物(亚砷或砷类化合物)生成。

关键词 N-氟代双苯磺酰胺; 氧化; 二硫化化合物

Metal-Free Oxidation of Thiols by N-Fluorobenzenesulfonimide: A Rapid and Efficient Method to Synthesize Disulfides

Jing, Li^{†,a} Jin, Hui^{†,a} Guan, Mei^b Wu, Xiaohua^a Wang, Qiantao^{*,a} Wu, Yong^{*,a}

(a Key Laboratory of Drug-Targeting of Education Ministry and Department of Medicinal Chemistry, West China School of Pharmacy, Sichuan University, Chengdu 61004)

(b West China Hospital, Sichuan University, Chengdu 610041)

Abstract A simple and rapid method is developed for the oxidation of thiols to the corresponding disulfides using N-fluorobenzenesulfonimide (NFSI) as the oxidant without any contamination by over oxidation. The synthetic protocol for disulfide bond formation proceeded efficiently under external base- and metal-free conditions with the advantages of simple operation, mild reaction conditions as well as short reaction times.

Keywords N-fluorobenzenesulfonimide; oxidation; disulfide

1 Introduction

Disulfide is an important moiety for the construction of various chemically and biologically significant molecules.^[1] Pharmaceutically important examples are biocefallin for cognition disorders and disulfiram for chronic alcoholism. For numerous native proteins, such as oxytocin, vasopressin, insulin and somatostatin, disulfide bridge plays a critical role in the formation of secondary and tertiary structures to maintain the biologically active conformation.^[2] Furthermore, it is also widely applied in the organic reactions, such as the sulfonylation of a variety of organic substrates including indoles,^[3] diazooxindoles,^[4] arenes,^[5] alkynes,^[6] ketones,^[7] pyrroles^[8] and sulfoximines.^[9] Moreover, disulfide is frequently used as vulcanizing agent for rubbers and elastomers.^[10]

Due to all these special characteristics and applications, numerous useful methods have been developed for the

Figure 1 Represented drugs with a disulfide motif

synthesis of disulfides, the most frequently used is the direct oxidation of thiols because a great many of thiols are commercially available or are easily synthesized. Peroxides,^[11] heterogeneous catalysts,^[12] photocatalysts,^[13] nitrates,^[14] halogenating agents,^[15] and other oxidizing reagents^[16] have been utilized for this transformation. Although these reagents can afford disulfides in good yields, they still suffer from several drawbacks. For instance, the use of bases or expensive metal-based oxidizing reagents, special treatment for the activation of reagents and low selectivity because of over oxidation of the final products into sulfoxides/sulfones. Therefore, the development of a

* Corresponding authors. E-mail: qwang@scu.edu.cn; wyong@scu.edu.cn

Received September 22, 2017; revised October 30, 2017; published online November 15, 2017.

Project supported by the National Natural Science Foundation of China (Nos. 81373259, 81573286).

国家自然科学基金(Nos. 81373259, 81573286)资助项目。

† 共同第一作者(These authors contributed equally to this work).

simple and efficient method for disulfides formation under mild reaction conditions remains highly desirable.

N-Fluorobenzenesulfonimide (NFSI) has been widely used as an electrophilic fluorinating agent since 1990s.^[17] Subsequently, it has been successfully applied as a strong oxidant for organometallic intermediates,^[18] an amination reagent,^[19] and a phenylsulfonyl group transfer reagent.^[20] In the course of our study on C—H fluorination, we found that it failed to provide the fluoro product, instead delivered a symmetrical disulfide when the substrate containing a thiol. To the best of our knowledge, oxidative coupling of thiols into disulfides with NFSI is rare in literature. In this context, a great deal of interest has been focused on this phenomenon. We find that NFSI is an ideal choice for the conversion of thiols to the corresponding disulfides at room temperature without any contamination by over oxidation. This new method has the advantages of straightforward workup, mild reaction conditions, high yields of the products, as well as short reaction times.

2 Results and discussion

Our initial experiment was carried out with benzenethiol (**1a**, 0.5 mmol), *N*-fluorobenzenesulfonimide (NFSI) (0.75 mmol) in *N,N*-dimethylformamide (DMF) (1 mL) stirred at room temperature for 1 h. Gratifyingly, the desired product **2a** was obtained in 90% yield under this condition (Table 1, Entry 1). Encouraged by this result, we further optimized the reaction conditions, and the results were showed in Table 1. Firstly, different solvents were examined, and ethyl acetate showed optimal influence on the reaction to give the product **2a** in 93% isolated yield (Entry 7). Other solvents, such as DMF, dichloromethane (DCM), acetone, MeCN, dimethyl sulfoxide (DMSO), EtOH were less effective (Entries 1~6). Then we focused our attention on the equivalent amounts of the oxidant. When the amount of NFSI was decreased from 1.2 equiv. to 1.0 equiv., an identical yield was still obtained, but further decrease resulted in only 86% of desired product (Entry 8). To our delight, shortening the reaction time from 1 h to 10 min did not reduce the efficiency of the reaction (Entry 9). Therefore, the optimized conditions for the oxidation of thiols to disulfides can be summarized as follows: 0.5 mmol of substrate, 0.5 mmol of NFSI in ethyl acetate stirred at room temperature for 10 min.

Once the efficient conditions had been established, we then performed further investigation to examine the application scope of the present protocol. As shown in Table 2, both electron donating (**1b**~**1d**, **1g**~**1j**) and electron withdrawing substituents (**1e**, **1f**, **1k**) on the aromatic ring were provided the corresponding disulfides in good to excellent yields. For example, thiosalicylic acid (**1k**) was smoothly derivatised to give the corresponding disulfide **2k** in 97% yield. The scope of this protocol was further extended to heteroaromatic thiols. As shown from the results, reaction of 2-mercaptobenzoxazole (**1m**) and 2-mercaptobenzothiazole (**1n**) proceeded smoothly to deliver the de-

Table 1 Optimization of reaction conditions^a

Entry	NFSI/equiv.	Solvent	Time/min	Yield ^b /%
1	1.2	DMF	60	90
2	1.2	DCM	60	55
3	1.2	Acetone	60	64
4	1.2	Acetonitrile	60	67
5	1.2	DMSO	60	69
6	1.2	Ethanol	60	81
7	1.0	Ethyl acetate	60	93
8	0.8	Ethyl acetate	60	86
9	1.0	Ethyl acetate	10	93
10	0	Ethyl acetate	60	Trace

^a Reaction conditions: benzenethiol (**1a**, 0.5 mmol), NFSI, solvent (1 mL), r.t.;

^b Isolated yield.

sired products **2m** and **2n** in 82% and 85% yields, respectively. Besides aromatic thiols, aliphatic thiols (**1o**~**1u**) also showed good reactivity in this novel transformation. For instance, 3-mercapto-propanoic acid (**1s**) gave the desired disulfide product **2s** in 96% yield under the optimum conditions. But the yield of di-*tert*-butyl disulfide (**2p**) was only 43%, presumably due to steric reasons. It is noteworthy that all these reactions showed high chemoselectivity in the case of bi-functional thiols, and a wide spectrum of synthetically variable functional group, such as halogen, methyl, amino and carboxyl were well tolerated, offering valuable synthetic handles for further derivatization of the disulfides. Moreover, conversion of the biologically important *L*-cysteine (**1t**) and *N*-acetylcysteine (**1u**) into the corresponding disulfides was also investigated. Although the oxidation in protic solvents required prolonged reaction times, the desired products **2t** and **2u** were obtained in good yields (97% and 84% respectively).

The mechanism for this reaction is still not clear. When radical scavenger, 2,2,6,6-tetramethyl-piperidine-1-oxyl (TEMPO), was added in the reaction of piperidine (**1a**) and NFSI under the optimized reaction conditions, the reaction was not evidently affected, which indicated that this transformation might not be a radical process. Based on the experimental results and the previous literature,^[21] a plausible mechanistic pathway is depicted in Scheme 1 with **1a** as the model substrate. The fluorosulfonium ion **A** is attacked by benzenethiol (**1a**) to deliver intermediate **B**, which undergo elimination to release intermediate **C**. Final nucleophilic addition of imide anion to **C** gives diaryl disulfide **2a** and byproduct **3** (compound **3** was confirmed by ¹H NMR and ¹³C NMR spectra).

3 Conclusions

In summary, we have demonstrated a simple and rapid method for oxidative coupling of thiols into the symmetrical disulfides using NFSI as an efficient oxidant for the first

$$\text{R-SH} \xrightarrow[\text{EtOAc, r.t., 10 min}]{\text{NFSI (1.0 equiv.)}} \text{R-S-S-R}$$

time. This protocol proceeded successfully under base- and metal-free conditions with the advantages of simple operation, high efficiency of transformation, broad functional group tolerance as well as short reaction times (as brief as 10 min). We believe that this method has potential applications in the synthesis of related bioactive compounds and intermediates.

4.1 General considerations

[illegible]

suppliers and used without further purification. All the solvents were treated according to general methods. Thin layer chromatography (TLC) was performed using pre-coated silica gel GF₂₅₄ (0.2 mm), while column chromatography was performed using silica gel (100~200 mesh). The melting point was measured on an YRT-3 melting point apparatus (Shantou Keyi instrument & Equipment Co. Ltd, Shantou, China). The optical rotation was measured on an Anton Paar MCP 200 polarimeter ($k = 589$ nm). NMR spectra were taken on a Varian INOVA 400 or 600 MHz (Varian, Palo Alto, CA, USA) using CDCl₃ or DMSO-*d*₆ as solvent. Chemical shifts were expressed in δ , with tetramethylsilane (TMS) functioning as the internal reference. Infrared (IR) spectra were recorded on a PerkinElmer Spectrum Two instrument. High resolution mass spectroscopy data of the products were collected on a Waters Micromass GCT or a Bruker Apex IV FTMS instrument.

In a flask, NFSI (0.5 mmol) was dissolved in ethyl acetate (1mL), then thiol **1** (0.5 mmol) was added, and the entire mixture was magnetically stirred at room temperature and monitored by TLC. After completion of reaction as indicated by TLC, the crude mixture was concentrated and directly purified by flash column chromatography [*V*(petroleum ether)/*V*(ethyl acetate)=20/1~5/1)] to give the desired product **2** with indicated yields.

1,2-Diphenyldisulfane (**2a**): Yield 93%. White solid, m.p. 60~62 °C (Lit.^[16c] m.p. 59~60 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.52 (d, *J*=4.0 Hz, 4H), 7.39 (t, *J*=4.0 Hz, 4H), 7.30 (t, *J*=4.0 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ : 137.1, 129.2, 127.6, 127.2; IR ν : 3072, 1552, 1385, 894 cm⁻¹; HRMS (EI) calcd for C₁₂H₁₀S₂ 218.0224, found 218.0222.

1,2-Di-*p*-tolylidysulfane (**2b**): Yield 95%. Yellow solid, m.p. 47~48 °C (Lit.^{11b} m.p. 47~48 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.39 (d, *J*=8.0 Hz, 4H), 7.19 (d, *J*=8.0 Hz, 4H), 2.28 (s, 6H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 137.6, 132.8, 130.2, 128.3, 20.8; IR ν: 3019, 2915, 2852, 1487, 1376, 1014, 816, 798 cm⁻¹; HRMS (EI) calcd for C₁₄H₁₄S₂ 246.0537, found 246.0537.

Chin. J. Org. Chem. **2018**, 38, 692~697

White solid, m.p. 86~87 °C (Lit.^[16e] m.p. 88~89 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.44 (d, *J*=8.0 Hz, 4H), 7.32 (d, *J*=8.0 Hz, 4H), 1.29 (s, 18H); ¹³C NMR (100 MHz, CDCl₃) δ: 150.5, 134.0, 127.7, 126.1, 34.5, 31.3; IR *v*: 2958, 2865, 2744, 1484, 1394, 1360, 838 cm⁻¹; HRMS (EI) calcd for C₂₀H₂₆S₂ 330.1476, found 330.1475.

1,2-Bis(4-methoxyphenyl)disulfane (**2d**): Yield 88%. White solid, m.p. 44~45 °C (Lit.²² m.p. 42~44 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.41 (d, *J*=8.0 Hz, 4H), 6.95 (d, *J*=8.0 Hz, 4H), 3.75 (s, 18H); ¹³C NMR (100 MHz, CDCl₃) δ: 159.9, 132.7, 128.4, 114.6, 55.4; IR *v*: 2967, 2840, 1489, 1236, 831 cm⁻¹; HRMS (EI) calcd for C₁₄H₁₄O₂S₂ 278.0435, found 278.0436.

1,2-Bis(4-chlorophenyl)disulfane (**2e**): Yield 89%. Yellow solid, m.p. 68~69 °C (Lit.^{11b} m.p. 71~73 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.40 (d, *J*=8.0 Hz, 4H), 7.28 (d, *J*=8.0 Hz, 4H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 134.7, 132.7, 129.7, 129.4; IR *v*: 3077, 1469; 812, 737 cm⁻¹; HRMS (EI) calcd for C₁₂H₈Cl₂S₂ 285.9444, found 285.9444.

1,2-Bis(4-bromophenyl)disulfane (**2f**): Yield 96%. White solid, m.p. 90~92 °C (Lit.^[16f] m.p. 94~95 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.42 (d, *J*=8.0 Hz, 4H), 7.33 (d, *J*=8.0 Hz, 4H); ¹³C NMR (100 MHz, CDCl₃) δ: 135.8, 132.3, 129.4, 121.6; IR *v*: 3073, 1465, 807, 493 cm⁻¹; HRMS (EI) calcd for C₁₂H₈Br₂S₂ 373.8434, found 373.8434.

N,N'-(Disulfanediybis(4,1-phenylene))diacetamide (**2g**): Yield 91%. White solid, m.p. 192~193 °C (Lit.^[23] m.p. 192~193 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ: 10.09 (s, 2H), 7.66 (d, *J*=8.0 Hz, 2H), 7.59 (d, *J*=8.0 Hz, 2H), 7.43 (d, *J*=8.0 Hz, 2H), 7.37 (d, *J*=8.0 Hz, 2H), 2.04 (s, 6H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 169.0, 139.9, 130.6, 129.9, 120.2, 24.5; IR *v*: 3292, 3111, 1646, 1534, 1395, 812 cm⁻¹; HRMS (EI) calcd for C₁₆H₁₆N₂O₂S₂ 332.0653, found 332.0650.

N,N'-(Disulfanediybis(4,1-phenylene))bis(2,2,2-trifluoroacetamide) (**2h**): Yield 89%. Yellow solid, m.p. 181~183 °C (Lit.^[24] m.p. 181~183 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ: 11.37 (br, 2H), 7.70 (d, *J*=8.0 Hz, 4H), 7.57 (d, *J*=8.0 Hz, 4H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 154.8 (q, *J*=30.0 Hz), 136.4, 132.6, 129.1, 128.3, 126.4, 122.1, 116.9; IR *v*: 3291, 3138, 1700, 1493, 1150, 817 cm⁻¹; HRMS (EI) calcd for C₁₆H₁₀F₆N₂O₂S₂ 440.0088, found 440.0089.

Di-*tert*-butyl(disulfanediybis(4,1-phenylene))dicarbamate (**2i**): Yield 52%. Yellow solid, ¹H NMR (400 MHz, DMSO-*d*₆) δ: 9.53 (s, 2H), 7.46 (d, *J*=8.0 Hz, 4H), 7.38 (d, *J*=8.0 Hz, 4H), 1.47 (s, 18H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 168.1, 139.0, 129.6, 128.9, 119.2, 79.6, 23.5; IR *v*: 3291, 3127, 1699, 1529, 1399, 1360, 1157, 817 cm⁻¹; HRMS (EI) calcd for C₂₂H₂₈N₂O₄S₂ 448.1490, found 448.1488.

N,N'-(Disulfanediybis(4,1-phenylene))bis(4-methylbenzenesulfonamide) (**2j**): Yield 74%. White solid, m.p. 171~173 °C (Lit.^[25] m.p. 171 °C); ¹H NMR (600 MHz, DMSO-*d*₆) δ: 10.43 (s, 2H), 7.64 (d, *J*=7.8 Hz, 4H), 7.32

(m, 8H), 7.07 (d, *J*=7.8 Hz, 4H), 2.33 (s, 6H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 143.9, 138.5, 136.9, 131.0, 130.7, 130.2, 127.2, 120.7, 21.4; IR *v*: 3271, 2921, 1511, 1334, 1187, 1175, 921, 810, 767 cm⁻¹; HRMS (EI) calcd for C₂₆H₂₄N₂O₄S₄ 556.0619, found 556.0619.

2,2'-Disulfanediyldibenzoic acid (**2k**): Yield 97%. White solid, m.p. 282~283 °C (Lit.^[26] m.p. 283~285 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ: 13.56 (br, 2H), 8.02 (dd, *J*=4.0, 0.8 Hz, 2H), 7.62 (d, *J*=4.0 Hz, 2H), 7.56 (dt, *J*=4.0, 0.8 Hz, 2H), 7.34 (t, *J*=4.0 Hz, 2H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 139.2, 133.5, 131.9, 128.3, 126.2, 125.3; IR *v*: 3091, 2814, 1676, 1561; 1415, 1310, 1258, 737, 696 cm⁻¹; HRMS (EI) calcd for C₁₄H₁₀O₄S₂ 306.0021, found 306.0022.

1,2-Di(naphthalen-2-yl)disulfane (**2l**): Yield 99%. White solid, m.p. 134~135 °C (Lit.^[16f] m.p. 135~138 °C); ¹H NMR (400 MHz, CDCl₃) δ: 7.98 (s, 2H), 7.78 (d, *J*=8.0 Hz, 4H), 7.13 (d, *J*=8.0 Hz, 2H), 7.62 (d, *J*=8.0 Hz, 2H), 7.45 (s, 4H); ¹³C NMR (100 MHz, CDCl₃) δ: 134.2, 133.4, 132.5, 128.9, 127.7, 127.4, 126.7, 126.5, 126.2, 125.6; IR *v*: 3050, 2918, 1584, 868, 735 cm⁻¹; HRMS (EI) calcd for C₂₀H₁₄S₂ 318.0537, found 318.0539.

1,2-Bis(benzo[*d*]oxazol-2-yl)disulfane (**2m**): Yield 82%. White solid, m.p. 110~111 °C (Lit.^[27] m.p. 111~113 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ: 8.07 (d, *J*=8.0 Hz, 2H), 7.94 (d, *J*=8.0 Hz, 2H), 7.53 (t, *J*=8.0 Hz, 2H), 7.44 (t, *J*=8.0 Hz, 2H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 159.7, 152.9, 142.5, 129.6, 128.9, 119.2, 111.2; IR *v*: 3214, 1657, 1458, 1234, 800, 739 cm⁻¹; HRMS (EI) calcd for C₁₄H₈N₂O₂S₂ 300.0027, found 300.0027.

1,2-Bis(benzo[*d*]thiazol-2-yl)disulfane (**2n**): Yield 85%. White solid, m.p. 181~182 °C (Lit.^[28] m.p. 177~179 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ: 8.07 (d, *J*=8.0 Hz, 2H), 7.94 (d, *J*=8.0 Hz, 2H), 7.53 (t, *J*=8.0 Hz, 2H), 7.44 (t, *J*=8.0 Hz, 2H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 160.4, 151.7, 135.2, 126.5, 124.3, 121.7, 121.0; IR *v*: 3109, 1651, 1458, 810, 797, 729 cm⁻¹; HRMS (EI) calcd for C₁₄H₈N₂S₄ 331.9570, found 331.9568.

1,2-Dibenzylidene disulfane (**2o**)^[16g]: Yield 82%. Colorless oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.28 (m, 10H), 3.60 (s, 4H); ¹³C NMR (100 MHz, CDCl₃) δ: 137.3, 129.4, 128.5, 127.4, 63.2; IR *v*: 3027, 2964, 2910, 1493, 1452, 757, 692 cm⁻¹; HRMS (EI) calcd for C₂₀H₁₄S₂ 318.0537, found 318.0539.

1,2-Di-*tert*-butylidene disulfane (**2p**)^[16g]: Yield 43%. Colorless oil. ¹H NMR (400 MHz, CDCl₃) δ: 1.41 (s, 18H); ¹³C NMR (100 MHz, CDCl₃) δ: 45.7, 30.5; IR *v*: 2960, 2922, 2895, 1455, 1388, 1360, 1160 cm⁻¹; HRMS (EI) calcd for C₂₀H₁₄S₂ 318.0537, found 318.0539.

1,2-Dioctylidene disulfane (**2q**): Yield 81%. White solid, m.p. 183~185 °C (Lit.^[29] m.p. 185~187 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ: 2.68 (t, *J*=7.2 Hz, 4H), 2.54 (m, 4H), 1.31 (m, 20H), 0.79 (m, 6H); ¹³C NMR (100 MHz, CDCl₃) δ: 39.2, 31.8, 29.3, 29.2, 29.2, 28.6, 22.7, 14.1; IR *v*: 2957, 2909, 2824, 1457, 1358 cm⁻¹; HRMS (EI) calcd for C₁₆H₃₄S₂ 290.2102, found 290.2103.

2,2'-Disulfanediybis(ethan-1-amine) (**2r**): Yield 95%.

Yellow solid, m.p. 133~135 °C; ^1H NMR (400 MHz, DMSO- d_6) δ : 8.26 (br, 4H), 3.09 (t, $J=4.0$ Hz, 4H), 3.01 (t, $J=4.0$ Hz, 4H); ^{13}C NMR (100 MHz, CDCl_3) δ : 37.9, 34.0; IR ν : 3112, 2952, 1601, 1402, 1243, 1096, 813 cm^{-1} ; HRMS (EI) calcd for $\text{C}_4\text{H}_{12}\text{N}_2\text{S}_2$ 152.0442, found 152.0442.

3,3'-Disulfanediylpropionic acid (**2s**): Yield 96%. White solid, m.p. 152~154 °C (Lit.^[30] m.p. 156 °C); ^1H NMR (400 MHz, DMSO- d_6) δ : 12.34 (br, 2H), 2.88 (t, $J=8.0$ Hz, 4H), 2.61 (t, $J=8.0$ Hz, 4H); ^{13}C NMR (100 MHz, CDCl_3) δ : 172.8, 33.7, 33.1; IR ν : 2968, 2916, 1620, 1483, 1404, 1296, 1193 cm^{-1} ; HRMS (EI) calcd for $\text{C}_6\text{H}_{10}\text{O}_4\text{S}_2$ 210.0021, found 210.0021.

(2*R*,2'*R*)-3,3'-Disulfanediylbis(2-aminopropanoic acid) (**2t**): Yield 97%. White solid, m.p. 202~204 °C (dec.); $[\alpha]_{\text{D}}^{20}$ -197.53 (c 1.00, 1 mol·L $^{-1}$ HCl) [Lit.^[31] m.p. 216~220 °C (dec.), $[\alpha]_{\text{D}}^{20}$ -205.17 (c 1.02, 1 mol·L $^{-1}$ HCl)]; ^1H NMR (400 MHz, DMSO- d_6) δ : 12.19 (br, 2H), 8.57 (br, 4H), 4.17 (t, $J=8.0$ Hz, 2H), 3.42 (d, $J=8.0$ Hz, 2H), 3.20 (d, $J=8.0$ Hz, 2H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 169.5, 51.4, 37.5; IR ν : 2880, 2633, 1918, 1726, 1590, 1431, 1362, 1090, 1133, 1082, 955, 868, 756, 541 cm^{-1} ; HRMS (EI) calcd for $\text{C}_6\text{H}_{12}\text{N}_2\text{O}_4\text{S}_2$ 240.0238, found 240.0235.

(2*R*,2'*R*)-3,3'-Disulfanediylbis(2-acetamidopropanoic acid) (**2u**): Yield 67%. White solid, m.p. 270~272 °C (Lit.^[13d,32] m.p. 273~275 °C); $[\alpha]_{\text{D}}^{24}$ -95.98 (c 1.00, D_2O) [Lit.^[13d] $[\alpha]_{\text{D}}^{24}$ -102.95 (c 1.06, D_2O)]; ^1H NMR (600 MHz, DMSO- d_6) δ : 12.91 (br, 2H), 8.30 (d, $J=4.0$ Hz, 4H), 4.46 (m, 2H), 3.12 (dd, $J=8.0$, 4.0 Hz, 2H), 2.90 (dd, $J=8.0$, 4.0 Hz, 2H), 1.85 (s, 6H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 172.5, 169.9, 51.7, 39.4, 22.8; IR ν : 3297, 2941, 2886, 1739, 1646, 1431, 1395, 957, 775 cm^{-1} ; HRMS (EI) calcd for $\text{C}_{10}\text{H}_{16}\text{N}_2\text{O}_6\text{S}_2$ 324.0450, found 324.0450.

Supporting Information The ^1H NMR and ^{13}C NMR spectra of products. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Kanda, Y.; Fukuyama, T. *J. Am. Chem. Soc.* **1993**, *115*, 8451.
(b) Schmidt, B.; Lindman, S.; Tong, W.; Lindeberg, G.; Gogoll, A.; Lai, Z.; Thornwall, M.; Synnergren, B.; Nilsson, A.; Welch, C. J.; Sohtell, M.; Westerlund, C.; Nyberg, F.; Karlen, A.; Hallberg, A. *J. Med. Chem.* **1997**, *40*, 903.
(c) Tam-Chang, S. W.; Stehouwer, J. S.; Hao, J. *J. Org. Chem.* **1999**, *64*, 334.
- [2] (a) Wu, L. F.; Meng, J.; Shen, Q.; Zhang, Y.; Pan, S.; Chen, Z.; Zhu, L.-Q.; Lu, Y.; Huang, Y.; Zhang, G. *Nat. Commun.* **2017**, *8*, 15904.
(b) Peters, S. T.; Bowen, M. T.; Bohrer, K.; McGregor, I. S.; Neumann, I. D. *Addict. Biol.* **2017**, *22*, 702.
(c) Ullian, M. E.; Beck, C. N.; Walker, L. P. *Am. J. Hypertens.* **2009**, *22*, 221.
(d) Pena, A.; Murat, B.; Trueba, M.; Ventura, M. A.; Wo, N. C.; Szeto, H. H.; Cheng, L. L.; Stoev, S.; Guillon, G.; Manning, M. *J. Med. Chem.* **2007**, *50*, 835.
(e) Anoop, A.; Ranganathan, S.; Dhaked, B. D.; Jha, N. N.; Pratihari, S.; Ghosh, S.; Sahay, S.; Kumar, S.; Das, S.; Kombrabail, M.; Agarwal, K.; Jacob, R. S.; Singru, P.; Bhaumik, P.; Padinhateeri, R.; Kumar, A.; Maji, S. K. *J. Biol. Chem.* **2014**, *289*, 16884.
- [3] (a) Ge, W. L.; Wei, Y. Y. *Green Chem.* **2012**, *14*, 2066.
(b) Sang, P.; Chen, Z. K.; Zou, J. W.; Zhang, Y. H. *Green Chem.* **2013**, *15*, 2096.
- [4] Khanal, H. D.; Kimb, S. H.; Lee, Y. R. *RSC Adv.* **2016**, *6*, 58501.
- [5] (a) Yang, Y. X.; Hou, W.; Qin, L. H.; Du, J. J.; Feng, H. J.; Zhou, B.; Li, Y. C. *Chem.-Eur. J.* **2014**, *20*, 416.
(b) Yan, G. B.; Borah, A. J.; Wang, L. G. *Org. Biomol. Chem.* **2014**, *12*, 9557.
(c) Tran, L. D.; Popov, I.; Daugulis, O. *J. Am. Chem. Soc.* **2012**, *134*, 18237.
- [6] (a) Xia, X.; Zou, K. B.; Li, R. X.; Zhang, Z. Y.; Shao, L. L.; Fang, D. W.; Xu, X. H. *Acta Chim. Sinica* **2008**, *66*, 1749 (in Chinese). (夏湘, 邹康兵, 李若信, 张尊英, 邵玲玲, 方大为, 许新华, 化学学报, **2008**, *66*, 1749.)
(b) Shi, Q.; Li, P. H.; Yan, Z. Y.; Wang, L. *Org. Chem. Front.* **2017**, *4*, 1322.
- [7] Devi, N.; Rahaman, R.; Sarma, K.; Khan, T.; Barman, P. *Eur. J. Org. Chem.* **2017**, *11*, 1520.
- [8] Alves, D.; Lara, R. G.; Contreira, M. E.; Radatz, C. S.; Duarte, L. F. B.; Perin, G. *Tetrahedron Letters* **2012**, *53*, 3364.
- [9] (a) Yang, L.; Feng, J.; Qiao, M.; Zeng, Q. L. *Org. Chem. Front.* **2017**, DOI: 10.1039/c7qo00766c.
(b) Zhu, H.; Yu, J. T.; Cheng, J. *Chem. Commun.* **2016**, *52*, 11908.
- [10] (a) Campbell, D. S. *J. Appl. Polym. Sci.* **1969**, *13*, 1201.
(b) Fettes, E. M.; Mark, H. J. *Appl. Polym. Sci.* **1963**, *7*, 2239.
- [11] (a) Hajjimi, M.; Shiri, L.; Jahanbakhshi, A. *Appl. Organomet. Chem.* **2015**, *29*, 668.
(b) He, Y.; Huang, D.; Lu, M. *Phosphorus, Sulfur, Silicon Relat. Elem.* **2012**, *187*, 1118.
(c) Madabhushi, S.; Jillell, R.; Sriramoju, V.; Sing, R. *Green Chem.* **2014**, *16*, 3125.
(d) Hajipour, A. R.; Mallakpour, S. E.; Adibi, H. *J. Org. Chem.* **2002**, *67*, 8666.
(e) Deng, S. R.; Hu, G. Q.; Yan, Y.; Guo, J. P.; Huang, W. G.; Zhou, Y. H.; Li, Z. Y. *Chin. J. Org. Chem.* **2005**, *25*, 815 (in Chinese). (邓仕任, 胡高强, 鄢勇, 郭建平, 黄伟国, 周运鸿, 李早英, 有机化学, **2005**, *25*, 815.)
- [12] (a) Corma, A.; Rodenas, T.; Sabater, M. J. *Chem. Sci.* **2012**, *3*, 398.
(b) Donck, S.; Gravel, E.; Li, A.; Prakash, P.; Shah, N.; Leroy, J.; Li, H. Y.; Namboothiri, I. N. N.; Doris, E. *Catal. Sci. Technol.* **2015**, *5*, 4542.
(c) Menini, L.; Pereira, M. C.; Ferreira, A. C.; Fabris, J. D.; Gusevskaya, E. V. *Appl. Catal. A: Gen.* **2011**, *392*, 151.
(d) Chauhan, D.; Kumar, P.; Joshi, C.; Labhsetwar, N.; Ganguly, S. K.; Jain, S. L. *New J. Chem.* **2015**, *39*, 6193.
(e) Dhakshinamoorthy, A.; Primo, A.; Esteve-Adell, I.; Alvaro, M.; Garcia, H. *ChemCatChem* **2015**, *7*, 776.
(f) Patra, A. K.; Dutta, A.; Pramanik, M.; Nandi, M.; Uyama, H.; Bhaumik, A. *ChemCatChem* **2014**, *6*, 220.
- [13] (a) Kumar, P.; Singh, G.; Tripathi, D.; Jain, S. L. *RSC Adv.* **2014**, *4*, 50331.
(b) Talla, A.; Driessen, B.; Straathof, N. J. W.; Milroy, L. G.; Brunsveldt, L.; Hessel, V.; Noel, T. *Adv. Synth. Catal.* **2015**, *357*, 2180.
(c) Dou, Y. C.; Huang, X.; Wang, H.; Yang, L. T.; Li, H.; Yuan, B. X.; Chang, G. Y. *Green Chem.* **2017**, *19*, 2491.
(d) Oba, M.; Tanaka, K.; Nishiyama, K.; Ando, W. *J. Org. Chem.* **2011**, *76*, 4173.
- [14] (a) Sharma, S.; Barooah, N.; Baruah, J. B. *J. Mol. Catal. A: Chem.* **2005**, *229*, 171.
(b) Nair, V.; Augustine, A. *Org. Lett.* **2003**, *5*, 543.
(c) Yi, S. L.; Li, M. C.; Hu, X. Q.; Mo, W. M.; Shen, Z. L. *Chin. Chem. Lett.* **2016**, *27*, 1505.
- [15] (a) Natarajan, P.; Sharma, H.; Kaur, M.; Sharma, P. *Tetrahedron Lett.* **2015**, *56*, 5578.
(b) Hashemi, M.; Ghafari, H.; Karimi-Jaberi, Z. *J. Sulfur Chem.* **2006**, *27*, 165.
(c) Huang, X. Y.; Wang, T.; Xiao, C. Q.; Yu, X. Q.; Xie, R. G. *Chin. J. Org. Chem.* **2004**, *24*, 1629 (in Chinese).

- (黄小毅, 王涛, 夏传琴, 余孝其, 谢如刚, 有机化学, **2004**, *24*, 1629.)
- [16] (a) Bagi, N.; Kaizer, J.; Speier, G. *RSC Adv.* **2015**, *5*, 45983.
(b) Vilaivan, T. T.; Poochampa, K.; Vilaivan, T.; Sukwattanasinitt, M.; Wacharasindhu, S. *Tetrahedron* **2016**, *72*, 788.
(c) Thurow, S.; Pereira, V. A.; Martinez, D. M.; Alves, D.; Perin, G.; Jacob, R. G.; Lenardao, E. J. *Tetrahedron Lett.* **2011**, *52*, 640.
(d) Shard, A.; Kumar, R.; Saima; Sharma, N.; Sinha, A. K. *RSC Adv.* **2014**, *4*, 33399.
(e) Zhu, R. H.; Shi, X. X. *Synth. Commun.* **2012**, *42*, 1108.
(f) Banfield, S. C.; Omori, A. T.; Leisch, H.; Hudlicky, T. *J. Org. Chem.* **2007**, *72*, 4989.
(g) Abbasi, M.; Sabet, A. *J. Organomet. Chem.* **2017**, *833*, 10.
(h) Wang, G.; Guo, Y.; Lv, Y.; Wang, X. C.; Quan, Z. *J. Chin. J. Org. Chem.* **2016**, *36*, 1375 (in Chinese).
(王刚, 郭燕, 吕颖, 王喜存, 权正军, 有机化学, **2016**, *36*, 1375.)
(i) Jiang, H.; Zhong, M. N.; Chen, J. S. *Chin. J. Org. Chem.* **1999**, *19*, 214 (in Chinese).
(蒋辉, 钟明鼎, 陈冀胜, 有机化学, **1999**, *19*, 214.)
- [17] (a) Rosewall, C. F.; Sibbald, P. A.; Liskin, D. V.; Michael, F. E. *J. Am. Chem. Soc.* **2009**, *131*, 9488.
(b) Sibbald, P. A.; Michael, F. E. *Org. Lett.* **2009**, *11*, 1147.
- [18] (a) Sun, K.; Li, Y.; Xiong, T.; Zhang, J. P.; Zhang, Q. *J. Am. Chem. Soc.* **2011**, *133*, 1694.
(b) Zheng, G. F.; Li, Y.; Han, J. J. *Nat. Commun.* **2015**, *6*, 7011.
- [19] (a) Roy, A.; Schneller, S. W. *Org. Lett.* **2005**, *7*, 3889.
(b) Boger, D. L.; Brunette, S. R.; Garbaccio, R. M. *J. Org. Chem.* **2001**, *66*, 5163.
- [20] Foye, W. O.; Speranza, J. J. *Pharm. Sci.* **1970**, *59*, 259.
- [21] Yang, Z. H.; Yang, S. Y.; Xu, J. X. *Tetrahedron* **2017**, *73*, 3240.
- [22] Shah, S. T. A.; Khan, K. M.; Fecker, M.; Voelter, W. *Tetrahedron Lett.* **2003**, *44*, 6789.
- [23] Johnson, R. N.; Smiles, S. J. *Chem. Soc., Trans.* **1923**, *123*, 2384.
- [24] Sengar, R. S.; Nemykin, V. N.; Basu, P. *New J. Chem.* **2003**, *27*, 1115.
- [25] Minakami, S.; Kono, M. *JP 42002425*, **1967** [*Chem. Abstr.* **1967**, *67*, 32484m].
- [26] Ghorbani-Choghamarani, A.; Nikoorazm, M.; Goudarziafshar, H.; Shokr, A.; Almasi, H. *J. Chem. Sci.* **2011**, *123*, 453.
- [27] Vandavasi, J. K.; Hu, W. P.; Chen, C. Y.; Wang, J. J. *Tetrahedron* **2011**, *67*, 8895.
- [28] Ghorbani-Choghamarani, A.; Nikoorazm, M.; Azadi, G. *J. Serb. Chem. Soc.* **2013**, *78*, 173.
- [29] Chauhan, S. M. S.; Kumar, A.; Srinivas, K. A. *Chem. Commun.* **2003**, *18*, 2348.
- [30] Danieli, B.; Sacchetti, A.; Silvani, A.; Pratesi, G.; Zunino, F. *J. Org. Chem.* **2006**, *71*, 2848.
- [31] (a) *Merck Index* **2001**, *13*, 2811.
(b) Data of commercial product (TCI C0519).
(c) Kirihaara, M.; Asai, Y.; Ogawa, S.; Noguchi, T.; Hatano, A.; Hirai, Y. *Synthesis* **2007**, *21*, 3286.
(d) Fles, D.; Markovac-Prpic, A.; Tomasic, V. *J. Am. Chem. Soc.* **1958**, *80*, 4654.
- [32] Morgan, L. R. *US 4827016*, **1989** [*Chem. Abstr.* **1989**, *111*, 225308m].

(Li, L.; Fan, Y.)