烯基叠氮与二异丙基黄原酸酯的自由基串联反应: 合成 6-巯甲基菲啶

陆露露 周丙伟 金红卫* 刘运奎*

(浙江工业大学化学工程学院 绿色化学与技术国家重点实验室培育基地 杭州 310014)

摘要 发展了偶氮二异丁腈诱发的烯基叠氮类化合物与二异丙基黄原酸酯的自由基串联反应, 一步构建了碳-硫和碳-氮键. 以 50%至 80%的收率方便和高区域选择性地合成了一系列官能化的 6-巯甲基菲啶类化合物. 机理研究表明该反应通过自由基路径进行.

关键词 烯基叠氮; 二硫化二异丙基黄原酸酯; 菲啶; 自由基反应

Radical-Triggered Tandem Reaction of Vinyl Azides with Isopropylxanthic Disulfide for the Synthesis of 6-Sulfanylmethyl Phenanthridines

Lu, Lulu Zhou, Bingwei Jin, Hongwei* Liu, Yunkui*

(State Key Laboratory Breeding Base of Green Chemistry-Synthesis Technology, College of Chemical Engineering, Zhejiang University of Technology, Hangzhou 310014)

Abstract An 2,2'-azobis(2-methylpropionitrile) (AIBN) initiated tandem reaction of vinyl azides with isopropylxanthic disulfide to construct C—S/C—N bonds was disclosed. A range of functionalized 6-sulfanylmethyl phenanthridines could be easily accessed in 50%~84% yields with a good regioselectivity. The mechanism study indicates a free radical pathway in this reaction.

Keywords vinyl azides; isopropylxanthic disulfide; phenanthridine; radical reaction

1 Introduction

Phenanthridines and their derivatives are of great interest in medicinal chemistry and material science due to their potential biological activities, optoelectronic properties and others.^[1,2] As a result, many useful methods for the synthesis of phenanthridines have been well developed.^[3~5] Among them, radical-initiated tandem cyclization of *ortho*-substituted diaryl compounds renders the formation of 6-functionalized phenanthridines in a one-pot procedure.^[4] For example, Nanni *et al.*^[4a] disclosed a sequential radical addition/cyclization reaction of biphenyl-2-ylisonitrile with 2,2'-azobis(2-methylpropionitrile) (AIBN) or dibenzoyl peroxide (DBP). Afterwards, alkyl, aryl, perfluoroalkyl, acyl, silyl, sulfonyl, arylsulfanyl, and thiocyanate substrates were all proved to be versatile radical precursors and were successfully applied to the reaction with biphenyl-2-ylisonitrile (Scheme 1). Alternatively, α -(biaryl-

2-yl)vinyl azides have also received much attention to access 6-methyl substituted phenanthridines via a radical pathway.^[5] Various functional groups such as alkyl, perfluoroalkyl, sulfonyl and phosphoryl can be incorporated into methyl group at the 6-position of phenanthridines. However, the tandem reaction of α -(biaryl-2-yl)vinyl azides with sulfanyl radicals derived from disulfides is still developed.

It is noteworthy that disulfides are recognized as one class of important sulfur reagents and bidentate ligand donors.^[6] They are widely used as synthetic intermediates in organic reactions for the synthesis of organosulfur compounds, such as in the transition-metal-catalyzed cross coupling reactions and addition reactions of S—S bond to alkynes.^[7] Other transformations of disulfides involving sulfanyl radicals have also been explored in the past few decades.^[8] To further extend the application of disulfides in the synthesis of organosulfur compounds,^[9] we herein

* Corresponding authors. E-mail: jhwei828@zjut.edu.cn; ykuiliu@zjut.edu.cn

Received July 16, 2018; revised September 10, 2018; published online September 14, 2018.

Project supported by the National Natural Science Foundation of China (Nos. 21772176, 21372201), the Opening Foundation of Zhejiang Key Course of Chemical Engineering and Technology, Zhejiang University of Technology.

国家自然科学基金(Nos. 21772176, 21372201)、浙江工业大学“省重中之重一级学科”开放基金资助项目。

(a) Synthesis of 6-functionalized phenanthridines (previous)

R = alkyl, aryl, perfluoroalkyl, acyl, thiocyanato, arylthio, etc.

(b) Synthesis of 6-methyl functionalized phenanthridines (previous)

R = alkyl, perfluoroalkyl, sulfonyl and phosphoryl

(c) Cyclization of vinyl azide with isopropylxanthic disulfide (this work)

Scheme 1 Radical-initiated cyclization of *ortho*-substituted diaryls

report an AIBN initiated tandem cyclization of vinyl azides with isopropylxanthic disulfide leading to 6-sulfanylmethyl phenanthridines in moderate to good yields. Notably, this protocol can serve as a supplementary strategy to previous methodologies for the synthesis of 6-functionalized phenanthridines.

2 Results and discussion

Our study began with the cyclization reaction using α -(biaryl-2-yl)vinyl azide (**1a**) and isopropylxanthic disulfide as model substrates, AIBN as a radical initiator in tetrahydrofuran (THF) for 8 h. To our delight, the desired product 6-sulfanylmethyl phenanthridine (**3a**) was obtained in 58% yield (Table 1, Entry 1). A control experiment showed that the radical initiator was indispensable to fulfill this transformation which implied a typical radical pathway (Entry 2). Next, dimethyl 2,2'-azobis(2-methylpropanoate) (AIBME) was used as a radical initiator instead of AIBN, whereas the yield of **3a** was not improved (Entry 3). We then examined the amount of isopropylxanthic disulfide on the reaction. It was found that a comparable yield of **3a** was obtained when the amount of disulfide was increased to 1.5 equiv. (Entries 4, 5). Increasing the amount of AIBN further improved the yield of **3a** to 84% (Entry 8). Screening various solvents revealed that THF remained the optimal one (Entry 10~12). Furthermore, either elevating or decreasing the temperature provided lower yields of **3a** (Entries 13, 14).

To evaluate the substrate scope of this cyclization reaction, a range of substituted vinyl azides were investigated under the optimized reaction conditions (Table 2). Different substituents R^1 on the phenyl ring showed unbiased electronic effect on the reaction outcome (**3a**~**3e**). The

Table 2 Screening for reaction parameters^a

Entry	Radical initiator (equiv.)	Solvent	Temp./°C	Yield ^b /%
1	AIBN (0.5)	THF	80	58
2	—	THF	80	0
3	AIBME (0.5) ^c	THF	80	50
4	AIBN (0.5)	THF	80	35 ^d
5	AIBN (0.5)	THF	80	56 ^e
6	AIBN (0.3)	THF	80	45
7	AIBN (0.75)	THF	80	63
8	AIBN (1.0)	THF	80	84 (71 ^f , 80 ^g)
9	AIBN (1.5)	THF	80	75
10	AIBN (1.0)	MeCN	80	24
11	AIBN (1.0)	DMSO	80	38
12	AIBN (1.0)	Toluene	80	Trace
13	AIBN (1.0)	THF	60	70
14	AIBN (1.0)	THF	100	52
15	AIBN (1.0)	2-MeTHF	80	65

^a Reaction conditions: **1a** (0.3 mmol), isopropylxanthic disulfide (0.3 mmol), radical initiator (0.3 mmol), solvent (3 mL) for 8 h under air. ^b Isolated yields were given. ^c AIBME = dimethyl 2,2'-azobis(2-methylpropanoate). ^d Isopropylxanthic disulfide (0.5 equiv.). ^e Isopropylxanthic disulfide (1.5 equiv.). ^f 2 mL of THF was used (0.15 mol/L for **1a**). ^g 4 mL of THF was used (0.075 mol/L for **1a**).

compatibility of **1** with various substituents R^2 on the phenyl ring was investigated and the corresponding products were obtained in moderate to good yields (**3f**~**3p**). Of note, substrates with electron-donating groups (EDGs) exhibited an inferior reactivity (**3j**~**3o**). When **1p** was used, the reaction preferred the formation of C—N bond at the less hindered 6'-position of **1p** and 2-chloro-phenanthridine derivative **3p** was isolated in 45% yield as a major isomer (Entry 16). The substrates that both of the two phenyl rings tethered with functional groups are also workable which provided multi-substituted phenanthridines in moderate yields (**3q**~**3t**).

In order to explore the reaction mechanism, the radical inhibiting experiments were carried out. The extra addition of 2,2,6,6-tetramethylpiperidin-1-oxyl (TEMPO)^[10] or 2,6-bis(1,1-dimethylethyl)-4-methylphenol^[11] (BHT) to the model reaction resulted in a drastic yield decrease, respectively (Scheme 2). It implied that a radical pathway might be involved. An attempt to run the reaction under the standard conditions except in argon atmosphere only resulted in low yield of **3a**, suggesting O_2 in open air plays important role in the reaction.

Based on the above investigations and literature clues,^[5,12,13] a possible reaction mechanism is illustrated in Scheme 3. A sulfanyl radical **A** is first generated by treatment of isopropylxanthic disulfide with AIBN.^[12] The addition of radical **A** to vinyl azide **1a** takes place to give an

Table 2 Substrate scope for vinyl azides **1**^a

Entry	R ¹	R ²	Product	Yield ^b /%
1	H	H	3a	84
2	5-F	H	3b	68
3	5-OCH ₃	H	3c	55
4	4-Cl	H	3d	56
5	4-Me	H	3e	53
6	H	4-F	3f	72
7	H	4-Cl	3g	70
8	H	4-Br	3h	74
9	H	4-OCF ₃	3i	76
10	H	4-CH ₃	3j	50
11	H	3,5-(Me) ₂	3k	52
12	H	4- <i>i</i> -Pr	3l	50
13	H	4- <i>t</i> -Bu	3m	57
14	H	4-OCH ₃	3n	52
15	H	2-OCH ₃	3o	53
16	H	3-Cl	3p	45 ^c
17	5-OCH ₃	4-Br	3q	70
18	5-OCH ₃	4- <i>i</i> -Pr	3r	63
19	5-F	4- <i>i</i> -P	3s	73
20	4-Cl	2-OCH ₃	3t	73

^a Reaction conditions: **1a** (0.3 mmol), isopropylxanthic disulfide (0.3 mmol), additive (0.3 mmol), solvent (3 mL) for 8 h under air. ^b Isolated yield. ^c The yield of regioisomer **3p'** is 7%.

Scheme 2 Mechanism experiments

imine radical **B** by extrusion of N₂.^[5] An intramolecular C—N cyclization of **B** produces an intermediate **C**.^[5] **C** undergoes an oxidation followed by an aromatization to afford the final product **3a**.^[13] In the reaction, the O₂ in the air may participate as the oxidant.^[13,14]

Scheme 3 Proposed mechanism for the formation of **3a**

3 Conclusions

In summary, we have developed a tandem reaction of vinyl azides with isopropylxanthic disulfide affording 6-sulfanylmethyl phenanthridines in moderate to good yields. The reaction has advantages of good functional group tolerance, high regioselectivity, simple operation, mild reaction conditions, and free-use of transition-metals. The development of novel reactions involving vinyl azides via a radical pathway is underway in our laboratory.

4 Experimental

4.1 General experimental information

Unless otherwise noted, all reactions were conducted in oven-dried sealed tubes with Teflon screw caps under air. All commercially available reagents and solvents were purchased from chemical suppliers and used as received. Flash column chromatography was performed on silica gel (100~200 mesh) with indicated solvent mixtures. Thin-layer chromatography (TLC) analysis was performed on pre-coated, glass-backed silica gel plates and visualized with UV light. The ¹H NMR and ¹³C NMR spectra were recorded on a Bruker 500 AV spectrometers. Chemical shifts (δ) were reported downfield from tetramethylsilane. Melting points were measured and uncorrected. The high resolution mass spectra (HRMS) were obtained on a TOF-MS instrument with ESI source. Vinyl azides were synthesized according to the previous literature.^[5a]

4.2 General synthesis procedure

Vinyl azide **1a** (0.3 mmol, 66.3 mg), isopropylxanthic disulfide **2** (0.3 mmol, 81 mg), azobisisobutyronitrile, AIBN (0.3 mmol, 49.2 mg), tetrahydrofuran (THF) (3 mL), and a stirrer bar were added to a 15 mL sealed tube with Teflon screw cap. The reaction mixture was then stirred at 80 °C for 8 h. After cooling, CH₂Cl₂ and H₂O were added to the mixture. The organic layer was separated and dried over anhydrous sodium sulfate and filtered through a short pad of Celite. After evaporation of the solvent under vacuum, the residue was purified by column chromatography on silica gel (100~200 mesh) using petrol ether (PE) and ethyl acetate (EA) (*V*: *V*=100:1) as an eluent to give

pure product **3a** in 84% yield as a yellow solid.

O-Isopropyl *S*-(phenanthridin-6-ylmethyl) carbonodithioate (**3a**): Yellow solid, m.p. 55~56 °C; ¹H NMR (500 MHz, CDCl₃) δ: 8.55 (d, *J*=8.5 Hz, 1H), 8.46 (dd, *J*=8.0, 1.0 Hz, 1H), 8.18 (d, *J*=8.0 Hz, 1H), 8.05 (dd, *J*=8.5, 1.0 Hz, 1H), 7.79~7.75 (m, 1H), 7.67~7.61 (m, 2H), 7.60~7.56 (m, 1H), 5.79~5.71 (m, 1H), 5.04 (s, 2H), 1.35 (d, *J*=6.0 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ: 213.0, 155.1, 143.3, 133.1, 130.9, 129.8, 128.9, 127.6, 127.3, 126.1, 124.7, 124.0, 122.6, 122.0, 78.3, 41.2, 21.36; HRMS (ESI) calcd for C₁₈H₁₈NOS₂ [M+H]⁺: 328.0830, found 328.0835.

S-((8-Fluorophenanthridin-6-yl)methyl) *O*-isopropyl carbonodithioate (**3b**): Yellow solid, m.p. 102~103 °C; ¹H NMR (500 MHz, CDCl₃) δ: 8.50 (d, *J*=8.0 Hz, 1H), 8.45 (dd, *J*=9.0, 5.5 Hz, 1H), 8.21 (d, *J*=8.0 Hz, 1H), 7.37~7.80 (m, 1H), 7.30 (dd, *J*=8.5, 2.5 Hz, 1H), 7.69~7.65 (m, 1H), 7.40~7.35 (m, 1H), 5.87~5.78 (m, 1H), 5.06 (s, 2H), 1.42 (d, *J*=6.0 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ: 212.6, 161.3 (d, ¹*J*_{C-F}=248.0 Hz), 154.2 (d, ⁴*J*_{C-F}=3.5 Hz), 143.0, 123.0, 129.7 (d, ⁴*J*_{C-F}=1.9 Hz), 128.1, 127.6, 125.8 (d, ³*J*_{C-F}=7.9 Hz), 125.1 (d, ³*J*_{C-F}=8.3 Hz), 123.4, 121.7, 120.0 (d, ²*J*_{C-F}=23.9 Hz), 110.6 (d, ²*J*_{C-F}=21.8 Hz), 78.5, 41.1, 21.3; HRMS (ESI) calcd for C₁₈H₁₇FNOS₂ [M+H]⁺: 346.0736, found 346.0739.

O-Isopropyl *S*-((8-methoxyphenanthridin-6-yl)methyl) carbonodithioate (**3c**): Yellow solid, m.p. 92~93 °C; ¹H NMR (500 MHz, CDCl₃) δ: 8.52 (d, *J*=9.0 Hz, 1H), 8.44 (dd, *J*=7.5, 1.0 Hz, 1H), 8.09 (dd, *J*=8.0, 1.0 Hz, 1H), 7.68~7.60 (m, 2H), 7.55 (d, *J*=2.5 Hz, 1H), 7.46 (dd, *J*=9.0, 2.5 Hz, 1H), 5.87~5.77 (m, 1H), 5.13 (s, 2H), 3.95 (s, 3H), 1.42 (d, *J*=6.2 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ: 213.7, 158.8, 154.4, 142.6, 129.8, 127.8, 127.5, 127.3, 125.9, 124.2, 124.2, 121.6, 121.5, 106.2, 78.6, 55.8, 41.8, 21.3; HRMS (ESI) calcd for C₁₉H₂₀NO₂S₂ [M+H]⁺: 358.0935, found 358.0930.

S-((9-Chlorophenanthridin-6-yl)methyl) *O*-isopropyl carbonodithioate (**3d**): Yellow solid, m.p. 90~91 °C; ¹H NMR (500 MHz, CDCl₃) δ: 8.45 (d, *J*=2.0 Hz, 1H), 8.33 (d, *J*=8.5 Hz, 1H), 8.10 (d, *J*=8.5 Hz, 1H), 8.06 (d, *J*=8.2, 1H), 7.72~7.69 (m, 1H), 7.62~7.56 (m, 2H), 5.82 (m, 1H), 5.01 (s, 2H), 1.42 (d, *J*=6.3 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ: 212.8, 154.5, 143.6, 137.2, 134.2, 129.9, 129.4, 128.0, 127.5, 122.4, 122.8, 122.7, 122.1, 121.9, 78.3, 41.2, 21.3; HRMS (ESI) calcd for C₁₈H₁₇ClNOS₂ [M+H]⁺: 362.0440, found 362.0436.

O-isopropyl *S*-((9-methylphenanthridin-6-yl)methyl) carbonodithioate (**3e**): Yellow solid, m.p. 89~90 °C; ¹H NMR (500 MHz, CDCl₃) δ: 8.36 (d, *J*=8.5 Hz, 1H), 8.21 (s, 1H), 8.04~8.02 (m, 1H), 7.98 (d, *J*=8.0 Hz, 1H), 7.65~7.61 (m, 1H), 7.55~7.52 (m, 1H), 7.36 (d, *J*=8.5 Hz, 1H), 5.84~5.76 (m, 1H), 4.97 (s, 2H), 2.51 (s, 3H), 1.39 (d, *J*=6.5 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ: 213.0, 154.5, 143.4, 141.1, 132.9, 129.6, 129.1, 128.5, 126.8, 125.8, 123.7, 122.5, 122.0, 121.8, 78.1, 41.2, 22.1, 21.3; HRMS (ESI) calcd for C₁₉H₂₀NOS₂ [M+H]⁺: 342.0986, found 342.0989.

S-((3-Fluorophenanthridin-6-yl)methyl) *O*-isopropyl carbonodithioate (**3f**): Yellow solid, m.p. 68~69 °C; ¹H NMR (500 MHz, CDCl₃) δ: 8.50 (d, *J*=8.0 Hz, 1H), 8.45 (dd, *J*=9.0, 5.5 Hz, 1H), 8.21 (d, *J*=8.0 Hz, 1H), 7.37~7.80 (m, 1H), 7.30 (dd, *J*=8.5, 2.5 Hz, 1H), 7.69~7.65 (m, 1H), 7.40~7.35 (m, 1H), 5.87~5.78 (m, 1H), 5.06 (s, 2H), 1.42 (d, *J*=6.0 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ: 212.9, 162.6 (d, ¹*J*_{C-F}=247.0 Hz), 156.5, 144.6 (d, ³*J*_{C-F}=12.1 Hz), 132.7, 131.1, 127.4, 126.0, 124.2, 123.9 (d, ³*J*_{C-F}=9.3 Hz), 122.3, 120.6, 116.2 (d, ²*J*_{C-F}=23.6 Hz), 114.3 (d, ²*J*_{C-F}=20.6 Hz), 78.3, 41.1, 21.32; HRMS (ESI) calcd for C₁₈H₁₇FNOS₂ [M+H]⁺: 346.0736, found 346.0740.

S-((3-Chlorophenanthridin-6-yl)methyl) *O*-isopropyl carbonodithioate (**3g**): Yellow solid, m.p. 86~87 °C; ¹H NMR (500 MHz, CDCl₃) δ: 8.48 (d, *J*=8.3 Hz, 1H), 8.35 (d, *J*=8.8 Hz, 1H), 8.18 (d, *J*=8.1 Hz, 1H), 8.02 (d, *J*=8.2 Hz, 1H), 7.81~7.77 (m, 1H), 7.67~7.63 (m, 1H), 7.51 (dd, *J*=8.8, 2.2 Hz, 1H), 5.75 (m, 1H), 5.01 (s, 2H), 1.35 (d, *J*=6.3 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ: 212.9, 156.5, 144.1, 134.5, 132.6, 131.2, 129.1, 127.9, 127.7, 126.1, 124.6, 123.3, 122.5, 122.5, 78.4, 41.2, 21.4; HRMS (ESI) calcd for C₁₈H₁₇ClNOS₂ [M+H]⁺: 362.0440, found 362.0437.

S-((3-Bromophenanthridin-6-yl)methyl) *O*-isopropyl carbonodithioate (**3h**): Yellow solid, m.p. 82~83 °C; ¹H NMR (500 MHz, CDCl₃) δ: 8.52 (d, *J*=8.0 Hz, 1H), 8.31 (d, *J*=8.5 Hz, 1H), 8.25 (d, *J*=2.0 Hz, 1H), 8.23 (d, *J*=8.1 Hz, 1H), 7.86~7.82 (m, 1H), 7.73~7.68 (m, 2H), 5.87~5.78 (m, 1H), 5.06 (s, 2H), 1.43 (d, *J*=6.3 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ: 212.9, 156.4, 144.2, 132.6, 132.2, 131.2, 130.3, 128.0, 126.1, 124.6, 123.4, 122.8, 122.5, 122.4, 78.4, 41.1, 21.4; HRMS (ESI) calcd for C₁₈H₁₇BrNOS₂ [M+H]⁺: 405.9935, found 405.9930.

O-Isopropyl *S*-((3-(trifluoromethoxy)phenanthridin-6-yl)methyl) carbonodithioate (**3i**): Yellow solid, m.p. 95~96 °C; ¹H NMR (500 MHz, CDCl₃) δ: 8.53 (d, *J*=8.5 Hz, 1H), 8.49 (d, *J*=9.0 Hz, 1H), 8.24 (d, *J*=8.0 Hz, 1H), 7.95 (dd, *J*=2.5, 1.0 Hz, 1H), 7.87~7.83 (m, 1H), 7.73~7.69 (m, 1H), 7.48 (dd, *J*=9.0, 2.0 Hz, 1H), 5.87~5.79 (m, 1H), 5.08 (s, 2H), 1.43 (d, *J*=6.0 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ: 212.8, 156.8, 149.2, 143.0, 132.4, 131.2, 128.0, 126.1, 124.6, 123.7, 122.5, 122.4, 120.6 (q, *J*=256.9 Hz), 120.4, 120.4, 78.4, 41.1, 21.3; HRMS (ESI) calcd for C₁₉H₁₇F₃NO₂S₂ [M+H]⁺: 412.0653, found 412.0659.

O-Isopropyl *S*-((3-methylphenanthridin-6-yl)methyl) carbonodithioate (**3j**): Yellow solid, m.p. 90~91 °C; ¹H NMR (500 MHz, CDCl₃) δ: 8.58 (d, *J*=8.5 Hz, 1H), 8.38 (d, *J*=8.5 Hz, 1H), 8.21 (d, *J*=8.0 Hz, 1H), 7.90 (s, 1H), 7.82~7.78 (m, 1H), 7.67~7.63 (m, 1H), 7.46 (dd, *J*=8.5, 1.5 Hz, 1H), 5.87~5.79 (m, 1H), 5.08 (s, 2H), 5.57 (s, 3H), 1.43 (d, *J*=6.3 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ: 231.2, 154.9, 143.5, 138.7, 133.1, 130.7, 129.4, 128.9, 127.1, 125.9, 124.3, 122.4, 121.7, 121.6, 78.2, 41.3, 21.5, 21.3; HRMS (ESI) calcd for C₁₉H₂₀NOS₂ [M+H]⁺: 342.0986, found 342.0980.

S-((2,4-Dimethylphenanthridin-6-yl)methyl) *O*-isopropyl carbonodithioate (**3k**): Yellow solid, m.p. 90~91 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.52 (d, $J=8.5$ Hz, 1H), 8.15 (d, $J=8.0$ Hz, 1H), 8.08 (s, 1H), 7.75 (t, $J=8.0$ Hz, 1H), 7.62 (t, $J=7.5$ Hz, 1H), 7.36 (s, 1H), 5.88~5.80 (m, 1H), 5.08 (s, 2H), 2.78 (s, 3H), 2.53 (s, 3H), 1.42 (d, $J=6.0$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 14.1, 152.0, 140.2, 137.5, 136.5, 133.0, 131.2, 130.3, 127.1, 125.6, 124.3, 123.6, 122.7, 119.3, 77.9, 42.0, 22.0, 21.4, 18.1; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{22}\text{NOS}_2$ $[\text{M} + \text{H}]^+$: 356.1143, found 356.1149.

O-Isopropyl *S*-((3-isopropylphenanthridin-6-yl)methyl) carbonodithioate (**3l**): Yellow solid, m.p. 85~87 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.57 (d, $J=8.0$ Hz, 1H), 8.44 (d, $J=8.5$ Hz, 1H), 8.23 (d, $J=8.2$ Hz, 1H), 7.98 (d, $J=1.5$ Hz, 1H), 7.83~7.79 (m, 1H), 7.67~7.63 (m, 1H), 7.54 (dd, $J=8.5$, 1.5 Hz, 1H), 5.87~5.79 (m, 1H), 5.16 (s, 2H), 3.18~3.09 (m, 1H), 1.42 (d, $J=6.0$ Hz, 6H), 1.38 (d, $J=7.0$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 213.1, 154.7, 149.9, 143.6, 133.1, 130.7, 127.1, 126.7, 126.6, 126.0, 124.4, 122.4, 122.0, 121.9, 78.2, 41.3, 34.1, 23.9, 21.4; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{24}\text{NOS}_2$ $[\text{M} + \text{H}]^+$: 370.1299, found 370.1293.

S-((3-*tert*-Butyl)phenanthridin-6-yl)methyl *O*-isopropyl carbonodithioate (**3m**): Yellow solid, m.p. 90~91 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.56 (d, $J=8.5$ Hz, 1H), 8.43 (d, $J=9.0$ Hz, 1H), 8.24~8.21 (m, 1H), 8.11 (d, $J=2.0$ Hz, 1H), 7.82~7.78 (m, 1H), 7.72 (dd, $J=8.5$, 2.0 Hz, 1H), 7.66~7.62 (m, 1H), 5.87~5.79 (m, 1H), 5.11 (s, 2H), 1.45 (s, 9H), 1.38 (d, $J=6.0$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 213.1, 154.8, 152.2, 143.4, 133.0, 130.7, 127.1, 126.0, 125.8, 125.5, 124.4, 122.4, 121.7, 121.6, 78.2, 41.3, 35.0, 31.3, 21.3; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{26}\text{NOS}_2$ $[\text{M} + \text{H}]^+$: 384.1456, found 384.1449.

O-Isopropyl *S*-((3-methoxyphenanthridin-6-yl)methyl) carbonodithioate (**3n**): Yellow solid, m.p. 85~86 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.53 (d, $J=8.3$ Hz, 1H), 8.42 (d, $J=9.0$ Hz, 1H), 8.24 (d, $J=8.2$ Hz, 1H), 7.84~7.80 (m, 1H), 7.63 (dd, $J=11.4$, 2.0 Hz, 1H), 7.53 (d, $J=2.6$ Hz, 1H), 7.29 (dd, $J=9.0$, 2.6 Hz, 1H), 5.88~5.79 (m, 1H), 5.11 (s, 2H), 3.98 (s, 3H), 1.43 (d, $J=6.3$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 213.0, 160.3, 155.5, 133.3, 131.0, 128.7, 126.5, 126.1, 123.8, 123.2, 122.1, 118.4, 118.1, 109.6, 78.3, 55.6, 41.3, 21.4; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{20}\text{NO}_2\text{S}_2$ $[\text{M} + \text{H}]^+$: 358.0935, found 358.0941.

O-Isopropyl *S*-((1-methoxyphenanthridin-6-yl)methyl) carbonodithioate (**3o**): Yellow solid, m.p. 56~57 °C; ^1H NMR (500 MHz, CDCl_3) δ : 9.58 (d, $J=9.0$ Hz, 1H), 8.25 (d, $J=8.0$ Hz, 1H), 7.84~7.76 (m, 2H), 7.70~7.62 (m, 2H), 7.14 (d, $J=8.0$ Hz, 1H); 5.87~5.79 (m, 1H), 5.10 (s, 2H), 4.12 (s, 3H), 1.42 (d, $J=6.0$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 213.2, 158.0, 155.4, 145.4, 133.1, 130.7, 128.4, 128.3, 126.9, 125.4, 125.0, 122.7, 114.7, 108.4, 78.2, 55.9, 41.4, 21.4; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{20}\text{NO}_2\text{S}_2$ $[\text{M} + \text{H}]^+$: 358.0935, found 358.0942.

S-((4-Chlorophenanthridin-6-yl)methyl) *O*-isopropyl carbonodithioate (**3p**): Yellow solid, m.p. 87~89 °C; ^1H

NMR (500 MHz, CDCl_3) δ : 8.59 (d, $J=8.5$ Hz, 1H), 8.43 (dd, $J=7.0$, 1.0 Hz, 1H), 8.28 (d, $J=8.0$ Hz, 1H), 7.89~7.85 (m, 1H), 7.82 (dd, $J=7.0$, 1.0 Hz, 1H), 7.77~7.73 (m, 1H), 7.55 (t, $J=8.0$ Hz, 1H), 5.90~5.82 (m, 1H), 5.17 (s, 2H), 1.45 (d, $J=6.0$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 213.3, 155.8, 139.7, 134.4, 132.8, 131.2, 129.2, 128.2, 127.0, 126.1, 125.6, 124.6, 122.9, 120.8, 78.3, 41.8, 21.3; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{17}\text{ClNOS}_2$ $[\text{M} + \text{H}]^+$: 362.0440, found 362.0433.

S-((2-Chlorophenanthridin-6-yl)methyl) *O*-isopropyl carbonodithioate (**3p'**): Yellow solid, m.p. 84~85 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.55 (d, $J=8.5$ Hz, 1H), 8.46 (dd, $J=8.0$, 1.0 Hz, 1H), 8.18 (d, $J=8.0$ Hz, 1H), 8.05 (dd, $J=8.5$, 1.0 Hz, 1H), 7.79~7.75 (m, 1H), 7.67~7.61 (m, 2H), 7.60~7.56 (m, 1H), 5.79~5.71 (m, 1H), 5.04 (s, 2H), 1.35 (d, $J=6.0$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 212.9, 155.5, 141.9, 133.2, 132.1, 131.4, 131.2, 129.4, 128.3, 126.2, 125.1, 124.9, 126.7, 121.7, 78.4, 41.2, 21.4; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{17}\text{ClNOS}_2$ $[\text{M} + \text{H}]^+$: 362.0440, found 362.0443.

S-((3-bromo-8-methoxyphenanthridin-6-yl)methyl) *O*-isopropyl carbonodithioate (**3q**): Yellow solid, m.p. 75~76 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.57 (d, $J=8.0$ Hz, 1H), 8.50 (d, $J=7.5$ Hz, 1H), 8.29 (d, $J=8.0$ Hz, 1H), 8.06 (d, $J=8.5$ Hz, 1H), 7.91~7.87 (m, 1H), 7.79~7.75 (m, 1H), 7.67 (dd, $J=9.0$, 2.5 Hz, 1H), 5.88~5.80 (m, 1H), 5.11 (s, 2H), 3.95 (s, 3H), 1.44 (d, $J=6.0$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 213.5, 159.0, 155.6, 144.3, 132.1, 130.3, 126.9, 125.8, 124.0, 122.9, 122.9, 121.9, 121.2, 106.2, 78.7, 55.8, 41.6, 21.3; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{19}\text{BrNO}_2\text{S}_2$ $[\text{M} + \text{H}]^+$: 436.0041, found 436.0037.

O-Isopropyl *S*-((3-isopropyl-8-methoxyphenanthridin-6-yl)methyl) carbonodithioate (**3r**): Yellow solid, m.p. 90~91 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.37 (d, $J=9.0$ Hz, 1H), 8.27 (d, $J=8.5$ Hz, 1H), 7.93 (d, $J=2.0$ Hz, 1H), 7.47 (dd, $J=8.0$, 1.5 Hz, 1H), 7.45 (d, $J=2.5$ Hz, 1H), 7.35 (dd, $J=9.0$, 2.0 Hz, 1H), 5.58~5.77 (m, 1H), 5.08 (s, 2H), 3.89 (s, 3H), 3.15~3.06 (m, 1H), 1.40 (d, $J=6.5$ Hz, 6H), 1.38 (d, $J=7.0$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 213.7, 158.3, 154.1, 148.7, 142.7, 127.4, 126.7, 126.3, 125.4, 123.9, 122.0, 121.4, 121.3, 105.9, 78.4, 55.6, 41.8, 33.9, 23.9, 21.4; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{26}\text{NO}_2\text{S}_2$ $[\text{M} + \text{H}]^+$: 400.1405, found 400.1412.

S-((8-Fluoro-3-isopropylphenanthridin-6-yl)methyl) *O*-isopropyl carbonodithioate (**3s**): Yellow solid; m.p. 77~78 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.57 (dd, $J=9.0$, 5.0 Hz, 1H), 8.38 (d, $J=8.5$ Hz, 1H), 7.97 (d, $J=1.5$ Hz, 1H), 7.87 (dd, $J=9.5$, 2.5 Hz, 1H), 7.59~7.54 (m, 2H), 5.88~5.80 (m, 1H), 5.04 (s, 2H), 3.19~3.10 (m, 1H), 1.44 (d, $J=6.5$ Hz, 6H), 1.38 (d, $J=7.0$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 213.7, 162.1, 160.2, 154.2 (d, $^4J_{\text{C-F}}=3.6$ Hz), 149.9, 143.3, 129.9, 127.2, 126.7, 125.6 (d, $^3J_{\text{C-F}}=8.0$ Hz), 125.0 (d, $^3J_{\text{C-F}}=8.3$ Hz), 121.6 (d, $^2J_{\text{C-F}}=10.4$ Hz), 120.0 (d, $^1J_{\text{C-F}}=23.9$ Hz), 110.7 (d, $^2J_{\text{C-F}}=21.5$ Hz), 78.5, 41.2, 34.1, 23.9, 21.4; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{23}\text{FNOS}_2$ $[\text{M} + \text{H}]^+$: 388.1205, found 388.1209.

S-((9-Chloro-1-methoxyphenanthridin-6-yl)methyl) *O*-isopropyl carbonodithioate (**3t**): Yellow solid, m.p. 87~89 °C; ¹H NMR (500 MHz, CDCl₃) δ: 9.54 (d, *J*=2.0 Hz, 1H), 8.15 (d, *J*=8.5 Hz, 1H), 7.75 (dd, *J*=8.5, 1.0 Hz, 1H), 7.65 (t, *J*=8.0 Hz, 1H), 7.61 (dd, *J*=9.0, 2.0 Hz, 1H), 7.13 (d, *J*=8.0 Hz, 1H), 5.87~5.64 (m, 1H), 5.04 (s, 2H), 4.13 (s, 3H), 1.42 (d, *J*=6.5 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ: 213.1, 160.2, 155.5, 145.1, 133.3, 130.9, 126.5, 126.0, 123.8, 123.2, 112.1, 118.4, 118.1, 109.6, 78.3, 55.6, 41.3, 21.4; HRMS (ESI) calcd for C₁₉H₁₉ClNO₂S₂ [M+H]⁺: 392.0546, found 392.0549.

Supporting Information The spectroscopic characterization of the products. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- (a) Keene, B. R. T.; Tislington, P. *Adv. Heterocycl. Chem.* **1971**, *13*, 315.
(b) Bernardo, P. H.; Wan, K. F.; Sivaraman, T.; Xu, J.; Moore, F. K.; Hung, A. W.; Mok, H. Y. K.; Yu, V. C.; Chai, C. L. *J. Med. Chem.* **2008**, *51*, 6699.
(c) Zhang, L.; Ang, G. Y.; Chiba, S. *Org. Lett.* **2010**, *12*, 3682.
- (a) Cappelli, A.; Anzini, M.; Vomero, S.; Mannuni, L.; Makovec, F.; Doucet, E.; Hamon, M.; Bruni, G.; Romeo, M. R.; Menziani, M. C.; Benedetti, P. G.; Langer, T. *J. Med. Chem.* **1998**, *41*, 728.
(b) Lynch, M. A.; Duval, O.; Sukhanova, A.; Devy, J.; MacKay, S. P.; Waigh, R. D.; Nabiev, I. *Bioorg. Med. Chem. Lett.* **2001**, *11*, 2643.
(c) Lewis, W. G.; Green, L. G.; Grynszpan, F.; Radic, Z.; Carlier, P. R.; Taylor, P.; Finn, M. G.; Sharpless, K. B. *Angew. Chem., Int. Ed.* **2002**, *41*, 1053.
(d) Cappoen, D.; Claes, P.; Jacobs, J.; Anthonissen, R.; Mathys, V.; Verschaeve, L.; Huygen, K.; Kimpe, N. D. *J. Med. Chem.* **2014**, *57*, 2895.
(e) Naidua, K. M.; Nagesha, H. N.; Singh, M.; Sriram, D.; Yogeeswarar, P.; Sekhara, K. V. G. C. *Eur. J. Med. Chem.* **2015**, *92*, 415.
(f) Riddell, I. A.; Johnstone, T. C.; Park, G. Y.; Lippard, S. J. *Chem.-Eur. J.* **2016**, *22*, 7574.
- (a) Alonso, R.; Campos, P. J.; García, B.; Rodríguez, M. A. *Org. Lett.* **2006**, *8*, 3521.
(b) Portela-Cubillo, F.; Lymer, J.; Scanlan, E. M.; Scott, J. S.; Walton, J. C. *Tetrahedron* **2008**, *64*, 11908.
(c) McBurney, R. T.; Slawin, A. M. Z.; Smart, L. A.; Yu, Y.; Walton, J. C. *Chem. Commun.* **2011**, *47*, 7974.
(d) McBurney, R. T.; Walton, J. C. *J. Am. Chem. Soc.* **2013**, *135*, 7349.
(e) McBurney, R. T.; Walton, J. C. *Beilstein J. Org. Chem.* **2013**, *9*, 1083.
(f) Jiang, H.; An, X.; Tong, K.; Zheng, T.; Zhang, Y.; Yu, S. *Angew. Chem., Int. Ed.* **2015**, *54*, 4055.
(g) Hofstra, J. L.; Grassbaugh, B. R.; Tran, Q. M.; Armada, N. R.; de Lijser, H. J. P. *J. Org. Chem.* **2015**, *80*, 256.
(h) Liu, X.; Qing, Z.; Cheng, P.; Zheng, X.; Zeng, J.; Xie, H. *Molecules* **2016**, *21*, 1690.
(i) Tang, J.; Sivaguru, P.; Ning, Y.; Zannoni, G.; Bi, X. *Org. Lett.* **2017**, *19*, 4026.
- (a) Nanni, D.; Pareschi, P.; Rizzoli, C.; Sgarabotto, P.; Tundo, A. *Tetrahedron* **1995**, *51*, 9045.
(b) Tobisu, M.; Koh, K.; Furukawa, T.; Chatani, N. *Angew. Chem., Int. Ed.* **2012**, *51*, 11363.
(c) Zhang, B.; Mück-Lichtenfeld, C.; Daniliuc, C. G.; Studer, A. *Angew. Chem., Int. Ed.* **2013**, *52*, 10972.
(d) Leifert, D.; Daniliuc, C. G.; Studer, A. *Org. Lett.* **2013**, *15*, 6286.
(e) Sha, W.; Yu, J.-T.; Jiang, Y.; Yang, H.; Cheng, J. *Chem. Commun.* **2014**, *50*, 9179.
(f) Fang, H.; Zhao, J.; Qian, P.; Han, J.; Pan, Y. *Asian J. Org. Chem.* **2014**, *3*, 1266.
(g) Xiao, T.; Li, L.; Lin, G.; Wang, Q.; Zhang, P.; Mao, Z.-W.; Zhou, L. *Green Chem.* **2014**, *16*, 2418.
(h) Fang, H.; Zhao, J. C.; Qian, P.; Han, J. L.; Pan, Y. *Asian J. Org. Chem.* **2014**, *12*, 1266.
(i) Zhang, Z.; Tang, X.; Dolbier, W. R., Jr. *Org. Lett.* **2015**, *17*, 4401.
(j) Lu, S.; Gong, Y.; Zhou, D. *J. Org. Chem.* **2015**, *80*, 9336.
(k) Zhang, H.; Shi, D.; Ren, S.; Jin, H.; Liu, Y. *Eur. J. Org. Chem.* **2016**, 4224.
(l) Wu, C.; Zhou, Y.; Dong, X.; Qu, J. *ARKIVOC* **2016**, 110.
(m) Yang, Z.; Song, X.; Wei, Z.; Cao, J.; Liang, D.; Duan, H.; Lin, Y. *Tetrahedron Lett.* **2016**, *57*, 2410.
(n) Singh, M.; Yadav, A. K.; Yadav, L. D. S.; Singh, R. K. P. *Synlett* **2018**, *29*, 176.
- (a) Wang, Y.-F.; Lonca, G. H.; Runigo, M. L.; Chiba, S. *Org. Lett.* **2014**, *16*, 4272.
(b) Yang, J.-C.; Zhang, J.-J.; Guo, L.-N. *Org. Biomol. Chem.* **2016**, *14*, 9806.
(c) Sun, X.; Yu, S. *Chem. Commun.* **2016**, *52*, 10898.
(d) Mackay, E. G.; Studer, A. *Chem. Eur. J.* **2016**, *22*, 13455.
(e) Mao, L.-L.; Zheng, D.-G.; Zhu, X.-H.; Zhou, A.-X.; Yang, S.-D. *Org. Chem. Front.* **2018**, *5*, 232.
(f) Li, Y.; Zhu, Y.; Yang, S.-D. *Org. Chem. Front.* **2018**, *5*, 822.
(g) Yang, J.-C.; Zhang, J.-Y.; Zhang, J.-J.; Duan, X.-H.; Guo, L.-N. *J. Org. Chem.* **2018**, *83*, 1598.
- (a) Casellato, U.; Vidali, M.; Vigato, P. A. *Coord. Chem. Rev.* **1979**, *28*, 231.
(b) Nief, F. *Coord. Chem. Rev.* **1998**, 178.
(c) Arda, M.; Ozturk, I. I.; Banti, C. N.; Kourkoulis, N.; Manoli, M.; Tasiopoulos, A. J.; Hadjikakou, S. K. *RSC Adv.* **2016**, *6*, 29026.
(d) Ephritikhine, M. *Coord. Chem. Rev.* **2016**, *319*, 35.
(e) Boreen, M. A.; Parker, B. F.; Hohloch, S.; Skeel, B. A.; Arnold, J. *Dalton Trans.* **2018**, *47*, 96.
- (a) Beletskaya, I. P.; Ananikov, V. P. *Chem. Rev.* **2011**, *111*, 1596.
(b) Dorosuk, J.; Musiejuk, M.; Demkowicz, S.; Rachon, J.; Witt, D. *RSC Adv.* **2016**, *6*, 105449.
(c) Jiao, J.; Wei, L.; Ji, X.-M.; Hu, M.-L.; Tang, R.-Y. *Adv. Synth. Catal.* **2016**, *358*, 268.
(d) Dong, Z.-B.; Liu, X.; Bolm, C. *Org. Lett.* **2017**, *19*, 5916.
(e) Cao, Q.; Peng, H.-Y.; Cheng, Y.; Dong, Z.-B. *Synthesis* **2018**, *50*, 1527.
- (a) Fuchigami, T.; Chen, C.-S.; Nonaka, T.; Yen, M.-Y.; Tien, H.-J. *Bull. Chem. Soc. Jpn.* **1986**, *59*, 487.
(b) Gueyraud, D.; Tatibouët, A.; Gareau, Y.; Rollin, P. *Org. Lett.* **1999**, *1*, 521.
(c) Enders, D.; Rembiak, A.; Liebich, J. X. *Synthesis* **2011**, 281.
(d) Camerel, F.; Jeannin, O.; Yzambart, G.; Fabre, B.; Lorcy, D.; Fourmigué, M. *New J. Chem.* **2013**, *37*, 992.
(e) Zhang, L.; Zhu, J.; Ma, J.; Wu, L.; Zhang, W.-H. *Org. Lett.* **2017**, *19*, 6308.
- (a) Tan, J.; Guo, Y.; Zeng, F.; Chen, G.; Xie, L.; He, W. *Chin. J. Org. Chem.* **2018**, *38*, 1740.
(b) Wu, C.; Lu, L.-H.; Peng, A.-Z.; Jia, G.-K.; Peng, C.; Cao, Z.; Tang, Z.; He, W.-M.; Xu, X. *Green Chem.* **2018**, *20*, 3683.
- (a) Sibbald, P. A.; Michael, F. E. *Org. Lett.* **2009**, *11*, 1147.
(b) Albéniz, A. C.; Espinet, P.; López-Fernández, R.; Sen, A. *J. Am. Chem. Soc.* **2002**, *124*, 11278.
- Winterie, J. S.; Mill, T. *J. Am. Chem. Soc.* **1980**, *102*, 6336.
- Gareau, Y.; Beauchemin, A. *Heterocycles* **1998**, *48*, 2003.
- Curran, D. P.; Keller, A. *J. Am. Chem. Soc.* **2006**, *128*, 13706.
- Liu, K.-J.; Jiang, S.; Lu, L.-H.; Tang, L.-L.; Tang, S.-S.; Tang, H.-S.; Tang, Z.; He, W.-M.; Xu, X. *Green Chem.* **2018**, *20*, 3038.

(Lu, Y.)