从 *D*-半乳糖或 *L*-阿拉伯糖合成叠氮鞘氨醇

高阳光^{*,a} 曹 周^a 韩忠享^a 张 强^a 胡 杰^a
郭 锐^b 贺贤然^a 丁 菲^a 尤庆亮^c 张勇民^{*,a,d}

(^a 江汉大学交叉学科研究院 武汉 430056)

(^b 江汉大学环境健康研究院 武汉 430056)

(^c 江汉大学光电化学材料与器件教育部重点实验室 武汉 430056)

(^d 索邦大学分子化学巴黎研究所 巴黎 75005)

摘要 分别以 *D*-半乳糖、*L*-阿拉伯糖为原料完成了叠氮鞘氨醇的合成, 总收率分别为 27% 和 20%。此策略主要特征是烯烃复分解反应、自由基诱导的异构化、Appel 反应和叠氮取代。

关键词 鞘氨醇; 合成; 糖; 烯烃复分解; 取代

Synthesis of Azidosphingosine from *D*-Galactose or *L*-Arabinose

Gao, Yangguang^{*,a} Cao, Zhou^a Han, Zhongxiang^a Zhang, Qiang^a Hu, Jie^a
Guo, Rui^b He, Xianran^a Ding, Fei^a You, Qingliang^c Zhang, Yongmin^{*,a,d}

(^a Institute for Interdisciplinary Research, Jiangnan University, Wuhan 430056)

(^b Institute of Environment and Health, Jiangnan University, Wuhan 430056)

(^c Key Laboratory of Optoelectronic Chemical Materials and Devices of Ministry of Education, Jiangnan University, Wuhan 430056)

(^d Institut Parisien de Chimie Moléculaire, Sorbonne Université, Paris 75005)

Abstract Synthesis of azidosphingosine has been described from *D*-galactose and *L*-arabinose with an overall yields of 27% and 20%, respectively. Our strategy features olefin cross-metathesis reaction, radical-induced isomerization, appel reaction, and azido replacement.

Keywords sphingosine; synthesis; carbohydrate; olefin cross-metathesis; replacement

1 Introduction

Sphingosine, named [(2*S*,3*R*,4*E*)-2-amino-3-hydroxyoctadec-4-en-1-ol] (Figure 1), which is the active component of many sphingolipids, was firstly isolated from human brain by Thudichum in 1884.^[1,2] While sphingolipids are ubiquitously found in cell membrane of eukaryotic cells, and serve as versatile physiological functions.^[3–6] Furthermore, sphingosine itself displays powerful inhibition of protein kinase C and plays critical role in signal pathway.^[7,8] Since the significance of sphingosine and its deficiency in nature, synthetic chemistry community has made a great endeavor to efficiently prepare optically pure sphingosine and its derivatives.

The main challenges towards synthesis of sphingosine

have two points: one is to exquisitely set stereogenic centers at C2 and C3, another is to selectively construct *E* carbon-carbon double bond. A great number of papers about synthesis of sphingosine have been published in the past decades.^[9–13] A typical synthesis of sphingosine was achieved by Zimmermann and Schmidt in 1988.^[14] The highlight of their scheme was that *D*-galactose and *D*-xylose were respectively used as chiral precursor to provide 2,4-*O*-protected *D*-threoses, upon which was treatment of ylide salt in the presence of an excess of lithium bromide to exclusively yield *E*-enetriols. More recently, Kitamura *et al.*^[15] described a novel methodology on construction of 1,2-amino alcohol via a catalytic asymmetric dehydrative allylation, which was applied to synthesize sphingosine within only five steps.

* Corresponding authors. E-mail: sunlt413@jhu.edu.cn; yongmin.zhang@upmc.fr

Received August 31, 2018; revised September 20, 2018; published online October 19, 2018.

Project supported by the National Natural Science Foundation of China (No. 21602082).

国家自然科学基金(No. 21602082)资助项目.

Figure 1 Structure of sphingosine

In a continuation of our review in 2016,^[13] we have been working to develop a concise and satisfactory protocol to synthesize sphingosine and its derivatives. Azidosphingosine is not only a stable and important precursor of sphingosine, but also an excellent acceptor of glycosylation.^[16] Consequently, synthesis of azidosphingosine counts for much. Herein, we described a modified synthesis of azidosphingosine from *D*-galactose. Besides, we developed a novel protocol for the synthesis of azidosphingosine from *L*-arabinose. Though *D*-arabinose has been used as chiral pool to synthesize *D*-erythro-sphingosine in the previous literature,^[17] in which *E*-selective Wittig olefination was employed as key step. To the best of our knowledge, this is the first report about the synthesis of azidosphingosine from *L*-arabinose.

2 Results and discussion

Synthesis of azidosphingosine (**7**) was summarized in Scheme 1. As shown in Scheme 1, firstly, *D*-galactose was protected as benzylidene ether at C4 and C6 with benzaldehyde dimethyl acetal in the presence of *p*-toluenesulfonic acid (PTSA) to provide compound **1** in a good yield (85%). Subsequent oxidative cleavage by sodium periodate in methanol was smoothly carried out to afford 2,4-benzylidene-*D*-threose (**2**), which was directly subjected to the following Wittig olefination using the *in situ* generated methyl Wittig reagent to give compound **3** in 81% yield for 2 steps. Then, elongation of compound **3** was successfully achieved by olefin cross-metathesis (OCM) between compound **3** and 1-pentadecene (**3a**) under the catalysis of Grubbs II catalyst in dichloromethane (DCM) at reflux temperature to afford compound **4a** in a yield of 62%, and albeit with 10% yield of compound **4b**. To note, conversion of compound **2** into compound **4a** underwent 2 steps other than one step in the previous literature, but the advantage of 2 steps operation was that it was unnecessary to use excess strong base such as PhLi^[18] or BuLi-LiBr.^[14] The undesired compound **4b** could be converted into the *E*-olefin **4b** by radical-induced isomerization^[19] in a moderate yield (64%). Furthermore, when we repeated the same Wittig olefination according to the literature, no exclusive *E*-selectivity was observed, and the best result of the mixture of *E*,*Z*-alkenes in a ratio of 10 : 1 was given. Similar report can also be found in literature.^[20] According to a literature procedure,^[14,18] compound **4a** was initially converted to its triflate followed by substitution with NaN₃ in dimethylformamide (DMF). To our surprise, beside the desired azido product **5**, 38% yield of elimination product compound **6** was also obtained. The possible reason for formation of elimination product can be explained as following: The triflate of compound **4a** may underwent E1

elimination in polar solvent (DMF) in the presence of weak base (azide anion) to give vinyl benzylidene ether, and one equivalent of triflic acid was released. At the same time, vinyl benzylidene ether was unstable under acidic condition. Thus, the benzylidene group was removed to give enol compound, which then tautomerize to its keto form, *i.e.* compound **6**. Because OTf usually works as a good leaving group, a less active group OMs was utilized instead. Unfortunately, subsequent replacement by NaN₃ proved to be problematic, which only gave 40% yield of compound **5** for 2 steps. That was in good accordance with the previous result.^[21] We were pleased to find when compound **4a** was conducted under Mitsunobu's condition,^[22] the above problems such as elimination or low yield could be circumvented, and compound **5** was obtained with 72% yield. Treatment of compound **5** with AcCl in methanol provided azidosphingosine (**7**) in 89% yield. The overall yield of 27% over 6 steps is superior to 7% overall yield over 6 steps in the previous literature.^[14]

Reagents and conditions: (a) PhCH(OMe)₂, PTSA, DMF (dry), 85%; (b) NaIO₄, MeOH, H₂O; (c) CH₃Ph₃P⁺·Br[−], *n*-BuLi, THF, 81% for 2 steps; (d) 1-pentadecene (**3a**), Grubbs II catalyst, 62% for **4a**, 10% for **4b**; (e) *p*-toluenethiol, AIBN, toluene, reflux, 64%; (f) Ph₃P, DEAD, DPPA, THF (dry), 72%; (g) (i) Tf₂O, DCM, pyridine (dry), −20 °C to r.t.; (ii) NaN₃, DMF (dry), 45% for **5**, 38% for **6**; (h) (i) MsCl, DCM, Et₃N, −30 °C to r.t.; (ii) NaN₃, DMF, 90 °C, 40%; (i) 1% AcCl in methanolic solution, 89%.

Scheme 1 Synthesis of azidosphingosine (**7**)

The second-generation synthesis of azidosphingosine (**7**) was depicted in Scheme 2. *L*-Arabinose was converted into

compound **8** according to the previously reported procedure.^[19] Unblocking of compound **8** in acidic solution followed by regioselective protection as benzylidene ether provided compound **9** in 85% yield for 2 steps. Initially, compound **9** was subjected to olefin cross-metathesis (OCM) with 1-pentadecene (**3a**) under the catalysis of Grubbs II catalyst. To our surprise, poor *E*-selectivity in CM reaction was observed, and the mixture of *Z,E*-alkenes was inseparable by routine column chromatography. Therefore, as a candidate route, Appel reaction^[23] of compound **9** was carried out with I_2 , PPh_3 , imidazole in tetrahydrofuran (THF) to give a 91% yield of compound **11**. Unfortunately, subsequent azido replacement of iodo product **11** by NaN_3 proved to be unsuccessful. Elimination product was obtained unexpectedly. To overcome the problem, we envisaged that elongation of the double bond might suppress the elimination due to the increasing electronic effect of the olefin. Thus, olefin cross-metathesis (OCM) between compound **11** and 1-pentadecene (**3a**) was executed to furnish compound **13** in 90% yield. To note, better *E*-selectivity (*E/Z*=10/1) was observed in OCM reaction than its counterpart compound **3** or **9**. Besides, compound **13** can be more easily separated from its *cis* byproduct. However, the following replacement by NaN_3 in DMF failed to provide compound **14**, and the elimination product compound **6** was still obtained instead. At this stage, we began to doubt that the presence of benzylidene group at C1 and C3 may promote the elimination of hydrogen at allylic position due to its conjugated effect. As expected, removal of benzylidene group followed by azido substitution gave azidosphingosine (**7**) successfully in a yield of 82%. The overall yield was 20% over 8 independent manipulations. The spectral and analytical data (1H NMR, ^{13}C NMR, HRMS and optical rotation values) of synthetic azidosphingosine were in good accordance with

the previously reported literatures.

3 Conclusions

In summary, a modified synthesis of azidosphingosine from *D*-galactose has been achieved, besides, a novel synthesis of azidosphingosine from *L*-arabinose has been developed. The overall yields were 27% and 20%, respectively, and both were better than that of the previous literature. Both routes underwent a similar benzylidene-protected enetriol, and OCM reaction was utilized as key step to elong the carbon-carbon double bond. The disfavored elimination during azido replacement was reported for the first time, and the reason for formation of elimination product was also presumed. Our protocol is flexible and effective, which could provide alternative syntheses of sphingosine and its diastereoisomers using chiral template. And these results will be reported in due course.

4 Experimental section

4.1 Instruments and reagents

All reagents were commercially available and used directly without further purification unless otherwise stated. Column chromatography was carried out by using silica gel (100~200 mesh). Routine monitoring of reactions was used silica gel60 F254 TLC plates. Dried THF was distilled from Na and benzophenone. DMF, DCM were treated with CaH_2 . All air-sensitive reactions were run under a nitrogen atmosphere. Purification by column chromatography was using 100~200 mesh silica gel. Reactions were monitored by TLC using silica gel 60 F254 plates. 1H NMR and ^{13}C NMR spectra were recorded with a Bruker Avance III spectrometer at 400 and 100 MHz respectively relative to Me_4Si ($\delta=0$) as internal standard. HRMS were measured with a ThermoFisher Q-Exactive mass spec-

Reagents and conditions: (a) (i) PTSA, MeOH; (ii) $PhCH(OMe)_2$, CH_3CN , 85% yield for 2 steps; (b) 1-pentadecene, Grubbs II catalyst, 65% yield for **10**, 90% yield for **13**; (c) I_2 , PPh_3 , imidazole, THF, 91% yield; (d) NaN_3 , DMF, 50 °C, 75% yield for **7**; (e) PTSA, MeOH, 82% yield.

Scheme 2 Second-generation synthesis of azidosphingosine (**7**)

trometer. IR spectra were recorded on neat sample with a NICOLET iS 10 infrared spectrometer. Optical rotations were recorded with a AUTOPOL IV automatic polarimeter.

4.2 Experimental method

4.2.1 Synthesis of 4,6-*O*-benzylidene- α/β -D-galactose (1)

D-Galactose (9.00 g, 0.05 mol) was dissolved in 100 mL of dried *N,N*-dimethyl formide (DMF). To the resulting solution was added benzaldehyde dimethyl acetal (12 mL, 0.08 mol), *p*-toluenesulfonic acid (PTSA) (1.00 g, 5.80 mmol) in sequence. The mixture was allowed to be heated at 50 °C, then 1 mL of Et₃N was added to quench the reaction until the starting material was thoroughly consumed (monitored by TLC). The solution was concentrated *in vacuo* and column chromatography of the residue on silica gel (EtOAc-MeOH, *V*: *V*=10:1) afforded a mixture of α/β -1 (α/β =5/1 analyzed by ¹H NMR) as a white solid (11.39 g, 85%). m.p. 180~183 °C (Lit.^[18] m.p. 171~173 °C); [α]_D²⁵ 122 (*c* 1.0, MeOH) [Lit.^[18] [α]_D²⁵ 132 (*c* 1.0, MeOH)]. For α -1: ¹H NMR (DMSO-*d*₆, 400 MHz) δ : 7.45 (d, *J*=5.2 Hz, 2H), 7.35 (d, *J*=5.6 Hz, 3H), 6.30 (d, *J*=4.4 Hz, 1H), 5.52 (s, 1H), 5.04 (t, *J*=3.6 Hz, 1H), 4.65 (d, *J*=6.4 Hz, 1H), 4.48 (d, *J*=6.4 Hz, 1H), 4.11 (d, *J*=3.2 Hz, 1H), 4.03 (m, 1H), 3.95 (d, *J*=15.6 Hz, 1H), 3.76~3.73 (m, 2H), 3.62~3.58 (m, 1H); ¹³C NMR (DMSO-*d*₆, 100 MHz) δ : 138.9, 128.7, 128.0, 126.4, 99.8, 93.2, 76.8, 69.0, 68.5, 67.8, 62.1; IR (neat) ν : 3392, 1400, 1167, 1093, 1048, 976 cm⁻¹. For β -1: ¹H NMR (DMSO-*d*₆, 400 MHz) δ : 7.45 (d, *J*=5.2 Hz, 2H), 7.35 (d, *J*=5.6 Hz, 3H), 6.61 (d, *J*=7.2 Hz, 1H), 5.54 (s, 1H), 4.82 (t, *J*=5.6 Hz, 1H), 4.76 (d, *J*=6.4 Hz, 1H), 4.33 (t, *J*=7.2 Hz, 1H), 4.00~4.10 (m, 3H), 3.45 (br, 1H), 3.40~3.46 (m, 1H), 3.28~3.34 (m, 1H); ¹³C NMR (DMSO-*d*₆, 100 MHz) δ : 138.9, 128.7, 128.0, 126.4, 99.8, 97.3, 76.2, 72.1, 71.7, 68.9, 65.9; HRMS (ESI, negative) calcd for C₁₅H₂₃O₆ [M-H]⁻ 299.1495, found 299.1480.

4.2.2 Synthesis of 2,4-*O*-benzylidene-D-threose (2)

Compound 1 (2.00 g, 7.46 mmol) was completely dissolved in 30 mL of methanol under sonication condition, then the solution was placed on an ice bath. To the solution was added sodium periodate (2.40 g, 11.22 mmol) in water (30 mL) via dropping funnel. The resulting solution was stirred at ambient temperature till the starting material thoroughly disappeared. The undissolved white solid was filtered, and the filtrate was extracted with DCM (50 mL $\times$ 3), combined the organic phase, concentrated, and the yellow residue was directly subjected to the next step without further purification.

4.2.3 Synthesis of (2*R*,3*R*)-1,3-*O*-benzylidene-4-pentene-1,2,3-triol (3)

To a suspension of methyltriphenylphosphonium bromide (5.00 g, 14.00 mmol) in anhydrous THF (30 mL) was added *n*-BuLi solution (7.8 mL, 2.2 mol·L⁻¹ in *n*-hexane, 17.16 mmol) dropwise at 0 °C under a nitrogen atmos-

phere. The mixture was turned to yellow immediately, 45 min later, to the yellow suspension was gradually added a solution of crude compound 2 (2.00 g, 9.60 mmol) in 10 mL of THF. Saturated ammonia chloride solution (20 mL) was added to quench the reaction until the completion of the reaction, then the organic layer was separated, and extracted with EtOAc (25 mL $\times$ 3), dried over anhydrous Na₂SO₄, and the organic layer was concentrated. The residue was purified by silica gel column chromatography (petroleum ether-EtOAc, *V*: *V*=6:1) to furnish compound 3 (1.60 g, 81% for 2 steps) as a white powder. m.p. 40~42 °C; [α]_D²⁵ +7.6 (*c* 0.96, MeOH); ¹H NMR (CDCl₃, 400 MHz) δ : 7.54 (d, *J*=7.6 Hz, 2H), 7.38 (d, *J*=6.8 Hz, 3H), 5.99 (ddd, *J*=16.8, 10.8, 5.2 Hz, 1H), 5.64 (s, 1H), 5.46 (d, *J*=17.6 Hz, 1H), 5.33 (d, *J*=10.8 Hz, 1H), 4.47 (d, *J*=4.8 Hz, 1H), 4.24 (d, *J*=12.0 Hz, 1H), 4.09 (d, *J*=11.6 Hz, 1H), 3.58 (s, 1H), 2.76 (br, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ : 137.8, 134.4, 129.1, 128.3, 126.1, 117.7, 101.3, 80.4, 72.4, 65.9; IR (neat) ν : 3281, 1451, 1397, 1132, 1092, 1022, 925 cm⁻¹; HRMS (ESI, negative) calcd for C₁₂H₁₃O₃ [M-H]⁻ 205.0865, found 205.0910.

4.2.4 Synthesis of 1-pentadecene (3a)

To a solution of 1-bromopentadecane (1.00 g, 3.43 mmol) in 10 mL of THF (dried) was added *t*-BuOK (1.12 g, 10.00 mmol), the solution was stirred at room temperature until the reaction completed (monitored by TLC). Saturated ammonia chloride solution was poured into the above solution to quench the reaction, then extracted with EtOAc (20 mL $\times$ 3), dried over anhydrous Na₂SO₄, and concentrated under reduced pressure. Column chromatography (pure petroleum ether) of the residue gave compound 3a as a colorless liquid (0.72 g, 99%). ¹H NMR (CDCl₃, 400 MHz) δ : 5.87~5.77 (m, 1H), 5.99 (dd, *J*=17.2, 2.0 Hz, 1H), 4.93 (dd, *J*=10.4, 0.8 Hz, 1H), 2.04 (q, *J*=7.2 Hz, 2H), 1.40~1.34 (m, 2H), 1.40~1.31 (m, 20H), 0.89 (t, *J*=7.2 Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz) δ : 139.4, 114.1, 33.9, 32.0, 29.8, 29.6, 29.4, 29.2, 29.0, 22.8, 14.2; IR (neat) ν : 3077, 2950, 1641, 1470, 1361 cm⁻¹; HRMS (ESI, positive) calcd for C₁₅H₃₀Na [M+Na]⁺ 233.2245, found 233.2239.

4.2.5 Synthesis of (2*R*,3*R*,4*E*)-1,3-*O*-benzylidene-4-octadecene-1,2,3-triol (4a) and (2*R*,3*R*,4*Z*)-1,3-*O*-benzylidene-4-octadecene-1,2,3-triol (4b)

To the mixture of compound 3 (600 mg, 2.91 mmol) and 3a (920 mg, 4.38 mmol) in dried DCM (10 mL) under a nitrogen atmosphere was added a catalyzed amount of Grubbs II catalyst (123 mg, 0.15 mmol). The solution was allowed to be heated at reflux temperature for 8 h, then concentrated under reduced pressure. The residue was purified by column chromatography (toluene-EtOAc, *V*: *V*=50:1) to give compound 4a as a white solid (701 mg, 62%), and compound 4b as a colorless liquid (113 mg, 10%). For 4a: m.p. 52~54 °C (Lit.^[18] m.p. 55~56 °C); [α]_D²⁵ -3.3 (*c* 0.487, CHCl₃) [Lit.^[17] [α]_D²² -3.8 (*c* 0.5, CHCl₃)]; ¹H NMR (CDCl₃, 400 MHz) δ : 7.52 (d, *J*=6.8 Hz, 2H), 7.37 (d, *J*=6.8 Hz, 3H), 5.87 (dt, *J*=15.6, 14 Hz,

1H), 5.66 (dd, $J=15.2, 6.0$ Hz, 1H), 5.62 (s, 1H), 4.41 (d, $J=6.0$ Hz, 1H), 4.25 (d, $J=12.0$ Hz, 1H), 4.08 (d, $J=11.6$ Hz, 1H), 3.54 (d, $J=10.4$ Hz, 1H), 2.64 (d, $J=10.4$ Hz, 1H), 2.08 (q, $J=6.8$ Hz, 2H), 1.40~1.38 (m, 2H), 1.26 (m, 20H), 0.88 (t, $J=6.0$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 137.9, 135.3, 129.1, 128.3, 126.1, 126.0, 101.5, 80.7, 72.5, 66.4, 32.5, 32.0, 29.7, 29.6, 29.5, 29.4, 29.3, 29.0, 22.7, 14.2; IR (neat) ν : 2916, 2849, 1464, 1395, 1364, 1127, 1082, 1035, 1012, 995 cm^{-1} ; HRMS (ESI, negative) calcd for $\text{C}_{25}\text{H}_{39}\text{O}_3$ $[\text{M}-\text{H}]^-$ 387.2899, found 387.2890. For **4b**: $[\alpha]_{\text{D}}^{25} +37.7$ (c 0.66, CHCl_3); ^1H NMR (CDCl_3 , 400 MHz) δ : 7.52 (d, $J=6.8$ Hz, 2H), 7.37 (d, $J=7.2$ Hz, 3H), 5.69 (dd, $J=9.6, 4.8$ Hz, 2H), 5.66 (s, 1H), 4.73 (d, $J=6.0$ Hz, 1H), 4.25 (d, $J=11.6$ Hz, 2H), 4.12 (d, $J=12.0$ Hz, 1H), 3.50 (s, 1H), 2.15~2.10 (m, 2H), 1.43~1.39 (m, 2H), 1.26 (m, 20H), 0.88 (t, $J=6.8$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 137.9, 135.0, 129.1, 128.4, 126.1, 125.7, 101.5, 76.5, 72.6, 66.6, 32.0, 29.8, 29.7, 29.7, 29.6, 29.5, 29.4, 28.4, 22.8, 14.2; IR (neat) ν : 3032, 2920, 2850, 1452, 1386, 1050, 698 cm^{-1} .

4.2.6 Isomerization of **4b** into **4a**

To a solution of compound **4b** (100 mg, 0.26 mmol) in anhydrous toluene (8 mL) was added *p*-toluenethiol (320 mg, 2.58 mmol), AIBN (424 mg, 2.58 mmol) under a nitrogen atmosphere. The mixture was heated at reflux for 8 h, then concentrated in vacuo. The residue was purified by column chromatography (toluene-EtOAc, $V:V=50:1$) to give compound **4a** (64 mg, 64%), and recovered 22 mg of compound **4b**.

4.2.7 Synthesis of (2*S*,3*R*,4*E*)-2-azido-1,3-*O*-benzylidene-4-octa-decene-1,3-diol (**5**)

Method A: To a solution of compound **4a** (1.00 g, 2.58 mmol) in 10 mL of dried THF under a nitrogen atmosphere was added PPh_3 (2.70 g, 10.30 mmol), diethyl azodiformate (DEAD) (1.55 mL, 10.3 mmol), diphenyl azidophosphate (DPPA) (0.22 mL, 10.3 mmol) in sequence. The mixture was stirred at room temperature until the completion of the reaction, concentrated. The residue was purified by column chromatography (petroleum ether-EtOAc, $V:V=50:1$) to give compound **5** as a white solid (766 mg, 72%). m.p. 41~43 °C (Lit.^[18] m.p. 43 °C); $[\alpha]_{\text{D}}^{25} -7.2$ (c 0.08, CHCl_3) [Lit.^[17] $[\alpha]_{\text{D}}^{20} -11.7$ (c 3.00, CHCl_3); ^1H NMR (CDCl_3 , 400 MHz) δ : 7.49~7.47 (m, 2H), 7.32~7.38 (m, 3H), 5.98 (dt, $J=15.4, 6.7$ Hz, 1H), 5.59 (dd, $J=15.4, 7.4$ Hz, 1H), 5.49 (s, 1H), 4.34 (dd, $J=11.1, 5.2$ Hz, 1H), 4.06 (dd, $J=9.5, 7.4$ Hz, 1H), 3.61 (dd, $J=11.1, 10.8$ Hz, 1H), 3.47 (ddd, $J=10.8, 9.5, 5.2$ Hz, 1H), 2.11 (dt, $J=7.2, 6.7$ Hz, 2H), 1.42 (q, $J=7.2$ Hz, 2H), 1.24~1.35 (m, 20H), 0.88 (t, $J=6.9$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 137.9, 137.4, 129.2, 128.4, 126.3, 125.9, 101.2, 81.9, 69.2, 57.6, 32.6, 32.0, 29.8, 29.7, 29.6, 29.5, 29.3, 28.8, 22.8, 14.2; IR (neat) ν : 2922, 2853, 2107, 1457, 1258, 1010, 788 cm^{-1} ; HRMS (ESI, positive) calcd for $\text{C}_{25}\text{H}_{40}\text{N}_3\text{O}_2$ $[\text{M}+\text{H}]^+$ 414.3121, found 414.3146.

Method B: To a solution of compound **4a** (150 mg, 0.39 mmol) in 6 mL of dried DCM was added 0.6 mL of dried

pyridine and Ti_2O (0.15 mL, 0.89 mmol) at -20 °C under a nitrogen atmosphere. After the disappearance of the starting material, the solution was concentrated in vacuo. NaN_3 (140 mg, 2.15 mmol) was added to the above residue followed by addition of DMF (3 mL). The mixture was allowed to stir at room temperature until the reaction completed, filtered the solid, the filtrate was concentrated in vacuo. Column chromatography (petroleum ether-EtOAc, $V:V=50:1$) of the residue provided compound **5** as a white powder (71.8 mg, 45%), compound **6** as a colorless liquid (41 mg, 38%). For **6**: ^1H NMR (CDCl_3 , 400 MHz) δ : 6.88 (dt, $J=15.6, 14.8$ Hz, 1H), 6.10 (d, $J=16.0$ Hz, 1H), 3.89 (t, $J=4.8$ Hz, 2H), 2.81 (t, $J=5.2$ Hz, 2H), 2.22 (q, $J=7.2$ Hz, 2H), 1.48~1.44 (m, 2H), 1.25 (m, 20H), 0.88 (t, $J=6.4$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 201.0, 149.2, 130.5, 58.2, 41.3, 32.6, 32.0, 29.8, 29.7, 29.7, 29.6, 29.5, 29.4, 29.3, 28.1, 22.8, 14.2; HRMS (ESI, positive) calcd for $\text{C}_{18}\text{H}_{34}\text{O}_2\text{Na}$ $[\text{M}+\text{Na}]^+$ 305.2457, found 305.2478.

Method C: To a solution of compound **4a** (350 mg, 0.90 mmol) in 8 mL of dried DCM was added 0.62 mL of Et_3N and MsCl (0.16 mL, 1.80 mmol) at -30 °C under a nitrogen atmosphere. After the disappearance of the starting material, the solution was concentrated under reduced pressure. The resulting white powder was dissolved in 10 mL of dried DMF, then NaN_3 (292 mg, 4.50 mmol) was added. The suspension was heated at 90 °C for 3 d till the starting material was consumed. 10 mL of water was poured into the suspension, extracted with petroleum ether, combined the organic layer, and evaporated. The residue was purified by column chromatography (petroleum ether-EtOAc, $V:V=50:1$) to furnish compound **5** (149 mg, 40%).

4.2.8 Synthesis of (2*S*,3*R*,4*E*)-2-azido-4-octadecene-1,3-diol (azidosphingosine) (**7**)

Method A: To a solution of compound **5** (800 mg, 1.93 mmol) in methanol (4 mL) was added AcCl (40 μL). The solution was allowed to stir at room temperature until the consumption of the starting material. The reaction was quenched by addition of 0.1 mL of Et_3N , then concentrated. Column chromatography (petroleum ether-EtOAc, $V:V=2:1$) of the residue provided compound **7** as a white solid (560 mg, 89%). m.p. 52~54 °C (Lit.^[18] m.p. 52~54 °C); $[\alpha]_{\text{D}}^{25} -28.6$ (c 0.30, CHCl_3) [Lit.^[18] $[\alpha]_{\text{D}}^{24} -33.0$ (c 4.00, CHCl_3); ^1H NMR (CDCl_3 , 400 MHz) δ : 5.82 (dt, $J=15.6, 6.4$ Hz, 1H), 5.53 (dd, $J=15.2, 7.2$ Hz, 1H), 4.25 (t, $J=6.4$ Hz, 1H), 3.78 (d, $J=5.1$ Hz, 2H), 3.50 (q, $J=5.2$ Hz, 1H), 2.07 (dt, $J=6.6$ Hz, 2H), 1.20~1.40 (m, 22H), 0.88 (t, $J=6.4$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 136.2, 128.1, 73.9, 66.8, 62.7, 32.4, 32.0, 29.8, 29.8, 29.7, 29.6, 29.5, 29.3, 29.0, 22.8, 14.2; IR (neat) ν : 3312, 2916, 2849, 2140, 2094, 1465, 1363, 1273, 998, 965 cm^{-1} . HRMS (ESI, negative) calcd for $\text{C}_{18}\text{H}_{34}\text{N}_3\text{O}_2$ $[\text{M}-\text{H}]^-$ 324.265, found 324.2657.

Method B: To a solution of compound **15** (100 mg, 0.24 mmol) in DMF (3 mL) was added NaN_3 (159 mg, 2.44

mmol). The mixture was heated at 50 °C till the starting material was thoroughly consumed. The solvent was concentrated in vacuo, the residue was added 10 mL of DCM, filtered. The filtrate was concentrated. Column chromatography (petroleum ether-EtOAc, $V:V=2:1$) of the residue provided compound **7** as a white solid (59 mg, 75%).

4.2.9 Synthesis of (2*S*,3*R*)-1,3-*O*-benzylidene-4-pentene-1,2,3-triol (**9**)

To a solution of compound **8** (500 mg, 3.16 mmol) in 10 mL of MeOH was added PTSA (60 mg, 0.32 mmol). The mixture was allowed to stir at room temperature, and the solvent was removed under reduced pressure till the completion of the reaction. The residue was dissolved in 10 mL of CH₃CN, then, benzaldehyde dimethyl acetal (0.6 mL, 4.00 mmol) was added. 2 mL of Et₃N was added to quench the reaction until the starting material completely disappeared (monitored by TLC). The mixture was concentrated, column chromatography of the residue (petroleum ether-EtOAc, $V:V=50:1$) gave compound **9** as a white solid (554 mg, 85%). m.p. 82~85 °C; $[\alpha]_D^{25} +50.9$ (c 0.37, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ : 7.52~7.49 (m, 2H), 7.37~7.35 (m, 3H), 5.99 (ddd, $J=17.2, 10.4, 6.8$ Hz, 1H), 5.55 (s, 1H), 5.49 (dt, $J=17.2, 1.2$ Hz, 1H), 5.37 (dd, $J=10.8, 1.2$ Hz, 1H), 4.36~4.34 (m, 1H), 4.03 (t, $J=7.6$ Hz, 1H), 3.69~3.63 (m, 2H), 1.84 (d, $J=3.2$ Hz, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ : 137.6, 134.6, 129.1, 128.4, 126.3, 119.4, 101.0, 83.5, 70.9, 65.3; IR (neat) ν : 3435, 2969, 2929, 1458, 1397, 1248, 1218, 1057, 1009, 918, 761, 698 cm⁻¹; HRMS (ESI, positive) calcd for C₁₂H₁₄O₃Na [M+Na]⁺ 229.084, found 229.0828.

4.2.10 Synthesis of (2*R*,3*R*)-1,3-*O*-benzylidene-2-iodo-4-pentene-1,3-diol (**11**)

To a solution of compound **9** (700 mg, 3.39 mmol) in 20 mL of THF (dried) was added I₂ (1.03 g, 4.06 mmol), PPh₃ (1.34 g, 5.10 mmol), imidazole (0.69 g, 10.13 mmol) in sequence. The mixture was heated at reflux temperature for 5 h. Saturated Na₂S₂O₃ solution was added to quench the reaction, extracted with EtOAc (20 mL×3), dried over anhydrous Na₂SO₄ and concentrated. The residue was purified by column chromatography (petroleum ether-EtOAc, $V:V=6:1$) to afford compound **11** as a white powder (976 mg, 91%). m.p. 66~73 °C; $[\alpha]_D^{25} -4$ (c 0.5, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ : 7.60~7.57 (m, 2H), 7.40~7.38 (m, 3H), 5.78 (ddd, $J=17.2, 10.4, 4.8$ Hz, 1H), 5.64 (s, 1H), 5.36 (dt, $J=17.2, 1.2$ Hz, 1H), 5.31 (dt, $J=10.4, 1.6$ Hz, 1H), 4.48 (dd, $J=12.8, 1.6$ Hz, 1H), 4.36 (dd, $J=12.8, 2.0$ Hz, 1H), 4.17 (q, $J=1.6$ Hz, 1H), 3.77 (ddd, $J=4.8, 3.2, 1.6$ Hz, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ : 138.3, 137.7, 129.3, 128.4, 126.6, 117.3, 102.3, 79.3, 74.1, 32.0; IR (neat) ν : 2834, 1448, 1391, 1358, 1296, 1112, 1079, 974, 750, 695 cm⁻¹; HRMS (ESI, positive) calcd for C₁₂H₁₃IO₂Na [M+Na]⁺ 338.9858, found 338.9839.

4.2.11 Synthesis of (2*R*,3*R*,4*E*)-1,3-*O*-benzylidene-2-iodo-4-octa-decene-1,3-diol (**13**)

To a mixture of compound **11** (600 mg, 1.90 mmol) and

1-pentadecene (**3a**) (800 mg, 3.80 mmol) was added Grubbs II catalyst (81 mg, 0.09 mmol). The solution was heated at reflux temperature. The solvent was removed until the starting material was thoroughly consumed. Column chromatography (petroleum ether-EtOAc, $V:V=10:1$) of the residue provided compound **13** as a white powder (852 mg, 90%). m.p. 52~57 °C; $[\alpha]_D^{25} -19.2$ (c 0.25, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ : 5.78 (dt, $J=14.8, 13.2$ Hz, 1H), 5.62 (s, 1H), 5.45 (ddt, $J=15.6, 5.6, 1.6$ Hz), 4.47 (dd, $J=12.8, 1.2$ Hz, 1H), 4.35 (dd, $J=12.8, 2.0$ Hz, 1H), 4.14 (d, $J=1.6$ Hz, 1H), 3.65 (d, $J=5.6$ Hz, 1H), 2.05 (q, $J=7.2$ Hz, 2H), 1.41~1.37 (m, 2H), 1.25 (m, 20 H), 0.88 (t, $J=6.8$ Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz) δ : 137.8, 134.7, 130.6, 129.3, 128.4, 126.7, 102.5, 79.5, 74.2, 33.5, 32.3, 32.0, 29.8, 29.7, 29.5, 29.4, 29.3, 28.8, 22.8, 14.2; IR (neat) ν : 2915, 2848, 1463, 1391, 1293, 1125, 1075, 1000, 956, 755, 699 cm⁻¹; HRMS (ESI, positive) calcd for C₂₅H₃₉IO₂Na [M+Na]⁺ 521.1893, found 521.1873.

4.2.12 Synthesis of (2*R*,3*R*,4*E*)-2-iodo-4-octa-decene-1,3-diol (**15**)

Compound **13** (500 mg, 1.00 mmol) was dissolved in 10 mL of MeOH. To the solution was added PTSA (10 mg, 0.05 mmol). 0.1 mL of Et₃N was added to neutralize the acid after the completion of the reaction, then the solvent was removed in vacuo. The residue was purified by column chromatography (petroleum ether-EtOAc, $V:V=2:1$) to afford compound **15** as an amorphous powder (337 mg, 82%). $[\alpha]_D^{25} -4.7$ (c 0.3, CHCl₃); ¹H NMR (CDCl₃, 400 MHz) δ : 5.78 (dt, $J=15.2, 14.0$ Hz, 1H), 5.46 (dd, $J=15.6, 6.0$ Hz, 1H), 4.25~4.21 (m, 1H), 3.94~3.89 (m, 3H), 2.35 (br, 1H), 2.21 (d, $J=5.6$ Hz, 1H), 2.05 (q, $J=7.2$ Hz, 2H), 1.40~1.34 (m, 2H), 1.25 (m, 20H), 0.87 (t, $J=6.8$ Hz, 3H); ¹³C NMR (CDCl₃, 100 MHz) δ : 134.8, 130.3, 73.2, 66.7, 53.5, 46.3, 32.7, 32.0, 29.8, 29.8, 29.7, 29.5, 29.2, 29.0, 22.8, 14.2; IR (neat) ν : 3259, 2916, 2848, 1473, 1117, 1067, 1038, 965, 719 cm⁻¹; HRMS (ESI, positive) calcd for C₁₈H₃₅IO₂Na [M+Na]⁺ 433.1580, found 433.1559.

Supporting Information ¹H NMR and ¹³C NMR spectra for compounds **1**, **3**, **3a**, **4a**, **4b**, **5**~**7**, **9**, **11**, **13** and **15**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] Thudichum, J. L. W. *A Treatise on the Chemical Constitution of the Brain*, Bailliere, Tindall, and Cox, London, 1884.
- [2] Du, H. W.; Chen, J. X.; Hao, J. S.; Pan, J. G.; Xia, C. Z. *Chem. J. Chin. Univ.* **1999**, 20, 590.
- [3] Chang, Y. T.; Choi, J.; Ding, S.; Prieschl, E. E.; Baumruker, T.; Lee, J. M.; Chung, S. K.; Schultz, P. G. *J. Am. Chem. Soc.* **2002**, 124, 1856.
- [4] Merrill, A. H. Jr.; Sandhoff, K. In *Biochemistry of Lipids, Lipoproteins and Membranes*, Eds.: Vance, D. E.; Vance, J. E., Elsevier, New York, **2002**, p. 373.
- [5] Lee, J. Y.; Hannun, Y. A.; Obeid, L. M. *J. Biol. Chem.* **1996**, 271, 13169.

- [6] Hannun, Y. A. *J. Biol. Chem.* **1994**, *269*, 3125.
[7] Hannun, Y. A.; Bell, R. M. *Science* **1989**, *243*, 500.
[8] Merrill, A. H. *J. Bioenerg. Biomembr.* **1991**, *23*, 83.
[9] Howell, A. R.; Ndakala, A. J. *Curr. Org. Chem.* **2002**, *6*, 365.
[10] Koskinen, P. M.; Koskinen, A. M. P. *Synthesis* **1998**, 1075.
[11] Morales-Serna, J. A.; Llaveria, J.; Díaz, Y.; Matheu, M. I.; Castillón, S. *Curr. Org. Chem.* **2010**, *14*, 2483.
[12] Liao, J. Y.; Tao, J. H.; Lin, G. Q.; Liu, D. G. *Tetrahedron* **2005**, *61*, 4715.
[13] Gao, Y. G.; He, X. R.; Ding, F.; Zhang, Y. M. *Synthesis* **2016**, *48*, 4017.
[14] Zimmermann, P.; Schmidt, R. R. *Liebigs Ann. Chem.* **1988**, 663.
[15] Vankar, Y. D.; Schmidt, R. R. *Chem. Soc. Rev.* **2000**, *29*, 201.
[16] Schmidt, R. R.; Zimmermann, P. *Tetrahedron Lett.* **1986**, *27*, 481.
[17] Tanaka, S.; Gunasekar, R.; Tanaka, T.; Iyoda, Y.; Suzuki, Y.; Kitamura, M. *J. Org. Chem.* **2017**, *82*, 9160.
[18] Duclos Jr, R. I. *Chem. Phys. Lipids* **2001**, *111*, 111.
[19] Gao, Y. G.; Cao, Z.; Su, C. Z.; Chen, Z. F.; He, X. R.; Ding, F.; Li, H.; Zhang, Y. M. *Synthesis* **2016**, *48*, 4471.
[20] Ohashi, K.; Yamagiwa, Y.; Kamikawa, T.; Kates, M. *Tetrahedron Lett.* **1988**, *29*, 1185.
[21] Compostella, F.; Franchini, L.; Panza, L.; Prosperi, D.; Ronchetti, F. *Tetrahedron* **2002**, *58*, 4425.
[22] (a) Niu, Y.; Cao, X.; Ye, X. S. *Helv. Chim. Acta* **2008**, *91*, 746.
(b) Jacqueline, E. M.; Krzysztof, J.; Philip, J. K.; Jorge, A. *Chem. Commun.* **2002**, 426.
[23] (a) Appel, R. *Angew. Chem.* **1975**, *87*, 863.
(b) Denton, R. M.; An, J.; Adeniran, B.; Blake, A. J.; Lewis, W.; Poulton, A. M. *J. Org. Chem.* **2011**, *76*, 6749.

(Li, L.; Fan, Y.)