一类基于 1,4,7,10-四氮杂环十二烷-1,4,7,10-四乙酸-酰肼结构的新型 含钆磁共振对比剂的设计、合成及性能表征

孙宏顺^{*,a,b} 周进^{a,b} 李玉龙^a 蒋蕪^a 张艳^c
王建强^b 郭成^b 沈临江^{b,d}

(^a 南京科技职业学院 江苏省磁共振靶向显像剂工程实验室 南京 210048)

(^b 南京工业大学化学与分子工程学院 南京 211816)

(^c 盐城工业职业技术学院纺织服装学院 盐城 224005)

(^d 南京工业大学数理科学学院 南京 211816)

摘要 磁共振成像技术被广泛应用于诊断医学和软组织成像,而磁共振对比剂有助于提高成像对比度。报道了一类共十二种基于钆-1,4,7,10-四氮杂环十二烷-1,4,7,10-四乙酸(DOTA)-酰肼结构的新型磁共振对比剂的设计、合成及性能表征。0.5 T 磁场下测得的弛豫率结果显示,对比剂 **5d**、**5h** 和 **5l** 的纵向弛豫率优于临床使用对比剂 Gd-DOTA,分别达到 4.67、4.85 和 5.33 L·mmol⁻¹·s⁻¹。进一步动物活体体内肝靶向磁共振成像研究显示,对比剂 **5d** 具有作为肝靶向磁共振对比剂的应用潜力。

关键词 磁共振成像; 对比剂; 肝靶向; DOTA-酰肼; 钆; 合成

Design, Synthesis and Evaluation of Novel Gd-Based 1,4,7,10-Tetraazacyclododecan-*N,N',N,N'*-tetraacetic Acid-Hydrazide Derived Contrast Agents for Magnetic Resonance Imaging

Sun, Hongshun^{*,a,b} Zhou, Jin^{a,b} Li, Yulong^a Jiang, Hong^a Zhang, Yan^c
Wang, Jianqiang^b Guo, Cheng^b Shen, Linjiang^{b,d}

(^a Targeted Magnetic Resonance Imaging Contrast Agents Laboratory of Jiangsu Province, Nanjing Polytechnic Institute, Nanjing 210048)

(^b College of Chemistry and Molecular Engineering, Nanjing Tech University, Nanjing 211816)

(^c College of Textiles and Clothing, Yancheng Vocational Institute of Industry Technology, Yancheng 224005)

(^d School of Physical and Mathematical Sciences, Nanjing Tech University, Nanjing 211816)

Abstract Magnetic resonance imaging (MRI) is widely used in diagnostic medicine and soft tissue imaging. Contrast agents (CAs) can improve the specificity of MRI enhancement. Herein, the design, synthesis and characterization of twelve novel Gd-based 1,4,7,10-tetraazacyclododecan-*N,N',N,N'*-tetraacetic acid (DOTA)-hydrazide derived contrast agents for MRI were reported. Among of them, **5d**, **5h** and **5l** exhibit higher longitudinal relaxivities than the clinical Gd-DOTA at 0.5 T. The relaxivities r_1 of **5d**, **5h** and **5l** are 4.67, 4.85 and 5.33 L·mmol⁻¹·s⁻¹ respectively. *In vivo* liver-target MRI shows that the potential of complex **5d** was used as a novel liver-target contrast agent for MRI.

Keywords magnetic resonance imaging (MRI); contrast agents; liver-target; DOTA-hydrazide; gadolinium; synthesis

* Corresponding author. E-mail: njutshs@126.com

Received August 7, 2018; revised September 30, 2018; published online October 26, 2018.

Project supported by the Natural Science Foundation of Jiangsu Province (No. BK20181486), the Natural Science Foundation of the Jiangsu Higher Education Institutions (Nos. 17KJB320001, 16KJB540006), the Overseas Training Program for Excellent Young Teachers and Principals of Jiangsu Province and the Qing Lan Project of Jiangsu Province.

江苏省自然科学基金(No. BK20181486)、江苏省高校自然科学基金(Nos. 17KJB320001, 16KJB540006)、江苏省高校优秀青年教师和校长境外研修计划和江苏省青蓝工程资助项目。

Molecular imaging plays an increasingly important role in bio-medical research, enabling a better understanding of fundamental molecular pathways in living organisms. Various different imaging techniques are used in molecular imaging, such as X-ray computer tomography (CT), positron emission tomography (PET), single photon emission tomography (SPECT), magnetic resonance imaging (MRI) and optical imaging (OI). Over the past few decades, the use of MRI has revolutionized diagnostic medicine and soft tissue imaging. In order to improve the specificity of MRI enhancement, contrast agents (CAs) are widely used to assist in the acquisition and interpretation of MRI images. About one-third of the 60 million MRI procedures done annually worldwide use contrast-enhancing agents, mostly containing the element gadolinium.^[1,2]

The first contrast agent approved for human use was $[\text{Gd}(\text{DTPA})(\text{H}_2\text{O})]^{2-}$ (DTPA: diethylenetriaminepentaacetate) (Chart 1).^[3,4] Up to now, the number of approved CAs has increased to nine, and all of them were gadolinium-based contrast agents (GBCAs) and T1-weighted MRI images. They can be divided into extracellular fluid (ECF) space, organ-specific (or targeted) and blood pool agents.^[5] $[\text{Gd}(\text{BOPTA})(\text{H}_2\text{O})]^{2-}$ and $[\text{Gd}(\text{EOB-DTPA})(\text{H}_2\text{O})]^{2-}$ (Chart 1) are liver-targeted agents with lipophilic side groups. The two targeted agents are all linear based on ligand DTPA.^[6,7] However, it has been proved that linear GBCAs have a risk of nephrogenic systemic fibrosis (NSF) in certain patients with kidney dysfunction. But macrocyclic GBCAs have a lower risk for the development of NSF.^[2,8] Because the macrocyclic agents are more stable than the linear agents. For this reason, a lot of macrocyclic GBCAs have been synthesized and researched.^[9–15]

Chart 1 Some clinical contrast agents

DOTA (1,4,7,10-tetraazacyclododecan- N,N',N'',N''' -tetraacetic acid) is the most commonly used macrocyclic chelating agent for the functionalization of peptides, proteins and antibodies with contrast agents, forming thermodynamically stable and kinetically inert complexes with

most trivalent metal ions.^[16,17] To enable efficient coupling to biomolecules, reactive groups need to be introduced into the DOTA complex. Amine and hydrazide groups have been established as versatile coupling sites in many different applications in the biomedical research. DOTA-hydrazide (DOTAH) is a good chelating agent for Gd^{3+} and has been researched in some literatures.^[18–20] Herein, we reported the synthesis and characterization of a series of Gd-based DOTA-hydrazide derived contrast agents for MRI. For these agents, we introduced a lipophilic group with benzene ring and hope that it can bind to liver. Then we can get a liver targeted agent based on macrocyclic ligand, which can overcome the drawback of NSF resulting from linear agents.

2 Results and discussion

2.1 Design and synthesis

In this study, DOTAH was chosen as the chelator for Gd^{3+} to build novel GBCAs. As a macrocyclic ligand, DOTAH has higher thermodynamic stability and slower dissociation kinetics to avoid transmetallation *in vivo* compared to the linear ligand like DTPA. At the same time, the Gd^{3+} binding neutralized the charges of the three acetic groups of DOTAH, leading to nonionic CAs, which is beneficial to low osmotic pressure. Then, kinds of lipophilic groups with benzene ring were introduced, which may make these new MRI contrast agents stay a longer time in liver so that they could get liver-target property.

The desired complexes **5a**~**5l** and their precursors were prepared following the procedures in Scheme 1. Raw material of 1,4,7-tris(tert-butoxycarbonylmethyl)-10-hydrazinylcarbonylmethyl-1,4,7,10-tetraazacyclododecane (*t*-Bu-DOTAH, **1**) was synthesized from cyclen according to the literature we have published before.^[20] The condensation reaction of the first step was accomplished using *t*-Bu-DOTAH and different benzaldehyde (**2**) in absolute ethanol refluxing for about 4 h. Deprotection of compounds **3a**~**3l** was achieved within 24 h by reacting with trifluoroacetic acid (TFA). Complexes **5a**~**5l** were obtained by combining **4a**~**4l** and gadolinium chloride in an aqueous solution at 50 °C, and the purification of **5a**~**5l** was accomplished by C-18 SPE cartridge.

2.2 In Vitro MRI

To evaluate the contrast enhancement of the new agents, longitudinal relaxivity r_1 and transverse relaxivity r_2 of them were calculated from the linear fitting of $1/T_1$ and $1/T_2$ plot versus metal concentrations. Commercially available Gd-DOTA was used as control. The values of r_1 and r_2 are estimated to be 2.94 to 5.33 and 3.63 to 6.32 $\text{L}\cdot\text{mmol}^{-1}\cdot\text{s}^{-1}$, respectively (Table 1). Among them, The relaxivities r_1 of **5d**, **5h** and **5l** are 4.67, 4.85 and 5.33 $\text{L}\cdot\text{mmol}^{-1}\cdot\text{s}^{-1}$ respectively, which are a little higher than that of the currently used clinical contrast agent Gd-DOTA with relaxivity r_1 of 4.41 $\text{L}\cdot\text{mmol}^{-1}\cdot\text{s}^{-1}$. It means that these three contrast agents may be used at lower dosage of agents and thus reducing the clinical toxicity. At the same

Table 1 r_1 and r_2 values of the synthesized compounds at 0.5 T, Gd-DOTA is as control

Entry	Compd.	$r_2/$ ($\text{L}\cdot\text{mmol}^{-1}\cdot\text{s}^{-1}$)	$r_1/$ ($\text{L}\cdot\text{mmol}^{-1}\cdot\text{s}^{-1}$)	r_2/r_1
1	Gd-DOTA	5.1289	4.4123	1.1624
2	5a	3.9125	3.2437	1.2062
3	5b	5.0427	4.1717	1.2088
4	5c	3.6309	2.9353	1.2370
5	5d	5.7423	4.6725	1.2290
6	5e	4.4016	3.7504	1.1736
7	5f	4.0837	3.5676	1.1447
8	5g	4.8747	3.9786	1.2252
9	5h	6.2079	4.8494	1.2801
10	5i	4.9583	3.9940	1.2414
11	5j	5.1293	4.2989	1.1932
12	5k	4.7857	3.9714	1.2050
13	5l	6.3216	5.3330	1.1854

time, the ratios of r_2/r_1 are calculated to be 1.14 to 1.28, the

low ratios of r_2/r_1 indicate that all the new agents can be used as T_1 -weighted MRI CAs. As shown in Figure 1, all new agents present excellent positive T_1 contrast enhancement, and the brightness of MR images enhance with the increasing concentrations of solutions, showing a clear dose-dependent color change, due to the relaxation of water proton increase with the increase of concentration dose.

2.3 *In vivo* liver-target MRI

For exploring the liver-target imaging property of these new synthesized contrast agents, we chose four of them with high relaxivity to do *in vivo* liver-target MRI test. Five adult male Sprague-Dawley (SD) rats weighing 280~300 g were randomly divided into five groups. All rats were scanned for livers with T_1 weighted imaging (T1WI). Then they were intravenously injected with **5d**, **5h**, **5j**, **5l** and Gd-DOTA as control at a dose of 0.1 mmol/kg in the tail vein, respectively. After administration, T1WI of liver was performed at 0 h and 6 h. As shown in Figure 2, immediately after administration of Gd-DOTA and four new

Scheme 1 Synthesis of **5a~5l** **Figure 1** T_1 -weighted images of **5a~5l** and Gd-DOTA at 0.5 T

Figure 2 MR images of liver at pre- and post-injection

Figure 3 Contrast ratio of liver vs normal muscle at pre- and post-injection

contrast agents, the signal intensities (SIs) of normal livers were all strongly enhanced. But after 6 h, only the SI administrated by **5d** was still enhanced. The contrast ratios of liver vs normal muscle at pre- and post-injection shown in Figure 3 also demonstrated that only the rat administrated by **5d** could get the continuous liver enhancement signal. All the results show the potential of compound **5d** used as a new liver-target contrast agent for MRI.

3 Conclusions

In summary, a series of novel Gd-based DOTA-hydrazide derived contrast agents have been designed and synthesized. And we explored their feasibilities as T_1 MRI CAs. The results showed that the thus-prepared agents have good positive imaging effect and low r_2/r_1 ratios suitable as T_1 -weighted MRI CAs. Moreover, three new

agents exhibited a little higher r_1 relaxivities and the ideal contrast enhancements in T_1 -weighted MR images, which were comparable to the clinical contrast agent Gd-DOTA. It means that they may be used at lower dosage of agents and thus reducing the clinical toxicity. *In vivo* liver-target MRI shows that complex **5d** may be used as a novel liver-target contrast agent for MRI.

4 Experimental section

4.1 General procedures

All chemicals and reagents were used as received without further purification. Glassware was dried in an oven at 130 °C and purged with a dry atmosphere prior to use. Unless otherwise mentioned, reactions were performed open to air. Reactions were monitored by thin-layer chromatography (TLC) and visualized by a dual short/long wavelength UV lamp. Flash column chromatography was performed using 200~400 mesh silica gel (Fisher). NMR spectra were recorded on a Bruker AVANCE AV-300 or a Bruker AVANCE AV-500 NMR spectrometer at room temperature. Chemical shifts for ^1H NMR and ^{13}C NMR were reported as δ . Spectra were referenced to internal standards (TMS in CDCl_3 and TMSP in D_2O). Multiplicity and coupling constants (J) were calculated automatically on MestReNova 10.0, a NMR processing software from Mestrelab Research. MS (ESI) was determined by using a microTOF-Q II HRMS/MS instrument (Bruker). Element analysis was determined by using a Perkin-Elmer 240c elemental analysis instrument. *In vitro* MRI testing and T_1 relaxation time measurements were tested at a 0.5 T NMR120-Analyst NMR Analyzing and Imaging system (Niuman Corporation, Shanghai, China). *In vivo* MRI was performed on a clinical 1.5 T MR magnet (Echo speed; GE Co., NY).

4.2 General method for the synthesis of **3a**~**3l**

To a 100 mL round-bottom flask fitted with a magnetic stir bar were added compound **1** (1.17 g, 2 mmol), different benzaldehyde (2 mmol) and 20 mL of absolute ethanol. The mixture was stirred at 75 °C for 4 h. After completion of the reaction, ethanol was removed *in vacuo*, and the resulting residue was recrystallized with ethyl acetate. Filtered, and washed with cold ethyl acetate, the resulting solid was used without further purification.

Tri-*tert*-butyl 2,2',2''-(10-(2-(2-(benzylidene)hydrazine-yl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)triacetate (**3a**): White solid, yield 88%. m.p. 204.5~205.8 °C; ^1H NMR (300 MHz, CDCl_3) δ : 12.99 (s, 1H), 8.76 (s, 1H), 7.83~7.70 (m, 2H), 7.39~7.25 (m, 3H), 3.62 (s, 2H), 3.29~2.81 (br, 6H), 2.74~1.97 (br, 16H), 1.49 (s, 15H), 1.46 (s, 12H); ^{13}C NMR (75 MHz, CDCl_3) δ : 171.9, 171.6, 168.3, 148.1, 134.6, 129.1, 128.2, 127.8, 127.3, 126.8, 81.7, 81.6, 55.6, 55.4, 55.2, 50.0, 27.7, 27.6; ESI-MS m/z : 675.5 $[\text{M}+\text{H}]^+$, 697.4 $[\text{M}+\text{Na}]^+$. Anal. calcd for $\text{C}_{35}\text{H}_{58}\text{N}_6\text{O}_7$: C 62.29, H 8.66, N 12.45; found C 62.23, H 8.68, N 12.47.

Tri-*tert*-butyl 2,2',2''-(10-(2-(2-(2-nitrobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)triacetate (**3b**): White solid, yield 90%. m.p. 194.7~195.2 °C; ¹H NMR (300 MHz, CDCl₃) δ: 13.31 (s, 1H), 8.90 (s, 1H), 8.73 (s, 1H), 8.14 (d, *J*=7.8 Hz, 1H), 7.95 (d, *J*=7.5 Hz, 1H), 7.50 (t, *J*=7.8 Hz, 1H), 3.62 (s, 2H), 3.44~2.92 (br, 6H), 2.86~1.99 (br, 16H), 1.51 (s, 15H), 1.47 (s, 12H); ¹³C NMR (75 MHz, CDCl₃) δ: 172.2, 168.9, 148.4, 145.3, 137.0, 133.9, 129.1, 123.5, 121.1, 82.2, 82.0, 55.8, 55.7, 55.4, 50.3, 28.0, 27.9; ESI-MS *m/z*: 720.6 [M+H]⁺. Anal. calcd for C₃₅H₅₇N₇O₉: C 58.40, H 7.98, N 13.62; found C 58.45, H 7.96, N 13.60.

Tri-*tert*-butyl 2,2',2''-(10-(2-(2-(4-methylbenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraaza-cyclododecane-1,4,7-triyl)triacetate (**3c**): White solid, yield 87%. m.p. 188.3~189.3 °C; ¹H NMR (300 MHz, CDCl₃) δ: 12.88 (s, 1H), 8.71 (s, 1H), 7.66 (d, *J*=7.8 Hz, 2H), 7.12 (d, *J*=7.8 Hz, 2H), 3.61 (s, 2H), 3.47 (s, 2H), 3.30~2.87 (br, 5H), 2.74~1.99 (br, 18H), 1.48 (s, 14H), 1.46 (s, 13H); ¹³C NMR (75 MHz, CDCl₃) δ: 172.1, 168.4, 148.7, 139.4, 132.1, 128.8, 127.6, 82.1, 82.0, 56.0, 55.7, 55.5, 50.3, 28.0, 27.9, 21.3; ESI-MS *m/z*: 689.5 [M+H]⁺. Anal. calcd for C₃₆H₆₀N₆O₇: C 62.77, H 8.78, N 12.20; found C 62.71, H 8.80, N 12.22.

Tri-*tert*-butyl 2,2',2''-(10-(2-(2-(4-nitrobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraaza-cyclododecane-1,4,7-triyl)triacetate (**3d**): Yellow solid, yield 87%. m.p. 138.9~140.5 °C; ¹H NMR (300 MHz, CDCl₃) δ: 13.26 (s, 1H), 8.87 (s, 1H), 8.18 (d, *J*=8.6 Hz, 2H), 7.93 (d, *J*=8.7 Hz, 2H), 3.61 (s, 2H), 2.39~2.87 (br, 6H), 2.81~2.15 (br, 16H), 1.49 (s, 14H), 1.47 (s, 13H); ¹³C NMR (75 MHz, CDCl₃) δ: 172.17, 169.2, 148.0, 145.5, 141.3, 128.0, 123.5, 82.2, 82.1, 55.9, 55.7, 55.5, 50.5, 28.0, 27.9; ESI-MS *m/z*: 720.5 [M+H]⁺, 742.5 [M+Na]⁺. Anal. calcd for C₃₅H₅₇N₇O₉: C 58.40, H 7.98, N 13.62; found C 58.44, H 7.97, N 13.60.

Tri-*tert*-butyl 2,2',2''-(10-(2-(2-(4-chlorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraaza-cyclododecane-1,4,7-triyl)triacetate (**3e**): White solid, yield 95%. m.p. 182.1~183.9 °C; ¹H NMR (300 MHz, CDCl₃) δ: 13.08 (s, 1H), 8.75 (s, 1H), 7.70 (d, *J*=8.3 Hz, 2H), 7.29 (d, *J*=8.0 Hz, 2H), 3.61 (s, 2H), 3.38~2.87 (br, 16H), 2.83~1.95 (br, 16H), 1.48 (s, 15H), 1.46 (s, 12H); ¹³C NMR (75 MHz, CDCl₃) δ: 172.1, 168.6, 147.1, 135.1, 133.5, 128.7, 128.3, 114.5, 82.1, 82.0, 55.9, 55.7, 55.5, 50.6, 28.6, 27.9; ESI-MS *m/z*: 709.5 [M+H]⁺, 731.4 [M+Na]⁺. Anal. calcd for C₃₅H₅₇ClN₆O₇: C 59.27, H 8.10, N 11.85; found C 59.21, H 8.11, N 11.87.

Tri-*tert*-butyl 2,2',2''-(10-(2-(2-(2,4-dichlorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraaza-cyclododecane-1,4,7-triyl)triacetate (**3f**): White solid, yield 90%. m.p. 164.6~166.1 °C; ¹H NMR (300 MHz, CDCl₃) δ: 13.26 (s, 1H), 9.08 (s, 1H), 8.02 (d, *J*=8.5 Hz, 1H), 7.33 (s, 1H), 7.17 (d, *J*=8.4 Hz, 1H), 3.62 (s, 2H), 3.40~2.88 (s, 6H), 2.84~2.15 (br, 16H), 1.48 (s, 15H), 1.46 (s, 12H). ¹³C NMR (126 MHz, CDCl₃) δ: 172.2, 169.1, 144.1, 135.4, 135.2, 131.5, 129.4, 128.4, 126.7, 82.1, 82.0, 56.0, 55.8,

55.6, 28.11, 28.0; ESI-MS *m/z*: 765.4 [M+Na]⁺. Anal. calcd for C₃₅H₅₆Cl₂N₆O₇: C 56.52, H 7.59, N 11.30; found C 56.57, H 7.58, N 11.28.

Tri-*tert*-butyl 2,2',2''-(10-(2-(2-(4-fluorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)triacetate (**3g**): Light yellow solid, yield 93%. m.p. 184.6~186.0 °C; ¹H NMR (300 MHz, CDCl₃) δ: 12.92 (s, 1H), 8.73 (s, 1H), 7.75 (s, 2H), 7.00 (t, *J*=8.2 Hz, 2H), 3.59 (s, 2H), 3.39~2.89 (br, 6H), 2.78~2.00 (br, 16H), 1.48 (s, 14H), 1.46 (s, 13H); ¹³C NMR (75 MHz, Chloroform-*d*) δ: 172.1, 168.6, 165.2, 161.8, 147.3, 131.1 (d), 129.4, 129.3, 115.3, 115.0, 82.1, 82.0, 55.9, 55.7, 55.4, 50.3, 28.0, 27.9; ESI-MS *m/z*: 693.5 [M+H]⁺, 715.5 [M+Na]⁺. Anal. calcd for C₃₅H₅₇FN₆O₇: C 60.67, H 8.29, N 12.13; found C 60.61, H 8.30, N 12.15.

Tri-*tert*-butyl 2,2',2''-(10-(2-(2-(3-fluorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraaza-cyclododecane-1,4,7-triyl)triacetate (**3h**): Light yellow solid, yield 91%. m.p. 185.3~186.7 °C; ¹H NMR (300 MHz, CDCl₃) δ: 12.97 (s, 1H), 8.56 (d, *J*=9.8 Hz, 1H), 7.45 (d, *J*=7.7 Hz, 1H), 7.28 (d, *J*=6.4 Hz, 1H), 7.00 (t, *J*=7.3 Hz, 1H), 3.59 (s, 2H), 3.36~2.83 (br, 6H), 2.77~1.96 (br, 16H), 1.49 (s, 14H), 1.46 (s, 13H); ¹³C NMR (75 MHz, CDCl₃) δ: 172.17, 169.2, 148.0, 145.5, 141.3, 128.0, 123.5, 82.2, 82.1, 55.9, 55.7, 55.5, 50.5, 28.0, 27.9; ESI-MS *m/z*: 720.5 [M+H]⁺, 742.5 [M+Na]⁺. Anal. calcd for C₃₅H₅₇FN₆O₇: C 60.67, H 8.29, N 12.13; found C 60.70, H 8.28, N 12.11.

Tri-*tert*-butyl 2,2',2''-(10-(2-(2-(2-fluorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraaza-cyclododecane-1,4,7-triyl)triacetate (**3i**): White solid, yield 89%. m.p. 193.4~195.0 °C; ¹H NMR (300 MHz, CDCl₃) δ: 13.10 (s, 1H), 8.93 (s, 1H), 8.00 (t, *J*=7.2 Hz, 1H), 7.28 (d, *J*=6.5 Hz, 1H), 7.13~6.91 (m, 2H), 3.63 (s, 2H), 3.41~2.93 (br, 6H), 2.83~1.95 (br, 16H), 1.48 (s, 14H), 1.46 (s, 13H); ¹³C NMR (75 MHz, CDCl₃) δ: 172.1, 168.9, 163.1, 159.7, 141.9, 141.9, 130.8, 127.4, 123.6, 123.5, 115.7, 115.4, 82.0, 81.9, 55.9, 55.7, 55.5, 50.2, 28.0, 27.9; ESI-MS *m/z*: 693.5 [M+H]⁺, 715.5 [M+Na]⁺. Anal. calcd for C₃₅H₅₇FN₆O₇: C 60.67, H 8.29, N 12.13; found C 60.62, H 8.30, N 12.14.

Tri-*tert*-butyl 2,2',2''-(10-(2-(2-(3,4-difluorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)triacetate (**3j**): White solid, yield 92%. m.p. 191.2~193.1 °C; ¹H NMR (300 MHz, CDCl₃) δ: 13.10 (s, 1H), 8.73 (s, 1H), 7.69 (t, *J*=9.0, 1H), 7.39 (s, 1H), 7.10 (dd, *J*=17.8, 8.5 Hz, 1H), 3.59 (s, 2H), 3.42~2.88 (br, 6H), 2.86~1.93 (br, 16H), 1.49 (s, 15H), 1.46 (s, 12H); ¹³C NMR (75 MHz, CDCl₃) δ: 172.1, 168.7, 145.9, 124.6, 117.0, 116.8, 115.2, 114.9, 82.1, 82.0, 55.94, 55.76, 55.49, 50.19, 28.04, 27.92; ESI-MS *m/z*: 733.5 [M+Na]⁺. Anal. calcd for C₃₅H₅₆F₂N₆O₇: C 59.14, H 7.94, N 11.82; found C 59.10, H 7.95, N 11.84.

Tri-*tert*-butyl 2,2',2''-(10-(2-(2-(4-methoxybenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)triacetate (**3k**): White solid, yield 85%. m.p. 177.5~178.9 °C; ¹H NMR (300 MHz, CDCl₃) δ: 12.80 (s, 1H), 8.70 (s, 1H), 7.71 (d, *J*=8.3 Hz, 2H), 6.84 (d, *J*=8.3

Hz, 2H), 3.81 (s, 3H), 3.59 (s, 2H), 3.36~2.87 (br, 6H), 2.81~1.91 (br, 16H), 1.49 (s, 14H), 1.46 (s, 13H); ^{13}C NMR (75 MHz, CDCl_3) δ : 172.1, 168.2, 160.7, 148.3, 129.1, 127.6, 114.5, 113.6, 82.0, 81.9, 56.0, 55.7, 55.5, 55.2, 50.3, 28.0, 27.9; ESI-MS m/z : 727.5 $[\text{M}+\text{Na}]^+$. Anal. calcd for $\text{C}_{36}\text{H}_{60}\text{N}_6\text{O}_8$: C 61.34, H 8.58, N 11.92; found C 61.39, H 8.57, N 11.90.

Tri-*tert*-butyl 2,2',2''-(10-(2-(2-(4-phenylbenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)triacetate (**3l**): White solid, yield 88%. m.p. 148.3~150.2 °C; ^1H NMR (300 MHz, CDCl_3) δ : 12.85 (s, 1H), 8.77 (s, 1H), 7.84 (d, $J=7.9$ Hz, 2H), 7.58 (t, $J=8.6$ Hz, 4H), 7.42 (t, $J=7.0$ Hz, 2H), 7.34 (d, $J=7.1$ Hz, 1H), 3.60 (s, 2H), 3.39~2.86 (br, 6H), 2.80~1.96 (br, 16H), 1.49 (s, 16H), 1.46 (s, 11H); ^{13}C NMR (75 MHz, CDCl_3) δ : 172.1, 168.6, 148.2, 142.0, 133.8, 128.6, 128.1, 127.6, 127.3, 127.0, 126.8, 82.1, 82.0, 56.0, 55.7, 55.5, 50.3, 28.0, 27.9. HRMS (ESI $^+$) calcd for $\text{C}_{41}\text{H}_{62}\text{N}_6\text{O}_7$ $[\text{M}+\text{H}]^+$ 751.4753, found 751.4752.

4.3 General method for the synthesis of 4a~4l

A solution of **3** (1.5 mmol) in trifluoroacetic acid (TFA) (10 mL) was stirred at room temperature for 24 h. TFA was removed under reduced pressure and the residue was dissolved in ethyl acetate (5 mL). After freezing for 12 h, the mixture was filtered to obtain solid compounds quantitatively.

2,2',2''-(10-(2-(2-(Benzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)triacetic acid (**4a**): White solid, yield 100%. m.p. 180.4~182.3 °C; ^1H NMR (300 MHz, D_2O) δ : 8.18 (s, 1H), 7.77 (s, 2H), 7.52 (s, 3H), 3.96~2.92 (m, 24H); ^{13}C NMR (75 MHz, D_2O) δ : 173.1, 172.6, 154.0, 135.7, 133.9, 133.6, 131.7, 130.5, 130.0, 59.1, 58.2, 56.6, 54.2, 53.5, 53.1, 51.2; ESI-MS m/z : 543.2 $[\text{M}+\text{K}-2\text{H}]^-$. Anal. calcd for $\text{C}_{23}\text{H}_{34}\text{N}_6\text{O}_7$: C 54.54, H 6.77, N 16.59; found C 54.50, H 6.78, N 16.62.

2,2',2''-(10-(2-(2-(2-Nitrobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)triacetic acid (**4b**): White solid, yield 100%. m.p. 169.9~172.0 °C; ^1H NMR (300 MHz, D_2O) δ : 8.47 (s, 1H), 8.25 (d, $J=9.0$ Hz, 1H), 8.14 (s, 1H), 8.03 (d, $J=8.5$ Hz, 1H), 4.06~2.87 (m, 24H); ^{13}C NMR (126 MHz, D_2O) δ : 148.5, 148.1, 148.0, 134.8, 133.8, 130.1, 125.2, 122.3, 117.6, 55.4, 54.1, 50.9, 48.6; ESI-MS m/z : 552.2 $[\text{M}+\text{H}]^+$, 550.0 $[\text{M}-\text{H}]^-$. Anal. calcd for $\text{C}_{23}\text{H}_{33}\text{N}_7\text{O}_9$: C 50.09, H 6.03, N 17.78; found C 50.14, H 6.02, N 17.76.

2,2',2''-(10-(2-(2-(4-Methylbenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)triacetic acid (**4c**): White solid, yield 100%. m.p. 185.7~187.3 °C; ^1H NMR (500 MHz, D_2O) δ : 8.14 (s, 1H), 7.66 (d, $J=7.0$ Hz, 2H), 7.33 (d, $J=6.4$ Hz, 2H), 3.96~2.96 (m, 24H), 2.38 (s, 3H); ^{13}C NMR (126 MHz, D_2O) δ : 170.4, 163.1, 162.8, 151.6, 142.3, 130.3, 129.7, 127.9, 127.4, 117.6, 115.3, 56.5, 54.1, 51.6, 50.9, 48.6, 48.0, 20.7; ESI-MS m/z : 521.2 $[\text{M}+\text{H}]^+$, 519.1 $[\text{M}-\text{H}]^-$. Anal. calcd for $\text{C}_{24}\text{H}_{36}\text{N}_6\text{O}_7$: C 55.37, H 6.97, N 16.14; found C 55.41, H 6.96, N 16.12.

2,2',2''-(10-(2-(2-(4-Nitrobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)triacetic acid (**4d**): Yellow solid, yield 100%. m.p. 183.8~185.6 °C; ^1H NMR (300 MHz, D_2O) δ : 8.23 (t, $J=8.0$ Hz, 2H), 8.14 (s, 1H), 7.87 (d, $J=8.2$ Hz, 1H), 7.81 (d, $J=8.2$ Hz, 1H), 4.04~2.97 (m, 24H); ^{13}C NMR (126 MHz, D_2O) δ : 163.0, 162.7, 148.3, 139.5, 128.5, 128.0, 124.0, 117.6, 115.2, 55.3, 53.8, 50.4, 48.6; ESI-MS m/z : 588.2 $[\text{M}+\text{K}-2\text{H}]^-$. Anal. calcd for $\text{C}_{23}\text{H}_{33}\text{N}_7\text{O}_9$: C 50.09, H 6.03, N 17.78; found C 50.05, H 6.04, N 17.80.

2,2',2''-(10-(2-(2-(4-Chlorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)triacetic acid (**4e**): Light yellow solid, yield 100%. m.p. 186.4~188.6 °C; ^1H NMR (500 MHz, D_2O) δ : 8.03 (s, 1H), 7.55 (d, $J=45.1$ Hz, 2H), 7.32 (d, $J=34.5$ Hz, 2H), 3.98~2.96 (m, 24H); ^{13}C NMR (126 MHz, D_2O) δ : 162.9, 162.6, 149.7, 136.2, 131.9, 129.2, 129.0, 117.6, 115.3, 55.3, 53.9, 50.9, 48.6, 48.1; ESI-MS m/z : 541.2 $[\text{M}+\text{H}]^+$, 539.0 $[\text{M}-\text{H}]^-$. Anal. calcd for $\text{C}_{23}\text{H}_{33}\text{ClN}_6\text{O}_7$: C 51.06, H 6.15, N 15.53; found C 51.10, H 6.14, N 15.50.

2,2',2''-(10-(2-(2-(2,4-Dichlorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)triacetic acid (**4f**): Light yellow solid, yield 100%. m.p. 184.5~186.4 °C; ^1H NMR (500 MHz, D_2O) δ : 8.34 (s, 1H), 8.11 (s, 1H), 7.16 (d, $J=24.4$ Hz, 1H), 7.02 (s, 1H), 3.97~2.85 (m, 24H); ^{13}C NMR (126 MHz, D_2O) δ : 162.9, 162.6, 136.4, 135.8, 134.8, 129.6, 127.3, 117.6, 115.3, 55.7, 50.9, 48.6, 48.1; ESI-MS m/z : 575.1 $[\text{M}+\text{H}]^+$, 573.0 $[\text{M}-\text{H}]^-$. Anal. calcd for $\text{C}_{23}\text{H}_{32}\text{Cl}_2\text{N}_6\text{O}_7$: C 48.01, H 5.61, N 14.60; found C 48.05, H 5.60, N 14.58.

2,2',2''-(10-(2-(2-(4-Fluorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)triacetic acid (**4g**): White solid, yield 100%. m.p. 163.6~166.5 °C; ^1H NMR (300 MHz, D_2O) δ : 8.13 (s, 1H), 8.00 (s, 1H), 7.78 (s, 2H), 7.22 (s, 1H), 4.00~2.95 (m, 24H); ^{13}C NMR (75 MHz, D_2O) δ : 168.4, 165.1, 152.9, 132.7, 132.6, 132.2, 132.0, 118.8, 118.5, 58.8, 57.7, 56.6, 53.43, 51.9, 50.7; ESI-MS m/z : 561.2 $[\text{M}+\text{K}-2\text{H}]^-$. Anal. calcd for $\text{C}_{23}\text{H}_{33}\text{FN}_6\text{O}_7$: C 52.66, H 6.34, N 16.02; found C 52.70, H 6.33, N 16.01.

2,2',2''-(10-(2-(2-(3-Fluorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)triacetic acid (**4h**): White solid, yield 100%. m.p. 170.0~171.5 °C; ^1H NMR (300 MHz, D_2O) δ : 8.12 (s, 1H), 7.96 (s, 1H), 7.49 (d, $J=14.8$ Hz, 3H), 7.23 (d, $J=7.5$ Hz, 1H), 3.99~3.04 (m, 24H); ^{13}C NMR (75 MHz, D_2O) δ : 173.0, 166.9, 163.7, 152.5, 138.0, 133.4, 126.9, 120.6, 116.1, 115.6, 58.9, 57.9, 56.6, 53.5, 51.2; ESI-MS m/z : 525.2 $[\text{M}+\text{H}]^+$, 523.1 $[\text{M}-\text{H}]^-$. Anal. calcd for $\text{C}_{23}\text{H}_{33}\text{FN}_6\text{O}_7$: C 52.66, H 6.34, N 16.02; found C 52.62, H 6.35, N 16.04.

2,2',2''-(10-(2-(2-(2-Fluorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)triacetic acid (**4i**): White solid, yield 100%. m.p. 170.1~172.5 °C; ^1H NMR (500 MHz, D_2O) δ : 8.34 (s, 1H), 7.90 (t, $J=7.1$ Hz, 1H), 7.56~7.46 (m, 1H), 7.32~7.17 (m, 2H), 3.95~3.09 (m, 24H); ^{13}C NMR (126 MHz, D_2O) δ : 170.5, 162.4, 160.4, 144.7, 133.1, 127.4, 124.8, 120.7,

117.6, 116.2, 55.5, 53.8, 50.9, 48.6; ESI-MS m/z : 525.2 $[M+H]^+$, 523.1 $[M-H]^-$. Anal. calcd for $C_{23}H_{33}FN_6O_7$: C 52.66, H 6.34, N 16.02; found C 52.69, H 6.33, N 16.00.

2,2',2''-(10-(2-(2-(3,4-Difluorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)tri-acetic acid (**4j**): White solid, yield 100%. m.p. 182.8~184.6 °C; 1H NMR (300 MHz, D_2O) δ : 8.04 (s, 1H), 7.72~7.22 (m, 3H), 4.01~3.06 (m, 24H); ^{13}C NMR (126 MHz, D_2O) δ : 173.3, 155.8, 154.4, 151.6, 148.1, 133.1, 127.9, 120.6, 118.3, 117.7, 58.8, 57.8, 56.5, 53.0, 51.5; ESI-MS m/z : 543.2 $[M+H]^+$, 541.1 $[M-H]^-$. Anal. calcd for $C_{23}H_{32}F_2N_6O_7$: C 50.92, H 5.95, N 15.49; found C 50.95, H 5.94, N 15.47.

2,2',2''-(10-(2-(2-(4-Methoxybenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)tri-acetic acid (**4k**): Yellow solid, yield 100%. m.p. 152.9~154.1 °C; 1H NMR (300 MHz, D_2O) δ : 8.09 (s, 1H), 7.71 (d, $J=7.8$ Hz, 1H), 7.65 (d, $J=7.9$ Hz, 1H), 7.05 (d, $J=7.2$ Hz, 2H), 3.98~2.98 (m, 27H); ^{13}C NMR (75 MHz, D_2O) δ : 197.5, 164.0, 163.7, 153.8, 135.4, 132.3, 131.8, 128.9, 128.6, 117.2, 58.9, 58.2, 56.6, 55.0, 53.5, 51.2; ESI-MS m/z : 537.2 $[M+H]^+$, 535.1 $[M-H]^-$. Anal. calcd for $C_{24}H_{36}N_6O_8$: C 53.72, H 6.76, N 15.66; found C 53.75, H 6.75, N 15.65.

2,2',2''-(10-(2-(2-(4-Phenylbenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)tri-acetic acid (**4l**): White solid, yield 100%. m.p. 180.5~181.7 °C; 1H NMR (300 MHz, D_2O) δ : 8.01 (s, 1H), 7.65~6.57 (m, 9H), 4.07~2.68 (m, 24H); ^{13}C NMR (75 MHz, D_2O) δ : 177.4, 173.1, 165.2, 152.1, 148.5, 144.5, 144.1, 142.0, 141.8, 135.2, 131.3, 129.2, 121.0, 117.2, 59.2, 55.7, 54.3, 53.4, 51.2, 50.7. HRMS (ESI $^+$) calcd for $C_{29}H_{38}N_6O_7$ $[M+H]^+$ 605.2694, found 605.2701.

4.4 General method for the synthesis of **5a**~**5l**

To a solution of **4a**~**4l** (1.5 mmol) in water (20 mL) was added a solution of $GdCl_3$ in water (0.05 mol/L, 36 mL, 1.8 mmol) while maintaining the pH at 6.5 by addition of 1 mol/L NaOH. The solution was stirred at 50 °C for 24 h. The pH was raised to 8.0. The solution was filtered through a 0.2 μm filter, and the pH was lowered to 7.0 by adding 1 mol/L HCl. The solution was loaded on a Hypersep C-18 cartridge, and the stationary phase was washed extensively with water. The product was recovered after gradient elution with MeOH/ H_2O from 1 : 9 to 3 : 2. The fractions containing the product were evaporated under reduced pressure to give **5a**~**5l** as solid compounds.

Gadolinium 2,2',2''-(10-(2-(2-(benzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)tri-acetic acetate (**5a**): White solid, yield 85%; ESI-MS m/z : 684.2 $[M+Na]^+$. A correct Gd isotope pattern was observed. Anal. calcd for $C_{23}H_{31}GdN_6O_7$: C 41.81, H 4.73, N 12.72; found C 41.74, H 4.74, N 13.19.

Gadolinium 2,2',2''-(10-(2-(2-(2-nitrobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)tri-acetic acetate (**5b**): Yellow solid, yield 90%; ESI-MS m/z : 729.1 $[M+Na]^+$. A correct Gd isotope pat-

tern was observed. Anal. calcd for $C_{23}H_{30}GdN_7O_9$: C 39.14, H 4.28, N 13.89; found C 39.19, H 4.27, N 13.87.

Gadolinium 2,2',2''-(10-(2-(2-(4-methylbenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)tri-acetic acetate (**5c**): White solid, yield 88%; ESI-MS m/z : 698.2 $[M+Na]^+$. A correct Gd isotope pattern was observed. Anal. calcd for $C_{24}H_{33}GdN_6O_7$: C 42.72, H 4.93, N 12.45; found C 42.66, H 4.94, N 12.47.

Gadolinium 2,2',2''-(10-(2-(2-(4-nitrobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)tri-acetic acetate (**5d**): Light yellow solid, yield 92%; ESI-MS m/z : 729.1 $[M+Na]^+$. A correct Gd isotope pattern was observed. Anal. calcd for $C_{23}H_{30}GdN_7O_9$: C 39.14, H 4.28, N 13.89; found C 39.18, H 4.27, N 13.86.

Gadolinium 2,2',2''-(10-(2-(2-(4-chlorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)tri-acetic acetate (**5e**): Light yellow solid, yield 95%; ESI-MS m/z : 718.2 $[M+Na]^+$. A correct Gd isotope pattern was observed. Anal. calcd for $C_{23}H_{30}ClGdN_6O_7$: C 39.74, H 4.35, N 12.09; found C 39.69, H 4.36, N 12.12.

Gadolinium 2,2',2''-(10-(2-(2-(2,4-dichlorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)tri-acetic acetate (**5f**): Light yellow solid, yield 90%; ESI-MS m/z : 729.2 $[M]^+$. A correct Gd isotope pattern was observed. Anal. calcd for $C_{23}H_{29}Cl_2GdN_6O_7$: C 37.86, H 4.01, N 11.52; found C 39.65, H 4.02, N 11.55.

Gadolinium 2,2',2''-(10-(2-(2-(4-fluorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)tri-acetic acetate (**5g**): Light yellow solid, yield 93%; ESI-MS m/z : 702.1 $[M+Na]^+$. A correct Gd isotope pattern was observed. Anal. calcd for $C_{23}H_{30}FGdN_6O_7$: C 40.70, H 4.46, N 12.38; found C 40.75, H 4.45, N 12.36.

Gadolinium 2,2',2''-(10-(2-(2-(3-fluorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)tri-acetic acetate (**5h**): Light yellow solid, yield 92%; ESI-MS m/z : 702.1 $[M+Na]^+$. A correct Gd isotope pattern was observed. Anal. calcd for $C_{23}H_{30}FGdN_6O_7$: C 40.70, H 4.46, N 12.38; found C 40.66, H 4.47, N 12.40.

Gadolinium 2,2',2''-(10-(2-(2-(2-fluorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)tri-acetic acetate (**5i**): White solid, yield 96%; ESI-MS m/z : 702.1 $[M+Na]^+$. A correct Gd isotope pattern was observed. Anal. calcd for $C_{23}H_{30}FGdN_6O_7$: C 40.70, H 4.46, N 12.38; found C 40.73, H 4.45, N 12.36.

Gadolinium 2,2',2''-(10-(2-(2-(3,4-difluorobenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)tri-acetic acetate (**5j**): White solid, yield 90%; ESI-MS m/z : 720.1 $[M+Na]^+$. A correct Gd isotope pattern was observed. Anal. calcd for $C_{23}H_{29}F_2GdN_6O_7$: C 39.65, H 4.20, N 12.06; found C 39.70, H 4.19, N 12.03.

Gadolinium 2,2',2''-(10-(2-(2-(4-methoxybenzylidene)hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4,7-triyl)tri-acetic acetate (**5k**): Light yellow solid, yield 88%; ESI-MS m/z : 714.1 $[M+Na]^+$. A correct Gd isotope pattern was observed. Anal. calcd for $C_{24}H_{33}GdN_6O_8$: C 41.73, H 4.82, N 12.17; found C 41.68, H 4.83, N 12.20.

Gadolinium 2,2',2''-(10-(2-(2-(4-phenylbenzylidene)-

hydrazinyl)-2-oxoethyl)-1,4,7,10-tetraazacyclododecane-1,4, 7-triyl)triacetic acetate (**5I**): Light yellow solid, yield 85%; ESI-MS m/z : 760.2 $[M+Na]^+$. A correct Gd isotope pattern was observed. Anal. calcd for $C_{29}H_{35}GdN_6O_7$: C 47.27, H 4.79, N 11.41; found C 47.33, H 4.78, N 11.38.

4.5 *In vitro* MRI

To obtain T_1 relaxation times of the aqueous solution of the new agents at different concentrations, following measurement parameters were used: repetition time (TR) is 6000 ms, number of data is 25, and number of averages (NA) is 2. While the T_2 relaxation times were determined with the following parameters: TR is 6000 ms, echo time (TE) is 1 ms, echo count is 6000 and NA is 2. Both r_1 and r_2 relaxivities were calculated by the linear fitting of $1/T_1$ or $1/T_2$ as a function of metal concentration. For T_1 -weighted MR images, the instrument parameters were set as follows: TR=100 ms, TE=18.2 ms, imaging matrix=192×256, slice thickness=5 mm, field of view (FOV)=100 mm×100 mm and NA=2.

4.6 Animals

Adult male Sprague-Dawley (SD) rats weighing 280~300 g were provided by the Experimental Animal Center, Jiangsu Province Academy of Traditional Chinese Medicine. All the animal experiments were implemented with the approval the institutional animal care and research advisory.

4.7 *In vivo* MRI and imaging analyses

MRI was performed on a clinical 1.5 T MR magnet (Echo speed; GE Co., NY) by using a rat coil. All animals were anaesthetized with isoflurane in a mixture of 80% room air and 20% oxygen. Spin-echo (SE) T_1 -weighted imaging (T1WI) was performed with the following parameters: TR=550 ms, TE=24 ms, FOV=100 mm×100 mm, imaging acquisition matrix=224×192, a total examination time of 185 s. Contrast agents of **5d**, **5h**, **5j**, **5I** and Gd-DOTA were used at the same dosage of 0.1 mmol/kg. MRI analyses were performed by using the built-in software of the system (GE, AW 4.3, GE Healthcare Bio-Sciences, Pittsburgh, PA). The signal intensities (SIs) of tissue were measured with an operator-defined circular region of interest (ROI) with 3 mm² on T1WI. Contrast ratio was used to express the visual conspicuity of adjacent component on the one image and calculated with the formula: $CR = SI_A/SI_B$.

Supporting Information ¹H NMR and ¹³C NMR spectra, mass spectra of the synthesized compounds. Longitu-

dinal and transverse relaxivities of Gd-DOTA and **5a**~**5I**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] Merbach, A.-E.; Helm, L.; Toth, E. *The Chemistry of Contrast Agents in Medical Magnetic Resonance Imaging*, Wiley, Chichester, **2013**, Preface.
- [2] He, W.; Oliver, T.-B.; Michael, G.-K.; Eric, C.-H.; Mariya, B.; Agata, W.; Ou, C.; Yue, C.; Nan, L.; Satoshi, O.; Jose, M. C.; Markus, H.; Christian, T.-F.; Daniel, M.-M.; Gerhard, A.; Harald, I.; Alan, J.; Peter, N.; Mounsi, G.-B. *Proc. Natl. Acad. Sci. U. S. A.* **2017**, *114*, 2325.
- [3] Carr, D.-H.; Brown, J.; Bydder, G.-M.; Steiner, R.-E.; Weinmann, H.-J.; Speck, U.; Hall, A.-S.; Young, I.-R. *Am. J. Roentgenol.* **1984**, *143*, 215.
- [4] Lohrke, J.; Frenzel, T.; Endrikat, J.; Alves, F.-C.; Grist, T.-M.; Law, M.; Lee, J.-M.; Leiner, T.; Li, K.-C.; Nikolaou, K.; Prince, M.-R.; Schild, H.-H.; Weinreb, J.-C.; Yoshikawa, K.; Pietsch, H. *Adv. Ther.* **2016**, *33*, 1.
- [5] Pierre, V.-C.; Allen, M.-J.; Caravan, P. *J. Biol. Inorg. Chem.* **2014**, *19*, 127.
- [6] Hermann, P.; Kotek, J.; Kubicek, V.; Lukes, I. *Dalton Trans.* **2008**, *23*, 3027.
- [7] Thian, Y.-L.; Riddell, A.-M.; Koh, D.-M. *Cancer Imaging* **2013**, *13*, 567.
- [8] Huckle, J.-E.; Altun, E.; Jay, M.-J.; Semelka, R.-C. *Invest. Radiol.* **2016**, *51*, 236.
- [9] McMahon, M.-T.; Chan, K.-W. *Adv. Cancer Res.* **2014**, *124*, 297.
- [10] Frullano, L.; Zhu, J.; Miller, R.-H.; Wang, Y.-J. *Med. Chem.* **2013**, *56*, 1629.
- [11] Song, Y.; Kohlmeir, E.-K.; Meade, T.-J. *J. Am. Chem. Soc.* **2008**, *130*, 6662.
- [12] Zhou, Y.; Kim, Y.-S.; Yan, X.; Jacobson, O.; Chen, X.; Liu, S. *Mol. Pharm.* **2011**, *8*, 1198.
- [13] Barge, A.; Tei, L.; Upadhyaya, D.; Fedeli, F.; Beltrami, L.; Stefania, R.; Aime, S.; Cravotto, G. *Org. Biomol. Chem.* **2008**, *6*, 1176.
- [14] Wang, S.-C.; Hsu, Y.-S.; Hsiao, C.-T.; Wu, C.-C.; Sue, Y.-C.; Alshehri, S.-M.; Ahamad, T.; Yamauchi, Y.; Chen, J.-E.; Wu, K.-C.-W.; Shieh, F.-K. *J. Inorg. Organomet. Polym.* **2016**, *26*, 165.
- [15] Zhang, W.; Lu, C.; Zhao, G.; Zhang, J.; Fang, X.; Wang, P.; Fang, X.; Xu, J.; Yang, W. *Z. Anorg. Allg. Chem.* **2015**, *641*, 578.
- [16] Cresens, E.; Ni, Y.-C.; Adriaens, P.; Verbruggen, A.; Marchal, G. *WO 2002038546*, **2002** [*Chem. Abstr.* **2002**, *136*, 379071].
- [17] Zhang, J.-D.; Curry, K. *WO 2016090491*, **2016** [*Chem. Abstr.* **2016**, *165*, 112684].
- [18] Frullano, L.; Tejerina, B.; Meade, T.-J. *Inorg. Chem.* **2006**, *45*, 8489.
- [19] Zhou, J.; Yang, H.-D.; Sun, H.-S.; Li, Y.-L.; Jiang, H. *Chin. J. Synth. Chem.* **2017**, *25*, 159 (in Chinese).
(周进, 杨海东, 孙宏顺, 李玉龙, 蒋蕪, 合成化学, **2017**, *25*, 159.)
- [20] Sun, H.-S.; Li, Y.-L.; Jiang, H.; Guo, C.; Shen, L.-J. *Chin. J. Org. Chem.* **2018**, *38*, 1779 (in Chinese).
(孙宏顺, 李玉龙, 蒋蕪, 郭成, 沈临江, 有机化学, **2018**, *38*, 1779.)

(Lu, Y.)