过渡金属催化的区域选择性芳烃 C—H 键氧化生成 C—O 键

杨帆致^{*,a} 张 晗^b 刘旭日^b 王 博^{*,b} Lutz Ackermann^{*,c}^a 北京理工大学前沿交叉科学研究院 北京 100081)^b 北京理工大学化学与化工学院 北京 100081)^c 哥廷根大学有机与生物分子化学研究所 哥廷根 德国 37077)

摘要 近年来过渡金属催化的芳烃直接 C—H 氧化反应取得了重要进展, 该策略被用于多种酚类化合物的制备. 反应体系使用的过渡金属催化剂包括钯、铜、钌、铱等; 氧化剂包括高价碘化合物、过硫酸盐、氧气等. 此前已有多篇综述就特定的过渡金属或氧化剂参与的 C—H 氧化反应进行了详尽的讨论. 本综述着重探讨过渡金属催化的邻、间、对位选择性的芳烃 C—H 氧化反应, 并试图阐释上述区域选择性的产生机制, 其中包括导向基团的螯合辅助作用、配体控制、底物自身因素等. 在讨论部分提出了过渡金属催化的芳烃 C—H 氧化反应中存在的问题以及影响该策略发展应用的可能限制因素.

关键词 过渡金属; 芳烃; C—H 氧化; 区域选择性; 导向基团

Transition Metal-Catalyzed Regio-selective Aromatic C—H Bond Oxidation for C—O Bond Formation

Yang, Fanzhi^{*,a} Zhang, Han^b Liu, Xuri^b Wang, Bo^{*,b} Ackermann, Lutz^{*,c}^a Advanced Research Institute of Multidisciplinary Science, Beijing Institute of Technology, Beijing 100081, China)^b School of Chemistry and Chemical Engineering, Beijing Institute of Technology, Beijing 100081 China)^c Institut für Organische und Biomolekulare Chemie, Georg-August-Universität Göttingen, Göttingen 37077 Germany)

Abstract Recent years the great progress in transition metal-catalyzed direct aromatic C—H oxidation has been witnessed, which has been utilized in the preparation of various phenolic compounds. These transformations employ inter alia palladium, copper, ruthenium, iridium, etc. as the transition metal catalysts, and hypervalent iodine, persulfate, or oxygen as the oxidants. There have been several reviews in which the C—H oxidations with specific transition metal or oxidant was discussed. This review focuses specifically on transition metal-catalyzed aromatic C—H oxidations with *ortho*-, *meta*-, or *para*-selectivity, and rationalizes the possible generation mechanism of regio-selectivities, which might be controlled by the directing group via chelation-assistance, the ligand, or intrinsic properties of the substrate. The discussion section indicated the existing problems of transition metal-catalyzed aromatic C—H oxidations, as well as the possible limiting factors for the development and application of this strategy.

Keywords transition metal; arene; C—H oxidation; regio-selectivity; directing group

1 Introduction

Phenols are important structural motifs in the research areas of pharmaceuticals, agrochemicals, polymers, and natural products. There has been a growing demand for developing versatile synthetic methodologies of phenol derivatives.^[1] Recently, transition metal-catalyzed direct aromatic C—H oxidation has emerged as a powerful tool

in the regio-selective synthesis of phenols. This strategy compares favorably to classical phenol synthesis with respect to the atom-economy as well as the step-economy.

There have already been several distinguished reviews on the research progress of transition metal-mediated aromatic C(sp²)—H oxygenation from different perspectives: Sanford *et al.* reviewed C—O bond formation as a section in the subject of palladium-catalyzed C—H functionaliza-

* Corresponding authors. E-mail: yangfanzhi@bit.edu.cn; bowang@bit.edu.cn; Lutz.Ackermann@chemie.uni-goettingen.de

Received August 16, 2018; revised October 22, 2018; published online October 26, 2018.

Dedicated to Professor Qingyun Chen on the occasion of his 90th birthday.

Project supported by the National Natural Science Foundation of China (No. 21801018) and the Beijing Institute of Technology Research Fund Program for Young Scholars (No. 1230011181807).

国家自然科学基金(No. 21801018)和北京理工大学青年教师学术启动计划(No. 1230011181807)资助项目.

tions.^[2] Later, Enthaler and Company summarized the progress in palladium-catalyzed hydroxylation and alkoxylation.^[3] Ackermann *et al.* reviewed the recent advances in ruthenium-catalyzed direct hydroxylation of C(sp²)—H and C(sp³)—H bonds.^[4] Baran *et al.* reviewed the oxidation of both C(sp²)—H and C(sp³)—H bonds, emphasizing the chemoselectivity imposed by the nature of the substrates, such as inductive effects, conjugation, hyper-conjugation, steric hindrance, and strain release.^[5] Regarding the oxidants, Jiao *et al.* summarized their research on the synthesis of oxygen-containing compounds via C—H or C—C bond activation with 101 kPa oxygen or air as the green oxidant.^[6] In the meanwhile, Saha and Laha *et al.* summarized the uses of K₂S₂O₈ as a cost-effective inorganic oxidant in metal-catalyzed and metal-free oxidative transformations.^[7]

The selectivity control of C—H functionalizations has been regarded as an essential section in the new “Holly Grails” of chemistry.^[8] Specific to the category of aromatic C—H bond oxidation, the selectivity mainly originates from electronic effect, steric effect, or chelation assistance. Very recently, several novel reaction systems have been developed for remote C—H bond oxidation. Given all of the above, we feel that it would be helpful to summarize the recent progress in transition metal-catalyzed aromatic C—H oxidations at *ortho*-, *meta*-, or *para*-position.

This review will focus on those aromatic oxidations where the oxygen atom on the newly formed C—O bond should originate from the terminal oxidant (Eq. 1). There are other pathways of aromatic C—H bond functionalizations for C—O bond generations: One way is the reaction between arenes and oxygen-containing nucleophiles via transition metal-catalyzed C—H activations,^[9] where the oxidation of transition metal core with an external oxidant is the essential step. It is noteworthy that the anode could also be regarded as an external oxidant, for instance, the groups of Ackermann, Mei, Lei *et al.* have recently made distinguished contributions to transition metal-catalyzed electrooxidative C—H functionalizations for C—O bond formations.^[10] Another way is C—H oxidation via radical aromatic substitution without transition metal catalyst.^[11] The above mentioned transformations will not be included in this review.

This review has compiled the research advances of transition metal-catalyzed regio-selective aromatic C—H oxidations up to spring of 2018. We hereby wish to give the readers an intuitive glance at the recent progress of this research area, as well as the generative mechanism of specific chemo- and regio-selectivity.

2 Direct C—H oxygenation of simple arenes

2.1 Palladium-catalyzed direct C—H oxidation

About 30 years ago, Jintoku, Fujiwara and co-workers

reported Pd(OAc)₂-catalyzed (OAc=acetate) oxidation of benzene to phenol under the atmosphere of oxygen (1.52 MPa) and carbon monoxide (1.52 MPa) at 180 °C. 1,10-Phenanthroline was recognized as the optimized ligand. Under optimized reaction conditions, phenyl acetate was monitored as a major side product, while benzoic acid was isolated as the major product in the absence of ligand.^[12]

The above systems have been suffering from high pressure, which could limit the applications of these strategies. In 2007, Ishii and co-workers utilized polyoxometalate H₅PMo₁₀V₂O₄₀•26H₂O (2 mol%) as the redox catalyst in palladium-catalyzed hydroxylation-carboxylation of biphenyl, and the reaction could proceed smoothly under an atmosphere of oxygen (50.7 kPa) and carbon monoxide (50.7 kPa) (Eq. 2).^[13]

Oxidants other than oxygen have proved applicable to palladium-catalyzed direct aromatic C—H oxidations. In 1996, Crabtree and co-workers employed PhI(OAc)₂ as the oxidant in Pd(OAc)₂-catalyzed C—H oxidation of arenes, delivering acetoxyated products. Mechanistic studies ruled out a radical route, and indicated that palladation should be the rate-determining step.^[14] Later, Sanford and co-workers performed a detailed mechanistic investigation on palladium-catalyzed PhI(OAc)₂ oxidized aromatic C—H acetoxylation. The results of NMR and kinetic studies suggested that the most active catalyst rests as a palladium dimer in solution (Scheme 1).^[15]

Scheme 1 Proposed catalytic cycle of palladium-catalyzed PhI(OAc)₂ oxidized aromatic C—H acetoxylation

Besides PhI(OAc)₂, K₂S₂O₈ has also been used as the terminal oxidant in palladium-catalyzed C—H oxidation.

In 2013, Sanford and co-workers reported palladium-catalyzed direct acetoxylation of benzene, using $K_2S_2O_8$ as the oxidant and a solvent mixture of $AcOH/Ac_2O$.^[16] In 2004, Nozaki and co-workers reported the palladium-catalyzed sequential hydroxylation-carboxylation using biphenyl using formic acid as the carbonyl source, generating a mixture of *meta*- and *para*-hydroxybiphenylcarboxylic acids in a combined yield of 45% (*m*-: *p*- = 17 : 83) (Eq. 3).^[17]

2.2 Manganese-catalyzed direct C—H oxidation

In 2013, Guo and co-workers reported tetrakis(pentafluorophenyl)porphyrin manganese(III) chloride ($F_{20}TPPMnCl$) catalyzed hydroxylation of 2-substituted naphthalene (Eq. 4). Depending on the electronic properties of the substituent on C2 position, the catalytic system showed different regio-selectivities. Naphthalene with an electron-donating substituent afforded C1-hydroxylated product, while with an electron-withdrawing substituent predominantly delivered C4-hydroxylated product.^[18]

R	Selectivity/%		
	6	7	8
H	93	—	—
Me/Et/ ⁱ Pr/OMe	92~93	Trace	7~8
Br/CO ₂ H	15~34	63~66	Trace~22

3 *ortho*-Selective aromatic C—H oxidation

In the category of transition metal-catalyzed *ortho*-selective aromatic C—H oxidation, the selectivity originated from the coordination of metal catalyst and the Lewis base substituent on the arene, which will tend to form a cyclized transition state via C—H metalation, followed by reductive elimination to afford the *ortho*-oxygenated product (Scheme 2).

3.1 Copper catalyzed *ortho*-C—H oxidation

Stoichiometric amount of copper has proved efficacy in

Scheme 2 *ortho*-Selective aromatic C—H oxidation

the *ortho*-selective hydroxylation of arenes, using molecular oxygen,^[19] iodosylbenzene,^[20] hydrogen peroxide,^[21] or trimethylamine *N*-oxide^[22] as the terminal oxidant. Besides these copper-catalyzed aromatic hydroxylations with external oxidants, $Cu(OAc)_2$ itself could also serve as the terminal oxidant in aromatic C—H oxidations. In 1991, Takizawa and co-workers reported stoichiometric amount of $Cu(OAc)_2$ mediated direct *ortho*-acetoxylation of 4-alkoxylated phenols. The configuration of the product was confirmed by nuclear Overhauser effect (NOE) experiment, and the possibility of transesterification involving the 1-hydroxy group was ruled out. However, this reaction system is not applicable to the acetoxylation of 4-methylphenols and 4-phenyl-phenols.^[23] In 2016, Jana and co-workers reported $Cu(OAc)_2 \cdot H_2O$ mediated hydroxylation via a bidentate directing group consisted of benzamide and 8-aminoquinoline. Radical trap experiment with (2,2,6,6-tetramethylpiperidin-1-yl)oxyl (TEMPO) found no significant effect on the reaction outcome, thus eliminating the radical pathway (Eq. 5).^[24]

In 2006, Yu and co-workers observed that an equimolar amount of $Cu(OAc)_2$ could mediate the *ortho*-hydroxylation of 2-phenylpyridine with 1 equiv of H_2O in CH_3CN under O_2 (101 kPa) at 130 °C. Labeling experiments using $H_2^{18}O$ demonstrated that the oxygen atom from $Cu(OAc)_2$ was incorporated into product. The author hypothesized the binding of the hydroxyl group and the nitrogen in the product to $Cu(II)$ might prevent catalytic turnover. By adding Ac_2O to the reaction system, the copper loading was successfully reduced to 10 mol % under O_2 atmosphere, albeit resulting in significant diacetoxylation (Eq. 6).^[25]

3.2 Palladium-catalyzed *ortho*-C—H oxidation

3.2.1 Palladium-catalyzed *ortho*-C—H oxidation via a 5-membered intermediate

Palladium-catalyzed *ortho*-C—H oxidation via a 5-membered intermediate is shown in Scheme 3.

Scheme 3 Palladium-catalyzed *ortho*-C—H oxidation via a 5-membered intermediate

CH₂NR₂ as DG In 1990s, Boersma, van Koten, and co-workers studied the generation of heteroleptic diarylpalladium compounds and the oxidation of Pd—Ar bond using *tert*-butyl hydroperoxide (TBHP) as the terminal oxidant. With the assistance of CH₂NR₂ group, this transformation regio-selectively delivered mono-hydroxylated compounds (Scheme 4).^[26]

Scheme 4 Oxidation of palladium complex **14** with TBHP

Carbonyl group as DG In 2009, Yu and co-workers described the Pd(OAc)₂-catalyzed regioselective *ortho*-hydroxylation of potassium benzoates or benzoic acids, employing the environmentally friendly molecular oxygen as the terminal oxidant (Eq. 7). The presence of benzoquinone (BQ) was found to significantly increase the reaction yield, however, it is not essential. Labeling experiment with ¹⁸O₂ proved that the oxygen-atom incorporated into the hydroxylated product is originated from molecular oxygen, while oxygen incorporation from H₂O was ruled out

via labeling experiment using H₂¹⁸O, indicating a direct oxygenation of the arylpalladium intermediates by molecular oxygen.^[27]

In 2012, Rao and co-workers developed a palladium-catalyzed *ortho*-hydroxylation system with selectfluor or K₂S₂O₈ as the oxidant and a TFA/TFAA (9 : 1) mixture as the solvent. This system has proved efficacy toward ketone-, ester-, amide-, acetanilide-, and sulfonamide-directed benzenes.^[28]

Imine, oxime or azo as DG Imine, oxime or azo as DG is shown in Scheme 5.

Scheme 5 Imine, oxime or azo as DG

In 2004, Sanford and co-workers proved the feasibility of Pd(OAc)₂-catalyzed *ortho*-selective C—H acetoxylation with imine as the directing group and PhI(OAc)₂ as the terminal oxidant, delivering a moderate yield (Scheme 6a).^[29] Later, Reddy and co-workers reported the acetoxylation of 1-aryl-3,4-dihydroisoquinolines under similar reaction conditions devoid of additional stoichiometric amount of Ac₂O (Scheme 6b).^[30]

Scheme 6 Pd(OAc)₂-catalyzed imine-directed *ortho*-selective C—H acetoxylation

Transient directing group (TDG) strategy has recently boomed in transition metal-catalyzed C—H functionalizations.^[31,32] To overcome the weak coordination between formyl group and transition metal, Sorensen and co-workers developed palladium-catalyzed *ortho*-hydroxylation of arylaldehyde utilizing anthranilic acid as the TDG (Scheme 7). The authors proposed that the reaction might go through a transient imine intermediate with catalytic amount of amines. This catalytic system could work with both electron-donating and electron-withdrawing groups substituted benzaldehydes.^[33]

In 2006, Sanford and co-workers reported Pd(OAc)₂-catalyzed acetoxylation of arenes with oxime ether as the

Scheme 7 $\text{Pd}(\text{OAc})_2$ -catalyzed *ortho*-C—H hydroxylation of arylaldehyde via a TDG strategy

directing group. They tested various oxidants, and found significant quantities of the *ortho*-acetoxyated product could be obtained with oxone, $\text{K}_2\text{S}_2\text{O}_8$, or $\text{PhI}(\text{OAc})_2$ as the oxidant (Eq. 8), however, the latter two oxidants could also induce the formation of small amount of di-*ortho*-acetoxyated product.^[34]

Oxidant	Isolated yield/%	Oxidant	Isolated yield/%
H_2O_2 •urea	10	$\text{CH}_3\text{CO}_3\text{H}$	34
50% aq. H_2O_2	11	Oxone	68
<i>m</i> -CPBA	14	$\text{K}_2\text{S}_2\text{O}_8$	76
70% aq. $t\text{BuOOH}$	18	$\text{PhI}(\text{OAc})_2$	81

With PPh_3 as the ligand and $\text{CHCl}_2\text{CHCl}_2$ as the solvent, Jiao and co-workers developed a strategy for $\text{Pd}(\text{OAc})_2$ -catalyzed *ortho*-C—H hydroxylation of aryl oxime ethers (Eq. 9).^[35] A $\text{Pd}^{\text{II}}/\text{Pd}^{\text{IV}}$ catalytic cycle was proposed as follows: firstly, oxime ether assisted *ortho*-C—H palladation formed a 5-membered palladacycle; secondly, oxidative addition of $\text{K}_2\text{S}_2\text{O}_8$ with Pd^{II} afforded a Pd^{IV} intermediate; finally, reductive elimination gave the hydroxylated product and regenerated Pd^{II} catalyst. The ligand was supposed to accelerate the C—O bond reductive elimination step (Scheme 8).

Scheme 8 Proposed catalytic cycle of $\text{Pd}(\text{OAc})_2$ -catalyzed *ortho*-hydroxylation of aryl oxime ethers

Azobenzene has been known as a functional group in among others Sudan Red B. Moreover, it could be used as the directing group in palladium-catalyzed C—H oxidations since the beginning of this century. In 2004, Sanford and co-workers disclosed $\text{Pd}(\text{OAc})_2$ -catalyzed azo-assisted *ortho*-C—H acetoxylation.^[29] Later, Joseph and co-workers developed $\text{Pd}(\text{OAc})_2$ -catalyzed *ortho*-C—H hydroxylation of symmetrical/unsymmetrical azobenzenes via a $\text{PhI}(\text{OTFA})_2/\text{DCE}$ system, generating unsymmetrical azophenols (Eq. 10).^[36]

In 2014, Zeng and co-workers reported $\text{Pd}(\text{OAc})_2$ -catalyzed *ortho*-esterification of azoarenes using aryl acylperoxide as the oxygen source (Eq. 11). The chemo-selectivity shifted with different solvents and temperatures: C—O functionalized product will be generated with CH_3CN as the solvent at 60 °C, while arylated product will be obtained with PhCl as the solvent at 130 °C. In contrast with the previously discussed $\text{Pd}^{\text{II}} \sim \text{Pd}^{\text{IV}}$ catalytic cycle, the authors proposed a radical mechanism, which might undergo a Pd^{III} or Pd^{IV} intermediate during the catalytic cycle (Scheme 9).^[37]

Scheme 9 Proposed catalytic cycle of $\text{Pd}(\text{OAc})_2$ -catalyzed *ortho*-esterification of azoarenes with aryl acylperoxide

In 2015, Cui, Wu and co-workers discovered the palladium-catalyzed *ortho*-esterification/etherification of azoxybenzenes at 100 °C or even at ambient temperature (Scheme 10). Based on mechanistic studies, a radical reaction pathway was proposed, including the formation of Pd-substrate complex via *ortho*-C—H activation, the generation of oxygen radical and its addition to the palladium center, and reductive elimination to afford the oxygenated product (Scheme 11).^[38]

N-Heterocycle as DG Nitrogen-containing heterocycles could be applied as the directing group in palladium-catalyzed *ortho*-selective aromatic oxidation. In early 2000s, Sanford and co-workers utilized the $\text{Pd}(\text{OAc})_2/\text{PhI}(\text{OAc})_2$ system for regio-selective C—H oxidation of arenes bearing pyridine or other *N*-heterocyclic substitu-

Scheme 10 $\text{Pd}(\text{O}_2\text{CCF}_3)_2$ -catalyzed *ortho*-esterification/etherification of azoxybenzenes

Scheme 11 Proposed catalytic cycle of $\text{Pd}(\text{O}_2\text{CCF}_3)_2$ -catalyzed *ortho*-oxidation of azoxybenzenes

ents, obtaining mono- or di-oxidized products. Various functional groups are well tolerated, including nitro, fluoride, bromide, and *etc.*^[29,39] Very recently, Correa and co-workers reported $\text{Pd}(\text{OAc})_2$ -catalyzed C—H acetoxylation/pivaloxylation of arenes with a triazole directing group, which could be easily prepared via click chemistry.^[40] Besides $\text{PhI}(\text{OAc})_2$, other oxidants like oxone, TBHP, and H_2O_2 have also been successfully employed in palladium-catalyzed aromatic C—H oxidation by the group of Kim,^[41] Sun,^[42] and Itoh,^[43] respectively, delivering numerous *ortho*-hydroxylated products (Eq. 12).

The detailed mechanism of palladium-catalyzed *ortho*-

C—H esterification on 2-phenylpyridine with hypervalent iodine as the oxidant was investigated by Sanford and Ritter, respectively. In 2005, Sanford and co-workers disclosed the Pd^{II}-Pd^{IV} catalytic cycle via mechanistic study. They designed and synthesized a series of stable Pd^{IV} complexes and presented detailed study of reductive elimination from this oxidation state which finally afforded oxygenated product (Eq. 13).^[44] Later, Sanford and Ritter *et al.* respectively proposed a Pd^{II}-Pd^{III} catalytic cycle in palladium-catalyzed C—H oxidation of 2-phenylpyridine as was substantiated by X-ray or kinetic isotope effect studies.^[45–47]

In 2015, Jiao and co-workers reported the palladium-catalyzed *ortho*-hydroxylation of 2-phenylpyridine. Interestingly, this system employed catalytic amount of NHPI under oxygen atmosphere (Scheme 12).^[48] Isotope labeling strategy proved that the oxygen atom in the hydroxy group originates from molecular oxygen. KIE study ($K_H/K_D=1.86$) indicated that C—H bond cleavage might be the rate-limiting step. Based on mechanistic studies, the catalytic cycle was as proposed, the radical stimulation and propagation should follow the sequence NHPI→PINO→benzyl radical→peroxide radical→hydroxyl radical, which will finally intervened in the palladium catalytic cycle (Scheme 12).

3.2.2 Palladium-catalyzed *ortho*-C—H oxidation via a 6-membered intermediate

Palladium-catalyzed *ortho*-C—H oxidation via a 6-membered intermediate is shown in Scheme 13.

CH₂-N-heterocycle as DG The diverse CH₂-N-heterocycle type directing group-assisted palladium-catalyzed aromatic C—H oxidation was respectively investigated by Sanford and Correa (Eq. 14). With a methylene spacer between the directing group and the aromatic C—H bond, the catalytic cycle would generate a 6-membered palladacycle. Individual kinetic isotope effect experiments were performed with both electron-donating and electron-withdrawing group substituted directing groups. The results demonstrated that reactions could be acceler-

Scheme 12 Palladium-catalyzed *ortho*-hydroxylation of 2-phenylpyridine with NHPI as the co-catalyst

Scheme 13 Palladium-catalyzed *ortho*-C—H oxidation via a 6-membered intermediate

ated by electron-withdrawing groups. A significant kinetic isotope effect was observed, indicating that the formation of the 6-membered palladacycle is the rate determining step.^[34,40,49]

Anilide or carbamate as DG During the past decades, several groups reported anilide-assisted palladium-catalyzed aromatic C—H oxidation with PhI(O₂CR)₂, oxone or K₂S₂O₈ as the terminal oxidant, delivering *ortho*-selective acetoxyated product (Eq. 15).^[34,39,50,51] The catalytic system could tolerate various functional groups, including nitro, fluoro, bromo, ketone, and *etc.*

In 2014, Rao and co-workers reported carbamate-assisted palladium-catalyzed *ortho*-C—H hydroxylation, using $K_2S_2O_8$ as the oxidant and a mixture of TFA/TFAA (1 : 1) as the solvent (Eq. 16). This catalytic system could proceed even at room temperature, however, delivering isolated yields of less than 50%.^[52]

Silicon-bridged DG Silicon-tethered directing groups have been used as easily installable and removable/modifiable auxiliaries for C—H functionalization reactions.^[53] In the early 2010s, Gevorgyan and co-workers applied the PyDipSi group as general directing group in palladium-catalyzed *ortho*-selective mono- or di-oxygenation of arenes, providing access to a variety of acetoxylated and pivaloxylated aromatic compounds in good yields (Scheme 14).^[54] It is notable that the palladium-catalyzed oxygenation system not only worked on simple arenes, but also on heterocycles like benzofuran and indole. After the reaction, PyDipSi group could be easily removed or converted into other functional groups.

Scheme 14 PyDipSi-directed palladium-catalyzed *ortho*-oxygenation of arenes

Silanol has also been employed as the directing group in palladium-catalyzed C—H oxidation, delivering a cyclic silicon-protected catechol (Scheme 15). The catechol product could be obtained via a routine desilylation of the silacyle with tetrabutylammonium fluoride (TBAF).^[55] To identify the source of oxygen of newly formed aryl silicon ether, labeling experiment was performed with ^{18}O -labelled silanol, after the reaction ^{18}O was detected only in HOAc rather than in the cyclized product. Based on the mechanistic studies, the authors proposed the catalytic cycle, in which reductive cyclization was forbidden, instead, acetoxylation followed by nucleophilic addition delivered the final product (Scheme 16).

Scheme 15 Palladium-catalyzed synthesis of cyclic silicon-protected catechol

Scheme 16 Proposed mechanism of palladium-catalyzed synthesis of cyclic silicon-protected catechol

3.3 Ruthenium-catalyzed *ortho*-C—H oxidation

3.3.1 Ruthenium-catalyzed *ortho*-C—H oxidation via a 5-membered intermediate

Ruthenium-catalyzed *ortho*-C—H oxidation via a 5-me-

numbered intermediate is shown is Scheme 17.

Scheme 17 Ruthenium-catalyzed *ortho*-C—H oxidation via a 5-membered intermediate

Carbonyl group as DG Recent years have witnessed a tremendous development in the *ortho*-selective C—H oxidation of arenes and heteroarenes with readily accessible ruthenium catalysts.^[5] In 2012, Rao and co-workers investigated $[\text{RuCl}_2(p\text{-cymene})]_2$ -catalyzed *ortho*-C—H hydroxylation of benzoates. The screening of oxidants demonstrated $\text{K}_2\text{S}_2\text{O}_8$ and selectfluor to be the optimized oxidants, meanwhile, a TFA/TFAA solvent mixture turned out to be critical. Notably, this catalytic system is valid towards heteroarenes. When using ethyl 1-naphthoate as the substrate, *peri*-hydroxylated product was found to be predominate isomer rather than *ortho*-hydroxylated product (Eq. 17).^[56]

In 2012, Ackermann and co-workers reported ruthenium-catalyzed amide-directed C—H hydroxylation on arenes. Several ruthenium catalysts were examined, out of which the well-defined ruthenium carboxylate complex $[\text{Ru}(\text{O}_2\text{CMes})_2(p\text{-cymene})]$ ^[57] was found to be the optimal catalyst at a remarkably low catalyst loading of only 1 mol% (Scheme 18).^[58] Later, Weinreb amide was applied as the directing group in ruthenium-catalyzed *ortho*-C—H hydroxylation of arenes with a similar reaction system but at a milder temperature. In comparison with simple amides, Weinreb amide exhibited more structural flexibility. After reaction, *ortho*-hydroxylated aryl Weinreb amide could be easily converted into salicylaldehyde.^[59]

In 2016, Zhang, Rao, and co-workers investigated the C—H oxidation of *N*-methyl-*N*-phenylbenzamide using $\text{K}_2\text{S}_2\text{O}_8$ as the oxidant and TFA/TFAA as the solvent (Scheme 19). It was found that ruthenium and palladium exhibited different regio-selectivities. They rationalized the

Scheme 18 Ruthenium-catalyzed *ortho*-C—H hydroxylation of benzamides and Weinreb amides

difference of regio-selectivities to the polarity of transient state via density functional theory (DFT) studies (Figure 1).^[51]

Scheme 19 Palladium-/ruthenium-catalyzed *ortho*-C—H hydroxylation of *N*-methyl-*N*-phenylbenzamide

Figure 1 Regio-selectivities to the polarity of transient state via DFT studies

In 2012 and 2013, Ackermann and Rao respectively reported ruthenium-catalyzed *ortho*-C—H bond oxygenations of arenes with weakly coordinating ketones as the directing group.^[60,61] In general, the reaction system employed oxone, K₂S₂O₈, Selectfluor, or PhI(OAc)₂ as the terminal oxidant and [RuCl₂(*p*-cymene)]₂, [RuCl₃(H₂O)_n] or [Ru(O₂CMe)₂(*p*-cymene)] as the catalyst, delivering mono-hydroxylated arylketones or mono/di-hydroxylated diarylketones with ample scope and excellent functional group tolerance. The products could be further converted into various functional molecules. Very recently, flavone and chromone derivatives have been applied in ruthenium-catalyzed site-selective hydroxylation, again reflecting the potential of this approach in the synthesis of bioactive molecules (Eq. 18).^[62]

Aldehyde is supposed to be the weakest carbonyl directing group. It is especially challenging to achieve aldehyde-assisted *ortho*-hydroxylation due to the inherent tendency of formyl group oxidation under oxidative environment, which will lead to the formation of benzoic acid. In 2014, Ackermann and co-workers developed the [RuCl₂(*p*-cymene)]₂/PhI(O₂CCF₃)₂/DCE system which could successfully convert arylaldehyde to salicylaldehyde with high chemo- and regio-selectivities. Intramolecular competition experiments were performed with benzaldehyde bearing an amide, ketone, or ester substituent, indicating that the coordinating ability of aldehyde is weaker than amide, ketone, and ester.⁶³ In contrast, all other transition metals fell short in providing the desired C—H oxygenation (Eq. 19).

R = N(<i>i</i> Pr) ₂ :	70%	0%
R = <i>t</i> -Bu:	76%	0%
R = OEt:	13%	0%
	+ 59% s.m.	

N-Heterocycle as DG Very recently, Singh and co-workers reported ruthenium-catalyzed *ortho*-hydroxylation of 2-phenylpyridine. The catalysts, naming SPS-Bpy and SPS-DB-Bpy, were derived from [Ru(O₂CMe)₂(*p*-cymene)] with bipyridine or dibromobipyridine as the respective substituents. These catalysts were also found to be highly chemo-selective in converting aromatic carboxylate esters. The authors presumed a Ru(II)-Ru(IV) catalytic cycle, including *ortho*-ruthelation, oxidative formation of C—Ru—O bond, and reductive elimination to afford the final product and regenerate the active catalyst (Scheme 20).^[64]

R = H, Br; L = *p*-cymene; L¹ = bipyridine, dibromobipyridine

Scheme 20 Proposed catalytic cycle of *ortho*-hydroxylation of 2-phenylpyridine with **81** as the pre-catalyst

3.3.2 Ruthenium-catalyzed *ortho*-C—H oxidation via a 6-membered ring transition state

Ruthenium-catalyzed *ortho*-C—H oxidation via a 6-membered ring transition state is shown in Scheme 21.

Scheme 21 Ruthenium-catalyzed *ortho*-C—H oxidation via a 6-membered ring transition state

Anilide or carbamate as DG In 2013, Rao and co-workers reported the synthesis of 2-aminophenols and 2-aminobenzene-1,3-diols via ruthenium-catalyzed *ortho*-selective mono- or di-hydroxylation with anilide⁶⁵ as a removable directing group.^[66] The reaction demonstrates excellent reactivity, regioselectivity, and good functional group tolerance. Besides removal from hydroxylated products, the *ortho*-substituted benzoyl group could be

further utilized in the synthesis of heterocycles, such as benzoxazoles and dibenzoxazepines (Scheme 22).

Scheme 22 Anilide-directed ruthenium-catalyzed *ortho*-hydroxylation of arenes

In the same year, Ackermann and co-workers independently reported the application of carbamate as the directing group in ruthenium-catalyzed *ortho*-hydroxylation. This catalytic system was widely applicable and could efficiently convert *para*-, *meta*-, and even more sterically hindered *ortho*-substituted aryl carbamates bearing various valuable electrophilic functional groups like chloro, bromo, or iodo substituents.^[67] Later, Rao and co-workers reported that the ruthenium-catalyzed carbamate assisted C—H hydroxylation system using $K_2S_2O_8$ as the oxidant and TFA/TFAA as the solvent.^[52] It is notable that carbamate directed *ortho*-hydroxylation is useful for the synthesis of catechol derivatives from easily accessible phenols (Scheme 23).

Scheme 23 Carbamate-directed ruthenium-catalyzed *ortho*-hydroxylation of arenes

3.4 Rhodium-catalyzed *ortho*-C—H oxidation

In 2013, Rao and co-workers reported $Rh(OAc)_2$ -catalyzed *ortho*-selective C—H hydroxylation on diarylketones with $K_2S_2O_8$ as the terminal oxidant and TFA/TFAA (9 : 1) as the solvent, delivering mono- or 2,2'-dihydroxylated

products in moderate to good yields. This method is also valid for the *ortho*-hydroxylation of benzoates and benzamides (Scheme 24).^[61]

Scheme 24 Ketone-directed rhodium-catalyzed *ortho*-hydroxylation of arenes

To understand the detailed mechanism of rhodium-catalyzed aromatic C—H oxidation, Jones, Templeton, and co-workers studied the oxygen insertion into the Cp^* -Rh-bpp half-sandwich complex using 2-*tert*-butylsulfonyliodosylbenzene (sPhIO) as the oxidant, forming a Rh(IV) complex. Treatment with acid generated 2-pyridyl-phenol (Scheme 25).^[68]

Scheme 25 Formation of Rh(IV) complex **90** via oxygen insertion

3.5 Iridium-catalyzed *ortho*-C—H oxidation

In 2017, Zhao and co-workers developed a redox-neutral strategy for catechol synthesis via iridium-catalyzed *ortho*-hydroxylation with an internal oxidant.^[69] The reaction was performed at room temperature and could tolerate various functional groups like halogens, carboxylates, heterocycles, and even chiral substituents. To understand the oxygen source, isotope labelling experiments were carried out, demonstrating an internal oxidant pathway. The proposed catalytic cycle started from iridium-catalyzed *ortho*-C—H activation, forming a 7-membered transition state, which underwent reductive elimination to form the C—O bond, oxidative insertion of Ir^I (Scheme 26) into the N—O bond and subsequently protonation afforded the original Ir^{III} catalyst and the imine compound, which would be converted to catechols via hydrolysis.

Scheme 26 Proposed catalytic cycle of iridium-catalyzed catechol synthesis with internal oxidant

4 meta-Selective aromatic C—H oxidation

4.1 Ligand controlled *meta*-C—H oxidation

In 2013, Sanford and co-workers disclosed that the regio-selectivity of palladium-catalyzed C—H acetoxylation could be altered with acridine as the ancillary ligand and $\text{MesI}(\text{OAc})_2$ as the terminal oxidant, delivering *meta*-acetoxylation arene (Scheme 27), thus overrode the substrate electronic bias that dominates the regio-selectivity observed in the reaction that used ligand-free $\text{Pd}(\text{OAc})_2$ as the catalyst and $\text{PhI}(\text{OAc})_2$ as the oxidant.^[70]

Scheme 27 Ligand-controlled regio-selectivity shift of palladium-catalyzed aromatic acetoxylation

4.2 Template assisted *meta*-C—H oxidation

Very recently, template-directed *meta*-selective $\text{C}(\text{sp}^2)$ —H functionalization has been recognized as a powerful tool in synthetic chemistry.^[71] In the category of transition metal-catalyzed *meta*-selective C—H oxidations, palladium-catalyzed *meta*-selective C—H oxygenation has been recently achieved with the assistance of a U-shaped template bearing a sulfonate or phosphonate linkage. The catalytic system has exhibited excellent *meta*-selectivity, delivering mono-hydroxylated/acetoxylation products in high yields (Scheme 28).^[72]

Based on mechanistic studies, the author proposed a $\text{Pd}^{\text{II}}/\text{Pd}^{\text{IV}}$ catalytic cycle: Firstly, the palladium catalyst coordinated with cyanide group, followed by the *meta*-C—H activation to form an 11-membered ring transition state. Pd^{II} was then oxidized to Pd^{IV} by hypervalent iodine rea-

Scheme 28 $\text{Pd}(\text{OAc})_2$ -catalyzed *meta*-selective aromatic hydroxylation/acetoxylation

gent. Finally, reductive elimination delivered the oxidized product and regenerated Pd^{II} catalyst (Scheme 29).

Scheme 29 Proposed catalytic cycle of palladium-catalyzed *meta*-acetoxylation

5 para-Selective aromatic C—H oxidation

5.1 Para-C—H oxidation of anisoles

The ruthenium-catalyzed *para*-C—H oxidation without a Lewis base directing group was achieved by Ackermann

and co-workers.^[67] Taking advantage of the $[\text{RuCl}_2(p\text{-cymene})]_2/\text{PhI}(\text{O}_2\text{CCF}_3)_2/\text{DCE}$ system, anisole derivatives were converted to the corresponding *para*-hydroxylated phenols. Oxidations of anisole in the presence of catalytic (10 mol%) or stoichiometric (1 equiv.) amounts of TEMPO furnished product in significantly reduced yields of 43% and 5%, respectively, indicating a single-electron-transfer (SET) pathway. However, the ruthenium catalyst's exact mode of action is as of yet not completely understood (Eq. 21).

5.2 Template assisted *para*-C—H oxidation

In 2015, Maiti and co-workers disclosed the palladium-catalyzed *para*-acetoxylation of substituted toluene derivatives (Scheme 30).^[73,74] High *para*-selectivity was achieved with the assistance of a U-shaped template. Installing two sterically hindered isopropyl groups at the Si atom might allow closer approach of the coordinating group to the *para*-C—H bond via the Thorpe-Ingold effect. In the control experiment without cyanide group, no *para*-acetoxylation product was generated.

6 Conclusions and perspectives

Recent years have witnessed tremendous progress in transition metal-catalyzed aromatic C—H oxidations. Enormous amount of phenols and phenolic derivatives have been synthesized via this strategy, in the meantime, huge advances have been made in the mechanistic study of these transformations. However, there are still many challenges to be addressed, such as developing cost-competitive transition metal catalysts, exploring switchable directing groups, expanding substrate scope, and reducing reaction temperature. In addition, there are two points deserve particular attention:

Point 1. Chelation-assisted aromatic C—H oxidation has been comprehensively studied over the past few years, delivering *ortho*-functionalized product. However, there have been only a few examples of *meta*- and *para*-selective aromatic C—H oxidations. Considering the diversity of functional molecules, the developing of synthetic methodologies for *meta*- and *para*-hydroxylated arenes are undoubtedly in great demand.

Point 2. Most of the oxidants are generally expensive and may cause environmental risks. The utilization of in-

Scheme 30 $\text{Pd}(\text{OAc})_2$ -catalyzed *para*-selective aromatic acetoxylation

expensive peroxides, oxygen or even air as a green oxidant in transition metal-catalyzed aromatic C—H oxidations is more advantageous, thus worth further research. However, the usually used high pressure reaction system could limit the development of this strategy. An additional redox catalyst might be the key solution to facilitate the reaction at ambient atmospheric pressure.

References

- [1] (a) Rappoport, Z. *The Chemistry of Phenols*, Wiley-VCH, Weinheim, **2003**.
 (b) Fiegel, H.; Voges, H. W.; Hamamoto, T.; Umemura, S.; Iwata, T.; Miki, H.; Fujita, Y.; Buysch, H. J.; Garbe, D.; Paulus, W. *Phenol Derivatives in Ullmann's Encyclopedia of Industrial Chemistry*, Wiley-VCH, New York, **2002**.
 (c) Hartwig, J. F. In *Handbook of Organopalladium Chemistry for Organic Synthesis*, Vol. 1, Ed.: Negishi, E.-I., Wiley-Interscience, New York, **2002**, p. 1097.
 (d) Tyman, J. H. P. *Synthetic and Natural Phenols*, Elsevier, New York, **1996**.
 (e) Liao, S.; Zhang, G.; Cao, H.; Chen, B.; Li, W.; Wu, X.; Feng, Y.

- Chin. J. Org. Chem.* **2018**, *38*, 1549.
- (f) Lin, W.; Cai, Q.; Zheng, C.; Huang, Z.; Shi, D. *Chin. J. Org. Chem.* **2017**, *37*, 2094.
- (g) Li, Z.; Kang, S.; Chen, L.; Wang, Y.; Li, J. *Chin. J. Org. Chem.* **2016**, *36*, 1143.
- [2] Lyons, T. W.; Sanford, M. S. *Chem. Rev.* **2010**, *110*, 1147.
- [3] Enthaler, S.; Company, A. *Chem. Soc. Rev.* **2011**, *40*, 4912.
- [4] Thirunavukkarasu, V. S.; Kozhushkov, S. I.; Ackermann, L. *Chem. Commun.* **2014**, *50*, 29.
- [5] Newhouse, T.; Baran, P. S. *Angew. Chem. Int. Ed.* **2011**, *50*, 3362.
- [6] Liang, Y.-F.; Jiao, N. *Acc. Chem. Res.* **2017**, *50*, 1640.
- [7] Mandal, S.; Bera, T.; Dubey, G.; Saha, J.; Laha, J. K. *ACS Catal.* **2018**, *8*, 5085.
- [8] (a) Huang, Z.; Dong, G. *Acc. Chem. Res.* **2017**, *50*, 465.
(b) Hartwig, J. F. *Acc. Chem. Res.* **2017**, *50*, 549.
(c) Kozłowski, M. C. *Acc. Chem. Res.* **2017**, *50*, 638.
- [9] For selected recent examples, see:
(a) Raghuvanshi, K.; Zell, D.; Ackermann, L. *Org. Lett.* **2017**, *19*, 1278.
(b) Raghuvanshi, K.; Rauch, K.; Ackermann, L. *Chem.-Eur. J.* **2015**, *21*, 1790.
(c) Hao, X.-Q.; Chen, L.-J.; Ren, B.; Li, L.-Y.; Yang, X.-Y.; Gong, J.-F.; Niu, J.-L.; Song, M.-P. *Org. Lett.* **2014**, *16*, 1104.
(d) Shi, S.; Kuang, C. J. *Org. Chem.* **2014**, *79*, 6105.
(e) Zhang, C.; Sun, P. J. *Org. Chem.* **2014**, *79*, 8457.
(f) Péron, F.; Fossey, C.; Sopkova-de Oliveira Santos, J.; Cailly, T.; Fabis, F. *Chem.-Eur. J.* **2014**, *20*, 7507.
(g) Bhadra, S.; Dzik, W. I.; Gooßen, L. J. *Angew. Chem., Int. Ed.* **2013**, *52*, 2959.
(h) Bhadra, S.; Matheis, C.; Katayev, D.; Gooßen, L. J. *Angew. Chem., Int. Ed.* **2013**, *52*, 9279.
(i) Suess, A. M.; Ertem, M. Z.; Cramer, C. J.; Stahl, S. S. *J. Am. Chem. Soc.* **2013**, *135*, 9797.
(j) Roane, J.; Daugulis, O. *Org. Lett.* **2013**, *15*, 5842.
(k) Eom, D.; Jeong, Y.; Kim, Y. R.; Lee, E.; Choi, W.; Lee, P. H. *Org. Lett.* **2013**, *15*, 5210.
(l) Zhao, J.; Wang, Y.; He, Y.; Liu, L.; Zhu, Q. *Org. Lett.* **2012**, *14*, 1078.
(m) Zhao, J.; Zhang, Q.; Liu, L.; He, Y.; Li, J.; Li, J.; Zhu, Q. *Org. Lett.* **2012**, *14*, 5362.
(n) Li, W.; Sun, P. J. *Org. Chem.* **2012**, *77*, 8362.
(o) Jiang, T.-S.; Wang, G.-W. *J. Org. Chem.* **2012**, *77*, 9504.
(p) Xiao, B.; Gong, T.-J.; Liu, Z.-J.; Liu, J.-H.; Luo, D.-F.; Xu, J.; Liu, L. *J. Am. Chem. Soc.* **2011**, *133*, 9250.
(q) Anand, M.; Sunoj, R. B. *Org. Lett.* **2011**, *13*, 4802.
(r) Wei, Y.; Yoshikai, N. *Org. Lett.* **2011**, *13*, 5504.
(s) Wang, X.; Lu, Y.; Dai, H.-X.; Yu, J.-Q. *J. Am. Chem. Soc.* **2010**, *132*, 12203.
(t) Wang, G.-W.; Yuan, T.-T. *J. Org. Chem.* **2010**, *75*, 476.
- [10] For selected reviews and recent examples, see:
(a) Tang, S.; Liu, Y.; Lei, A. *Chem* **2018**, *4*, 27.
(b) Tang, S.; Zeng, L.; Lei, A. *J. Am. Chem. Soc.* **2018**, *140*, 13128.
(c) Ma, C.; Fang, P.; Mei, T.-S. *ACS Catal.* **2018**, *8*, 7179.
(d) Sauermann, N.; Meyer, T. H.; Qiu, Y.; Ackermann, L. *ACS Catal.* **2018**, *8*, 7086.
(e) Sauermann, N.; Meyer, T. H.; Ackermann, L. *Chem.-Eur. J.* **2018**, *24*, 16209.
(f) Jiao, K.-J.; Li, Z.-M.; Xu, X.-T.; Zhang, L.-P.; Li, Y.-Q.; Zhang, K.; Mei, T.-S. *Org. Chem. Front.* **2018**, *5*, 2244.
(g) Shao, A.; Li, N.; Gao, Y.; Zhan, J.; Chiang, C.-W.; Lei, A. *Chin. J. Chem.* **2018**, *36*, 619.
(h) Yang, Q.-L.; Fang, P.; Mei, T.-S. *Chin. J. Chem.* **2018**, *36*, 338.
(i) Jiao, K.-J.; Zhao, C.-Q.; Fang, P.; Mei, T.-S. *Tetrahedron Lett.* **2017**, *58*, 797.
(j) Yuan, Y.; Chen, Y.; Tang, S.; Huang, Z.; Lei, A. *Sci. Adv.* **2018**, *4*, eaat5312.
(k) Shrestha, A.; Lee, M.; Dunn, A. L.; Sanford, M. S. *Org. Lett.* **2018**, *20*, 204.
(l) Sauermann, N.; Meyer, T. H.; Tian, C.; Ackermann, L. *J. Am. Chem. Soc.* **2017**, *139*, 18452.
(m) Yang, Q.-L.; Li, Y.-Q.; Ma, C.; Fang, P.; Zhang, X.-J.; Mei, T.-S. *J. Am. Chem. Soc.* **2017**, *139*, 3293.
(n) Li, Y.-Q.; Yang, Q.-L.; Fang, P.; Mei, T.-S.; Zhang, D. *Org. Lett.* **2017**, *19*, 2905.
- [11] A recent example: Zhang, Z.-J.; Quan, X.-J.; Ren, Z.-H.; Wang, Y.-Y.; Guan, Z.-H. *Org. Lett.* **2014**, *16*, 3292.
- [12] (a) Jintoku, T.; Taniguchi, H.; Fujiwara, Y. *Chem. Lett.* **1987**, 1865.
(b) Jintoku, T.; Takaki, K.; Fujiwara, Y.; Fuchita, Y.; Hiraki, K. *Bull. Chem. Soc. Jpn.* **1990**, *63*, 438.
(c) Jintoku, T.; Nishimura, K.; Takaki, K.; Fujiwara, Y. *Chem. Lett.* **1990**, 1687.
- [13] Yamada, S.; Sakaguchi, S.; Ishii, Y. *J. Mol. Catal. A Chem.* **2007**, *262*, 48.
- [14] Yoneyama, T.; Crabtree, R. H. *J. Mol. Catal. A: Chem.* **1996**, *108*, 35.
- [15] Cook, A. K.; Sanford, M. S. *J. Am. Chem. Soc.* **2015**, *137*, 3109.
- [16] Gary, J. B.; Cook, A. K.; Sanford, M. S. *ACS Catal.* **2013**, *3*, 700.
- [17] Shibahara, F.; Kinoshita, S.; Nozaki, K. *Org. Lett.* **2004**, *6*, 2437.
- [18] Chen, C.-D.; Sheng, W.-B.; Shi, G.-J.; Guo, C.-C. *J. Phys. Org. Chem.* **2013**, *26*, 23.
- [19] Casella, L.; Rigoni, L. *J. Chem. Soc., Chem. Commun.* **1985**, 1668.
- [20] Réglier, M.; Amadeï, E.; Tadayoni, R.; Waegell, B. *J. Chem. Soc., Chem. Commun.* **1989**, 447.
- [21] Cruse, R. W.; Kaderli, S.; Meyer, C. J.; Zuberbühler, A. D.; Karlin, K. D. *J. Am. Chem. Soc.* **1988**, *110*, 5020.
- [22] Reinaud, O.; Capdevielle, P.; Maumy, M. *J. Chem. Soc., Chem. Commun.* **1990**, 566.
- [23] Takizawa, Y.; Tateishi, A.; Sugiyama, J.; Yoshida, H.; Yoshihara, N. *J. Chem. Soc., Chem. Commun.* **1991**, 104.
- [24] Singh, B. K.; Jana, R. *J. Org. Chem.* **2016**, *81*, 831.
- [25] Chen, X.; Hao, X.-S.; Goodhue, C.-E.; Yu, J.-Q. *J. Am. Chem. Soc.* **2006**, *128*, 6790.
- [26] (a) Valk, J.-M.; van Belzen, R.; Boersma, J.; Spek, A. L.; van Koten, G. *J. Chem. Soc., Dalton Trans.* **1994**, 2293.
(b) Valk, J.-M.; Boersma, J.; van Koten, G. *Organometallics* **1996**, *15*, 4366.
- [27] Zhang, Y.-H.; Yu, J.-Q. *J. Am. Chem. Soc.* **2009**, *131*, 14654.
- [28] Shan, G.; Yang, X.; Ma, L.; Rao, Y. *Angew. Chem., Int. Ed.* **2012**, *51*, 13070.
- [29] Dick, A. R.; Hull, K. L.; Sanford, M. S. *J. Am. Chem. Soc.* **2004**, *126*, 2300.
- [30] Subba Reddy, B. V.; Umadevi, N.; Narasimulu, G.; Yadav, J. S. *Tetrahedron Lett.* **2012**, *53*, 6091.
- [31] Zhao, Q.; Poisson, T.; Pannecoucke, X.; Besset, T. *Synthesis* **2017**, *49*, 4808.
- [32] Gandeepan, P.; Ackermann, L. *Chem* **2018**, *4*, 199.
- [33] Chen, X.-Y.; Ozturk, S.; Sorensen, E. *J. Org. Lett.* **2017**, *19*, 6280.
- [34] Desai, L. V.; Malik, H. A.; Sanford, M. S. *Org. Lett.* **2006**, *8*, 1141.
- [35] Liang, Y.-F.; Wang, X.; Yuan, Y.; Liang, Y.; Li, X.; Jiao, N. *ACS Catal.* **2015**, *5*, 6148.
- [36] Nguyen, T. H. L.; Gigant, N.; Delarue-Cochin, S.; Joseph, D. *J. Org. Chem.* **2016**, *81*, 1850.
- [37] Qian, C.; Lin, D.; Deng, Y.; Zhang, X.-Q.; Jiang, H.; Miao, G.; Tang, X.; Zeng, W. *Org. Biomol. Chem.* **2014**, *12*, 5866.
- [38] Zhang, D.; Cui, X.; Yang, F.; Zhang, Q.; Zhu, Y.; Wu, Y. *Org. Chem. Front.* **2015**, *2*, 951.
- [39] Kalyani, D.; Sanford, M. S. *Org. Lett.* **2005**, *7*, 4149.
- [40] Irastorza, A.; Aizpurua, J. M.; Correa, A. *Org. Lett.* **2016**, *18*, 1080.
- [41] Kim, S. H.; Lee, H. S.; Kim, S. H.; Kim, J. N. *Tetrahedron Lett.* **2008**, *49*, 5863.
- [42] Dong, J.; Liu, P.; Sun, P. *J. Org. Chem.* **2015**, *80*, 2925.
- [43] Yamaguchi, T.; Yamaguchi, E.; Tada, N.; Itoh, A. *Adv. Synth. Catal.* **2015**, *357*, 2017.
- [44] Dick, A. R.; Kampf, J. W.; Sanford, M. S. *J. Am. Chem. Soc.* **2005**, *127*, 12790.
- [45] Stowers, K. J.; Sanford, M. S. *Org. Lett.* **2009**, *11*, 4584.

- [46] Powers, D. C.; Geibel, M. A. L.; Klein, J. E. M. N.; Ritter, T. *J. Am. Chem. Soc.* **2009**, *131*, 17050.
- [47] Powers, D. C.; Xiao, D. Y.; Geibel, M. A. L.; Ritter, T. *J. Am. Chem. Soc.* **2010**, *132*, 14530.
- [48] Yan, Y.; Feng, P.; Zheng, Q.-Z.; Liang, Y.-F.; Lu, J.-F.; Cui, Y.; Jiao, N. *Angew. Chem., Int. Ed.* **2013**, *52*, 5827.
- [49] Desai, L. V.; Stowers, K. J.; Sanford, M. S. *J. Am. Chem. Soc.* **2008**, *130*, 13285.
- [50] Wang, G.-W.; Yuan, T.-T.; Wu, X.-L. *J. Org. Chem.* **2008**, *73*, 4717.
- [51] Sun, Y.-H.; Sun, T.-Y.; Wu, Y.-D.; Zhang, X.; Rao, Y. *Chem. Sci.* **2016**, *7*, 2229.
- [52] Yang, X.; Sun, Y.; Chen, Z.; Rao, Y. *Adv. Synth. Catal.* **2014**, *356*, 1625.
- [53] A review: Parasram, M.; Gevorgyan, V. *Acc. Chem. Res.* **2017**, *50*, 2038.
- [54] (a) Chernyak, N.; Dudnik, A. S.; Huang, C.; Gevorgyan, V. *J. Am. Chem. Soc.* **2010**, *132*, 8270.
(b) Huang, C.; Chernyak, N.; Dudnik, A. S.; Gevorgyan, V. *Adv. Synth. Catal.* **2011**, *353*, 1285.
(c) Gulevich, A. V.; Melkonyan, F. S.; Sarkar, D.; Gevorgyan, V. *J. Am. Chem. Soc.* **2012**, *134*, 5528.
- [55] (a) Huang, C.; Ghavtadze, N.; Chattopadhyay, B.; Gevorgyan, V. *J. Am. Chem. Soc.* **2011**, *133*, 17630.
(b) Huang, C.; Ghavtadze, N.; Godoi, B.; Gevorgyan, V. *Chem.-Eur. J.* **2012**, *18*, 9789.
- [56] Yang, Y.; Lin, Y.; Rao, Y. *Org. Lett.* **2012**, *14*, 2874.
- [57] Ackermann, L.; Vicente, R.; Potukuchi, H. K.; Pirovano, V. *Org. Lett.* **2010**, *12*, 5032.
- [58] Thirunavukkarasu, V. S.; Hubrich, J.; Ackermann, L. *Org. Lett.* **2012**, *14*, 4210.
- [59] Yang, F.; Ackermann, L. *Org. Lett.* **2013**, *15*, 718.
- [60] Thirunavukkarasu, V. S.; Ackermann, L. *Org. Lett.* **2012**, *14*, 6206.
- [61] Shan, G.; Han, X.; Lin, Y.; Yu, S.; Rao, Y. *Org. Biomol. Chem.* **2013**, *11*, 2318.
- [62] Kim, K.; Choe, H.; Jeong, Y.; Lee, J. H.; Hong, S. *Org. Lett.* **2015**, *17*, 2550.
- [63] Yang, F.; Rauch, K.; Kettelhoit, K.; Ackermann, L. *Angew. Chem., Int. Ed.* **2014**, *53*, 11285.
- [64] Shome, S.; Singh, S. P. *Tetrahedron Lett.* **2017**, *58*, 3743.
- [65] Ackermann, L.; Wang, L.; Wolfram, R.; Lygin, A. V. *Org. Lett.* **2012**, *14*, 728.
- [66] Yang, X.; Shan, G.; Rao, Y. *Org. Lett.* **2013**, *15*, 2334.
- [67] Liu, W.; Ackermann, L. *Org. Lett.* **2013**, *15*, 3484.
- [68] Turlington, C. R.; Morris, J.; White, P. S.; Brennessel, W. W.; Jones, W. D.; Brookhart, M.; Templeton, J. L. *Organometallics* **2014**, *33*, 4442.
- [69] Wu, Q.; Yan, D.; Chen, Y.; Wang, T.; Xiong, F.; Wei, W.; Lu, Y.; Sun, W.-Y.; Li, J. J.; Zhao, J. *Nat. Commun.* **2017**, *8*, 14227.
- [70] Cook, A. K.; Emmert, M. H.; Sanford, M. S. *Org. Lett.* **2013**, *15*, 5428.
- [71] For selected recent views on template-directed *meta*-selective C(sp²)-H functionalizations, see:
(a) Mihai, M. T.; Genov, G. R.; Phipps, R. J. *Chem. Soc. Rev.* **2017**, *47*, 149.
(b) Ping, L.; Chung, D. S.; Bouffard, J.; Lee, S.-G. *Chem. Soc. Rev.* **2017**, *46*, 4299.
(c) Mihai, M. T.; Phipps, R. J. *Synlett* **2017**, *28*, 1011.
(d) Dey, A.; Agasti, S.; Maiti, D. *Org. Biomol. Chem.* **2016**, *14*, 5440.
(e) Yang, G.; Butt, N.; Zhang, W. *Chin. J. Catal.* **2016**, *37*, 98.
(f) Yang, G.; Butt, N.; Zhang, W. *Chin. J. Catal.* **2016**, *37*, 98.
(g) Ackermann, L.; Li, J. *Nat. Chem.* **2015**, *7*, 686.
(h) Yang, J. *Org. Biomol. Chem.* **2015**, *13*, 1930.
- [72] (a) Bera, M.; Sahoo, S. K.; Maiti, D. *ACS Catal.* **2016**, *6*, 3575.
(b) Maji, A.; Bhaskararao, B.; Singha, S.; Sunoj, R. B.; Maiti, D. *Chem. Sci.* **2016**, *7*, 3147.
- [73] Bag, S.; Patra, T.; Modak, A.; Deb, A.; Maity, S.; Dutta, U.; Dey, A.; Kancherla, R.; Maji, A.; Hazra, A.; Bera, M.; Maiti, D. *J. Am. Chem. Soc.* **2015**, *137*, 11888.
- [74] A review on template-directed *para*-selective C(sp²)-H functionalizations: Dey, A.; Maity, S.; Maiti, D. *Chem. Commun.* **2016**, *52*, 12398.

(Li, L.; Fan, Y.)