新型查尔酮衍生物的合成及抗肿瘤活性评价

盛琦威 赵婉秋 曾明 谢中袍 夏雅平 崔冬梅*

(浙江工业大学药学院 杭州 310014)

摘要 主要通过 Claisen-Schmidt 缩合反应设计合成了一系列含有哌啶、吗啡啉、*N*-甲基哌嗪取代的查尔酮衍生物, 并通过噻唑蓝(MTT)法对 MCF-7(人乳腺癌细胞系)、A549(人肺癌细胞系)、HL-60(人白血病细胞系)、Hela(人宫颈癌细胞系)和 Bewo(人绒毛膜癌细胞系)等五种癌细胞系进行了体外抗癌活性研究. 研究表明, 化合物 **4a**, **4e**, **4f**, **4j**, **4m**, **4o** 对 MCF-7, A549, HL-60 三种癌细胞系抗癌活性最佳, IC₅₀ 值均在 10 μmol/L 以下.

关键词 查尔酮; 缩合; 抗癌; MTT 法

Synthesis and Evaluation of Chalcone Derivatives as Novel Anticancer Agents

Sheng, Qiwei Zhao, Wanqiu Zeng, Ming Xie, Zhongpao
Xia, Yaping Cui, Dongmei*

(College of Pharmaceutical Science, Zhejiang University of Technology, Hangzhou 310014)

Abstract Three series of chalcones bearing a piperidino, morpholino, and 1-methylpiperazino moiety were synthesized in two steps with the key step being Claisen-Schmidt condensation and tested for the activity against five cell lines, MCF-7 (human breast adenocarcinoma cell line), A549 (human lung adenocarcinoma epithelial cell line), HL-60 (human leukemia cell line), Hela (human cervical cancer cell line), and Bewo (human chorionic tumor cell line) by thiazolyl blue tetrazolium bromide (MTT) assay. Some chalcones exhibited good anticancer activity, and among them **4a**, **4e**, **4f**, **4j**, **4m**, and **4o** displayed the best anticancer activity for MCF-7 breast cancer cells, A549 lung cancer cells, and HL-60 leukemia cancer cells with IC₅₀ values below 10 μmol/L, respectively.

Keywords chalcone; condensation; anticancer agents; MTT assay

1 Introduction

Chalcones, analogs of 1,3-diaryl prop-2-ene-1-one containing two aromatic rings bound with vinyl ketone fragment form a wide class of compounds, represent an interesting group of natural compounds that are abundant in fruits, vegetables, spices, tea, and soy based foodstuff and possess multifarious pharmacological potentials, such as anti-fungal,^[1] anti-hypertensive,^[2] hypnotic,^[3] antispasmodic,^[4] immunosuppressant,^[5] anti-platelet,^[6] anti-diabetic,^[7] anti-tubercular,^[8] anti-neoplastic,^[9] antiangiogenic,^[10] anti-oxidant,^[11] hypolipidemic,^[12] anti-filarial,^[13] anti-retroviral,^[14] anti-malarial,^[15] antibacterial,^[16] anti-ulcer,^[17] anti-arrhythmic,^[18] anti-invasive,^[19] anti-histaminic,^[20] etc. The potential anticancer activities of chalcone derivatives has been explored. Recently, the anticancer activity of chalcones have been observed on inhib-

iting various molecular targets, like ABCG2/P-gp/BCRP, 5α-reductase, aromatase, 17-β-hydroxysteroid dehydrogenase, HDAC/Sirtuin-1, proteasome, VEGF, VEGFR-2 kinase, MMP-2/9, JAK/STAT signaling pathways, CDC-25B, tubulin, cathepsin-K, topoisomerase-II, Wnt, NF-κB, B-Raf and mTOR.^[9] Heterocyclic nitrogen-containing substrates are common constituents of natural products, agrochemicals, and pharmaceuticals. Recently, the synthesis of hybrid compounds of chalcone and heterocycles, such as chalcone substituted piperazine, and their anticancer activity have been reported, but few chalcone substituted other heterocycles, for example piperidine, morpholine, have been documented in the literature.^[21] Based on the above observations, we have synthesized chalcone derivatives containing heterocyclic nitrogen. The synthesized compounds act as potential anticancer agents.

* Corresponding author. E-mail: cuidongmei@zjut.edu.cn

Received August 29, 2018; revised October 23, 2018; published online November 25, 2018.

2 Results and discussion

2.1 Synthesis

The synthetic route of the proposed compounds **4a~4s** is shown in Scheme 1. First, the acetophenones derivatives **3a~3c** were obtained by the reaction of 4-fluorophenylethanone with piperidine, morpholine, and 1-methylpiperazine with stirring at reflux according to literature procedures. In the present investigation, substituted chalcones were prepared by the Claisen-Schmidt condensation of substituted 1-phenylethanone and substituted benzaldehyde with NaOH (aq.) as base in high yield by a known literature method. All chalcones were prepared from the corresponding reactants in 38%~87% yields. Comparative data for novel derivatives with respective to structures, melting point and yield are provided in Table 1. A range of benzaldehydes, with electronrich or electron-poor substituents on the aryl rings at different positions, were well-tolerated for the condensation reaction, giving the corresponding products. The heterocyclic aldehydes were also found to be suitable reaction partners for oxygen- and nitrogen-containing aromatics, and the corresponding products were obtained.

Scheme 1 Synthesis of chalcone derivatives **4a~4s**

Table 1 Physical data of compounds **4a~4s**

Compd.	X	Ar	Yield/%	m.p./°C
4a	CH ₂	3,4-(OCH ₂ O)C ₆ H ₃	76	170~173
4b	CH ₂	3-CH ₃ O-4-HOCH ₂ CH ₂ OC ₆ H ₃	65	167~170
4c	CH ₂	3-CH ₃ OC ₆ H ₄	44	160~163
4d	CH ₂	4-(CH ₃) ₂ NC ₆ H ₄	58	214~218
4e	CH ₂	2-Furyl	69	179~180
4f	CH ₂	2-Pyridyl	72	167~170
4g	O	3,4-(OCH ₂ O)C ₆ H ₃	76	167~171
4h	O	3-CH ₃ O-4-HOCH ₂ CH ₂ OC ₆ H ₃	41	165~168
4i	O	3-CH ₃ OC ₆ H ₄	87	130~133
4j	O	4-(CH ₃) ₂ NC ₆ H ₄	77	226~226
4k	O	4-BrC ₆ H ₄	79	213~216
4l	O	2,4-Cl ₂ C ₆ H ₃	40	147~148
4m	O	2-Furyl	38	195~197
4n	O	2-Pyridyl	52	150~153
4o	CH ₃ N	3,4-(OCH ₂ O)C ₆ H ₃	52	155~158
4p	CH ₃ N	3-CH ₃ O-4-HOCH ₂ CH ₂ OC ₆ H ₃	54	185~188
4q	CH ₃ N	3-CH ₃ OC ₆ H ₄	46	147~149
4r	CH ₃ N	4-(CH ₃) ₂ NC ₆ H ₄	57	198~201
4s	CH ₃ N	2-Furyl	39	179~182

Reaction conditions: **3** (1.0 mmol), aldehyde (1.0 mmol), 50% KOH (aq.) (1.0 mL), MeOH (3 mL), isolated yields.

2.2 Spectra

The structures of the obtained compounds **4a~4s** were confirmed by ¹H NMR, ¹³C NMR and HRMS spectral data. In the ¹H NMR spectra of the compounds, the CH protons of the olefin resonated as a doublet at δ 7.58~8.17 and 7.41~8.09, respectively. The large coupling constant of the olefinic protons confirms the *trans* configuration. In general, IR spectra of these compounds exhibited the presence of absorption bands for C=O between 1639~1649 cm⁻¹.

2.3 Biological studies

All the synthesized chalcone derivatives bearing a heterocyclic nitrogen moiety **4a~4s** were evaluated for their anticancer activity *in vitro* against five cancer cell lines, MCF-7 (human breast adenocarcinoma cell line), A549 (human lung adenocarcinoma epithelial cell line), HL-60 (human leukemia cell line), Hela (human cervical cancer cell line), and Bewo (Human chorionic tumor cell line), using the 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) assay method and compared with the well-known anticancer drug cisplatin. The anticancer activity results are shown in Table 2. The results showed that the structures of compounds had an obvious influence on anticancer activities. Among all compounds, there were three series of substituents of core benzene ring, including piperidino, morpholino, and 1-methylpiperazino. In general, derivatives containing piperidino were more potent than compounds with other substituted groups. For example, compound **4a** displayed the best anticancer activity against five cancer cell lines [for MCF-7 (IC₅₀ = 8.70 μ mol/L), A549 (IC₅₀ = 9.72 μ mol/L), HL-60 (IC₅₀ = 7.15 μ mol/L), Hela (IC₅₀ = 12.88 μ mol/L), and Bewo (IC₅₀ = 18.81 μ mol/L)]. Among all synthesized derivatives, compound **4j** showed the highest anticancer activity against MCF-7 with an IC₅₀ value of 4.63 μ mol/L, and better than cisplatin against MCF-7 cells. Compound **4j** also displayed the best anticancer activity for HL-60 cells (IC₅₀ = 6.00 μ mol/L), compounds **4e**, **4f**, **4m**, and **4o** also showed selected anticancer activity against HL-60 cell lines. According to the result, it is found that most chalcone derivatives were non-sensitive to Hela cells. Compound **4a** showed the most anticancer activity (IC₅₀ = 12.88 μ mol/L), and compounds **4c**, **4d**, **4h**, **4j**, and **4n** had weak anticancer activity for Hela (IC₅₀ < 50 μ mol/L). All compounds, except for **4a**, showed anticancer activity against Bewo cells at an IC₅₀ level of > 50 μ mol/L.

The compounds **4a~4e** containing piperidine group have significant potent anticancer activity (IC₅₀ = 8.70~10.78 μ mol/L) against MCF-7 as compared to the other compounds **4g~4i**, **4m** and **4o~4s** containing morpholine or methylpiperazine groups as shown in Table 2. These data indicated that the activity of compounds **4a~4e** was considerably attributed to the presence of piperidine group. The compound **4a** with substituted C-3 phenyl ring [3,4-(OCH₂O)] showed to have good activity (IC₅₀ = 8.70 μ mol/L). The change of 3,4-(OCH₂O) group to 3-CH₃O

4-HOCH₂CH₂O, 3-CH₃O, and 4-(CH₃)₂N did not show good activity. The activity of compound **4e** which C-3 phenyl ring was changed to furanyl ring was similar to that of compound **4a**. In the case of HL-60, investigation of structure-activity relationship (SAR) revealed that compounds **4d**, **4e**, **4f**, **4j**, **4m** and **4o** (IC₅₀ < 10.0 μmol/L) have a significant potent activity against HL-60 as compared to the standard drug cisplatin (IC₅₀ = 2.90 μmol/L) as shown in Table 2. This potency due to the presence of 3,4-(OCH₂O) and 4-(CH₃)₂N group at C-3 phenyl ring in compounds **4a**, **4j** and **4o** (IC₅₀ = 7.15, 6.00 and 8.15 μmol/L), or furyl and pyridyl ring at C-3 in compounds **4e** and **4f** (IC₅₀ = 8.77 and 9.62 μmol/L). In the case of A549, Hela, and Bewo, compound **4a** also retained the activity (IC₅₀ = 9.72, 12.88 and 18.81 μmol/L), and the other compounds showed lower or no anticancer activity in three cell lines. It indicated the importance of the type of piperidine group and 3,4-(OCH₂O)-substituted group.

Table 2 *In vitro* anticancer activity of the title compounds **4a**~**4s** against MCF-7, A549, HL-60, Hela, and Bewo cancer cell lines

Compd.	IC ₅₀ ^a /(μmol·L ⁻¹)				
	MCF-7	A549	HL-60	Hela	Bewo
4a	8.70	9.72	7.15	12.88	18.81
4b	14.99	>50	22.51	>50	>50
4c	21.50	>50	12.62	42.92	>50
4d	17.59	44.99	49.35	45.37	>50
4e	10.78	>50	8.77	>50	>50
4f	26.73	>50	9.62	>50	>50
4g	>50	>50	>50	>50	>50
4h	19.39	43.63	26.63	32.43	>50
4i	>50	>50	14.98	>50	>50
4j	4.63	35.62	6.00	39.24	>50
4k	>50	>50	>50	>50	>50
4l	>50	>50	>50	>50	>50
4m	15.69	>50	9.11	>50	>50
4n	16.58	44.19	11.37	25.29	>50
4o	39.46	>50	8.15	>50	>50
4p	>50	>50	>50	>50	>50
4q	23.52	45.49	>50	>50	>50
4r	>50	>50	24.86	>50	>50
4s	24.43	>50	13.67	>50	>50
Cisplatin	8.75	2.65	2.90	0.59	2.24

^a Anticancer activity was assayed by exposure for 48 h to the tested substances and expressed as the concentration required to inhibit tumour cell proliferation by 50% (IC₅₀).

3 Conclusions

In summary, a series of chalcone derivatives have been synthesized and the effects of all the compounds on anticancer activity of five cancer cell lines (MCF-7, A549, HL-60, Hela, and Bewo) were investigated. Compounds **4a**, **4e**, **4f**, **4j**, **4m**, and **4o** displayed the best anticancer activity for MCF-7 breast cancer cells, A549, lung cancer cells, and HL-60 leukemia cancer cells, respectively (IC₅₀ < 10 μmol/L). Moreover, compound **4a** had inhibitory effects

on the growth of five cancer cell lines.

4 Experimental

4.1 Experimental details

Under otherwise noted, materials were obtained from Aladdin Co., Ltd. and used without further purification. Thin layer chromatography (TLC) was performed using silica gel 60 F254 and visualized using UV light. Column chromatography was performed with silica gel (mesh 300 ~ 400). NMR spectra were obtained on a Bruker Avance 500 spectrometer (¹H NMR at 500 Hz, ¹³C NMR at 125 Hz) in CDCl₃ or DMSO-*d*₆ using TMS as an internal standard. High-resolution mass spectra (HRMS) were measured on a high resolution mass spectrometer (GCT Premier). Infrared spectra (IR) were obtained on a 370 FT-IR spectrometer.

4.2 4-Substituted acetophenone **3a**~**3c**

These were synthesized as described in previous work.^[22]

4.3 General procedure for synthesis of (*E*)-chalcones (**4**)

To a mixture of acetophenone (1.0 mmol) and aldehyde (1.0 mmol) in methanol (10 mL) was added 50% KOH (aq.) (1 mL). The resulting mixture was stirred for 10 h at reflux. The reaction mixture was cooled to room temperature. After evaporation of methanol and added water, the mixture was extracted with ethyl acetate. The organic phase was dried over anhydrous Na₂SO₄. The crude residue was obtained after evaporation of the solvent in vacuum, and the residue was purified by flash chromatography with petroleum and EtOAc as the eluent to give the pure product.

(*E*)-3-(1,3-Benzodioxol-5-yl)-1-(4-piperidinyl)phenyl-2-propen-1-one (**4a**): 1-(4-(Piperidin-1-yl)phenyl)ethan-1-one (204.1 mg, 1.0 mmol) and benzo[*d*][1,3]dioxole-5-carbaldehyde (149.9 mg, 1.0 mmol) in methanol (10 mL) was reacted according to the general procedure for compounds **4** to give the product **4a** (253.2 mg, 76% yield). *R*_f = 0.45 [*V*(petroleum ether) : *V*(ethyl acetate) = 3 : 1]. Yellow solid, m.p. 170 ~ 173 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.98 (d, *J* = 9.0 Hz, 2H), 7.72 (d, *J* = 15.5 Hz, 1H), 7.42 (d, *J* = 15.5 Hz, 1H), 7.18 (d, *J* = 1.3 Hz, 1H), 7.12 (dd, *J* = 8.0, 1.3 Hz, 1H), 6.91 (d, *J* = 9.0 Hz, 2H), 6.85 (d, *J* = 8.0 Hz, 1H), 6.03 (s, 2H), 3.39 (t, *J* = 4.9 Hz, 2H), 1.68 ~ 1.70 (m, 6H); ¹³C NMR (100 MHz, CDCl₃) δ: 187.4, 154.0, 149.2, 148.1, 142.4, 130.5, 129.7, 127.3, 124.6, 120.0, 113.3, 108.5, 106.5, 101.4, 48.6, 25.5, 24.5; IR (KBr) ν: 2935, 1651, 1600, 1499, 1360, 1196, 1124, 1034 cm⁻¹. HRMS (ESI) calcd for C₂₁H₂₂NO₃ [*M* + *H*]⁺: 226.1600, found 226.1603.

(*E*)-3-[(3-Methoxyphenyl-4-(2-hydroxyethoxy)]-1-(4-piperidinyl)phenyl-2-propen-1-one (**4b**): *R*_f = 0.2 [*V*(petroleum ether) : *V*(ethyl acetate) = 1 : 1]. Yellow solid, m.p. 167 ~ 170 °C; ¹H NMR (500 MHz, CDCl₃) δ: 7.99 (d, *J* = 8.9 Hz, 2H), 7.73 (d, *J* = 15.5 Hz, 1H), 7.45 (d, *J* = 15.5 Hz,

1H), 7.21 (dd, $J=8.3, 1.4$ Hz, 1H), 7.16 (d, $J=1.4$ Hz, 1H), 6.93 (d, $J=8.3$ Hz, 1H), 6.90 (d, $J=8.9$ Hz, 2H), 4.18 (t, $J=4.4$ Hz, 2H), 4.00~3.99 (m, 2H), 3.93 (s, 3H), 3.39 (t, $J=5.4$ Hz, 4H), 2.79 (s, 1H), 1.69~1.67 (m, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 187.9, 150.1, 149.9, 142.8, 130.8, 129.4, 122.5, 120.6, 114.2, 113.5, 110.9, 71.1, 61.3, 56.0, 48.7, 25.4, 24.4; IR (KBr) ν : 3537, 2939, 1643, 1609, 1511, 1268, 1207, 1033 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{28}\text{NO}_4$ $[\text{M}+\text{H}]^+$: 382.2018; found 382.2014.

(*E*)-3-(3-Methoxyphenyl)-1-(4-piperidinyl)phenyl-2-propen-1-one (**4c**): $R_f=0.25$ [$V(\text{petroleum ether}) : V(\text{ethyl acetate})=10 : 1$]. Yellow solid, m.p. 160~163 $^{\circ}\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 8.00 (d, $J=8.9$ Hz, 2H), 7.76 (d, $J=15.6$ Hz, 1H), 7.56 (d, $J=15.6$ Hz, 1H), 7.34 (t, $J=7.8$ Hz, 1H), 7.25 (d, $J=7.8$ Hz, 1H), 7.17 (d, $J=2.1$ Hz, 1H), 6.96~6.92 (m, 3H), 3.87 (s, 3H), 3.41 (t, $J=4.9$ Hz, 4H), 1.66~1.72 (m, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 187.9, 159.9, 142.8, 136.9, 130.9, 129.9, 122.6, 120.9, 115.8, 113.5, 113.4, 55.4, 48.7, 25.4, 24.4; IR (KBr) ν : 3420, 2945, 2849, 1649, 1592, 1384, 1251, 1127, 1040 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{24}\text{NO}_2$ $[\text{M}+\text{H}]^+$: 322.1807, found 322.1809.

(*E*)-3-[(4-Dimethylamino)phenyl]-1-(4-piperidinyl)phenyl-2-propen-1-one (**4d**): $R_f=0.23$ [$V(\text{petroleum ether}) : V(\text{ethyl acetate})=1 : 1$]. Yellow solid, m.p. 214~218 $^{\circ}\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 8.00 (d, $J=8.8$ Hz, 2H), 7.79 (d, $J=15.4$ Hz, 1H), 7.56 (d, $J=8.8$ Hz, 2H), 7.41 (d, $J=15.4$ Hz, 1H), 6.91 (d, $J=8.8$ Hz, 2H), 6.71 (d, $J=8.8$ Hz, 2H), 3.38 (t, $J=5.1$ Hz, 4H), 3.04 (s, 6H), 1.74~1.62 (m, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 188.2, 151.7, 143.8, 130.4, 130.0, 123.3, 117.1, 113.6, 111.9, 48.8, 40.2, 25.4, 24.3; IR (KBr) ν : 3408, 2933, 2854, 1732, 1641, 1599, 1228, 1029 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{27}\text{N}_2\text{O}$ $[\text{M}+\text{H}]^+$: 335.2123, found 335.2118.

(*E*)-3-(3-Furan-2-yl)-1-(4-piperidinyl)phenyl-2-propen-1-one (**4e**): $R_f=0.32$ [$V(\text{petroleum ether}) : V(\text{ethyl acetate})=5 : 1$]. Yellow solid, m.p. 179~180 $^{\circ}\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 8.01 (d, $J=8.9$ Hz, 2H), 7.58 (d, $J=15.5$ Hz, 1H), 7.52~7.48 (m, 2H), 6.98~6.84 (m, 2H), 6.68 (d, $J=3.4$ Hz, 1H), 6.52~6.50 (m, 1H), 3.41 (t, $J=4.1$ Hz, 4H), 1.69~1.62 (m, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 187.3, 152.1, 144.4, 130.7, 129.1, 119.6, 115.1, 113.5, 112.5, 48.7, 25.4, 24.3; IR (KBr) ν : 3429, 2940, 2922, 2854, 1647, 1603, 1387, 1234, 1194, 1123, 1020 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{20}\text{NO}_2$ $[\text{M}+\text{H}]^+$: 282.1494; found 282.1490.

(*E*)-3-(Pyridin-2-yl)-1-(4-piperidinyl)phenyl-2-propen-1-one (**4f**): $R_f=0.25$ [$V(\text{petroleum ether}) : V(\text{ethyl acetate})=3 : 1$]. Yellow solid, m.p. 167~170 $^{\circ}\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 8.70 (d, $J=4.0$ Hz, 1H), 8.17 (d, $J=15.2$ Hz, 1H), 8.07 (d, $J=9.0$ Hz, 2H), 7.76 (d, $J=15.2$ Hz, 1H), 7.76~7.72 (m, 1H), 7.47 (d, $J=7.8$ Hz, 1H), 7.29~7.27 (m, 1H), 6.90 (d, $J=9.0$ Hz, 2H), 3.41 (t, $J=4.8$ Hz, 4H), 1.70~1.68 (m, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ : 187.6, 154.4, 153.7, 150.0, 140.8, 136.8, 131.1, 126.9, 125.8, 125.1, 123.9, 113.2, 48.4, 25.3, 24.3; IR (KBr) ν : 3421, 2934, 1647, 1588, 1392, 1192, 1126, 994 cm^{-1} . HRMS

(ESI) calcd for $\text{C}_{19}\text{H}_{21}\text{N}_2\text{O}$ $[\text{M}+\text{H}]^+$: 293.1654; found 293.1655.

(*E*)-3-(1,3-Benzodioxol-5-yl)-1-[4-(4-morpholinyl)phenyl]-2-propen-1-one (**4g**) (CAS: 305859-40-9): $R_f=0.19$ [$V(\text{petroleum ether}) : V(\text{ethyl acetate})=3 : 1$]. Yellow solid, m.p. 167~171 $^{\circ}\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 8.01 (d, $J=9.0$ Hz, 2H), 7.74 (d, $J=15.5$ Hz, 1H), 7.42 (d, $J=15.5$ Hz, 1H), 7.19 (d, $J=1.5$ Hz, 1H), 7.13 (dd, $J=8.0, 1.5$ Hz, 1H), 6.93 (d, $J=9.0$ Hz, 2H), 6.86 (d, $J=8.0$ Hz, 1H), 6.04 (s, 2H), 3.89 (t, $J=4.9$ Hz, 4H), 3.35 (t, $J=4.9$ Hz, 4H).

(*E*)-3-[(3-Methoxyphenyl-4-(2-hydroxyethoxy))-1-[4-(4-morpholinyl)phenyl]-2-propen-1-one (**4h**): $R_f=0.13$ [$V(\text{petroleum ether}) : V(\text{ethyl acetate})=1 : 1$]. Yellow solid, m.p. 165~168 $^{\circ}\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 8.02 (d, $J=8.8$ Hz, 2H), 7.75 (d, $J=15.6$ Hz, 1H), 7.44 (d, $J=15.6$ Hz, 1H), 7.23 (dd, $J=8.2, 1.6$ Hz, 1H), 7.17 (d, $J=1.6$ Hz, 1H), 6.96~6.92 (m, 3H), 4.19 (t, $J=4.4$ Hz, 2H), 4.01~3.99 (m, 2H), 3.95 (s, 3H), 3.88 (t, $J=4.8$ Hz, 4H), 3.35 (t, $J=4.8$ Hz, 4H), 2.54 (t, $J=6.1$ Hz, 1H); ^{13}C NMR (125 MHz, CDCl_3) δ : 188.2, 154.0, 150.2, 149.9, 143.3, 130.6, 129.2, 122.5, 120.4, 114.1, 113.6, 110.9, 71.1, 66.6, 61.2, 56.0, 47.7; IR (KBr) ν : 3539, 2966, 2858, 1641, 1607, 1508, 1266, 1032 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{26}\text{NO}_5$ $[\text{M}+\text{H}]^+$: 384.1811; found 384.1808.

(*E*)-3-(3-Methoxyphenyl)-1-[4-(4-morpholinyl)phenyl]-2-propen-1-one (**4i**) (CAS: 850351-83-6): $R_f=0.2$ [$V(\text{petroleum ether}) : V(\text{ethyl acetate})=3 : 1$]. Yellow solid, m.p. 130~133 $^{\circ}\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 8.03 (d, $J=8.9$ Hz, 2H), 7.77 (d, $J=15.6$ Hz, 1H), 7.55 (d, $J=15.6$ Hz, 1H), 7.34 (t, $J=7.8$ Hz, 1H), 7.26 (d, $J=7.8$ Hz, 1H), 7.17 (s, 1H), 6.97~6.94 (m, 1H), 6.93 (d, $J=8.9$ Hz, 2H), 3.89~3.88 (m, 4H), 3.87 (s, 3H), 3.35 (t, $J=5.0$ Hz, 4H).

(*E*)-3-[(4-Dimethylamino)phenyl]-1-[4-(4-morpholinyl)phenyl]-2-propen-1-one (**4j**) (CAS: 1004038-07-6): $R_f=0.35$ [$V(\text{petroleum ether}) : V(\text{ethyl acetate})=2 : 1$]; yellow solid, m.p. 226~228 $^{\circ}\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 8.02 (d, $J=8.9$ Hz, 2H), 7.80 (d, $J=15.4$ Hz, 1H), 7.56 (d, $J=8.8$ Hz, 2H), 7.39 (d, $J=15.4$ Hz, 1H), 6.93 (d, $J=8.9$ Hz, 2H), 6.71 (d, $J=8.8$ Hz, 2H), 3.88 (t, $J=4.9$ Hz, 4H), 3.35 (t, $J=4.9$ Hz, 4H), 3.05 (s, 6H).

(*E*)-3-(4-Bromophenyl)-1-[4-(4-morpholinyl)phenyl]-2-propen-1-one (**4k**)^[23]: $R_f=0.26$ [$V(\text{petroleum ether}) : V(\text{ethyl acetate})=3 : 1$]. Yellow solid, m.p. 213~216 $^{\circ}\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 8.02 (d, $J=9.0$ Hz, 2H), 7.73 (d, $J=15.6$ Hz, 1H), 7.57~7.50 (m, 5H), 6.93 (d, $J=9.0$ Hz, 2H), 3.88 (t, $J=4.9$ Hz, 4H), 3.55 (t, $J=4.9$ Hz, 4H).

(*E*)-3-(2,4-Dichlorophenyl)-1-[4-(4-morpholinyl)phenyl]-2-propen-1-one (**4l**) (CAS: 796989-79-2): $R_f=0.33$ [$V(\text{petroleum ether}) : V(\text{ethyl acetate})=3 : 1$]. Yellow solid, m.p. 147~148 $^{\circ}\text{C}$; ^1H NMR (500 MHz, CDCl_3) δ : 8.09 (d, $J=15.7$ Hz, 1H), 8.01 (d, $J=8.9$ Hz, 2H), 7.69 (d, $J=8.5$ Hz, 1H), 7.51 (d, $J=15.7$ Hz, 1H), 7.48 (d, $J=2.0$ Hz, 1H), 7.31 (dd, $J=8.5, 2.0$ Hz, 1H), 6.93 (d, $J=8.9$ Hz, 2H), 3.89 (t, $J=5.0$ Hz, 4H), 3.36 (t, $J=5.0$ Hz, 4H).

(*E*)-3-(3-Furan-2-yl)-1-[4-(4-morpholinyl)phenyl]-2-propen-1-one (**4m**): $R_f=0.3$ [*V*(petroleum ether) : *V*(ethyl acetate) = 3 : 1]. Yellow solid, m.p. 195 ~ 197 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.03 (d, $J=9.0$ Hz, 2H), 7.59 (d, $J=15.3$ Hz, 1H), 7.53 (d, $J=1.7$ Hz, 1H), 7.50 (d, $J=15.3$ Hz, 1H), 6.94 (d, $J=9.0$ Hz, 2H), 6.70 (d, $J=3.4$ Hz, 1H), 6.52 (dd, $J=3.4, 1.7$ Hz, 1H), 3.89 (t, $J=4.9$ Hz, 4H), 3.35 (t, $J=4.9$ Hz, 4H); ^{13}C NMR (125 MHz, CDCl_3) δ : 187.6, 154.1, 152.0, 144.5, 130.6, 129.5, 128.9, 119.4, 115.4, 113.5, 112.5, 66.6, 47.6; IR (KBr) ν : 3457, 2974, 2850, 1647, 1604, 1384, 1231, 1193, 1130, 1025 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{18}\text{NO}_3$ [$\text{M}+\text{H}$] $^+$: 284.1287; found 284.1287.

(*E*)-3-(Pyridin-2-yl)-1-[4-(4-morpholinyl)phenyl]-2-propen-1-one (**4n**): $R_f=0.41$ [*V*(petroleum ether) : *V*(ethyl acetate) = 3 : 1]. Yellow solid, m.p. 150 ~ 153 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.69 (d, $J=3.9$ Hz, 1H), 8.15 (d, $J=15.2$ Hz, 1H), 8.09 (d, $J=9.0$ Hz, 2H), 7.76 (d, $J=15.2$ Hz, 1H), 7.75 ~ 7.72 (m, 1H), 7.47 (d, $J=7.7$ Hz, 1H), 7.30 ~ 7.28 (m, 1H), 6.91 (d, $J=9.0$ Hz, 2H), 3.87 (t, $J=4.9$ Hz, 4H), 3.35 (t, $J=4.9$ Hz, 4H); ^{13}C NMR (125 MHz, CDCl_3) δ : 188.0, 154.3, 153.6, 150.1, 141.3, 136.9, 130.9, 128.6, 125.7, 125.3, 124.1, 113.3, 66.6, 47.4; IR (KBr) ν : 3463, 2960, 1656, 1591, 1382, 1193, 1117, 1049 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{19}\text{N}_2\text{O}_2$ [$\text{M}+\text{H}$] $^+$: 295.1447; found 295.1448.

(*E*)-3-(1,3-Benzodioxol-5-yl)-1-[4-(4-methylpiperazin-1-yl)phenyl]-2-propen-1-one (**4o**): $R_f=0.45$ (ethyl acetate). Yellow solid, m.p. 155 ~ 158 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.00 (d, $J=8.9$ Hz, 2H), 7.73 (d, $J=15.5$ Hz, 1H), 7.41 (d, $J=15.5$ Hz, 1H), 7.18 (d, $J=1.6$ Hz, 1H), 7.13 (dd, $J=8.0, 1.6$ Hz, 1H), 6.93 (d, $J=8.9$ Hz, 2H), 6.85 (d, $J=8.0$ Hz, 1H), 6.03 (s, 2H), 3.42 (t, $J=4.9$ Hz, 4H), 2.60 (t, $J=4.9$ Hz, 4H), 2.39 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 187.6, 153.7, 149.3, 148.1, 142.8, 130.4, 129.7, 128.3, 124.7, 119.9, 113.5, 108.5, 106.5, 101.4, 54.7, 47.3, 46.1; IR (KBr) ν : 3418, 2904, 1647, 1609, 1384, 1250, 1198, 1037, 810 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{23}\text{N}_2\text{O}_3$ [$\text{M}+\text{H}$] $^+$: 351.1709; found 351.1711.

(*E*)-3-[(3-Methoxyphenyl)-4-(2-hydroxyethoxy)]-1-[4-(4-methylpiperazin-1-yl)phenyl]-2-propen-1-one (**4p**): $R_f=0.18$ (ethyl acetate). Yellow solid, m.p. 185 ~ 188 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.01 (d, $J=8.9$ Hz, 2H), 7.75 (d, $J=15.6$ Hz, 1H), 7.44 (d, $J=15.6$ Hz, 1H), 7.23 (dd, $J=8.3, 1.7$ Hz, 1H), 7.17 (d, $J=1.7$ Hz, 1H), 6.95 ~ 6.92 (m, 3H), 4.19 (t, $J=4.5$ Hz, 2H), 4.00 (t, $J=4.5$ Hz, 2H), 3.94 (s, 3H), 3.42 (t, $J=4.9$ Hz, 4H), 2.59 (t, $J=4.9$ Hz, 4H), 2.38 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 188.2, 154.1, 150.2, 149.9, 143.2, 130.7, 129.3, 128.6, 122.5, 120.5, 114.1, 113.7, 110.9, 71.1, 61.3, 56.0, 54.8, 47.3, 46.1; IR (KBr) ν : 3414, 2939, 2844, 1644, 1599, 1511, 1269, 1141, 1032 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{29}\text{N}_2\text{O}_4$ [$\text{M}+\text{H}$] $^+$: 397.2127; found 397.2123.

(*E*)-3-(3-Methoxyphenyl)-1-[4-(4-methylpiperazin-1-yl)phenyl]-2-propen-1-one (**4q**): $R_f=0.21$ (ethyl acetate). Yellow solid, m.p. 147 ~ 149 °C; ^1H NMR (400 MHz,

CDCl_3) δ : 7.97 (d, $J=8.9$ Hz, 2H), 7.73 (d, $J=15.6$ Hz, 1H), 7.52 (d, $J=15.6$ Hz, 1H), 7.31 (t, $J=7.7$ Hz, 1H), 7.22 (d, $J=7.7$ Hz, 1H), 7.13 (t, $J=1.9$ Hz, 1H), 6.93 ~ 6.92 (m, 1H), 6.90 (d, $J=8.9$ Hz, 2H), 3.85 (s, 3H), 3.39 (t, $J=4.9$ Hz, 4H), 2.56 (t, $J=4.9$ Hz, 4H), 2.36 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 187.7, 159.7, 153.8, 142.8, 136.5, 130.5, 129.7, 128.1, 122.2, 120.8, 115.6, 113.4, 113.2, 55.4, 54.7, 47.3, 46.2; IR (KBr) ν : 3473, 2936, 1660, 1603, 1382, 1250, 1203, 1053 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{25}\text{N}_2\text{O}_2$ [$\text{M}+\text{H}$] $^+$: 337.1916; found 337.1919.

(*E*)-3-[(4-Dimethylamino)phenyl]-1-[4-(4-methylpiperazin-1-yl)phenyl]-2-propen-1-one (**4r**): $R_f=0.19$ (ethyl acetate). Yellow solid, m.p. 198 ~ 201 °C; ^1H NMR (500 MHz, CDCl_3) δ : 8.01 (d, $J=8.9$ Hz, 2H), 7.79 (d, $J=15.4$ Hz, 1H), 7.56 (d, $J=8.8$ Hz, 2H), 7.39 (d, $J=15.4$ Hz, 1H), 6.93 (d, $J=8.9$ Hz, 2H), 6.71 (d, $J=8.8$ Hz, 2H), 3.40 (t, $J=5.0$ Hz, 4H), 3.04 (s, 6H), 2.58 (t, $J=5.0$ Hz, 4H), 2.37 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 188.0, 153.5, 151.5, 143.9, 130.2, 129.9, 129.0, 122.9, 116.8, 113.5, 111.7, 54.8, 47.4, 46.2, 40.2; IR (KBr) ν : 3455, 2936, 2795, 1636, 1598, 1518, 1231, 1180, 1033 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{28}\text{N}_3\text{O}$ [$\text{M}+\text{H}$] $^+$: 350.2232; found 350.2231.

(*E*)-3-(3-Furan-2-yl)-1-[4-(4-methylpiperazin-1-yl)phenyl]-2-propen-1-one (**4s**): $R_f=0.15$ (ethyl acetate). Yellow solid, m.p. 179 ~ 182 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.98 (d, $J=8.9$ Hz, 2H), 7.55 (d, $J=15.3$ Hz, 1H), 7.49 ~ 7.44 (m, 2H), 6.89 (d, $J=8.9$ Hz, 2H), 6.65 (d, $J=3.3$ Hz, 1H), 6.48 (dd, $J=3.3, 1.7$ Hz, 1H), 3.39 (t, $J=4.9$ Hz, 4H), 2.56 (t, $J=4.9$ Hz, 4H), 2.36 (s, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ : 187.4, 154.0, 152.1, 144.4, 130.6, 129.3, 128.3, 119.5, 115.2, 113.5, 112.5, 54.7, 47.2, 46.1; IR (KBr) ν : 3416, 2845, 1647, 1602, 1383, 1233, 1187, 1019, 923 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{21}\text{N}_2\text{O}_2$ [$\text{M}+\text{H}$] $^+$: 297.1603; found 297.1606.

4.4 Cytotoxic activity

For observing the anticancer activity of the target compounds *in vitro*, MTT assay was used. MCF-7, A549, HL-60, Hela, and Bewo cells were harvested in the logarithmic growth phase and seeded in 96-well plates, and cultured at 37 °C in humidified atmosphere containing 5% CO_2 in Dulbecco's Modified Eagle Medium (DMEM) with 10% FBS for 24 h before any treatments. Tested compounds were dissolved in dimethyl sulfoxide (DMSO) and diluted in the culture fluid to get various concentrations. The cells were treated with target compounds subsequently and incubated 24 h. Then 20 μL of MTT (5 mg/mL) was added in a 37 °C, 5% CO_2 incubator for 4 h, the medium was removed immediately and MTT formazan was solubilized in 150 μL of DMSO. Cell survival was assayed by enzyme-labeled instrument. Inhibitory effects were determined as IC_{50} value.^[23]

Supporting Information The electronic supplementary information (ESI) available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] Lahtchev, K. L.; Batovska, D. I.; Parushev, S. P.; Ubiyovkov, V. M.; Sibimy, A. A. *Eur. J. Med. Chem.* **2008**, *43*, 2220.
- [2] (a) Kantevari, S.; Addla, D.; Bagul, P. K.; Sridhar, B.; Banerjee, S. K. *Bioorg. Med. Chem.* **2011**, *19*, 4772.
(b) Yan, Y. K.; Xu, Q.; Gao, Y.; Liu, H.; Tang, X. R. *Chin. J. Org. Chem.* **2018**, *38*, 1763 (in Chinese).
(严映坤, 徐侨, 高扬, 刘辉, 唐孝荣, 有机化学, **2018**, *38*, 1763.)
- [3] (a) Cho, S.; Kim, S.; Jin, Z.; Yang, H.; Han, D.; Baek, N. I.; Jo, J.; Cho, C. W.; Park, J. H.; Shimizu, M.; Jin, Y. H. *Res. Commun.* **2011**, *413*, 637.
(b) Zhang, E.; Wang, M. M.; Xu, S. M.; Wang, S.; Zhao, D.; Bai, P. Y.; Cui, D. Y.; Hua, Y. G.; Wang, Y. N.; Qin, S. S.; Liu, H. M. *Chin. J. Org. Chem.* **2017**, *37*, 959 (in Chinese).
(张恩, 王铭铭, 徐帅民, 王上, 赵娣, 白鹏燕, 崔得运, 化永刚, 王亚娜, 秦上尚, 刘宏民, 有机化学, **2017**, *37*, 959.)
- [4] Sato, Y.; He, J.; Nagai, H.; Tani, T.; Akao, T. *Biol. Pharm. Bull.* **2007**, *30*, 145.
- [5] Luo, Y.; Song, R.; Li, Y.; Zhang, S.; Liu, Z. J.; Fu, J.; Zhu, H. L. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 3039.
- [6] Zhao, L.; Jin, H.; Sun, L.; Piao, H.; Quan, Z. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 5027.
- [7] Mahapatra, D. K.; Asati, V.; Bharti, S. K. *Eur. J. Med. Chem.* **2015**, *92*, 839.
- [8] Mascarello, A.; Chiaradia, L. D.; Vernal, J.; Villarino, A.; Guido, R. V.; Perizzolo, P.; Poirier, V.; Wong, D.; Martins, P. G.; Nunes, R. J.; Yunes, R. A.; Andricopulo, A. D.; Av-Gay, Y.; Terenzi, H. *Bioorg. Med. Chem.* **2010**, *18*, 3783.
- [9] Mahapatra, D. K.; Bharti, S. K.; Asati, V. *Eur. J. Med. Chem.* **2015**, *98*, 69.
- [10] Lee, Y. S.; Lim, S. S.; Shin, K. H.; Kim, Y. S.; Ohuchi, K.; Jung, S. H. *Biol. Pharm. Bull.* **2006**, *29*, 1028.
- [11] Aoki, N.; Muko, M.; Ohta, E.; Ohta, S. *J. Nat. Prod.* **2008**, *71*, 1308.
- [12] Sashidhara, K. V.; Palnati, G. R.; Sonkar, R.; Avula, S. R.; Awasthi, C.; Bhatia, G. *Eur. J. Med. Chem.* **2013**, *64*, 422.
- [13] Sashidhara, K. V.; Rao, K. B.; Kushwaha, V.; Modukuri, R. K.; Verma, R.; Murthy, P. K. *Eur. J. Med. Chem.* **2014**, *81*, 473.
- [14] Rizvi, S. U. F.; Siddiqui, H. L.; Johns, M.; Dettori, M.; Schinazi, R. F. *Med. Chem. Res.* **2012**, *21*, 3741.
- [15] Tomar, V.; Bhattacharjee, G.; Kamaluddin; Rajakumar, S.; Srivastava, K.; Puri, S. K. *Eur. J. Med. Chem.* **2010**, *45*, 745.
- [16] Abdullah, M. I.; Mahmood, A.; Madni, M.; Masood, S.; Kashif, M. *Bioorg. Chem.* **2014**, *54*, 31.
- [17] Sashidhara, K. V.; Avula, S. R.; Mishra, V.; Palnati, G. R.; Singh, L. R.; Singh, N.; Chhonker, Y. S.; Swamy, P.; Bhatta, R. S.; Palit, G. *Eur. J. Med. Chem.* **2015**, *89*, 638.
- [18] Yarishkin, O. V.; Ryu, H. W.; Park, J.; Yang, M. S.; Hong, S.; Park, K. H. *Bioorg. Med. Chem. Lett.* **2008**, *18*, 137.
- [19] Wang, L.; Chen, G.; Lu, X.; Wang, S.; Hans, S.; Li, Y.; Ping, G.; Jiang, X.; Li, H.; Yang, J.; Wu, C. *Eur. J. Med. Chem.* **2015**, *89*, 88.
- [20] Yamamoto, T.; Yoshimura, M.; Yamaguchi, F.; Kouchi, T.; Tsuji, R.; Saito, M.; Obata, A.; Kikuchi, M. *Biosci. Biotechnol. Biochem.* **2004**, *68*, 1706.
- [21] Gao, H.; Zheng, X.; Qi, Y.; Wang, S.; Wan, C. P.; Rao, G. X.; Mao, Z. W. *Chin. J. Org. Chem.* **2018**, *38*, 648 (in Chinese).
(高慧, 郑喜, 祁燕, 王斯, 万春平, 饶高雄, 毛泽伟, 有机化学, **2018**, *38*, 648.)
- [22] Thanusu, J.; Kanagarajan, V.; Gopalakrishnan, M. *J. Enzyme Inhib. Med. Chem.* **2010**, *25*, 347.
- [23] Ramya, V.; Vembu, S.; Ariharasivakumar, G.; Gopalakrishnan, M. *Pharma Chem.* **2017**, *9*, 46.
- [24] Arasakumar, T.; Mathusalini, S.; Ata, A.; Shankar, R.; Gopalan, S.; Lakshmi, K.; Sakthivel, P.; Mohan, P. S. *Mol. Diversity* **2017**, *21*, 37.
- [25] Zhang, X.; Shi, G.; He, L.; Tao, K.; Hou, T. *Pesticides* **2012**, *51*, 301 (in Chinese).
(张新刚, 史冠莹, 何利钦, 陶科, 侯太平, 农药, **2012**, *51*, 301.)

(Lu, Y.)