苯硫酚还原 α,α,α -三溴甲基酮类化合物的研究

杨莹 哈斯木江·巴拉提 阿布都热西提·阿布力克木*

(新疆师范大学化学系 乌鲁木齐 830054)

摘要 以苯硫酚为还原剂, 在绿色溶剂、室温下将 α,α,α -三溴甲基酮类化合物还原成 α,α -二溴甲基酮类化合物。此反应适用于芳香族、脂肪族和杂环类 α,α,α -三溴甲基酮类化合物, 产率高达 97%。进一步的机理研究表明, 该反应是自由基反应。

关键词 苯硫酚; α,α -二溴甲基酮; 绿色合成; 还原反应

Research on Reduction of α,α,α -Tribromomethyl Ketones via Thiophenol

Yang, Ying Balati, Hasimujiang Abulikemu, Abudu Rexit*

(Department of Chemistry, Xinjiang Normal University, Urumqi 830054)

Abstract α,α -Dibromomethyl ketones were synthesized with high yields through a thiophenol-promoted reduction of α,α,α -tribromomethyl ketones under mild conditions within one hour. A further mechanistic study showed that the reaction proceeded via a radical process. This is an example that uses thiophenol as the radical stimulator in the reduction of multi-halogenated compounds.

Keywords thiophenol; α,α -dibromomethyl ketones; green synthesis; reduction

1 Introduction

Halogenated organic compounds are widely used as starting materials and intermediates in organic and organometallic synthesis. This includes irreplaceable applications of α,α -dibromoacetophenones in the construction of heterocyclic structures^[1] or other small rings,^[2] the preparation of compounds with a double^[3] or triple^[4] bonds, the synthesis of unsaturated acids and ynols,^[5] and the preparation of cyclic compounds.^[6]

The synthesis of 2,2-dibromo-1-arylethanones is traditionally based on the bromination of acetophenones with molecular bromine,^[7] with *N*-halosuccinimides (NXS),^[8] hexabromocyclopentadiene,^[9] and the H_2O_2 -HBr system,^[10] or through an oxyhalogenation of alkynes using oxone-KX,^[11] Selectfluor/KBr,^[12] 48% HBr (aq.),^[13] and 1,3-dibromo-5,5-dimethylhydantoin (DBH)^[14] in an oxy-1,1-dihalogenation of alkynes with NaX mediated by $\text{K}_2\text{S}_2\text{O}_8$ in an aqueous system.^[15] The widely-known synthetic methods of α,α -dibromoacetophenones do not meet the requirements for green chemistry because of several

elements. They use low-atom-efficiency reagents or non-environmentally friendly organic solvents. In addition, the critical reaction conditions require high temperatures or a complex operation. Finally, they only provide low yields.

Recent research^[16] showed that radical reactions can be easily realized under visible light with a catalytic amount of thiophenols or mercaptans. On the basis of the earlier work of the project group,^[17] our results showed that a similar situation occurred for the reaction of α,α,α -tribromomethyl ketones in the presence of thiophenols or mercaptans. Furthermore, among various possible radical stimulators (AIBN, RSH, NBS), thiophenol was the optimal choice.^[18] The preparation of dibromomethyl ketones from tribromomethyl ketones to produce sulfur radicals was simpler and more environment-friendly in the presence of thiophenol in a sodium bromide solution (0.1 mmol/mL).

Another difficulty arose from the intense and increasing interest in developing "greener" chemistry in the laboratory and in the industry. Green synthesis is difficult to achieve for halogenated organic compounds because of the unavoidable use of organic solvents because of their insolubil-

* Corresponding author. E-mail: aarexit@xjnu.edu.cn

Received October 5, 2018; revised November 7, 2018; published online November 25, 2018.

Project supported by the National Natural Science Foundation of China (No. 21662035).

国家自然科学基金(No. 21662035)资助项目.

ity in aqueous solutions. To meet this requirement, aqueous alcohol solvent mixtures were tested (Scheme 1).

Scheme 1 Strategies for synthesis of α,α -dibromoacetophenones

2 Results and discussion

Using α,α,α -tribromomethyl ketone **1a** and thiophenol compound **2a** as the starting materials in a 1 : 2 molar ratio, various solvents were tested at room temperature (Table 1). Studying the effects of different solvents such as ethanol, CH_2Cl_2 , CH_3CN , PhCH_3 and $\text{H}_2\text{O}:\text{EtOH}$ (Table 1, Entries 2~6), solvent screening revealed that ethanol was the best reaction media since it produced **3a** with 87% yield (Entry 2).

Environment-friendly aqueous solutions are preferred over organic solvents because they are safe, benign, non-toxic, inexpensive, and easily separable.

Another optimization of the $\text{H}_2\text{O}/\text{EtOH}$ system was investigated (Table 1, Entries 6~18). The concentrations of inorganic salts were also tested in $\text{H}_2\text{O}:\text{EtOH}$ systems (Entries 7~18). Among NaCl , KBr , Na_2CO_3 , K_2CO_3 , NH_4Cl

and NaBr (Entries 7~12), the saturated solution of sodium bromide (Entry 12) corresponded to the best choice by producing α,α -tribromomethyl ketone **3a** with 83% yield. A detailed study on the concentration of sodium bromide showed that a 0.1 mol/L solution resulted in the highest yield for producing α,α -tribromomethyl ketone **3a** with 96% (Entry 17).

Under the optimized reaction conditions established in Table 1, various substrates of tribromomethyl ketones were screened (Table 2). The results showed that this methodology could be used for a variety of aromatic, aliphatic, and heterocyclic tribromomethyl ketones. Meanwhile, changing the position of the substituent in the substrate had no influence on the reaction time and the yield. As shown in Entries 2~4, the corresponding α,α -dibromo acetophenones **3b**, **3c**, and **3d** were obtained with 96%, 97% and 95%, respectively. When an electron-withdrawing group like bromine was introduced in the benzene ring in **1e** and **1f**, 93% and 95% yields were obtained (Entries 5 and 6). Additionally, an electron-donating group like methyl also resulted in good yields of 96%, 90% and 91% (Entries 7~9). The introduction of strong electron-absorption groups had a negative effect and resulted in lower yields (Entry 11). Aliphatic tribromomethyl ketones also performed well in the same conditions, producing the relevant products with 83%~94% yields (Entries 12~14). For the heterocyclic tribromomethyl ketone **3o**, the yield could reach up to 92% (Entry 15). Good isolation yields were also obtained for the synthesis of **3p** and **3q** (Entries 16 and 17).

Table 1 Optimization of the reaction conditions^a

Entry	2a	Solvent	Time/h	Yield ^b / % of 3a
1	PhCH_2SH	EtOH	10	73
2	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	EtOH	1/4	87
3	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	CH_2Cl_2	1/4	88
4	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	CH_3CN	1	82
5	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	PhCH_3	1	82
6	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	$\text{H}_2\text{O}/\text{EtOH}$ (V : V=9 : 1)	1/4	64
7	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	NaCl (saturated)/EtOH (V : V=9 : 1)	1/4	73
8	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	KBr (saturated)/EtOH (V : V=9 : 1)	1/4	68
9	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	Na_2CO_3 (saturated)/EtOH (V : V=9 : 1)	1/4	77
10	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	K_2CO_3 (saturated)/EtOH (V : V=9 : 1)	1/4	79
11	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	NH_4Cl (saturated)/EtOH (V : V=9 : 1)	1/4	82
12	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	NaBr (saturated)/EtOH (V : V=9 : 1)	1/4	83
13	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	NaBr (1.0 mmol/mL)/EtOH (V : V=9 : 1)	1/4	86
14	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	NaBr (0.5 mmol/mL)/EtOH (V : V=9 : 1)	1/4	90
15	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	NaBr (0.2 mmol/mL)/EtOH (V : V=9 : 1)	1/4	93
16	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	NaBr (0.1 mmol/mL)/EtOH (V : V=9 : 1)	1/4	95
17	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	NaBr (0.1 mmol/mL)/EtOH (V : V=9 : 1)	1/4	96 ^c
18	$p\text{-CH}_3\text{C}_6\text{H}_4\text{SH}$	NaBr (0.1 mmol/mL)/EtOH (V : V=9 : 1)	1/2	65 ^d

^a The reactions of **1a** (0.10 mmol) and **2a** (0.2 mmol) were carried out in 2 mL of solvent at room temperature. ^b Isolated yield. ^c **2a** (2.05 equiv.). ^d **2a** (1.5 equiv.).

Table 2 Synthesis of α,α -dibromoacetophenones from tribromomethyl ketones^a

Entry	Substrate	Product	Time/min	Yield ^b /%
1	1a	3a	15	96
2	1b	3b	60	96 ^c
3	1c	3c	25	97
4	1d	3d	15	95
5	1e	3e	60	93 ^c
6	1f	3f	40	95
7	1g	3g	35	96
8	1h	3h	40	90
9	1i	3i	15	91

Continued

Entry	Substrate	Product	Time/min	Yield ^b /%
10	1j	3j	35	93
11	1k	3k	60	75
12	1l	3l	40	94
13	1m	3m	60	94
14	1n	3n	30	83
15	1o	3o	30	92
16	1p	3p	45	85
17	1q	3q	30	85

^a The reactions of **1a** (0.10 mmol) and **2a** (2.05 mmol) were carried out in 2 mL of NaBr (0.1 mmol/mL)/EtOH ($V : V = 9 : 1$) at the room temperature. ^b Isolated yield. ^c At 0 °C.

2.1 Mechanism study

Controlled experiments were designed to study the reaction mechanism. As an initial step, 3 equiv. 2,2,6,6-tetramethyl-1-pyrroline *N*-oxide (TEMPO) was added to the reaction system, the yield of **3a** was suppressed to 10%, and

most of the starting substrate **1a** was recovered. In addition, this reaction was conducted to 3 equiv. benzoyl peroxide gave a similar result. It indicates that this reaction involves free radical initiated mechanism (Scheme 2).

Scheme 2 Proposed reaction mechanism

3 Conclusions

An efficient method that easily convert tribromomethyl ketones to α,α -dibromo acetophenones at room temperature was developed and produced a series of α,α -dibromo acetophenones with high yields. Investigating the reaction mechanism showed that it was a radical reaction and it could be used in the reduction of aliphatic, aromatic, and heterocyclic tribromomethyl ketones. Our research innovatively used thiophenols as the radical reaction stimulator and tested aqueous/alcohol solvent systems. To our best knowledge, this is the first time attempt to reduce multi-halogenated compounds that use radical reactions.

4 Experimental section

4.1 General information

Visualization on thin-layer chromatography (TLC) was achieved by use of UV light (254 nm) or iodine. ^1H NMR (400 MHz) and ^{13}C NMR (100 MHz) spectra were recorded on a Varian-400 MHz spectrometer. The spectra were recorded in CDCl_3 as solvent at room temperature, ^1H NMR and ^{13}C NMR chemical shifts are reported relative to either the residual solvent peak (^{13}C) (δ 77.00 or 52.00) or TMS (^1H) (δ 0) as an internal standard. ESI-MS spectra were recorded on a BioTOF Q. Infrared spectra were recorded on a Nicolet MX-1E FT-IR spectrometer.

4.2 General procedure for the reaction

α,α,α -Tribromomethyl ketone (0.1 mmol, 1 equiv.) was added to 0.2 mL of ethanol and mixed at room temperature. After 5 min, 1.8 mL of sodium bromide solution (0.1 mmol/mL) and thiophenol (2.05 equiv.) were introduced. The reaction mixture was then stirred at room temperature and monitored by TLC. The reaction mixture was extracted with EtOAc, washed with brine, dried with Na_2SO_4 and then concentrated. A further purification by column chromatography ethyl acetate and petroleum ether ($V:V=1:9$) as the eluent afforded pure products.

2,2-Dibromo-1-phenyl-ethanone (3a): Reaction time 15 min, white solid (0.0264 g, 96% yield), m.p. 36 °C (Ref.^[15] yellow oil); ^1H NMR (CDCl_3 , 400 MHz) δ : 8.07 (t, $J=4.0$ Hz, 2H), 7.63 (t, $J=8$ Hz, 1H), 7.50 (t, $J=8$ Hz, 2H), 6.70 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 185.92, 134.41, 130.84, 129.68, 128.91, 39.64; IR (KBr) ν : 3064, 3015, 2975, 2850, 1697, 1579, 1418, 1270, 1192, 1161, 981, 801, 782, 684 cm^{-1} .

2,2-Dibromo-1-(4-chloro-phenyl)-ethanone (3b): Reaction time 60 min, white solid (0.0288 g, 96% yield). m.p. 91~92 °C (Ref.^[10a] 92.5~93.5 °C); ^1H NMR (CDCl_3 , 400 MHz) δ : 8.07 (d, $J=12.0$ Hz, 2H) 7.50 (d, $J=8.0$ Hz, 2H), 6.60 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 184.85, 141.04, 131.17, 129.26, 129.05, 39.20; IR (KBr) ν : 3059, 3037, 2926, 2853, 1695, 1587, 1567, 1487, 1401, 1275, 1206, 1093, 989, 845, 728 cm^{-1} .

2,2-Dibromo-1-(3-chloro-phenyl)-ethanone (3c): Reaction time 25 min; colorless oil (0.0302 g, 97% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 8.05 (d, $J=4.0$ Hz, 1H), 7.98

(d, $J=8.0$ Hz, 1H), 7.61~7.58 (m, 1H), 7.45 (t, $J=8.0$ Hz, 1H), 6.60 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 184.77, 135.26, 134.32, 132.37, 130.13, 129.72, 127.74, 39.05; IR (KBr) ν : 3068, 3021, 2927, 2850, 1703, 1572, 1421, 1280, 1192, 990, 803, 749, 677 cm^{-1} ; ESI HRMS calcd for $\text{C}_8\text{H}_6\text{Br}_2\text{ClO}$ [$\text{M} + \text{H}$] $^+$ 310.84684, found 310.84644.

2,2-Dibromo-1-(2-chloro-phenyl)-ethanone (3d): Reaction time 15 min, colorless oil (0.0282 g, 95% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.62 (d, $J=8.0$ Hz, 1H), 7.50~7.44 (m, 2H), 7.40~7.36 (m, 1H), 6.76 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 188.72, 134.02, 132.96, 131.07, 130.87, 130.43, 127.18, 42.06; IR (KBr) ν : 3019, 2925, 2849, 1720, 1588, 1434, 1282, 1184, 980, 782, 748, 672 cm^{-1} ; ESI HRMS calcd for $\text{C}_8\text{H}_6\text{Br}_2\text{ClO}$ [$\text{M} + \text{H}$] $^+$ 310.84684, found 310.84601.

2,2-Dibromo-1-(4-bromo-phenyl)-ethanone (3e): Reaction time 60 min, white solid (0.0332 g, 93% yield). m.p. 90~91 °C (Ref.^[10a] 92~93 °C); ^1H NMR (CDCl_3 , 400 MHz) δ : 7.98 (d, $J=8.0$ Hz, 2H), 7.67 (d, $J=8.0$ Hz, 2H), 6.60 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 185.07, 132.26, 131.19, 129.86, 129.48, 39.19; IR (KBr) ν : 3034, 1693, 1582, 1484, 1395, 1273, 1203, 987, 820, 719 cm^{-1} .

2,2-Dibromo-1-(2-bromo-phenyl)-ethanone (3f): Reaction time 40 min, colorless oil (0.0339 g, 95% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.65 (d, $J=8.0$ Hz, 1H), 7.57 (d, $J=8.0$ Hz, 1H), 7.45~7.36 (m, 2H), 6.73 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 189.32, 136.38, 133.48, 132.79, 130.72, 127.60, 119.04, 41.69; IR (KBr) ν : 3012, 2920, 2850, 2310, 1721, 1564, 1464, 1428, 1278, 1250, 1184, 1132, 1053, 1207, 978, 781, 724, 668 cm^{-1} ; ESI HRMS calcd for $\text{C}_8\text{H}_6\text{Br}_3\text{O}$ [$\text{M} + \text{H}$] $^+$ 354.79633, found 354.79553.

2,2-Dibromo-1-*p*-tolyl-ethanone (3g): Reaction time 35 min, white solid (0.0316 g, 96% yield). m.p. 95~96 °C (Ref.^[10b] 97~98 °C); ^1H NMR (CDCl_3 , 400 MHz) δ : 7.98 (d, $J=8.0$ Hz, 2H), 7.30 (d, $J=8.0$ Hz, 2H), 6.68 (s, 1H), 2.43 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 185.61, 145.69, 129.79, 129.62, 128.18, 39.48, 21.81; IR (KBr) ν : 3028, 2920, 2851, 1685, 1603, 1409, 1319, 1274, 1224, 1195, 1124, 960, 833, 752, 697 cm^{-1} .

2,2-Dibromo-1-*o*-tolyl-ethanone (3h): Reaction time 40 min, colorless oil (0.0288 g, 90% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.69 (d, $J=8.0$ Hz, 1H), 7.47~7.43 (m, 1H), 7.33~7.26 (m, 2H), 6.67 (s, 1H), 2.51 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 188.52, 140.21, 132.66, 132.25, 132.22, 128.05, 125.72, 42.18, 21.06; IR (KBr) ν : 3064, 3021, 2927, 1704, 1599, 1569, 1456, 1291, 1255, 1209, 1178, 970, 769, 738 cm^{-1} ; ESI HRMS calcd for $\text{C}_9\text{H}_9\text{Br}_2\text{O}$ [$\text{M} + \text{H}$] $^+$ 290.90147, found 290.90076.

2,2-Dibromo-1-(4-tert-butyl-phenyl)-ethanone (3i): Reaction time 15 min, colorless oil (0.0280 g, 91% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 8.03 (d, $J=8.0$ Hz, 2H), 7.52 (d, $J=8.0$ Hz, 2H), 6.68 (s, 1H), 1.34 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ : 185.55, 158.55, 129.70, 128.06, 125.92, 39.85, 35.33, 30.97; IR (KBr) ν : 2964, 2905, 2869, 1697, 1676, 1603, 1564, 1462, 1408, 1364, 1271, 1194,

1107, 982, 850 cm^{-1} ; ESI HRMS calcd for $\text{C}_{12}\text{H}_{15}\text{Br}_2\text{O}$ $[\text{M}+\text{H}]^+$ 332.94842, found 332.9476.

2,2-Dibromo-1-(4-fluoro-phenyl)-ethanone (3j): Reaction time 35 min, white solid (0.0265 g, 93% yield). m.p. 36 °C (Ref.^[15] yellow oil); ^1H NMR (CDCl_3 , 400 MHz) δ : 8.17~8.14 (m, 2H), 7.21~7.17 (m, 2H), 6.62 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 184.51, 167.59~165.02 (d, $J=257$ Hz), 132.71~132.61 (d, $J=10.0$ Hz), 127.07~127.05 (d, $J=2.0$ Hz), 116.30~116.08 (d, $J=22.0$ Hz), 39.31; IR (KBr) ν : 3016, 2927, 2851, 1697, 1597, 1507, 1270, 1241, 1189, 1160, 984, 850, 765 cm^{-1} .

4-(2,2-Dibromo-acetyl)-benzonitrile (3k): Reaction time 60 min, white solid (0.0255 g, 75% yield). m.p. 97~98 °C; ^1H NMR (CDCl_3 , 400 MHz) δ : 8.23 (d, $J=4.0$ Hz, 2H), 7.83 (d, $J=8.0$ Hz, 2H), 6.58 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 184.65, 134.06, 132.56, 130.26, 117.51, 117.47, 38.74; IR (KBr) ν : 3092, 3049, 3008, 2924, 2334, 1692, 1605, 1566, 1408, 1266, 1171, 1113, 991, 857, 766, 704, 678, 554, 523 cm^{-1} ; ESI HRMS calcd for $\text{C}_9\text{H}_6\text{Br}_2\text{NO}$ $[\text{M}+\text{H}]^+$ 300.87324, found 300.87356.

1,1-Dibromo-4-phenyl-butan-2-one (3l): Reaction time 40 min, colorless oil (0.0271 g, 94% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.31~7.28 (m, 2H), 7.23~7.20 (m, 3H), 5.74 (s, 1H), 3.24 (t, $J=8.0$ Hz, 2H), 2.98 (t, $J=8.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 195.88, 139.95, 128.58, 128.34, 126.44, 42.68, 36.68, 30.44; IR (KBr) ν : 3027, 2927, 2852, 2378, 2310, 1721, 1604, 1584, 1496, 1453, 1362, 1154, 1076, 751, 700 cm^{-1} ; ESI HRMS calcd for $\text{C}_{10}\text{H}_{11}\text{Br}_2\text{O}$ $[\text{M}+\text{H}]^+$ 303.90929, found 303.90955.

1,1-Dibromo-nonan-2-one (3m): Reaction time 60 min, colorless oil (0.0278 g, 94% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 5.72 (s, 1H), 2.94~2.88 (m, 1H), 2.82~2.76 (m, 1H), 2.21~2.14 (m, 1H), 1.30~1.14 (m, 4H), 1.10 (d, $J=8.0$ Hz, 3H), 0.92 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ : 196.15, 50.57, 44.34, 43.40, 29.95, 25.86, 22.41; IR (KBr) ν : 2927, 2856, 1723, 1465, 1405, 1376, 1149, 1094, 1056, 722, 659, 596, 507 cm^{-1} ; ESI HRMS calcd for $\text{C}_9\text{H}_{17}\text{Br}_2\text{O}$ $[\text{M}+\text{H}]^+$ 298.96407, found 298.96357.

1,1-Dibromo-4,6,6-trimethyl-heptan-2-one (3n): Reaction time 30 min, colorless oil (0.0259 g, 83% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 5.78 (s, 1H), 2.94 (t, $J=8.0$ Hz, 2H), 1.71~1.64 (m, 2H), 1.32~1.29 (m, 9H), 0.90~0.87 (m, 4H); ^{13}C NMR (100 MHz, CDCl_3) δ : 197.01, 42.96, 34.09, 31.61, 29.92, 28.85, 24.34, 22.57, 14.05; IR (KBr) ν : 2957, 2956, 2360, 1722, 1472, 1364, 1111, 668 cm^{-1} ; ESI HRMS calcd for $\text{C}_{10}\text{H}_{19}\text{Br}_2\text{O}$ $[\text{M}+\text{H}]^+$ 312.97972, found 312.97997.

2,2-Dibromo-1-thiophen-2-yl-ethanone (3o): Reaction time 30 min, colorless oil (0.0260 g, 92% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 8.00 (t, $J=4.0$ Hz, 1H), 7.78 (t, $J=4.0$ Hz, 1H), 7.20 (t, $J=4.0$ Hz, 1H), 6.48 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 179.71, 136.51, 136.27, 134.64, 128.48, 38.91; IR (KBr) ν : 3099, 3013, 2918, 2850, 2378, 2320, 1673, 1515, 1410, 1354, 1279, 1181, 1065, 860, 728 cm^{-1} ; ESI HRMS calcd for $\text{C}_6\text{H}_5\text{Br}_2\text{SO}$ $[\text{M}+\text{H}]^+$ 281.83441, found 281.83438.

2,2-Dibromo-1-naphthalen-1-yl-ethanone (3p): Reaction

time 45 min, colorless oil (0.0277 g, 85% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 8.42 (d, $J=8.0$ Hz, 1H), 8.07 (d, $J=8.0$ Hz, 1H), 7.94~7.89 (m, 2H), 7.65 (t, $J=4.0$ Hz, 1H), 7.58 (t, $J=4.0$ Hz, 1H), 7.51 (t, $J=4.0$ Hz, 1H), 6.83 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 188.59, 134.14, 133.92, 130.93, 130.31, 128.67, 127.50, 127.48, 127.04, 125.41, 124.10, 42.35; IR (KBr) ν : 3013, 2961, 2852, 1693, 1592, 1461, 1367, 1276, 1214, 1045, 1023, 940, 865, 760, 731, 643, 497, 441 cm^{-1} ; ESI HRMS calcd for $\text{C}_{12}\text{H}_9\text{Br}_2\text{O}$ $[\text{M}+\text{H}]^+$ 325.89364, found 325.89387.

2,2-Dibromo-1-naphthalen-2-yl-ethanone (3q): Reaction time 30 min, white solid (0.0269 g, 85% yield). m.p. 97~98 °C (Ref.^[10b] 100~101.5 °C); ^1H NMR (CDCl_3 , 400 MHz) δ : 8.63 (s, 1H), 8.09 (d, $J=4.0$ Hz, 1H), 7.99 (d, $J=4.0$ Hz, 1H), 7.94~7.89 (m, 2H), 7.65 (t, $J=4.0$ Hz, 1H), 7.59 (t, $J=4.0$ Hz, 1H), 6.86 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 185.98, 136.01, 132.27, 131.77, 129.82, 129.43, 128.92, 128.07, 127.86, 127.23, 124.68, 39.79; IR (KBr) ν : 3025, 2953, 2056, 2269, 1684, 1625, 1541, 1469, 1388, 1277, 1185, 1043, 917, 774, 732, 688, 625, 591, 471, 419 cm^{-1} .

Supporting Information ^1H NMR and ^{13}C NMR spectra of all products **3a**~**3q**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Kim, K.; Cho, J.; Yoon, S. C. *J. Chem. Soc., Perkin Trans. 1* **1995**, 253.
(b) Maglioli, P.; de, L. O.; Delogu, G.; Valle, G. *Tetrahedron* **1992**, 3, 365.
(c) Nemeryuk, M. P.; Likhovidova, M. M.; Mitsev, I. D.; Safonova, T. S. *Chem. Heterocycl. Compd. (Engl. Transl.)* **1986**, 22, 468.
- [2] Araki, S.; Hirashita, T.; Shimizu, K.; Ikeda, T.; Butsugan, Y. *Tetrahedron* **1996**, 52, 2803.
- [3] Kowalski, C. J.; Reddy, R. E. *J. Org. Chem.* **1992**, 57, 7194.
- [4] Zhdankin, V. V.; Stang, P. J. *Tetrahedron Lett.* **1993**, 34, 1461.
- [5] Kowalski, C. J.; Fields, K. W. *J. Am. Chem. Soc.* **1982**, 104, 321.
- [6] (a) Fujiwara, J.; Matsumura, A.; Matsuoka, Y. K. *J. Bull. Chem. Soc. Jpn.* **1976**, 49, 829.
(b) Kawabata, N.; Fujii, T.; Naka, M.; Yamashita, S. *J. Bull. Chem. Soc. Jpn.* **1977**, 50, 1005.
- [7] (a) Kowalski, C. J.; Fields, K. W. *J. Am. Chem. Soc.* **1982**, 104, 321.
(b) Ding, Y.; Huang, X. *Heteroat. Chem.* **2003**, 14, 304.
(c) Ghiaci, M.; Asgsarl, J. *Bull. Chem. Soc. Jpn.* **2001**, 74, 1151.
(d) Taylor, W. *J. Chem. Soc.* **1937**, 304.
- [8] (a) Pravst, I.; Zupan, M.; Stavber, S. *Tetrahedron* **2008**, 64, 5191.
(b) Liu, J.-H.; Li, W.-J.; Wang, C.; Li, Y.; Li, Z. *Tetrahedron* **2011**, 52, 4320.
(c) Kotek, V.; Polak, P.; Tobrman, T. *Monatsh. Chem.* **2016**, 147, 405.
(d) Catano, B.; Lee, J.; Kim, C.; Farrell, D.; Petersen, J. L.; Xin, Y. *Tetrahedron Lett.* **2015**, 56, 4124.
- [9] Magen, S.; Oren, J.; Fuchs, B. *Tetrahedron Lett.* **1984**, 25, 3369.
- [10] (a) Terent'ev, A. O.; Khodykin, S. V.; Krylov, I. B.; Ogibin, Y. N.; Nikishin, G. I. *Synthesis* **2006**, 1087.
(b) Terent'ev, A. O.; Krylov, I. B.; Vil, V. A.; Yu, Z.; Nikishin, G. I. *Cent. Eur. J. Chem.* **2012**, 10, 360.
- [11] Madabhushi, S.; Jillella, R.; Mallu, K. K. R.; Godala, K. R.; Vangipuram, V. S. *Tetrahedron Lett.* **2013**, 54, 3993.

- [12] Ye, C.-F.; Shreeve, J. M. *J. Org. Chem.* **2004**, *69*, 8561.
- [13] (a) Nobuta, T.; Hirashima, S.; Tada, N.; Miura, T.; Itoh, A. *Tetrahedron Lett.* **2010**, *51*, 4576.
(b) Tada, N.; Ban, K.; Ishigami, T.; Nobuta, T.; Miura, T.; Itoh, A. *Tetrahedron Lett.* **2011**, *52*, 3821.
- [14] Wu, P.; Xu, S.-H.; Xu, H.; Hu, H.-Y.; Zhang, W. *Tetrahedron Lett.* **2017**, *58*, 618.
- [15] Wang, J.-Y.; Jiang, Q.; Guo, C.-C. *Synth. Commun.* **2014**, 3130.
- [16] (a) Qvortrup, K.; Rankic, D. A.; Mac Millan, D. W. C. *J. Am. Chem. Soc.* **2014**, *136*, 626.
(b) Hager, D.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2014**, *136*, 16986.
(c) Cuthbertson, J. D.; MacMillan, D. W. C. *Nature* **2015**, *519*, 74.
- [17] Balati, H.; Zeng, J.; Zhang, Y.-H.; Abulikemu, A. R. *Synth. Commun.* **2018**, 887.
- [18] (a) Dhakshinamoorthy, A.; Alvaro, M.; Garcia, H. *Chem. Commun.* **2010**, *46*, 6476.
(b) Liu, Y.; Wang, H.; Wang, C.-P.; Wan, J.-P.; Wen, C.-P. *RSC Adv.* **2013**, *3*, 21368.

(Cheng, F.)