苯胺基于钛硅分子筛的选择性氧化合成氧化偶氮苯

宋玉婉 冯韶静 韩 维* 肖亚平*

(南京师范大学化学与材料科学学院 南京 210023)

摘要 报道了一种通过调节 TS-1 的硅钛摩尔比来提高苯胺高效、高选择性转化为相应氧化偶氮苯的方法. 在 TS-1 的硅钛摩尔比为 80 时, 以 H_2O_2 为氧化剂, 芳基苯胺以良好的产率以及高选择性转化为相应的氧化偶氮苯类化合物. 还研究了 H_2O_2 摩尔比、溶剂、催化剂回收率等反应参数对反应的影响.

关键词 苯胺; 氧化; 氧化偶氮苯; H_2O_2

Synthesis of Azoxybenzenes by Selective Oxidation of Anilines on Titanium Silicate Molecular Sieves

Song, Yuwan Feng, Shaojing Han, Wei* Xiao, Yaping*

(School of Chemistry and Materials Science, Nanjing Normal University, Nanjing 210023)

Abstract A convenient method for improving the conversion and selectivity of aniline to azoxybenzene by adjusting the ratio of silicon to titanium of TS-1 is reported. Over TS-1 (80), aryl anilines are transformed into the corresponding azoxybenzenes in good yields with H_2O_2 as the oxidant. The effect of different reaction parameters like molar ratio of H_2O_2 to aniline, solvent and the activity in catalyst recycling has been studied.

Keywords anilines; oxidation; azoxybenzene; H_2O_2

1 Introduction

The oxidation of anilines to azoxybenzenes is an important transformation that has been widely applied in the synthesis of their oxygenated derivatives such as hydroxylamine, oxime, azoxy, nitro, nitroso and azo compounds (Scheme 1).^[1] Additionally, azoxybenzenes are used as reducing agents, polymerization inhibitors, chemical stabilizers, and dyes. Therefore, many methods have been reported for the synthesis of azoxybenzene. Traditional methods heavily rely on stoichiometric reduction and oxidation such as reduction of nitro and nitroso aromatic compounds by using sodium bis(trimethylsilyl)amide^[2] and potassium borohydride,^[3] and oxidation of anilines with stoichiometric oxidants of sulfonic peracids,^[4] $\text{Bu}_4\text{N-OSO}_3$,^[5] and *m*-chloroperoxybenzoic acid.^[6] However, the use of stoichiometric reagents makes these processes environmentally unfriendly.

Various attempts were made for the oxidative coupling of aniline to azoxybenzene by catalytic protocols. For instance, homogeneous ruthenium-catalyzed oxidation of

aniline to provide azoxybenzene with the aid of environmentally clean and cheaper oxidant H_2O_2 , has been reported.^[7] However, homogeneous catalysis has a common problem of recycling of the catalyst. In order to address this drawback, heterogeneous catalysts like zeolites, vanadium silicates, and titanium containing molecular sieves, and cobalt oxide supported mesoporous silica for the synthesis of azoxybenzene from aniline are described in the literature.^[8] Additionally, silver supported tungsten oxide nanostructured catalyst^[9] and CuCr_2O_4 spinel nano catalyst^[10] were found to be suitable for the oxidation of aniline to azoxybenzene with 91% and 92% selectivities, respectively. Recently, Zhang *et al.* reported the synthesis of azoxybenzenes using mesoporous titania microspheres.^[11] Although various catalysts have been efficiently used for the preparation of azoxybenzene, a successful catalytic system with very high selectivity^[12] and yield of azoxybenzene using lower hydrogen peroxide concentration under mild conditions has not been achieved. Herein, we develop a general and high yielding method for the

* Corresponding authors. E-mail: hanwei@njnu.edu.cn; xiaoyaping@njnu.edu.cn

Received August 28, 2018; revised November 13, 2018; published online November 30, 2018.

Project supported by the National Natural Science Foundation of China (Nos. 21776139, 21302099), the Natural Science Foundation of Jiangsu Province (No. BK20161553), the Natural Science Foundation of Jiangsu Provincial Colleges and Universities (No. 16KJB150019).

国家自然科学基金(Nos. 21776139, 21302099)、江苏省自然科学基金(No. BK20161553)、江苏省高效自然科学基金(No. 16KJB150019)资助项目.

synthesis of various azoxybenzenes via titanium silicate molecular sieves-catalyzed oxidation of anilines using H_2O_2 as an oxidant. Notably, we observed that the molar ratios of silicon to titanium can adjust the selectivity of the reaction.^[13]

Scheme 1 Pathways for the synthesis of azoxybenzene

2 Results and discussion

2.1 Scanning electron microscope images of TS-1 (80)

The scanning electron microscope (SEM) images of TS-1 (80) are shown in Figure 1. From the images, it is observed that all the particles are spherical in shape with size of 150~200 nm.

Figure 1 Scanning-electron micrographs of TS-1 (80)

2.2 Influence of catalysts

The influence of several catalysts on the oxidation of aniline to azoxybenzene was studied. As shown in Table 1 are the results for the reactions performed in acetone and at 60 °C for 5 h with aniline to H_2O_2 molar ratio of 1 : 1. The catalyst plays a decisive role in this reaction according to the blank experimental result (Table 1, Entry 1). As can be seen from Entries 2~5 in Table 1, TS-1 gave higher reactivity than any of the other catalysts. Under the same conditions, the influence of different Si/Ti ratios was also explored. When the ratio of Si/Ti was decreased from 100 to 80, the yield would increase from 49.1% to 52.6%. However, the Si/Ti ratio further decreased to 33 and even to 10, leading to an apparent decrease in the yield. This is likely due to higher titanium content producing more white precipitate from the hydrolysis of the titanium source dur-

ing the catalyst synthesis that adheres to the surface of the catalyst and prevents the incorporation of the titanium into the TS-1 framework.

Table 1 Optimization of reaction conditions^a

Entry	Catalyst	H_2O_2 /Substrate	Solvent	Yield/%
1	No catalyst	1.0	Acetone	No product
2	TS-1 (33)	1.0	Acetone	36.6
3	Mg-TS-1	1.0	Acetone	28.1
4	Al-TS-1	1.0	Acetone	31.1
5	Fe-TS-1	1.0	Acetone	1.4
6	TS-1 (80)	1.0	Acetone	52.6
7	TS-1 (10)	1.0	Acetone	35.0
8	TS-1 (100)	1.0	Acetone	49.1
9	TS-1 (80)	1.5	Acetone	84.9
10	TS-1 (80)	2.0	Acetone	96.6
11	TS-1 (80)	2.5	Acetone	96.6
12	TS-1 (80)	2.0	Methanol	85.0
13	TS-1 (80)	2.0	Acetonitrile	95.3

^a Reaction conditions: substrate (1 mmol), catalyst (0.1 g), solvent (2 mL), water bath 60 °C, 5 h.

2.3 Dose of H_2O_2

Next, we have chosen TS-1 (80) as the catalyst for investigating the influence of the molar ratios of H_2O_2 to aniline on the oxidation reaction (Table 1, Entries 6 and 9~11). With increasing the ratio of H_2O_2 to aniline in the reaction mixture, the yield was found to increase. When the ratio of H_2O_2 to aniline reaches 2.0, the optimal yield of azoxybenzene was obtained (96.6%). High concentration of H_2O_2 didn't lead to significant change in the yield.

2.4 Influence of solvent

In a series of reactions with titanium silicate as the catalyst, the solvent was explored for the activity and selectivity.^[14] Table 1 (Entries 10 and 12~13) shows the effect of three different solvents on the oxidation of aniline. The reactions were proceeded at 60 °C for 5 h with keeping molar ratio of H_2O_2 to aniline. Acetone gave a higher yield (96%) for the desired product **2a** than acetonitrile (95%) or methanol (85%).

2.5 Reaction applicability

The optimized reaction conditions were then tested with a wide range of anilines (Table 2). Varying the substituents on the aromatic unit showed that electron-withdrawing (**2e~2h** and **2k~2l**) and electron-rich (**2b~2d**, **2i~2j**, **2m~2t**) aniline were competent substrates. The results indicate that the electronic effects of the substituents have little influence on the transformation. In contrast, the steric effect of the substituents was observed. For instance, 2-methylaniline (**1d**) gave the desired product **2d** in moderate yield, and extremely bulky 2,6-dimethyl aniline (**1p**) only led to a poor yield of **2p**.

Table 2 Oxidation of different anilines to azoxybenzenes^a

Entry	Aniline	Product	Yield/%	Entry	Aniline	Product	Yield/%
1			96	11			90
2			98	12			98
3			60	13			98
4			70	14			90
5			80	15			92
6			97	16			50
7			70	17			85
8			96	18			96
9			98	19			85

续表

Entry	Aniline	Product	Yield/%	Entry	Aniline	Product	Yield/%
10			96	20			88

^a Reaction conditions: **1** (1 mmol), H₂O₂ (2 mmol), catalyst (0.1 g), acetone (2 mL), water bath 60 °C, 5 h. ^b Conversion was determined by separation.

2.6 Reusability test

The reusability of TS-1 (80) was tested in the oxidation system for subsequent cycles. At the end of reaction, the catalyst was centrifuged from reaction mixture. The catalyst was calcined in a muffle furnace at 550 °C for 6 h and then reused in the next reaction. Three runs were performed under the same conditions. There is no significant loss of catalyst activity and the yield for each run was 94%, 93%, and 93%, respectively.

3 Conclusions

In summary, we achieved the selective oxidation of anilines to azoxybenzenes by adjusting the titanium content. It has been demonstrated that TS-1 with the Si/Ti molar ratio of 80 is superior to other ratios of titanium silicon molecular sieves in the oxidation of aniline to azoxybenzene. The catalyst also has excellent performance for the selective oxidation of a wide scope of substituted anilines to the corresponding azoxybenzenes. The reactions worked well (yields: 50%~98%) with different anilines at 60 °C with environmentally benign H₂O₂ as the oxidation and acetone as the solvent. In addition, the catalyst maintained its catalytic activity even after three recycles. These features make the protocol attractive for broad applications.

4 Experimental section

The TS-1 was prepared by hydrothermal method reported in Ref. [13, 15]. The TS-1 samples with different Si/Ti molar ratios were synthesized by adjusting the molar ratio of silicon source (tetraethyl orthosilicate, TEOS) to titanium source (titanium isopropoxide, TTIP). In a synthesis of Si/Ti=80, as an example, TPAOH (32 g, 25% aq.) was dissolved in deionized water (8 g) completely. TEOS (21 g, 98.5%) was added to the solution under stirring, and then the stirring was continued for 1 h. To a solution of TTIP (0.4 g) in dry isopropanol (2 g) was slowly dropped into the reaction mixture with stirring for 1 h. Note that in the above process the formation of white solid titanium dioxide by hydrolysis of isopropyl titanate has to be avoided. Finally, phosphoric acid (0.3 g, 85%, aq.) in distilled water (9.7 g) was added to the above mixed system and the reaction mixture was stirred for one more hour. The molar composition of the mixture was $n(\text{TEOS}) : n(\text{TTIP}) : n(\text{TPAOH}) : n(\text{IPA}) : n(\text{H}_3\text{PO}_4) : n(\text{H}_2\text{O}) = 80 : 1 : 27.9 : 23.6 : 1.8 : 1646$.

The above mixture was transferred to the autoclave and

crystallized for 6 h at 433 K. The solid product was separated from the mother liquor by centrifugation, followed by washing thoroughly with deionized water, 1 wt% aqueous solution of ammonium acetate, and deionized water. The precipitate was collected, dried in the oven at 393 K, and calcined in a muffle furnace at 823 K for 16 h.

The supported metal (Al, Fe, and Mg) catalyst was synthesized by an impregnation method.^[16] First, the above TS-1 sample was kept dry in dryer. A 5 mL of solution of corresponding metal salts [Al₂(NO₃)₃, FeCl₂•4H₂O, and Mg(OAc)₂] with 0.2 mol•L⁻¹ was used, and the TS-1 (1 g) was impregnated into the support under room temperature for 2 h. Then, the catalyst was dried in an oven for 6 h at 120 °C to remove water. Finally, the catalyst was calcined at 540 °C for 6 h in a muffle furnace.

All the catalytic reactions were carried out in an oven dried reaction tube with a magnetic stir bar. The reaction tube was charged with TS-1 (Si/Ti=80, 0.1 g), and a solution of substrate (1 mmol) in solvent (acetone, 2 mL). H₂O₂ (30%, 2 mmol) was added successively. Then, the reaction mixture was heated in a water bath at 60 °C for 5 h. Conversion was calculated on the basis of the substrate. The progress of the reaction was monitored by TLC. After that, the catalyst was centrifuged from the reaction mixture for recycling. The mixture was washed with saturated brine and extracted with ethyl acetate (15 mL×3). The organic solution was dried over anhydrous Na₂SO₄ and concentrated to 1~2 mL by rotary evaporator. Then, by means of column chromatography, the target products **2a**~**2t** were obtained.

1,2-Diphenyldiazene oxide (**2a**):^[17] 99.1 mg, 96% yield. Yellow liquid, m.p. 34~35 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.32 (d, *J*=8.0 Hz, 2H), 8.17 (d, *J*=7.5 Hz, 2H), 7.57~7.40 (m, 6H); ¹³C NMR (100 MHz, CDCl₃) δ: 148.3, 143.9, 131.6, 129.6, 128.8, 128.7, 125.5, 122.3.

1,2-Di-*p*-tolylidiazene oxide (**2b**):^[17] 113.1 mg, 99% yield. Orange solid, m.p. 67~68 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.18 (d, *J*=8.6 Hz, 2H), 8.12 (d, *J*=8.4 Hz, 2H), 7.28 (d, *J*=8.1 Hz, 4H), 2.44 (s, 3H), 2.41 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 146.1, 141.9, 141.7, 140.0, 129.3, 129.3, 125.6, 122.1, 21.6, 21.3.

1,2-Di-*m*-tolylidiazene oxide (**2c**):^[6] 113.1 mg, 60% yield. Orange solid, m.p. 35~36 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.10 (d, *J*=12.8 Hz, 2H), 7.98 (d, *J*=7.2 Hz, 2H), 7.41~7.35 (m, 3H), 7.22 (d, *J*=7.8 Hz, 1H), 2.47 (s, 3H), 2.44 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 148.4, 144.0, 139.0, 138.4, 132.3, 130.4, 128.6, 126.0, 122.7,

122.5, 119.5, 21.5, 21.4.

1,2-Di-*o*-tolylidiazene oxide (**2d**):^[6] 113.1 mg, 75% yield. Orange solid, m.p. 57~58 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.03 (d, *J*=7.9 Hz, 2H), 7.67 (d, *J*=7.7 Hz, 2H), 7.40~7.28 (m, 6H), 2.52 (s, 3H), 2.38 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 149.4, 142.7, 134.1, 131.8, 131.2, 130.8, 128.6, 126.6, 126.0, 123.5, 121.5, 18.4, 18.4.

1,2-Bis(3-bromophenyl)diazene oxide (**2e**):^[18] 178.0 mg, 80% yield. Yellow solid, m.p. 110~111 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.48 (t, *J*=2.0 Hz, 1H), 8.42 (t, *J*=1.9 Hz, 1H), 8.26~8.23 (m, 1H), 8.07~8.04 (m, 1H), 7.73~7.70 (m, 1H); 7.56~7.53 (m, 1H), 7.43~7.35 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 148.8, 144.6, 134.9, 132.8, 130.2, 130.0, 128.3, 125.7, 124.5, 122.4, 122.3, 121.1.

1,2-Bis(4-bromophenyl)diazene oxide (**2f**):^[17] 178.0 mg, 99% yield. Yellow solid, m.p. 169~170 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.18 (d, *J*=8.9 Hz, 2H), 8.08 (d, *J*=8.8 Hz, 2H), 7.65 (d, *J*=9.0 Hz, 2H), 7.61 (d, *J*=8.8 Hz, 2H); ¹³C NMR (100 MHz, CDCl₃) δ: 147.0, 142.6, 132.0, 132.0, 127.2, 126.5, 123.9, 123.6.

1,2-Bis(3-chlorophenyl)diazene oxide (**2g**):^[18] 133.6 mg, 70% yield. Yellow solid, m.p. 95~96 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.32 (s, 1H), 8.27 (s, 1H), 8.21 (d, *J*=8.2 Hz, 1H), 8.01 (d, *J*=7.6 Hz, 1H), 7.56 (d, *J*=8.0 Hz, 1H); 7.49~7.38 (m, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 153.0, 148.8, 144.5, 134.8, 134.4, 132.0, 129.9, 129.7, 125.4, 124.1, 122.8, 120.6.

1,2-Bis(4-chlorophenyl)diazene oxide (**2h**):^[18] 133.6 mg, 96% yield. Yellow solid, m.p. 151~152 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.26 (d, *J*=9.0 Hz, 2H), 8.16 (d, *J*=9.0 Hz, 2H), 7.50~7.44 (m, 4H); ¹³C NMR (100 MHz, CDCl₃) δ: 146.5, 142.2, 138.1, 135.2, 129.0, 129.0, 127.1, 123.7.

1,2-Bis(4-methoxyphenyl)diazene oxide (**2i**):^[17] 129.1 mg, 98% yield. Yellow solid, m.p. 115~116 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.29~8.24 (m, 4H), 6.99~6.95 (m, 4H), 3.89 (s, 3H), 3.88 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 161.8, 160.2, 141.7, 138.0, 127.8, 123.7, 113.7, 113.6, 55.7, 55.5.

1,2-Bis(3-methoxyphenyl)diazene oxide (**2j**):^[19] 129.1 mg, 96% yield. Yellow solid, m.p. 50~51 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.91~7.89 (s, 1H), 7.84 (s, 1H), 7.80 (m, 1H), 7.75~7.73 (m, 1H), 7.41~7.39 (m, 2H), 7.12~7.09 (m, 1H), 6.99~6.96 (m, 1H), 3.90 (s, 3H), 3.87 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 159.8, 159.5, 149.4, 144.9, 129.4, 129.3, 118.4, 116.3, 114.6, 110.0, 107.4, 55.7, 55.4.

1,2-Bis(3-fluorophenyl)diazene oxide (**2k**):^[6] 117.1 mg, 95% yield. Yellow solid, m.p. 50~51 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.13 (dd, *J*=8.2 Hz, 1H), 8.09~8.02 (m, 2H), 7.86~7.84 (m, 1H), 7.53~7.43 (m, 2H), 7.32~7.28 (m, 1H), 7.16~7.11 (m, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.6 (d, *J*=7.7 Hz), 161.1 (d, *J*=5.2 Hz), 149.3 (d, *J*=9.7 Hz), 144.8 (d, *J*=9.2 Hz), 130.1 (d, *J*=8.3 Hz), 129.8 (d, *J*=8.7 Hz), 122.1 (d, *J*=3.0 Hz), 119.0 (d, *J*=21.3 Hz), 118.1 (d, *J*=3.2 Hz), 116.9 (d, *J*=21.5

Hz), 112.2 (d, *J*=24.5 Hz), 110.3 (d, *J*=26.6 Hz).

1,2-Bis(4-fluorophenyl)diazene oxide (**2l**):^[17] 117.1 mg, 98% yield. Yellow solid, m.p. 89~90 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.34~8.30 (m, 2H), 8.28~8.24 (m, 2H), 7.19~7.15 (m, 4H); ¹³C NMR (100 MHz, CDCl₃) δ: 163.8 (d, *J*=251.3 Hz), 162.5 (d, *J*=250.9 Hz), 144.2 (d, *J*=2.1 Hz), 140.2 (d, *J*=3.3 Hz), 128.0 (d, *J*=8.4 Hz), 124.5 (d, *J*=9.2 Hz), 115.8 (d, *J*=5.8 Hz), 115.6 (d, *J*=4.9 Hz).

1,2-Bis(4-ethylphenyl)diazene oxide (**2m**):^[18] 127.2 mg, 98% yield. Orange liquid, ¹H NMR (400 MHz, CDCl₃) δ: 8.22 (d, *J*=8.5 Hz, 2H), 8.16 (d, *J*=8.4 Hz, 2H), 7.32 (d, *J*=8.4 Hz, 4H), 2.77~2.69 (m, 4H), 1.29 (t, *J*=15.6 Hz, 6H); ¹³C NMR (100 MHz, CDCl₃) δ: 148.1, 146.3, 146.2, 142.0, 128.1, 128.0, 125.7, 122.2, 28.8, 28.6, 15.3, 15.3.

1,2-Bis(4-isopropylphenyl)diazene oxide (**2n**): 141.2 mg, 90% yield. Orange liquid, ¹H NMR (400 MHz, CDCl₃) δ: 8.18~8.01 (m, 4H), 7.44~7.40 (m, 3H), 7.28 (d, *J*=7.7 Hz, 1H), 3.08~2.97 (m, 2H), 1.32 (dd, *J*=6.9 Hz, 6.9 Hz, 12H); ¹³C NMR (100 MHz, CDCl₃) δ: 150.0, 149.4, 148.5, 144.1, 129.7, 128.7, 128.5, 127.9, 123.8, 122.7, 120.3, 119.9, 34.1, 23.9, 23.8.

1,2-Bis(3-isopropylphenyl)diazene oxide (**2o**): 141.2 mg, 92% yield. Orange liquid, ¹H NMR (400 MHz, CDCl₃) δ: 8.17~8.00 (m, 4H), 7.43~7.39 (m, 3H), 7.28~7.23 (m, 1H), 3.06~2.95 (m, 2H), 1.32~1.29 (m, 12H); ¹³C NMR (100 MHz, CDCl₃) δ: 150.0, 149.4, 148.5, 144.1, 129.7, 128.6, 128.5, 127.9, 123.8, 122.7, 120.3, 119.8, 34.1, 23.9, 23.8.

1,2-Bis(2,6-dimethylphenyl)diazene oxide (**2p**):^[20] 127.2 mg, 50% yield. Orange solid, m.p. 79~81 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.18~7.13 (m, 6H), 2.47 (s, 6H), 2.32 (s, 6H); ¹³C NMR (100 MHz, CDCl₃) δ: 149.5, 141.7, 130.8, 130.3, 129.0, 129.0, 128.5, 127.5, 19.4, 18.0.

1,2-Bis(4-ethoxyphenyl)diazene oxide (**2q**):^[17] 143.2 mg, 85% yield. orange solid known. m.p. 134~135 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.28~8.22 (m, 4H), 6.97~6.93 (m, 4H), 4.11 (q, *J*=6.9 Hz, 4H), 1.47~1.43 (m, 6H); ¹³C NMR (100 MHz, CDCl₃) δ: 161.2, 159.6, 141.5, 137.8, 127.8, 123.7, 114.1, 114.0, 63.9, 63.7, 14.8, 14.7.

1,2-Bis(4-(tert-butyl)phenyl)diazene oxide (**2r**):^[6] 155.2 mg, 96% yield. Yellow solid, m.p. 141~142 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.21 (d, *J*=8.9 Hz, 2H), 8.15 (d, *J*=8.8 Hz, 2H), 7.50 (dd, *J*=8.9 Hz, 8.8 Hz, 4H), 1.37 (s, 9H), 1.36 (s, 9H); ¹³C NMR (100 MHz, CDCl₃) δ: 155.0, 153.0, 146.0, 141.7, 125.7, 125.6, 121.9, 35.0, 31.2, 31.2; m.p. 141~142 °C.

1,2-Bis(3-chloro-4-methylphenyl)diazene oxide (**2s**):^[19] 147.6 mg, 85% yield. Yellow solid, m.p. 116~117 °C; ¹H NMR (400 MHz, CDCl₃) δ: 8.31 (q, *J*=2.5 Hz, 2H), 8.08 (dd, *J*=8.4 Hz, 8.4 Hz, 1H), 7.97 (dd, *J*=8.3 Hz, 8.3 Hz, 1H), 7.37~7.32 (m, 2H), 2.46 (s, 3H), 2.43 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ: 146.7, 142.5, 140.3, 138.2, 134.6, 134.3, 130.9, 130.8, 126.0, 124.2, 123.0, 120.4, 20.3, 20.1.

1,2-Bis(3,5-dimethylphenyl)diazene oxide (**2t**): 127.2 mg, 88% yield. Yellow solid, m.p. 106~107 °C; ¹H NMR

(400 MHz, CDCl₃) δ : 7.9 (s, 2H), 7.78 (s, 2H), 7.17 (s, 1H), 7.04 (s, 1H), 2.42 (6H, s), 2.39 (6H, s); ¹³C NMR (100 MHz, CDCl₃) δ : 148.5, 144.1, 138.7, 138.2, 133.1, 131.2, 123.0, 120.0, 21.4, 21.3.

Supporting Information ¹H NMR and ¹³C NMR spectra for compounds **2a** ~ **2t**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] Suresh, S.; Joseph, R.; Jayachandran, B.; Pol, A. V.; Vinod, M. P.; Sudalai, A.; Sonawane, H. R.; Ravindranathan, T. *Tetrahedron Lett.* **1995**, 51, 11305.
- [2] Hwu, J. R.; Das, A. R.; Yang, C. W.; Jiannjyh Huang, A.; Hsu, M. H. *Org. Lett.* **2005**, 7, 3211.
- [3] Liu, Y.; Liu, B.; Guo, A.; Dong, Z.; Jin, S.; Lu, Y. *Molecules* **2011**, 16, 3563.
- [4] Kluge, R.; Schulz, M.; Liebsch, S. *Tetrahedron Lett.* **1996**, 52, 5773.
- [5] Rezaeifard, A.; Jafarpour, M.; Naseri, M. A.; Shariati, R. *Dyes Pigm.* **2008**, 76, 840.
- [6] Yang, F.; Wang, Z.; Zhang, X.; Jiang, L.; Li, Y.; Wang, L. *Chem-CatChem* **2015**, 7, 3450.
- [7] Barak, G.; Sasson, Y. *J. Org. Chem.* **1989**, 54, 3484.
- [8] (a) Sonawane, H. R.; Pol, A. V.; Moghe, P. P.; Biswas, S. S.; Shariati, R. *Chem. Commun.* **1994**, 25, 1215.
(b) Gontier, S.; Tuel, A. *J. Catal.* **1995**, 157, 124.
(c) Chang, C. F.; Liu, S. T. *J. Mol. Catal. A: Chem.* **2009**, 299, 121.
(d) Das, D. R.; Talukdar, A. K. *ChemistrySelect* **2017**, 2, 8983.
- [9] Ghosh, S.; Acharyya, S. S.; Sasaki, T.; Bal, R. *Green Chem.* **2015**, 17, 1867.
- [10] Acharyya, S. S.; Ghosh, S.; Bal, R. *ACS Sustainable Chem. Eng.* **2014**, 2, 584.
- [11] Yang, L.; Shi, G.; Ke, X.; Shena, R.; Zhang, L. *CrystEngComm* **2014**, 16, 1620.
- [12] For aniline forming polyaniline during the oxidation process, see:
(a) Yu, L.; Huang, Y.; Wei, Z.; Ding, Y.; Su, C.; Xu, Q. *ChemInform* **2016**, 47, 8677.
(b) Yu, L.; Han, Z. *Mater. Lett.* **2016**, 184, 312.
(c) Yu, L.; Han, Z.; Ding, Y. *Org. Process Res. Dev.* **2016**, 20, 2124.
(d) Wang, Q.; Jing, X.; Han, J.; Yu, L.; Xu, Q. *Mater. Lett.* **2018**, 215, 65.
(e) Liu, Y.; Tang, D.; Cao, K.; Yu, L.; Han, J.; Xu, Q. *J. Catal.* **2018**, 360, 250.
- [13] For an example of the molar ratios of silicon to titanium affecting the selectivity of oxidation of aromatic compounds, see: Mukherjee, P.; Bhaumik, A.; Kumar, R. *Ind. Eng. Chem. Res.* **2007**, 46, 8657.
- [14] Barbera, D.; Cavani, F.; D'Alessandro, T.; Fornasari, G.; Guidetti, S.; Aloise, A.; Giordano, G.; Piumetti, M.; Bonelli, B.; Zanzottera, C. *J. Catal.* **2010**, 275, 158.
- [15] Zuo, Y.; Wang, X.; Guo, X. *Ind. Eng. Chem. Res.* **2011**, 50, 8485.
- [16] Ahmed, M. H. M.; Muraza, O.; Jamil, A. K.; Shafei, E. N.; Yamani, Z. H.; Choi, K. H. *Energy Fuels* **2017**, 31, 5482.
- [17] Voutyritsa, E.; Theodorou, A.; Kokotou, M. G.; Kokotos, C. G. *Green Chem.* **2017**, 19, 1291.
- [18] Chen, Y. F.; Chen, J.; Lin, L. J.; Chuang, G. J. *J. Org. Chem.* **2017**, 82, 11626.
- [19] Long, Z.; Wang, Z.; Zhou, D.; Wan, D.; You, J. *Org. Lett.* **2017**, 19.
- [20] Stowell, J. C.; Lau, C. M. *J. Org. Chem.* **2002**, 51, 1614.

(Li, L.; Fan, Y.)