碘诱导的醛糖与 *o*-氨基苯甲酰胺的氧化缩合合成糖基喹唑啉酮

琚欢欢 孙佳婧 李小六* 陈 华*

(河北大学化学与环境科学学院 河北省化学生物学重点实验室 保定 071002)

摘要 以不保护的单/双糖和邻氨基苯甲酰胺为原料, 通过碘诱导的氧化缩合一锅法反应, 合成了系列糖基喹唑啉酮衍生物, 产率 30%~80%, 为该类化合物的合成提供了一个有效方法. 碘的用量对该反应有重要的影响.

关键词 糖基喹唑啉酮; 氧化缩合; 碘诱导

Synthesis of Aldo-quinazolinones by Iodine-Induced Oxidative Condensation of Aldoses and *o*-Aminobenzamides

Ju, Huanhuan Sun, Jiajing Li, Xiaoliu* Chen, Hua*

(Key Laboratory of Chemical Biology of Hebei Province, College of Chemistry and Environmental Science, Hebei University, Baoding 071002)

Abstract A series of aldo-quinazolinones were synthesized by a one-pot iodine-induced oxidative condensation of the unprotected mono-/di-saccharides and *o*-aminobenzamides in the yields of 30%~80%, providing an effective protocol for the preparation of aldo-quinazolinones. The amount of iodine has an important effect on the reaction.

Keywords aldo-quinazolinone; oxidative condensation; iodine-induced

1 Introduction

4-(3*H*)-Quinazolinone and its derivatives have attracted many attentions due to their diverse biological activities,^[1] such as anti-cancer,^[2] anti-bacterial,^[3] anti-virus,^[4] anti-diabetes,^[5] and anti-glucosidase activities.^[6] As an important nitrogen-containing pharmacophore, 4-(3*H*)-quinazolinone has been made many efforts to develop the synthetic strategies and structural diversities to meet the ever-growing requirements for the new drug discovery.^[7,8] So far, a vast number of 4-(3*H*)-quinazolinone derivatives have been prepared and evaluated for their various biological activities.^[9] Owing to their widespread applications, the development of novel quinazolinones has been a long-standing interest for chemists.^[10~12]

There are two main strategies for constructing the skeleton of 4-(3*H*)-quinazolinone. One is the direct condensation of aldehyde and *o*-aminobenzamide,^[13,14] the other is the ammonolysis reaction of aromatic amine and isatoic anhydride derived from anthranilic acid.^[15,16] The former is the most common protocol as its simple and efficient.^[17,18] The condensation reactions were generally catalyzed by

Lewis acids, such as FeCl₃,^[19] CuCl₂,^[20] and I₂.^[21~23] The starting reactants could be the precursors of aldehyde and *o*-aminobenzamide, respectively, such as benzyl alcohol,^[24] benzyl amine,^[25] and *o*-aminobenzonitrile.^[26] Although there are several literatures to describe the preparation of 4-(3*H*)-quinazolinone from the aliphatic aldehydes,^[27~29] aldoses as reactants weren't studied yet. As a continuation of our researches on novel 4-(3*H*)-quinazolinone derivatives,^[30~32] herein, we would like to report the synthesis of the aldo-quinazolinone by a one-pot iodine-induced reaction.

2 Results and discussion

We initiated our experiment by examining the reaction of *D*-ribose (**1a**) with *o*-aminobenzamide (**2a**) in dimethyl sulfoxide (DMSO) solution at 60 °C (Table 1, Entry 1) according to the method for the aromatic aldehydes.^[13] However, the reaction didn't occur. Subsequently, the reaction was carried out in methanol solution at 40 °C in the presence of some Lewis acids FeCl₃^[19] or CuCl₂^[20] (Table 1, Entries 2, 3), but the reactions still didn't proceed. When using AcOH as the catalyst,^[33] the reaction could yield

* Corresponding authors. E-mail: lixl@hbu.cn; hua-todd@163.com

Received January 17, 2019; revised March 4, 2019; published online March 21, 2019.

Project supported by the National Natural Science Foundation of China (Nos. 21772031, 21372060).

国家自然科学基金(Nos. 21772031, 21372060)资助项目.

small amount of the intermediate **A** (Scheme 1, two pairs of inseparable amino glycosides including α/β -furanosides and α/β -pyranosides) (Table 1, Entry 4). Finally, iodine (I_2) was added to promote the reaction (Table 1, Entry 5).^[21] The reaction could succeed in generating the target ribo-quinazolinone **3a** with a low yield of 12%, although the iodine-induced reaction was very complicated.

The condensation conditions were further optimized. The results showed that the amount of iodine had an important effect on the reaction. Adjusting the amount of iodine could control the synthesis in its different stages to give the different products. For example, adding the catalytic amount of iodine (0.02 equiv.) would lead to the reaction which only afford the intermediate **A** (Table 1, Entry

6). When the amount of I_2 increased to 0.1 equiv., intermediate **A** decreased slowly, another key intermediate **B** (ribo-dihydroquinazolinones, a pair of inseparable isomers) was greatly produced, and **3a** was synthesized a little simultaneously (Table 1, Entry 7). Further improving the amount of I_2 to 0.23 equiv., the intermediate **B** could be oxidized to **3a** completely (Table 1, Entry 8).

The possible mechanism for the synthesis of **3a** was summarized in Scheme 1.^[21,34] It was likely that the Schiff base was firstly formed between *D*-ribose and *o*-aminobenzamide. Theoretically, 4-/5-hydroxyl on the ribose could attack the imine to generate the intermediate **A**, while the amino on the amide could attack the imine to produce the intermediate **B**, respectively. In the presence of iodine, the

Table 1 Optimization of the condensation of *D*-ribose and *o*-aminobenzamide

Entry	Condition ^a	I_2 /equiv.	Product	Yield/%
1	DMSO, 60 °C, 24 h		— ^b	
2	$FeCl_3$ (2 mmol), CH_3OH , 40 °C, 24 h		—	
3	$CuCl_2$ (2 mmol), CH_3OH , 40 °C, 24 h		—	
4	$AcOH$ (0.1 mL), CH_3OH , 40 °C, 24 h		Intermediate A	28
5	I_2 , CH_3OH , 40 °C, 24 h	0.15	Intermediate B and 3a	12 ^c
6	I_2 , CH_3OH , 40 °C, 8 h	0.02	Intermediate A	62
7	I_2 , CH_3OH , 40 °C, 8 h	0.1	Intermediate B and 3a	6 ^c
8	I_2 , CH_3OH , 40 °C, 24 h	0.23	3a	46

^a The molar ratio of *D*-ribose (180 mg, 1 mmol) and *o*-aminobenzamide is 1.0 : 1.2, the solvent is 5 mL. ^b No reaction. ^c Yield of **3a**.

Figure 1 X-ray crystallographic structure of **3a**
 synthesis of **B** was predominant, and **A** could transform to

B at the same time. Finally, the ribo-dihydroquinazolinone **B** was oxidized to **3a** by iodine, which was identified by its X-ray crystallographic structure (Figure 1).^[35] To confirm the rationality of the mechanism, the reaction was carried out using the pure intermediate **A**, which afforded **B** and **3a** as expected in the presence of I_2 . Additionally, the pure intermediate **B** could greatly convert to **3a** (70% yield) by the I_2 -catalyzed reaction at 70 °C.

Table 2 Synthesis of the aldo-quinazolinones **3a**~**3g** through the condensation of the unprotected saccharides and *o*-aminobenzamide

Entry	Saccharide	I_2 /equiv.	Product	Yield/%
1	 1a <i>D</i> -ribose	0.23	 3a <i>ribo</i> -quinazolinone	46
2	 1b <i>D</i> -xylose	0.65	 3b <i>xylo</i> -quinazolinone	43
3	 1c <i>D</i> -arabinose	0.78	 3c <i>arabino</i> -quinazolinone	48
4	 1d <i>D</i> -glucose	0.49	 3d <i>gluco</i> -quinazolinone	45
5	 1e <i>D</i> -galactose	0.82	 3e <i>galacto</i> -quinazolinone	41
6	 1f <i>D</i> -mannose	0.42	 3f <i>manno</i> -quinazolinone	59
7	 1g <i>D</i> -maltose	2.42	 3g <i>malto</i> -quinazolinone	30

To simplify the synthetic progress, we could directly add 0.23 equiv. of I_2 into the methanol solution of *D*-ribose and *o*-aminobenzamide at 40 °C for 24 h. After simple workup and silica gel column chromatography, aldo-quinazolinone **3a** would precipitate from the eluent in the yield of 46% (Table 2, Entry 1). Under the similar conditions using different equivalents of I_2 , the corresponding aldo-quinazolinones **3b**~**3g** (Table 2, Entries 2~7) were successfully obtained from seven saccharides with *o*-aminobenzamides in the satisfactory yields of 30%~59%. The used saccharides included five monosaccharides, *D*-xylose (**1b**), *D*-arabinose (**1c**), *D*-glucose (**1d**), *D*-mannose (**1e**) and *D*-galactose (**1f**), and one disaccharide, *D*-maltose (**1g**), respectively. To demonstrate the generality of the reaction, the substituted *o*-aminobenzamides were also used to react with *D*-arabinose (**1c**), *D*-mannose (**1e**), and *D*-maltose (**1g**) to prepare the corresponding aldo-quinazolinones **3c-1**~**3c-3**, **3f-1**~**3f-3**, and **3g-1**~**3g-2** (Table 3, Entries 1~8) in the moderate yields of 38%~80%, respectively. However, the production of **3g-2** was accompanied by a side reaction leading to an inseparable I-substituted byproduct **3g-2-1** (Table 3, Entry 8), as the same as the literature described.^[36] Its molecular formula was deduced to be $C_{20}H_{27}IN_2O_{11}$ by using HRESIMS.

It was worth noting that the fluorescent characteristics of the quinazolinone analogs were interesting. For example, the intermediate **A** exhibited blue fluorescence, the intermediate **B** appeared green fluorescence, while the aldo-quinazolinones showed no fluorescence in the fluorescent examination. The significant change of the fluorescence color of the compounds could help us to easily monitor the one-pot multistep transformation in the thin layer chromatography (TLC) detection.

2 Conclusions

In conclusion, a series of novel aldo-quinazolinones were synthesized by the iodine-induced condensation of the monosaccharides and *o*-aminobenzamides, providing an effective protocol for the preparation of aldo-quinazolinones.

3 Experimental

3.1 Materials

Melting points were measured on an SGW[®] X-4 micro melting point apparatus and were uncorrected. Optical rotations were determined on an SGW[®]-1 automatic polarimeter. 1H NMR, and ^{13}C NMR spectra were measured on a RT-NMR Bruker AVANCE 600 (600 MHz) and AVANCE 400 (400 MHz), NMR spectrometer using tetramethylsilane (Me_4Si) as an internal standard. Mass Spectra (MS) and High Resolution Mass Spectra (HRMS) were carried out on a FTICR-MS (Ionspec 7.0T) mass spectrometer with electrospray ionization (ESI). X-Ray crystallographic measurements were made on a Bruker SMART CCD Diffractometer. Thin-layer chromatography (TLC) was performed on precoated plates (Qingdao GF₂₅₄) with detection with

phosphomolybdic acid in EtOH/H₂O followed by heating. Column chromatography was performed using SiO₂ (Qingdao 300~400 mesh).

3.2 Methods

3.2.1 Synthesis of the intermediate A

A mixture of *D*-ribose (1.8 g, 12.0 mmol) and *o*-aminobenzamide (2.0 g, 1.2 equiv.) in 60 mL methanol was stirred at 40 °C under N₂ atmosphere for 10 min, then iodine (0.1 g, 0.02 equiv.) was added into the solution. The mixture was continued to stir at 40 °C for 8 h, the reaction finished (monitored by TLC). After the solution cooled down, methanol was evaporated under vacuum to afford the crude product. The residue was purified using flash column chromatography (dichloromethane-methanol *V*:*V*=2:1) to afford the intermediate **A** (two pairs of inseparable isomers, four compounds). The 1H and ^{13}C NMR spectra of **A** could be seen in the supporting materials. HR-ESI-MS calcd for $C_{12}H_{16}N_2O_5Na$ ($[M+Na]^+$) 291.0957, found 291.0956.

3.2.2 Synthesis of the intermediate B

A mixture of *D*-ribose (1.8 g, 12.0 mmol) and 2-aminobenzamide (2.0 g, 1.2 equiv.) in 60 mL methanol was stirred at 40 °C under N₂ atmosphere for 10 min, then iodine (0.5 g, 0.1 equiv.) was added into the solution. The mixture was stirred for 8 h, the reaction completed (the intermediate **A** disappeared on TLC). After the solution cooled down, methanol was evaporated under vacuum to afford the crude product. The residue was purified using flash column chromatography (dichloromethane-methanol *V*:*V*=6:1) to afford the intermediate **B**.

((2*S*,3*R*)-1,2,3,4-tetrahydroxybutyl)-2,3-dihydroquinazolin-4(1*H*)-one (intermediate **B**, a pair of inseparable isomers): White solid, yield 17%; m.p. 165.9~169.6 °C, 1H NMR (600 MHz, CD₃OD) δ : 7.69~7.67 (m, 1H), 7.28~7.25 (m, 1H), 6.78 (d, *J*=8.4 Hz, 0.6H), 6.74~6.71 (m, 1.4H), 5.03 (d, *J*=4.8 Hz, 0.4H), 4.97 (d, *J*=5.4 Hz, 1H), 3.88 (t, *J*=4.2 Hz, 0.5H), 3.84~3.81 (m, 1H), 3.79~3.76 (m, 1.5H), 3.67 (dd, *J*=11.4, 6.0 Hz, 1H); ^{13}C NMR (150 MHz, CD₃OD) δ : 167.4, 167.3, 149.9, 149.5, 135.2, 128.7, 128.6, 118.9, 116.1, 115.8, 115.7, 74.7, 74.4, 73.9, 73.8, 73.7, 68.0, 67.9, 64.6, 64.5; HR-ESI-MS calcd for $C_{12}H_{17}N_2O_5$ ($[M+H]^+$) 269.1137, found 269.1132.

3.2.3 General synthesis of the aldo-quinazolinones 3

A mixture of *D*-ribose (1.8 g, 12.0 mmol) and 2-aminobenzamide (2.0 g, 1.2 equiv.) in 60 mL methanol was stirred at 40 °C under N₂ atmosphere for 10 min, then iodine (0.9 g, 0.23 equiv.) was added into the solution. The mixture was stirred for 24 h till the reaction completed. After the solution cooled down, methanol was evaporated under vacuum to afford the crude product. The residue was purified using flash column chromatography (dichloromethane-methanol *V*:*V*=6:1). White solid could precipitate from the eluent and was washed with ethyl acetate (5 mL×3) to afford compound **3a**.

Table 3 Synthesis of the aldo-quinazolinones **3c-1**~**3c-3**, **3f-1**~**3f-3**, and **3g-1**~**3g-2** through the condensation of the unprotected saccharides and the substituted *o*-aminobenzamides

Entry	Saccharide	Substituted <i>o</i> -amino benzamide	I ₂ /equiv.	Product	Yield/%
1	1c <i>D</i> -arabinose	2b	1.37	3c-1	57
2	1c <i>D</i> -arabinose	2c	0.62	3c-2	50
3	1f <i>D</i> -mannose	2d	0.80	3c-3	45
4	1f <i>D</i> -mannose	2b	1.16	3f-1	78
5	1f <i>D</i> -mannose	2c	0.75	3f-2	57
6	1f <i>D</i> -mannose	2d	0.81	3f-3	80
7	1g <i>D</i> -maltose	2b	2.75	3g-1	38
8	1g <i>D</i> -maltose	2d	1.87	3g-2 3g-2-1	35 ^a

^a Total yields of **3g-2** and **3g-2-1**.

Under the similar conditions with different amounts of I_2 , the corresponding aldo-quinazolinones **3** were obtained in satisfactory yields.

2-((1*S*,2*S*,3*R*)-1,2,3,4-Tetrahydroxybutyl)quinazolin-4(3*H*)-one (**3a**): White solid, yield 46%. m.p. 183.9 ~ 185.6 °C; $[\alpha]_D^{25}$ -8.0 (*c* 0.01, H_2O); 1H NMR (600 MHz, $DMSO-d_6$) δ : 11.36 (s, 1H), 8.11 (d, *J*=7.8 Hz, 1H), 7.79 (t, *J*=7.2 Hz, 1H), 7.64 (d, *J*=7.8 Hz, 1H), 7.48 (t, *J*=7.2 Hz, 1H), 5.92 (s, 1H), 5.08 (s, 1H), 4.71 (s, 1H), 4.64 (s, 1H), 4.40 (t, *J*=5.4 Hz, 1H), 3.86 (t, *J*=5.4 Hz, 1H), 3.61 ~ 3.58 (m, 2H), 3.45 ~ 3.41 (m, 1H); ^{13}C NMR (150 MHz, $DMSO-d_6$) δ : 161.1, 157.7, 148.5, 134.3, 126.9, 126.2, 125.8, 121.3, 74.2, 72.7, 72.0, 62.9; HR-ESI-MS calcd for $C_{12}H_{14}N_2O_5Na$ ($[M+Na]^+$) 289.0800, found 289.0802.

2-((1*S*,2*R*,3*R*)-1,2,3,4-Tetrahydroxybutyl)quinazolin-4(3*H*)-one (**3b**): White solid, yield 43%. m.p. 115.4 ~ 117.0 °C; $[\alpha]_D^{25}$ +36.8 (*c* 0.04, H_2O); 1H NMR (600 MHz, CD_3OD) δ : 8.15 (d, *J*=7.8 Hz, 1H), 7.71 (t, *J*=7.8 Hz, 1H), 7.63 (d, *J*=7.8 Hz, 1H), 7.41 (t, *J*=7.8 Hz, 1H), 4.783 (s, 1H), 4.045 (s, 1H), 3.88 (d, *J*=4.2 Hz, 1H), 3.75 ~ 3.69 (m, 2H); ^{13}C NMR (150 MHz, CD_3OD) δ : 167.8, 162.2, 150.3, 135.1, 127.1, 127.1, 127.1, 122.2, 74.4, 74.4, 74.0, 64.2; HR-ESI-MS calcd for $C_{12}H_{14}N_2O_5Na$ ($[M+Na]^+$) 289.0800, found 289.0804.

2-((1*R*,2*S*,3*R*)-1,2,3,4-tetrahydroxybutyl)quinazolin-4(3*H*)-one (**3c**): White solid, yield 48%, m.p. 206.0 ~ 207.7 °C, $[\alpha]_D^{25}$ -15.6 (*c* 0.04, H_2O); 1H NMR (600 MHz, $DMSO-d_6$) δ : 11.20 (s, 1H, NH), 8.11 (d, *J*=7.8 Hz, 1H), 7.79 (t, *J*=7.2 Hz, 1H), 7.63 (d, *J*=7.2 Hz, 1H), 7.48 (t, *J*=7.8 Hz, 1H), 5.56 (d, *J*=6.0 Hz, 1H), 4.81 (d, *J*=5.4 Hz, 2H), 4.73 (d, *J*=4.8 Hz, 1H), 4.44 (s, 1H), 3.76 ~ 3.73 (m, 1H), 3.64 (d, *J*=10.8 Hz, 1H), 3.57 (s, 1H), 3.48 ~ 3.46 (m, 1H); ^{13}C NMR (150 MHz, $DMSO-d_6$) δ : 161.1, 159.1, 148.6, 134.4, 126.8, 126.1, 125.1, 121.1, 73.4, 70.8, 70.5, 63.3; HR-ESI-MS calcd for $C_{12}H_{14}N_2O_5Na$ ($[M+Na]^+$) 289.0800, found 289.0802.

2-((1*S*,2*R*,3*R*,4*R*)-1,2,3,4,5-pentahydroxypentyl)quinazolin-4(3*H*)-one (**3d**): White solid, yield 45%, m.p. 166.9 ~ 168.7 °C, $[\alpha]_D^{25}$ 16.5 (*c* 0.04, H_2O); 1H NMR (600 MHz, $DMSO-d_6$) δ : 11.50 (s, 1H), 8.11 (d, *J*=7.8 Hz, 1H), 7.80 (t, *J*=7.2 Hz, 1H), 7.65 (d, *J*=7.8 Hz, 1H), 7.49 (t, *J*=7.8 Hz, 1H), 4.71 ~ 4.37 (m, 5H), 4.09 (d, *J*=2.4 Hz, 1H), 3.58 (dd, *J*=10.8, 3.0 Hz, 1H), 3.53 ~ 3.50 (m, 1H), 3.47 (d, *J*=8.4, 1H), 3.39 (q, *J*=5.4 Hz, 1H); ^{13}C NMR (150 MHz, $DMSO-d_6$) δ : 161.2, 157.8, 148.4, 134.4, 126.9, 126.3, 125.8, 121.2, 73.3, 71.6, 71.4, 71.3, 63.2; HR-ESI-MS calcd for $C_{13}H_{16}N_2O_6Na$ ($[M+Na]^+$) 319.0906, found 319.0908.

2-((1*S*,2*R*,3*S*,4*R*)-1,2,3,4,5-pentahydroxypentyl)quinazolin-4(3*H*)-one (**3e**): White solid, yield 41%, m.p. 211.4 ~ 212.3 °C, $[\alpha]_D^{25}$ 22.6 (*c* 0.04, H_2O); 1H NMR (600 MHz, $DMSO-d_6$) δ : 11.21 (s, 1H), 8.11 (d, *J*=7.8 Hz, 1H), 7.79 (t, *J*=7.8 Hz, 1H), 7.64 (d, *J*=7.8 Hz, 1H), 7.48 (t, *J*=7.2 Hz, 1H), 5.52 (s, 1H), 4.84 (s, 1H), 4.74 (s, 1H), 4.52 (s, 1H), 4.28 (s, 2H), 3.88 (d, *J*=9.0 Hz, 1H), 3.74 (t, *J*=6.0 Hz, 1H), 3.59 (d, *J*=9.0 Hz, 1H), 3.47 ~ 3.41 (m, 3H); ^{13}C NMR (150 MHz, $DMSO-d_6$) δ : 161.1, 159.5, 148.5, 134.4, 126.6, 126.0, 125.8, 121.0, 72.6, 70.5, 69.6, 69.0, 63.1; HR-ESI-

MS calcd for $C_{13}H_{16}N_2O_6Na$ ($[M+Na]^+$) 319.0906, found 319.0903.

2-((1*R*,2*R*,3*R*,4*R*)-1,2,3,4,5-pentahydroxypentyl)quinazolin-4(3*H*)-one (**3f**): White solid, yield 59%, m.p. 209.9 ~ 211.5 °C, $[\alpha]_D^{25}$ -16.6 (*c* 0.04, H_2O); 1H NMR (600 MHz, $DMSO-d_6$) δ : 11.93 (s, 1H), 8.10 (d, *J*=7.8 Hz, 1H), 7.79 (t, *J*=7.8 Hz, 1H), 7.64 (d, *J*=7.8 Hz, 1H), 7.48 (t, *J*=7.2 Hz, 1H), 5.60 (d, *J*=4.8 Hz, 1H), 4.50 (s, 1H), 4.46 (q, *J*=6.0 Hz, 1H), 4.40 (s, 3H), 4.10 (t, *J*=6.6 Hz, 1H), 3.67 ~ 3.62 (m, 2H), 3.51 (s, 1H), 3.45 ~ 3.42 (m, 1H); ^{13}C NMR (150 MHz, $DMSO-d_6$) δ : 161.5, 158.9, 148.6, 134.2, 126.9, 126.1, 125.7, 121.4, 71.9, 71.2, 70.9, 69.5, 63.8; HR-ESI-MS calcd for $C_{13}H_{16}N_2O_6Na$ ($[M+Na]^+$) 319.0906, found 319.0907.

2-((1*S*,2*R*,3*R*,4*R*)-1,2,4,5-tetrahydroxy-3-(((2*S*,3*R*,4*S*,5*S*,6*R*)-3,4,5-trihydroxy-6-(hydroxymethyl)tetrahydro-2*H*-pyran-2-yl)oxy)pentyl)quinazolin-4(3*H*)-one (**3g**): White solid, yield 30%, m.p. 172.6 ~ 173.8 °C, $[\alpha]_D^{25}$ 114.3 (*c* 0.04, H_2O); 1H NMR (400 MHz, $DMSO-d_6$) δ : 11.31 (s, 1H), 8.11 (d, *J*=7.6 Hz, 1H), 7.81 (t, *J*=7.2 Hz, 1H), 7.64 (d, *J*=8.0 Hz, 1H), 7.49 (t, *J*=7.2 Hz, 1H), 5.75 (s, 1H), 5.56 (d, *J*=4.8 Hz, 1H), 5.06 (s, 1H), 4.95 ~ 4.92 (m, 3H), 4.69 (s, 1H), 4.60 (d, *J*=5.2 Hz, 1H), 4.54 (t, *J*=5.2 Hz, 1H), 4.48 (t, *J*=5.2 Hz, 1H), 4.14 (s, 1H), 3.72 (s, 2H), 3.65 ~ 3.61 (m, 3H), 3.50 ~ 3.39 (m, 3H), 3.27 (d, *J*=8.8 Hz, 1H), 3.13 ~ 3.08 (m, 1H); ^{13}C NMR (150 MHz, $DMSO-d_6$) δ : 161.0, 157.8, 148.4, 134.5, 126.8, 126.3, 125.8, 121.1, 100.8, 82.2, 73.4, 73.2, 73.1, 72.3, 72.2, 71.6, 69.8, 62.6, 60.6; HR-ESI-MS calcd for $C_{19}H_{26}N_2O_{11}Na$ ($[M+Na]^+$) 481.1434, found 481.1437.

6-Chloro-2-((1*R*,2*S*,3*R*)-1,2,3,4-tetrahydroxybutyl)-quinazolin-4(3*H*)-one (**3c-1**): White solid, yield 57%, m.p. 201.9 ~ 203.1 °C, $[\alpha]_D^{25}$ -14.5 (*c* 0.04, H_2O); 1H NMR (400 MHz, $DMSO-d_6$) δ : 11.41 ~ 11.36 (m, 1H), 8.02 (d, *J*=2.4 Hz, 1H), 7.81 (dd, *J*=8.8 Hz, 2.4 Hz, 1H), 7.65 (d, *J*=8.4 Hz, 1H), 5.53 (d, *J*=0.4 Hz, 1H), 4.77 ~ 4.68 (m, 3H), 4.39 (t, *J*=4.8 Hz, 1H), 3.72 ~ 3.54 (m, 3H), 3.43 (m, 1H); ^{13}C NMR (150 MHz, $DMSO-d_6$) δ : 160.1, 159.8, 147.4, 134.5, 130.3, 129.0, 124.8, 122.4, 73.4, 70.8, 70.6, 63.3; HR-ESI-MS calcd for $C_{12}H_{13}ClN_2O_5Na$ ($[M+Na]^+$) 323.0410, found 323.0407.

7-Chloro-2-((1*R*,2*S*,3*R*)-1,2,3,4-tetrahydroxybutyl)-quinazolin-4(3*H*)-one (**3c-2**): White solid, yield 50%, m.p. 168.9 ~ 169.8 °C, $[\alpha]_D^{25}$ -32.2 (*c* 0.04, H_2O); 1H NMR (600 MHz, $DMSO-d_6$) δ : 11.33 (s, 1H), 8.10 (d, *J*=8.4 Hz, 1H), 7.68 (s, 1H), 7.50 (d, *J*=8.4 Hz, 1H), 5.55 (s, 1H), 4.79 (s, 2H), 4.70 (s, 1H), 4.40 (s, 1H), 3.73 (d, *J*=7.2 Hz, 1H), 3.64 (dd, *J*=11.4, 2.4 Hz, 1H), 3.56 (s, 1H), 3.45 (dd, *J*=11.4, 6.0 Hz, 1H); ^{13}C NMR (150 MHz, $DMSO-d_6$) δ : 160.9, 160.5, 149.8, 139.0, 127.9, 126.3, 125.9, 120.0, 73.4, 70.8, 70.7, 63.3; HR-ESI-MS calcd for $C_{12}H_{13}ClN_2O_5Na$ ($[M+Na]^+$) 323.0410, found 323.0415.

7-Methyl-2-((1*R*,2*S*,3*R*)-1,2,3,4-tetrahydroxybutyl)-quinazolin-4(3*H*)-one (**3c-3**): Yellow solid, yield 45%, m.p. 211.0 ~ 212.0 °C, $[\alpha]_D^{25}$ -72.5 (*c* 0.04, H_2O); 1H NMR (400 MHz, $DMSO-d_6$) δ : 11.05 ~ 11.03 (m, 1H), 7.98 (d, *J*=8.0 Hz, 1H), 7.43 (s, 1H), 7.30 (d, *J*=8.0 Hz, 1H), 5.54

(d, $J=1.2$ Hz, 1H), 4.77~4.69 (m, 3H), 4.41 (t, $J=5.2$ Hz, 2H), 3.70~3.55 (m, 3H), 3.47~3.40 (m, 1H), 2.44 (s, 3H); ^{13}C NMR (150 MHz, DMSO- d_6) δ : 160.9, 159.1, 148.8, 144.8, 127.5, 126.4, 125.7, 118.7, 73.4, 70.8, 70.5, 63.3, 21.3; HR-ESI-MS calcd for $\text{C}_{13}\text{H}_{16}\text{N}_2\text{O}_5\text{Na}$ ($[\text{M}+\text{Na}]^+$) 303.0957, found 303.0962.

6-Chloro-2-((1*R*,2*R*,3*R*,4*R*)-1,2,3,4,5-pentahydroxypentyl)quinazolin-4(3*H*)-one (**3f-1**): White solid, yield 78%, m.p. 232.6~233.4 °C, $[\alpha]_{\text{D}}^{25} -13.5$ (c 0.04, H_2O), ^1H NMR (400 MHz, DMSO- d_6) δ : 8.03 (d, $J=2.4$ Hz, 1H), 7.79 (d, $J=2.4$ Hz, 1H), 7.66 (d, $J=8.8$ Hz, 1H), 4.50~4.33 (m, 3H), 4.06 (d, $J=8.8$ Hz, 1H), 3.65~3.60 (m, 2H), 3.51~3.42 (m, 2H); ^{13}C NMR (150 MHz, DMSO- d_6) δ : 160.8, 159.9, 147.2, 134.3, 130.3, 129.1, 124.7, 122.6, 71.9, 71.1, 70.9, 69.5, 63.7; HR-ESI-MS calcd for $\text{C}_{13}\text{H}_{15}\text{ClN}_2\text{O}_6\text{Na}$ ($[\text{M}+\text{Na}]^+$) 353.0516, found 353.0515.

7-Chloro-2-((1*R*,2*R*,3*R*,4*R*)-1,2,3,4,5-pentahydroxypentyl)quinazolin-4(3*H*)-one (**3f-2**): White solid, yield 57%, m.p. 182.9~183.8 °C, $[\alpha]_{\text{D}}^{25} -33.5$ (c 0.04, H_2O), ^1H NMR (400 MHz, DMSO- d_6) δ : 8.09 (d, $J=8.4$ Hz, 1H), 7.68 (d, $J=2.0$ Hz, 1H), 7.51 (dd, $J=8.4$ Hz, 2.0 Hz, 1H), 4.53~4.41 (m, 4H), 4.07 (d, $J=8.8$ Hz, 1H), 3.65~3.60 (m, 2H), 3.49 (d, $J=5.2$ Hz, 1H); ^{13}C NMR (150 MHz, DMSO- d_6) δ : 161.0, 160.7, 149.7, 138.8, 127.9, 126.4, 126.1, 120.2, 71.9, 71.1, 70.8, 69.4, 63.7; HR-ESI-MS calcd for $\text{C}_{13}\text{H}_{15}\text{ClN}_2\text{O}_6\text{Na}$ ($[\text{M}+\text{Na}]^+$) 353.0516, found 353.0518.

7-Methyl-2-((1*R*,2*R*,3*R*,4*R*)-1,2,3,4,5-pentahydroxypentyl)quinazolin-4(3*H*)-one (**3f-3**): Yellow solid, yield 80%, m.p. 191.8~192.7 °C, $[\alpha]_{\text{D}}^{25} -66.0$ (c 0.04, H_2O), ^1H NMR (400 MHz, DMSO- d_6) δ : 7.98 (d, $J=8.0$ Hz, 1H), 7.43 (s, 1H), 7.30 (d, $J=8.0$ Hz, 1H), 4.48~4.37 (m, 4H), 4.07 (d, $J=8.8$ Hz, 1H), 3.65~3.60 (m, 2H), 3.48 (d, $J=6.0$ Hz, 1H), 3.44~3.35 (m, 1H), 2.45 (s, 3H); ^{13}C NMR (150 MHz, DMSO- d_6) δ : 161.4, 158.9, 148.7, 144.6, 127.6, 126.6, 125.6, 119.0, 71.9, 71.1, 70.9, 69.5, 63.7, 21.3; HR-ESI-MS calcd for $\text{C}_{14}\text{H}_{18}\text{N}_2\text{O}_6\text{Na}$ ($[\text{M}+\text{Na}]^+$) 333.1062, found 333.1064.

6-Chloro-2-((1*S*,2*R*,3*R*,4*R*)-1,2,4,5-tetrahydroxy-3-(((2*R*,3*R*,4*S*,5*S*,6*R*)-3,4,5-trihydroxy-6-(hydroxymethyl)tetrahydro-2*H*-pyran-2-yl)oxy)pentyl)quinazolin-4(3*H*)-one (**3g-1**): White solid, yield 38%, m.p. 198.4~198.8 °C, $[\alpha]_{\text{D}}^{25} 97.6$ (c 0.04, H_2O), ^1H NMR (600 MHz, DMSO- d_6) δ : 11.55 (d, $J=0.8$ Hz, 1H), 8.04 (s, 1H), 7.83 (dd, $J=8.8$, 2.0 Hz, 1H), 7.67 (d, $J=8.4$ Hz, 1H), 5.77 (d, $J=0.8$ Hz, 1H), 5.55 (s, 1H), 5.06 (s, 1H), 4.92 (d, $J=2.8$ Hz, 3H), 4.69 (s, 1H), 4.60~4.46 (m, 3H), 4.13 (s, 1H), 3.72 (s, 2H), 3.63~3.61 (m, 3H), 3.49~3.38 (m, 3H), 3.28 (s, 1H), 3.13~3.07 (m, 1H); ^{13}C NMR (150 MHz, DMSO- d_6) δ : 160.1, 158.6, 147.1, 134.6, 130.5, 129.1, 124.8, 122.4, 100.8, 82.2, 73.3, 73.2, 73.0, 72.3, 72.2, 71.7, 69.8, 62.6, 60.6; HR-ESI-MS calcd for $\text{C}_{19}\text{H}_{25}\text{ClN}_2\text{O}_{11}\text{Na}$ ($[\text{M}+\text{Na}]^+$) 515.1044, found 515.1049.

7-Methyl-2-((1*S*,2*R*,3*R*,4*R*)-1,2,4,5-tetrahydroxy-3-(((2*R*,3*R*,4*S*,5*S*,6*R*)-3,4,5-trihydroxy-6-(hydroxymethyl)tetrahydro-2*H*-pyran-2-yl)oxy)pentyl)quinazolin-4(3*H*)-one (**3g-2**) and 6-iodo-7-methyl-2-((1*S*,2*R*,3*R*,4*R*)-1,2,4,5-

tetrahydroxy-3-(((2*R*,3*R*,4*S*,5*S*,6*R*)-3,4,5-trihydroxy-6-(hydroxymethyl)tetrahydro-2*H*-pyran-2-yl)oxy)pentyl)quinazolin-4(3*H*)-one (**3g-2-1**): White solid (mixture), total yields 35%, ^1H NMR (400 MHz, DMSO- d_6) δ : 11.34 (s, 0.5H), 11.14 (s, 0.8H), 8.46 (s, 0.5H), 8.00~7.99 (m, 0.8H), 7.62 (s, 0.5H), 7.45 (s, 0.8H), 7.33~7.31 (m, 0.8H), 5.71 (s, 1.5H), 5.49 (s, 1.5H), 5.03~4.85 (m, 6.0H), 4.68 (s, 1.7H), 4.55~4.42 (m, 4.7H), 4.14 (s, 1.4H), 3.73~3.63 (m, 7.8H), 3.54~3.37 (m, 5.0H), 3.18~3.11 (m, 2.0H), 2.45 (s, 3.0H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 162.5, 162.1, 159.3, 159.2, 149.2, 148.8, 146.6, 145.1, 135.1, 127.9, 127.3, 126.6, 126.2, 121.4, 119.2, 101.3, 97.4, 82.8, 82.7, 73.9, 73.8, 73.7, 73.6, 73.0, 72.9, 72.7, 72.6, 72.3, 70.0, 63.0, 61.0, 28.4, 21.9; HR-ESI-MS calcd for $\text{C}_{20}\text{H}_{29}\text{N}_2\text{O}_{11}$ ($[\text{M}+\text{H}]^+$), 473.1766, found 473.1772; and HR-ESI-MS calcd for $\text{C}_{20}\text{H}_{28}\text{IN}_2\text{O}_{11}$ ($[\text{M}+\text{H}]^+$) 599.0732, found 599.0738.

Supporting Information The NMR characterization data for compounds **A**, **B**, **3a**~**3g**, **3c-1**~**3c-3**, **3f-1**~**3f-3**, and **3g-1**~**3g-2** are available. Supplementary data associated with this article can be found, in the online version, at doi:

References

- [1] Alagarsamy, V.; Chitra, K.; Saravanan, G.; Raja Solomon, V.; Sulthana, M. T.; Narendhar, B. *Eur. J. Med. Chem.* **2018**, *151*, 628.
- [2] Dohle, W.; Jourdan, F. L.; Menchon, G.; Protta, A. E.; Foster, P. A.; Mannion, P.; Hamel, E.; Thomas, M. P.; Kasprzyk, P. G.; Ferrandis, E.; Steinmetz, M. O.; Leese, M. P.; Potter, B. V. L. *J. Med. Chem.* **2018**, *61*, 1031.
- [3] Bouley, R.; Ding, D. R.; Peng, Z. H.; Bastian, M.; Lastochkin, E.; Song, W.; Suckow, M. A.; Schroeder, V. A.; Wolter, W. R.; Mobashery, S.; Chang, M. *J. Med. Chem.* **2016**, *59*, 5011.
- [4] Leivers, A. L.; Tallant, M.; Brad Shotwell, J.; Dickerson, S.; Leivers, M. R.; McDonald, O. B.; Gobel, J.; Creech, K. L.; Strum, S. L.; Mathis, A.; Rogers, S.; Moore, C. B.; Botyanszki, J. *J. Med. Chem.* **2014**, *57*, 2091.
- [5] Kerru, N.; Singh-Pillay, A.; Awolade, P.; Singh, P. *Eur. J. Med. Chem.* **2018**, *152*, 436.
- [6] Wei, M. K.; Chai, W. M.; Wang, R.; Yang, Q.; Deng, Z. H.; Peng, Y. *Bioorg. Med. Chem.* **2017**, *25*, 1303.
- [7] Khan, I.; Ibrar, A.; Ahmed, W.; Saeed, A. *Eur. J. Med. Chem.* **2015**, *90*, 124.
- [8] Khan, I.; Ibrar, A.; Abbas, N.; Saeed, A. *Eur. J. Med. Chem.* **2014**, *76*, 193.
- [9] Khan, I.; Zaib, S.; Batool, S.; Abbas, N.; Ashraf, Z.; Iqbal, J.; Saeed, A. *Bioorg. Med. Chem.* **2016**, *24*, 2361.
- [10] Hou, J.; Kovacs, M. S.; Dhanvantari, S.; Luyt, L. G. *J. Med. Chem.* **2018**, *61*, 1261.
- [11] Wang, J. P.; Chang, Y. J.; Dong, X. Y.; Zhang, R. S.; Tang, Y.; Zhang, M. *Carbohydr. Res.* **2018**, *463*, 4.
- [12] Ma, C. C.; Zhang, C. M.; Tang, L. Q.; Liu, Z. P. *Eur. J. Med. Chem.* **2018**, *151*, 9.
- [13] Kim, N. Y.; Cheon, C. H. *Tetrahedron Lett.* **2014**, *55*, 2340.
- [14] Wang, Z. Z.; Tang, Y. *Tetrahedron* **2016**, *72*, 1330.
- [15] Yu, C. W.; Chang, P. T.; Hsin, L. W.; Chern, J. W. *J. Med. Chem.* **2013**, *56*, 6775.
- [16] Che, T.; Chen, S. B.; Tu, J. L.; Wang, B.; Wang, Y. Q.; Zhang, Y.; Wang, J.; Wang, Z. Q.; Zhang, Z. P.; Ou, T. M.; Zhao, Y.; Tan, J. H.; Huang, Z. S. *J. Med. Chem.* **2018**, *61*, 3436.
- [17] Saleeb, M.; Sundin, C.; Aglar, Ö.; Pinto, A. F.; Ebrahimi, M.; Forsberg, Å.; Schüller, H.; Eloffsson, M. *Eur. J. Med. Chem.* **2018**,

- 143, 568.
- [18] Sales, Z. S.; Mani, N. S.; Allison, B. D. *Tetrahedron Lett.* **2018**, 59, 1623.
- [19] Shen, G. S.; Zhou, H. F.; Sui, Y. B.; Liu, Q. X.; Zou, K. *Tetrahedron Lett.* **2016**, 57, 587.
- [20] Khan, K. M.; Saad, S. M.; Shaikh, N. N.; Hussain, S.; Fakhri, M. I.; Perveen, S.; Taha, M.; Choudhary, M. I. *Bioorg. Med. Chem.* **2014**, 22, 3449.
- [21] Ge, W.; Zhu, X.; Wei, Y. *RSC Adv.* **2013**, 3, 10817.
- [22] Yan, Y. Z.; Xu, Y.; Niu, B.; Xie, H. F.; Liu, Y. Q. *J. Org. Chem.* **2015**, 80, 5581.
- [23] Tian, X. H.; Song, L. N.; Li, E. T.; Wang, Q.; Yu, W. Q.; Chang, J. B. *RSC Adv.* **2015**, 76, 62194.
- [24] Sun, J. W.; Tao, T.; Xu, D.; Cao, H.; Kong, Q. G.; Wang, X. Y.; Liu, Y.; Zhao, J. L.; Wang, Y.; Pan, Y. *Tetrahedron Lett.* **2018**, 59, 2099.
- [25] Qi, X. X.; Song, Z. Z.; Gong, J. L.; Fang, Z. Y.; Wu, X. F. *Chin. Chem. Lett.* **2016**, 27, 21.
- [26] Kabri, Y.; Croze, M. D.; Terme, T.; Vanelle, P. *Eur. J. Org. Chem.* **2015**, 3806.
- [27] Adib, M.; Ansari, S.; Mohammadi, A.; Bijanzadeh, H. R. *Tetrahedron Lett.* **2010**, 51, 30.
- [28] Ren, Z. L.; Kong, H. H.; Lu, W. T.; Sun, M.; Ding, M. W. *Tetrahedron* **2018**, 74, 184.
- [29] Ramesh, K.; Karnakar, K.; Satish, G.; Anil Kumar, B. S. P.; Nageswar, Y. V. D. *Tetrahedron Lett.* **2012**, 53, 6936.
- [30] Wang, S. X.; Gao, M. Y.; Tan, G. H.; Ma, H. X.; Zhao, Y. Y.; Du, H. Y.; Wang, Z. S.; Chen, H.; Li, X. L. *Chin. J. Org. Chem.* **2017**, 37, 385 (in Chinese).
(王淑霞, 高梦颖, 谭官海, 马海霞, 赵莹莹, 杜鸿源, 王总帅, 陈华, 李小六, 有机化学, **2017**, 37, 385.)
- [31] Feng, Y. X.; Tan, G. H.; Zhou, L. K.; Wang, S. X.; Chen, H.; Li, X. L. *Chin. J. Org. Chem.* **2017**, 37, 429 (in Chinese).
(冯玉欣, 谭官海, 周利凯, 王淑霞, 陈华, 李小六, 有机化学, **2017**, 37, 429.)
- [32] Sun, J. J.; Zhou, L. K.; Tan, G. H.; Li, S.; Wang, S. X.; Chen, H.; Li, X. L. *Chin. J. Org. Chem.* **2017**, 37, 455 (in Chinese).
(孙佳婧, 周利凯, 谭官海, 李帅, 王淑霞, 陈华, 李小六, 有机化学, **2017**, 37, 455.)
- [33] Cheng, R.; Tang, L.; Guo, T.; Zhang, N.; Zhang, N. D.; Du, Y. *RSC Adv.* **2015**, 5, 26434.
- [34] Lin, C.; Lai, P. T.; Liao, K. S.; Hung, W. T.; Yang, W. B.; Fang, J. M. *J. Org. Chem.* **2008**, 73, 3848.
- [35] CCDC-1851215 (for **3a**) contain the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [36] Rambabu, D.; Kumar, S. K.; Sreenivas, B. Y.; Sandra, S.; Kandale, A.; Misra, P.; Basaveswara Rao, M. V.; Pal, M. *Tetrahedron Lett.* **2013**, 54, 495.

(Lu, Y.)