B₂pin₂ 参与的含硼酸酯的芳香胺和醛的一锅两步还原胺化反应

刘雪滢^a 刘振伟^a 郭圆圆^a 李敬亚^b
 邹大鹏^{*,a} 吴豫生^{*,b} 吴养洁^{*,a}
 (^a 郑州大学化学与分子工程学院 郑州 450052)
 (^b 郑州泰基鸿诺医药股份有限公司 郑州 450052)

摘要 官能团化的仲胺广泛存在于生物和天然产物, 在有机化学中具有重要的地位. 此外, 含硼酸频哪醇酯的芳香仲胺是一种重要的有机合成中间体. 研究了在氢氧化钠和乙醇的存在下, 联硼酸频哪醇酯作为还原剂, 含硼酸频哪醇酯的芳香伯胺和醛的一锅两步还原胺化反应. 通过对碱、温度、溶剂、时间、醛的用量等进行筛选, 在较优的反应条件下以中等到较好的收率得到了一系列相应的含硼酸频哪醇酯的芳香仲胺. 该方法产率较高, 具有良好的官能团容忍性, 并且合成的芳香仲胺中硼酸频哪醇酯取代基还可以进一步官能团化.

关键词 还原胺化; 含硼的芳香仲胺; 联硼酸频哪醇酯; 醛

One-Pot, Two-Step Reductive Amination of Boronate Ester Containing Aromatic Amines and Aldehydes Using B₂pin₂ as Reductant

Liu, Xueying^a Liu, Zhenwei^a Guo, Yuanyuan^a Li, Jingya^b
 Zou, Dapeng^{*,a} Wu, Yusheng^{*,b} Wu, Yangjie^{*,a}
 (^a College of Chemistry and Molecular Engineering, Zhengzhou University, Zhengzhou 450052)
 (^b Tetranov Biopharm, Limited Liability Company, Zhengzhou 450052)

Abstract The aromatic amine functionality occupies a very important role in organic chemistry due to its prominence in biological and naturally occurring molecules. In addition, the synthesized secondary aromatic amines with pendant boronate ester are versatile intermediates in several organic transformations. The one-pot, two-step reductive amination of boron-containing primary aromatic amines and aldehydes has been achieved in the presence of NaOH in ethanol using B₂pin₂ as reductant. After extensive screening of various reaction parameters, such as base, reaction temperature, solvent, reaction time and protective gas, a series of secondary aromatic amines with pendant boronate ester and various functional groups were obtained in moderate to good yields under the optimal reaction conditions. This system features generally high yields and broad functional group tolerance. The boronate ester substituent is a very good handle to be further functionalized.

Keywords reductive amination; boron-containing secondary aromatic amines; bis(pinacolato)diboron; aldehydes

1 Introduction

The aromatic amine functionality occupies a very important role in organic chemistry due to its prominence in biological and naturally occurring molecules.^[1] Reductive amination is one of the most efficient approaches to obtain highly substituted amines.^[2] Various strategies have been developed to achieve this transformation. Reductive amination has been generally accomplished by two protocols according to the nature of the reducing agent. (1) Met-

al-based catalytic hydrogenation: Both heterogeneous hydrogenation and homogeneous reductive amination use Ir,^[3] Rh,^[4] Fe,^[5] Ag,^[6] Ni^[7] and Pd^[8] complexes as catalysts. These methods have been applied in industry because of their economical and effective property.^[9] There are still some drawbacks, such as lack of chemoselectivity and not compatible with reducible groups.^[10] (2) Hydride reducing agents^[11-14] involved reductive amination.^[11] Borohydrides reagents have been employed in these methods, such as NaBH₃CN^[12] and NaBH(OAc)₃^[13] etc. The selectivity of

* Corresponding authors. E-mail: yusheng.wu@tetranovglobal.com; wyj@zzu.edu.cn; zdp@zzu.edu.cn

Received February 13, 2019; revised March 15, 2019; published online March 29, 2019.

Project supported by the National Natural Science Foundation of China (Nos. 21172200, 21702191).

国家自然科学基金(Nos. 21172200, 21702191)资助项目.

these reagents allows for a convenient access to reductive amination. Although various reductive amination systems have been well developed and applied, it is still necessary to develop economical and efficient methods for the highly chemoselective reduction amination under user-friendly conditions.

Bis(pinacolato)diboron (B_2pin_2), which is an air stable, green and inexpensive borylation reagent, has been frequently used in organic synthesis.^[15] The new reactivities of B_2pin_2 which are used for reduction reaction have been developed in recent years.^[16–21] A number of elegant works disclosed their abilities to reduce the C—C,^[17] C—O,^[18] N—O^[19] and S—O^[20] unsaturated bonds. Nevertheless, the reduction of C—N unsaturated bonds using B_2pin_2 as reductant is rarely reported. The only one research about Pd-catalyzed transfer hydrogenation of various N-hetero-aromatic compounds using B_2pin_2 as a mediator was reported by Song group.^[21] As a part of our ongoing efforts on the synthesis of organoborane reagents, as well as synthesis of various valuable compounds by using organoborane reagents.^[23] Herein, we report a novel one-pot, two-step reductive amination protocol for the synthesis of boron-containing secondary aromatic amines using B_2pin_2 as reductant. In addition, the synthesized secondary aromatic amines with pendant boronate ester are versatile intermediates in several organic transformations.^[24]

2 Results and discussion

The model reaction of 3-aminophenylboronic acid pinacol ester (**1a**, 1.0 equiv.) and benzaldehyde (**2a**, 1.0 equiv.) was chosen to optimize the reaction parameters and the results are summarized in Table 1. Firstly, the reaction was kept in *i*PrOH at 25 °C for 1 h under argon to obtain the intermediate (*E*)-*N*-benzylidene-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline. Secondly, B_2pin_2 (3.0 equiv.) and ^tBuOK (2.0 equiv.) were added to the reaction and the mixture was stirred for 4 h at 110 °C under argon (Table 1, Entry 1). The desired reductive amination product **3a** was observed in 32% yield in HPLC detection. Encouraged by this preliminary result, various reaction parameters, including bases, solvents and temperature were screened to optimize the reaction conditions. When a series of bases were studied, NaOH was found to be the best choice (Table 1, Entries 1~6). The reaction did not work without base. It indicated that the base is essential for the reaction (Table 1, Entry 7). When the solvent was changed to EtOH, the yield improved to 77%. Whereas, other solvents such as MeOH, ⁿPrOH, THF, dioxane and H₂O were found to be less effective for the reaction (Table 1, Entries 8~13). The effect of temperature was also studied, and the best yield was obtained under 110 °C (Table 1, Entries 14, 15). When the amount of B_2pin_2 decreased to 2.0 equiv., the yield of the desired product reduced to 66% (Table 1, Entry 16). Increasing the amount of **2a** to 1.2 equiv., the yield of the target product **3a** improved to 82% (Table 1, Entry 17). Under air atmosphere, the desired product **3a** was obtained

in 76% yield (Table 1, Entry 18). Lowering the amount of base, the yield of **3a** was slightly decreased (Table 1, Entries 19, 20). Consequently, the optimized reaction conditions for the one-pot, two-step reductive amination were determined as follows: firstly, 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**1a**, 1.0 equiv.) and benzaldehyde (**2a**, 1.2 equiv.) were kept in EtOH at 25 °C for 1 h under argon. Secondly, B_2pin_2 (3.0 equiv.) and NaOH (2.0 equiv.) were added to the reaction and the mixture was stirred for 4 h at 110 °C oil bath under argon.

Table 1 Optimization of the reaction conditions^a

Entry	Base	Solvent	Temp./°C	Yield ^b /%
1	^t BuOK	ⁱ PrOH	25/110	32
2	CH ₃ ONa	ⁱ PrOH	25/110	40
3	Cs ₂ CO ₃	ⁱ PrOH	25/110	20
4	KOH	ⁱ PrOH	25/110	50
5	NaOH	ⁱ PrOH	25/110	71
6	Et ₃ N	ⁱ PrOH	25/110	nr
7	—	ⁱ PrOH	25/110	nr
8	NaOH	MeOH	25/110	35
9	NaOH	ⁿ PrOH	25/110	47
10	NaOH	EtOH	25/110	77
11	NaOH	THF	25/110	4
12	NaOH	Dioxane	25/110	8
13	NaOH	H ₂ O	25/110	8
14	NaOH	EtOH	25/80	73
15	NaOH	EtOH	25/r.t.	nr
16 ^c	NaOH	EtOH	25/110	66
17 ^d	NaOH	EtOH	25/110	82/75 ^e
18 ^f	NaOH	EtOH	25/110	76
19 ^{d,g}	NaOH	EtOH	25/110	75
20 ^{d,h}	NaOH	EtOH	25/110	78

^a Reaction conditions: (1) **1a** (0.5 mmol) and **2a** (0.5 mmol), solvent (5.0 mL), r.t., 1 h. (2) B_2pin_2 (1.5 mmol), base (1.0 mmol), at 110 °C oil bath for 4 h, under Ar. ^b HPLC yield. ^c 2.0 equiv. of B_2pin_2 . ^d 1.2 equiv. of **2a**. ^e Isolated yield. ^f Under air. n.r. = no reaction. ^g 1.0 equiv. of base. ^h 1.5 equiv. of base.

With the optimal reaction conditions in hand, the scope of substrates were examined (Table 2). Initially, a variety of benzaldehyde derivatives were investigated. It was found that benzaldehydes with electron-donating groups worked comparatively well (**3ab**~**3ag**) except for (**3af**). Benzaldehyde possessing a strong electron-withdrawing group resulted in slightly lower yields (**3ah**~**3ai**). Trace amount of the desired product was observed when 4-nitrobenzaldehyde was employed (**3aj**). The halogen-containing benzaldehydes were also compatible and gave the corresponding products in good yields (**3ak**~**3as**). When benzaldehyde bearing a *meta*- or *para*-substitute boronate ester group was checked, the desired products **3at** and **3au** were obtained in 83% and 60% yields, respectively. Moreover,

the reaction seemed to be sensitive to steric effects. When acetophenone was used as substrate, only trace amount of target product was obtained (**3av**). Generally, arylamine proceeded well in the optimal reaction conditions (**3aa**~**3au**). In comparison, aliphatic amine gave the desired product in lower yields (**3aw**~**3ax**). The possible reason is about volatilization of acetaldehyde or propionaldehyde, as well as the difference of reactivities between arylamine and aliphatic amine.

To establish the scope of this reductive amination protocol, we subsequently tested the *ortho*- and *para*-boronate ester group substitute anilines (Table 3). The reaction of 4-aminophenylboronic acid pinacol ester and substituted benzaldehydes gave the desired products (**4a**, **4b**, **4d** and **4e**) in satisfactory yields except for **4c** due to volatilization of 4-(trifluoromethyl)benzaldehyde. Whereas, 2-amino-phenylboronic acid pinacol ester gave the target products in lower yields, indicated that the steric hindrance possibly had

a negative effect on this reaction (**4f**, **4g**). Only trace amount of the desired product was obtained when 2-aminopyridine-4-boronic acid pinacol ester was used as substrate (**4h**). The results indicated that the property of the aromatic ring in the substrates might influence the reaction.

Based on previous reports^[18d,25] and the data presented here, a reaction pathway was proposed as shown in Scheme 1. Initially, the aromatic amine **1** reacted with aldehyde **2** to generate imine **3** and water.^[25] Next, B—B bond in B₂pin chelated with H₂O in the presence of base, forming species **A** under the electronic field effect.^[18d] In the species **A**, the O atom of H₂O was inserted into the polarized B—B bond, forcing H-atom to leave as a hydride species. C=N bond in imine **3** chelated with the polarized B—B bond to form species **B**. Then, species **B** was attacked by the newly generated hydride to obtain compound **C**.^[18d] Finally, compound **C** was protonated to give the desired product **4**.

Table 2 Scope of substrate^{a,b}

^a Reaction conditions: (1) **1a** (0.5 mmol) and **2** (0.6 mmol), EtOH (5.0 mL), r.t., 1 h. (2) B₂pin₂ (1.5 mmol), NaOH (1.0 mmol), at 110 °C oil bath for 4 h, under Ar.

^b Determined by LC-MS analysis of the crude reaction mixture, with the isolated yield in parentheses. ^c The yield was determined by LC-MS analysis of the crude reaction mixture.

Table 3 Scope of substrate^{a,b}

^a Reaction conditions: (1) **1** (0.5 mmol) and **2** (0.6 mmol), EtOH (5.0 mL), r.t., 1 h. (2) B₂pin₂ (1.5 mmol), NaOH (1.0 mmol), at 110 °C oil bath for 4 h, under Ar.

^b Determined by LC-MS analysis of the crude reaction mixture, with the isolated yield in parentheses. ^c The yield was determined by LC-MS analysis of the crude reaction mixture. ^d Isolated yield.

Scheme 1 Plausible mechanism

To further investigate the importance and practical utility of this method, a gram-scale reaction of 3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**1a**, 1.1 g, 5.0 mmol) and benzaldehyde (**2a**, 0.6 g, 6.0 mmol) was performed under the optimal reaction conditions. As a result, the reductive amination reaction proceeded smoothly to afford *N*-benzyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3a**) in 70% isolated yield (Scheme 2).

Scheme 2 Scalability of the reductive amination reaction

3 Conclusions

In summary, a simple and highly efficient protocol for the synthesis of various boron-containing secondary aromatic amines was developed using commercially available B₂pin₂

as reductant. This system features generally high yields and broad functional group tolerance. The boronate ester substituent is a very good handle to be further functionalized. We believe that this method will be practical for further explorations in synthetic applications.

4 Experimental section

3.1 General information

All reactions were performed in dried glass reaction tube equipped with a magnetic stir bar under argon atmosphere. Unless otherwise noted, solvents and reagents were purchased from commercial sources and used as received. Flash column chromatography was performed using silica gel (60-Å pore size, 32~63 μm, standard grade). Analytical thin-layer chromatography was performed using glass plates pre-coated with 0.25 mm of 230~400 mesh silica gel impregnated with a fluorescent indicator (254 nm). Thin layer chromatography plates were visualized by exposure to ultraviolet light. Mass spectra and HPLC yields were measured on a LC-MSD-Trap-XCT instrument. NMR spectra were obtained on a Bruker AVANCE III systems using CDCl₃ or (CD₃)₂SO as solvent, TMS as internal standard substance at 400 MHz for ¹H NMR, 100

MHz for ^{13}C NMR, and 376 MHz for ^{19}F NMR. Melting points were measured with an XT4A microscopic apparatus. The high resolution mass spectrum was received via an Agilent Technologies 6540 UHD Accurate-mass Q-ToF LC/MS, with ESI as ion source. Preparative TLC was performed on silica gel plates and developed with ethyl acetate/hexane.

3.2 General experimental procedures

A glass reaction tube equipped with a magnetic stir bar was charged with compound **1** (0.5 mmol, 1.0 equiv.), compound **2** (0.6 mmol, 1.2 equiv.), EtOH (5 mL). The reaction mixture was stirred at 25 °C under argon atmosphere for 1 h. B_2pin_2 (1.5 mmol, 3.0 equiv.), and NaOH (1.0 mmol, 2.0 equiv.) were added and then reacted at 110 °C oil bath under argon atmosphere for 4 h. The reaction progress was monitored by TLC. The reaction mixture was concentrated *in vacuo*, and then extracted with ethyl acetate three times. The combined organic layers were washed with brine, dried over anhydrous MgSO_4 and concentrated under reduced pressure. The residue was purified by silica gel flash chromatography to produce the desired product.

N-Benzyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3aa**):^[26] 115.9 mg, yield 75%. White solid, m.p. 87~88 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.38~7.27 (m, 5H), 7.19~7.14 (m, 3H), 6.73~6.70 (m, 1H), 4.35 (s, 2H), 4.00 (s, 1H), 1.33 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 147.6, 139.5, 128.8, 128.6, 127.6, 127.2, 124.0, 119.3, 115.5, 83.7, 48.4, 24.9; GC-MS (EI) m/z : 309.1 $[\text{M}]^+$.

N-(2-Methylbenzyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3ab**): 134.0 mg, yield 83%. White solid, m.p. 89~90 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.33~7.31 (m, 1H), 7.21~7.16 (m, 6H), 6.74 (s, 1H), 4.29 (s, 2H), 2.36 (s, 3H), 1.34 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 147.7, 137.1, 136.5, 130.4, 128.8, 128.5, 127.5, 126.2, 124.0, 119.0, 115.4, 83.7, 46.5, 24.9, 19.0. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{27}\text{O}_2\text{BN}$ $[\text{M}+\text{H}]^+$ 324.2129, found: 324.2134.

N-(3-Methylbenzyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3ac**): 130.8 mg, yield 81%. White solid, m.p. 88~89 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.25~7.21 (m, 1H), 7.20~7.14 (m, 5H), 7.10~7.08 (m, 1H), 6.74~6.71 (m, 1H), 4.31 (s, 2H), 4.00 (s, 1H), 2.35 (s, 3H), 1.34 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 147.7, 139.4, 138.3, 128.8, 128.6, 128.4, 128.0, 124.7, 124.0, 119.3, 115.5, 83.7, 48.4, 24.9, 21.5. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{27}\text{O}_2\text{BN}$ $[\text{M}+\text{H}]^+$ 324.2129, found 324.2134.

N-(4-Methylbenzyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3ad**): 138.9 mg, yield 86%. White solid, m.p. 85~86 °C; ^1H NMR (400 MHz, DMSO) δ : 7.23~7.21 (m, 2H), 7.12~7.10 (m, 2H), 7.04~7.01 (m, 1H), 6.97~6.96 (m, 1H), 6.84~6.82 (m, 1H), 6.63~6.61 (m, 1H), 6.20~6.17 (m, 1H), 4.21~4.20 (m, 2H), 2.26 (s, 3H), 1.26 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 147.6, 136.9, 136.4, 129.3, 128.8, 127.6, 124.0, 119.3, 115.5, 83.7, 48.1, 24.9, 21.1; HRMS (ESI) calcd for

$\text{C}_{20}\text{H}_{27}\text{O}_2\text{BN}$ $[\text{M}+\text{H}]^+$ 324.2129, found 324.2136.

N-(4-Methoxybenzyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3ae**): 135.6 mg, yield 80%. White solid, m.p. 97~98 °C (lit.^[26] 105 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.29~7.27 (m, 2H), 7.21~7.16 (m, 2H), 7.13~7.12 (m, 1H), 6.89~6.85 (m, 2H), 6.74~6.71 (m, 1H), 4.27 (s, 2H), 3.80 (s, 3H), 1.33 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 159.0, 147.7, 131.6, 129.0, 128.9, 124.1, 119.4, 115.7, 114.1, 83.8, 55.4, 48.0, 25.0; LC-MS (ESI) m/z : 340.2 $[\text{M}+\text{H}]^+$.

2-(((3-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-phenyl)amino)methyl)phenol (**3af**): 81.3 mg, yield 50%. White solid, m.p. 90~91 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.40~7.38 (m, 1H), 7.35~7.34 (m, 1H), 7.29~7.27 (m, 1H), 7.25~7.20 (m, 1H), 7.16~7.14 (m, 1H), 6.98~6.96 (m, 1H), 6.93~6.91 (m, 1H), 6.88~6.84 (m, 1H), 4.43 (s, 2H), 1.34 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 156.7, 146.5, 129.2, 128.9, 127.2, 123.0, 122.2, 122.1, 120.0, 118.7, 116.6, 83.9, 48.7, 24.9; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{25}\text{O}_3\text{BN}$ $[\text{M}+\text{H}]^+$ 326.1922, found 326.1933.

N,N-Dimethyl-4-(((3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)amino)methyl)aniline (**3ag**): 114.4 mg, yield 65%. White solid, m.p. 112~114 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.25~7.22 (m, 2H), 7.21~7.15 (m, 2H), 7.13~7.12 (m, 1H), 6.74~6.72 (m, 3H), 4.22 (s, 2H), 3.84 (s, 1H), 2.94 (s, 6H), 1.34 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 150.0, 147.9, 128.8, 128.7, 127.2, 123.8, 119.1, 115.6, 112.8, 83.7, 48.0, 40.8, 24.9; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{30}\text{O}_2\text{BN}_2$ $[\text{M}+\text{H}]^+$ 353.2395, found 353.2394.

3-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-*N*-(4-(trifluoromethyl)benzyl)aniline (**3ah**): 43.4 mg, yield 23%. White solid, m.p. 71~72 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.59~7.57 (m, 2H), 7.48~7.46 (m, 2H), 7.21~7.16 (m, 2H), 7.14~7.13 (m, 1H), 6.69~6.66 (m, 1H), 4.43 (s, 2H), 1.33 (s, 12H); ^{19}F NMR (376 MHz, CDCl_3) δ : -62.4 (s); ^{13}C NMR (100 MHz, CDCl_3) δ : 147.1, 143.8, 129.5 (q, $J=32.0$ Hz), 128.8, 127.7, 125.5 (q, $J=3.8$ Hz), 124.5, 124.3 (q, $J=270.3$ Hz), 119.4, 115.4, 83.7, 47.8, 24.9; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{24}\text{O}_2\text{BF}_3\text{N}$ $[\text{M}+\text{H}]^+$ 378.1847, found 378.1850.

4-(((3-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-phenyl)amino)methyl)benzonitrile (**3ai**): 93.5 mg, yield 56%. White solid, m.p. 128~129 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.63~7.61 (m, 2H), 7.48~7.46 (m, 2H), 7.21~7.15 (m, 2H), 7.11~7.10 (m, 1H), 6.64~6.61 (m, 1H), 4.44 (s, 2H), 4.13 (s, 1H), 1.33 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 146.8, 145.4, 132.4, 128.9, 127.8, 124.6, 119.4, 118.9, 115.3, 110.9, 83.8, 47.8, 24.9; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{24}\text{O}_2\text{BN}_2$ $[\text{M}+\text{H}]^+$ 335.1926, found 335.1931.

N-(2-Bromobenzyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3ak**): 150.9 mg, yield 78%. White solid, m.p. 133~134 °C; ^1H NMR (400 MHz, DMSO) δ : 7.63~7.61 (m, 1H), 7.37~7.30 (m, 2H), 7.21~7.17 (m, 1H), 7.08~7.04 (m, 1H), 6.98~6.97 (m, 1H), 6.88~6.86 (m, 1H), 6.58~6.55 (m, 1H), 6.34~6.31

(m, 1H), 4.29~4.27 (m, 2H), 1.26 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 147.1, 138.3, 132.8, 129.4, 128.8, 128.7, 127.6, 124.2, 123.4, 119.7, 115.3, 83.7, 48.4, 24.9; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{24}\text{O}_2\text{BBrN}$ $[\text{M} + \text{H}]^+$ 388.1078, found 388.1087.

N-(3-Bromobenzyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3al**): 160.6 mg, yield 83%. White solid, m.p. 73~74 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.52 (s, 1H), 7.40~7.39 (m, 1H), 7.30~7.28 (m, 1H), 7.21~7.18 (m, 3H), 7.13~7.12 (m, 1H), 6.70~6.67 (m, 1H), 4.34 (s, 2H), 4.10 (s, 1H), 1.34 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 147.2, 142.0, 130.5, 130.3, 130.2, 128.8, 126.0, 124.3, 122.7, 119.4, 115.4, 83.7, 47.7, 24.9; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{24}\text{O}_2\text{BBrN}$ $[\text{M} + \text{H}]^+$ 388.1078, found 388.1087.

N-(4-Bromobenzyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3am**): 90.9 mg, yield 47%. White solid, m.p. 92~93 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.46~7.44 (m, 2H), 7.24~7.22 (m, 2H), 7.19~7.17 (m, 2H), 7.12~7.11 (m, 1H), 6.70~6.65 (m, 1H), 4.31 (s, 2H), 1.33 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 147.2, 138.6, 131.7, 129.2, 128.8, 124.3, 120.9, 119.4, 115.5, 83.7, 47.7, 24.9; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{24}\text{O}_2\text{BBrN}$ $[\text{M} + \text{H}]^+$ 388.1078, found 388.1089.

N-(2-Fluorobenzyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3an**): 137.3 mg, yield 84%. White solid, m.p. 67~68 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.40~7.36 (m, 1H), 7.25~7.21 (m, 1H), 7.20~7.16 (m, 3H), 7.10~7.02 (m, 2H), 6.77~6.74 (m, 1H), 4.42 (s, 2H), 1.33 (s, 12H); ^{19}F NMR (376 MHz, CDCl_3) δ : -119.0 (s). ^{13}C NMR (100 MHz, CDCl_3) δ : 161.0 (d, $J=244.4$ Hz), 147.2, 129.5 (d, $J=4.4$ Hz), 128.9, 128.8, 126.3 (d, $J=14.6$ Hz), 124.3, 124.2 (d, $J=3.6$ Hz), 119.6, 115.5, 115.3 (d, $J=21.2$ Hz), 83.7, 41.9 (d, $J=4.2$ Hz), 24.9. HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{24}\text{O}_2\text{BFN}$ $[\text{M} + \text{H}]^+$ 328.1879, found 328.1885.

N-(3-Fluorobenzyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3ao**): 145.5 mg, yield 89%. White solid, m.p. 79~80 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.31~7.27 (m, 1H), 7.20~7.18 (m, 2H), 7.15~7.13 (m, 2H), 7.09~7.06 (m, 1H), 6.97~6.92 (m, 1H), 6.71~6.69 (m, 1H), 4.36 (s, 2H), 1.33 (s, 12H); ^{19}F NMR (376 MHz, CDCl_3) δ : -113.0 (s); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.1 (d, $J=244.6$ Hz), 147.2, 142.3 (d, $J=6.7$ Hz), 130.1 (d, $J=8.2$ Hz), 128.8, 124.3, 122.9 (d, $J=2.7$ Hz), 119.4, 115.5, 114.3 (d, $J=22.5$ Hz), 114.1 (d, $J=21.4$ Hz), 83.7, 47.8 (d, $J=1.4$ Hz), 24.9; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{24}\text{O}_2\text{BFN}$ $[\text{M} + \text{H}]^+$ 328.1879, found 328.1886.

N-(2-Chlorobenzyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3ap**): 128.6 mg, yield 75%. White solid, m.p. 115~116 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.42~7.35 (m, 2H), 7.22~7.17 (m, 4H), 7.15~7.14 (m, 1H), 6.70~6.67 (m, 1H), 4.46 (s, 2H), 1.34 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 147.1, 136.7, 133.4, 129.5, 129.2, 128.8, 128.4, 126.9, 124.3, 119.7, 115.3, 83.7, 45.9, 24.9; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{24}\text{O}_2\text{BClN}$ $[\text{M} + \text{H}]^+$ 344.1583, found 344.1590.

N-(4-Chlorobenzyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3aq**): 137.2 mg, yield 80%. White solid, m.p. 90~91 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.29 (s, 4H), 7.21~7.16 (m, 2H), 7.14~7.13 (m, 1H), 6.72~6.69 (m, 1H), 4.33 (s, 2H), 1.33 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 147.3, 138.0, 132.9, 128.8, 128.7, 124.3, 119.4, 119.3, 115.5, 83.7, 47.6, 24.9; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{24}\text{O}_2\text{BClN}$ $[\text{M} + \text{H}]^+$ 344.1583, found 344.1593.

N-(3-Chloro-4-fluorobenzyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3ar**): 169.7 mg, yield 94%. White solid, m.p. 125~126 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.42~7.40 (m, 1H), 7.24~7.16 (m, 3H), 7.12~7.07 (m, 2H), 6.68~6.65 (m, 1H), 4.31 (s, 2H), 1.33 (s, 12H); ^{19}F NMR (376 MHz, CDCl_3) δ : -117.9 (s); ^{13}C NMR (100 MHz, CDCl_3) δ : 157.2 (d, $J=246.5$ Hz), 146.9, 136.6 (d, $J=3.7$ Hz), 129.5, 128.8, 127.0 (d, $J=7.0$ Hz), 124.5, 121.1 (d, $J=17.9$ Hz), 119.4, 116.6 (d, $J=21.0$ Hz), 115.5, 83.7, 47.2, 24.9; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{23}\text{O}_2\text{BClFN}$ $[\text{M} + \text{H}]^+$ 362.1489, found 362.1498.

N-(4-Chloro-3-(trifluoromethyl)benzyl)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3as**): 168.5 mg, yield 82%. White solid, m.p. 91~92 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.69 (s, 1H), 7.49~7.44 (m, 2H), 7.21~7.16 (m, 2H), 7.12~7.11 (m, 1H), 6.67~6.64 (m, 1H), 4.39 (s, 2H), 4.17 (s, 1H), 1.33 (s, 12H); ^{19}F NMR (376 MHz, CDCl_3) δ : -62.6 (s); ^{13}C NMR (100 MHz, CDCl_3) δ : 146.8, 139.0, 131.7, 130.8, 128.8, 128.5 (q, $J=31.1$ Hz), 126.5 (q, $J=5.1$ Hz), 124.6, 122.8 (q, $J=27.1$ Hz), 119.4, 115.5, 83.8, 47.4, 24.9. HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{24}\text{O}_2\text{BF}_3\text{ClN}$ $[\text{M} + \text{H}]^+$ 412.1457, found 412.1465.

3-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-*N*-(3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzyl)aniline (**3at**): 130.5 mg, yield 60%. White solid, m.p. 165~166 °C (lit.^[26] 166 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.80 (s, 1H), 7.73~7.71 (m, 1H), 7.48~7.46 (m, 1H), 7.35~7.32 (m, 1H), 7.20~7.14 (m, 3H), 6.73~6.71 (m, 1H), 4.34 (s, 2H), 4.01 (s, 1H), 1.35 (s, 12H), 1.33 (s, 12H). ^{13}C NMR (100 MHz, CDCl_3) δ : 147.6, 138.7, 134.0, 133.7, 130.6, 128.7, 128.0, 123.9, 119.3, 115.5, 83.9, 83.7, 48.3, 24.9, 24.8; MS (ESI) m/z : 436.4 $[\text{M} + \text{H}]^+$.

3-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-*N*-(4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzyl)aniline (**3au**): 180.5 mg, yield 83%. White solid, m.p. 128~129 °C (lit.^[26] 131 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.79~7.77 (m, 2H), 7.38~7.36 (m, 2H), 7.18~7.13 (m, 3H), 6.71~6.68 (m, 1H), 4.37 (s, 2H), 4.02 (s, 1H), 1.34 (s, 12H), 1.33 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 147.4, 142.7, 135.1, 128.8, 126.9, 124.2, 119.4, 115.5, 83.8, 83.7, 48.5, 24.9; MS (ESI) m/z : 436.4 $[\text{M} + \text{H}]^+$.

N-Propyl-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**3ax**): 11.7 mg, yield 9%. Brown oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.23~7.16 (m, 2H), 7.08 (d, $J=2.3$ Hz, 1H), 6.74~6.72 (m, 1H), 3.30 (s, 1H), 3.11 (t, $J=7.0$ Hz, 2H), 1.64 (q, $J=7.3$ Hz, 2H), 1.35 (s, 12H), 1.00 (t,

$J=7.4$ Hz, 3 H); ^{13}C NMR (100 MHz, CDCl_3) δ : 147.9, 128.7, 123.7, 119.1, 115.6, 83.6, 45.9, 24.9, 22.8, 11.7. HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{25}\text{O}_2\text{BN}$ $[\text{M}+\text{H}]^+$ 262.1973, found 262.1985.

N-(4-Methylbenzyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**4a**): 121.1 mg, yield 75%. White solid, m.p. 101~102 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.64 (d, $J=8.5$ Hz, 2H), 7.25 (d, $J=8.0$ Hz, 2H), 7.15 (d, $J=7.9$ Hz, 2H), 6.62 (d, $J=8.5$ Hz, 2H), 4.31 (s, 2H), 2.34 (s, 3H), 1.31 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 150.7, 137.0, 136.4, 136.0, 129.3, 127.5, 111.9, 83.2, 47.5, 24.9, 21.1; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{27}\text{O}_2\text{BN}$ $[\text{M}+\text{H}]^+$ 324.2129, found 324.2137.

N-(4-Methoxybenzyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**4b**): 149.2 mg, yield 88%. White solid, m.p. 119~120 °C (lit.^[27] 116~118 °C); ^1H NMR (400 MHz, CDCl_3) δ : 7.64 (d, $J=8.5$ Hz, 2H), 7.28 (d, $J=7.4$ Hz, 2H), 6.89~6.85 (m, 2H), 6.62 (d, $J=8.5$ Hz, 2H), 4.28 (s, 2H), 3.80 (s, 3H), 1.31 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 158.9, 150.7, 136.4, 131.0, 128.8, 114.1, 111.9, 83.2, 55.3, 47.2, 24.9; LC-MS (ESI) m/z : 340.2 $[\text{M}+\text{H}]^+$.

4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)-*N*-(4-(trifluoromethyl)benzyl)aniline (**4c**): 41.5 mg, yield 22%. White solid, m.p. 153~154 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.64 (d, $J=8.5$ Hz, 2H), 7.59 (d, $J=8.1$ Hz, 2H), 7.47 (d, $J=8.1$ Hz, 2H), 6.59 (d, $J=8.6$ Hz, 2H), 4.45 (s, 2H), 4.38 (s, 1H), 1.31 (s, 12H); ^{19}F NMR (376 MHz, CDCl_3) δ : -62.4 (s); ^{13}C NMR (100 MHz, CDCl_3) δ : 150.1, 143.4, 136.4, 129.5 (q, $J=32.2$ Hz), 127.4, 125.6 (q, $J=3.7$ Hz), 124.2 (q, $J=270.4$ Hz), 112.0, 83.3, 47.2, 24.8; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{24}\text{O}_2\text{BF}_3\text{N}$ $[\text{M}+\text{H}]^+$ 378.1847, found 378.1850.

4-(((4-(4,4,5,5-Tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)amino)methyl)benzonitrile (**4d**): 116.9 mg, yield 70%. White solid, m.p. 157~158 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.63~7.60 (m, 4H), 7.46~7.44 (m, 2H), 6.56~6.54 (m, 2H), 4.47~4.46 (m, 2H), 4.39 (s, 1H), 1.31 (s, 12H); ^{13}C NMR (100 MHz, CDCl_3) δ : 149.8, 145.0, 136.4, 132.5, 127.7, 118.8, 112.0, 110.0, 83.3, 47.2, 24.8; HRMS (ESI) calcd for $\text{C}_{20}\text{H}_{24}\text{O}_2\text{BN}_2$ $[\text{M}+\text{H}]^+$ 335.1925, found 335.1936.

N-(3-Fluorobenzyl)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)aniline (**4e**): 130.8 mg, yield 80%. White solid, m.p. 99~100 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.63~7.61 (m, 2H), 7.24~7.19 (m, 1H), 7.07~7.05 (m, 1H), 7.01~6.99 (m, 1H), 6.93~6.88 (m, 1H), 6.55~6.53 (m, 2H), 4.30 (s, 2H), 1.28 (s, 12H); ^{19}F NMR (376 MHz, CDCl_3) δ : -112.9 (s); ^{13}C NMR (100 MHz, CDCl_3) δ : 163.2 (d, $J=244.6$ Hz), 150.5, 142.2 (d, $J=6.6$ Hz), 136.5, 130.2 (d, $J=8.2$ Hz), 122.8 (d, $J=2.5$ Hz), 114.2, 114.0, 112.1, 83.4, 47.1 (d, $J=1.5$ Hz), 24.9; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{24}\text{O}_2\text{BFN}$ $[\text{M}+\text{H}]^+$ 328.1879, found 328.1883.

Supporting Information Optimization of the reaction conditions. The LC-MS data of the products **3aw**, **4f**~**4g**. Copies of ^1H NMR, ^{13}C NMR and ^{19}F NMR spectra of the

products. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- Pham, P. D.; Bertus, P.; Legoupy, S. *Chem. Commun.* **2009**, 6207.
- For literature pertaining to the origins and definition of reductive amination, see: (a) Emerson, W. S. *Org. React.* **1948**, *4*, 174. (b) Moore, M. L. *Org. React.* **1949**, *5*, 301. (c) Nugenta, T. C.; El-Shazly, M. *Adv. Synth. Catal.* **2010**, *352*, 753.
- Chi, Y.; Zhou, Y.; Zhang, X. *J. Org. Chem.* **2003**, *68*, 4120.
- (a) Skucas, E.; Kong, J.; Krische, M. J. *J. Am. Chem. Soc.* **2007**, *129*, 7242. (b) Park, J. W.; Chung, Y. K. *ACS Catal.* **2015**, *5*, 4846.
- Pagnoux-Ozherelyeva, A.; Pannetier, N.; Mbaye, M. D.; Gaillard, S.; Renaud, J.-L. *Angew. Chem., Int. Ed.* **2012**, *51*, 4976.
- Nasrollahzadeh, M. *New J. Chem.* **2014**, *38*, 5544.
- Huang, L.; Wang, Z.; Geng, L.; Chen, R.; Xing, W.; Wang, Y.; Huang, J. *RSC Adv.* **2015**, *5*, 56936.
- Gao, G.; Sun, P.; Li, Y.; Wang, F.; Zhao, Z.; Qin, Y.; Li, F. *ACS Catal.* **2017**, *7*, 4927.
- Abdel-Magid, A. F.; Carson, K. G.; Harris, B. D.; Maryanoff, C. A.; Shah, R. D. *J. Org. Chem.* **1996**, *61*, 3849.
- Roe, A.; Montgomery, J. A. *J. Am. Chem. Soc.* **1953**, *75*, 910.
- Tripathi, R. P.; Verma, S. S.; Pandey, J.; Tiwari, V. K. *Curr. Org. Chem.* **2008**, *12*, 1093.
- (a) Borch, R. F.; Bernstein, M. D.; Durst, H. D. *J. Am. Chem. Soc.* **1971**, *93*, 2897. (b) Borch, R. F.; Hassid, A. I. *J. Org. Chem.* **1972**, *37*, 1673. (c) Marchini, P.; Liso, G.; Reho, A.; Liberatore, F.; Moracci, F. M. *J. Org. Chem.* **1975**, *40*, 3453. (d) Lane, C. F. *Synthesis* **1975**, 135.
- (a) Abdel-Magid, A. F.; Maryanoff, C. A.; Carson, K. G. *Tetrahedron Lett.* **1990**, *31*, 5595. (b) Kim, H. O.; Carrol, B.; Lee, M. S. *Synth. Commun.* **1997**, *27*, 2505. (c) Tarasevich, V. A.; Kozlov, N. G. *Russ. Chem. Rev.* **1999**, *68*, 55.
- Borch, R. F.; Durst, H. D. *J. Am. Chem. Soc.* **1969**, *91*, 3996.
- (a) Ros, A.; Fernandez, R.; Lassaletta, J. M. *Chem. Soc. Rev.* **2014**, *43*, 3229. (b) Hartwig, J. F. *Acc. Chem. Res.* **2012**, *45*, 864. (c) Hartwig, J. F. *Chem. Soc. Rev.* **2011**, *40*, 1992. (d) Mkhali, I. A. I.; Barnard, J. H.; Marder, T. B.; Murphy, J. M.; Hartwig, J. F. *Chem. Rev.* **2010**, *110*, 890.
- (a) Xu, H.; Zhao, C.; Qian, Q.; Deng, W.; Gong, H. *Chem. Sci.* **2013**, *4*, 4022. (b) Yu, X.; Wang, S.; Xu, H.; Gong, H. *Org. Lett.* **2011**, *13*, 2138. (c) Liang, Z.; Xue, W.; Lin, K.; Gong, H. *Org. Lett.* **2014**, *16*, 5620. (d) Zhang, G.; Xie, Y.; Wang, Z.; Liu, Y.; Huang, H. *Chem. Commun.* **2015**, *51*, 1850. (e) Ke, M.; Song, Q. *J. Org. Chem.* **2016**, *81*, 3654. (f) Doi, R.; Ohashi, M.; Ogoshi, S. *Angew. Chem., Int. Ed.* **2016**, *55*, 341. (g) Ke, M.; Song, Q. *Chem. Commun.* **2017**, *53*, 2222. (h) Chen, Z.; Wang, X. *Org. Biomol. Chem.* **2017**, *15*, 5790. (i) Ke, M.; Song, Q. *Adv. Synth. Catal.* **2017**, *359*, 384. (j) Lu, X.; Wang, Y.; Zhang, B.; Pi, J.; Wang, X.; Gong, T.; Xiao, B.; Fu, Y. *J. Am. Chem. Soc.* **2017**, *139*, 12632. (k) Kuang, Z.; Li, B.; Song, Q. *Chem. Commun.* **2018**, *54*, 34.
- (a) Ojha, D. P.; Gadde, K.; Prabhu, K. R. *Org. Lett.* **2016**, *18*, 5062. (b) Ding, W.; Song, Q. *Org. Chem. Front.* **2016**, *3*, 14. (c) Wang, Q.; Yang, J.; Fang, D.; Ren, J.; Dong, B.; Zhou, B.; Zeng, B. *Tetrahedron Lett.* **2016**, *57*, 2587.
- (a) Laitar, D. S.; Müller, P.; Sadighi, J. P. *J. Am. Chem. Soc.* **2005**, *127*, 17196. (b) Bae, S.; Lakshman, M. K. *J. Org. Chem.* **2008**, *73*, 1311.

- (c) Kokatla, H. P.; Thomson, P. F.; Bae, S.; Doddi, V. R.; Lakshman, M. K. *J. Org. Chem.* **2011**, *76*, 7842.
- (d) Xuan, Q.; Zhao, C.; Song, Q. *Org. Biomol. Chem.* **2017**, *15*, 5140.
- [19] (a) Lu, H.; Geng, Z.; Li, J.; Zou, D.; Wu, Y. S.; Wu, Y. *J. Org. Lett.* **2016**, *18*, 2774.
- (b) Yang, K.; Zhou, F.; Kuang, Z.; Gao, G.; Driver, T. *Org. Lett.* **2016**, *18*, 4088.
- [20] Enthaler, S. *Catal. Lett.* **2012**, *142*, 1306.
- [21] Xuan, Q.; Song, Q. *Org. Lett.* **2016**, *18*, 4250.
- [22] (a) Lu, H.; Wang, S.; Li, J.; Zou, D.; Wu, Y. S.; Wu, Y. *J. Tetrahedron Lett.* **2017**, *58*, 839.
- (b) Zhi, W.; Li, J.; Zou, D.; Wu, Y.; Wu, Y. *J. Org. Chem.* **2017**, *82*, 12286.
- (c) Ren, X.; Han, S.; Gao, X.; Li, J.; Zou, D.; Wu, Y.; Wu, Y. *Tetrahedron Lett.* **2018**, *59*, 1065.
- (d) Zhi, W.; Li, J.; Zou, D.; Wu, Y.; Wu, Y. *Tetrahedron Lett.* **2018**, *59*, 537.
- (e) Zhi, W.; Li, J.; Zou, D.; Wu, Y.; Wu, Y. *Tetrahedron Lett.* **2018**, *59*, 2736.
- [23] (a) Geng, Z.; Zhang, Y.; Zheng, L.; Li, J.; Zou, D.; Wu, Y. J.; Wu, Y. S. *Tetrahedron Lett.* **2016**, *57*, 3063.
- (b) Zhang, Y.; Geng, Z.; Li, J.; Zou, D.; Wu, Y.; Wu, Y. *Adv. Synth. Catal.* **2017**, *359*, 390.
- (c) Zhu, M.; Qiu, Z.; Zhang, Y.; Du, H.; Li, J.; Zou, D.; Wu, Y.; Wu, Y. *Tetrahedron Lett.* **2017**, *58*, 2255.
- (d) Zhu, M.; Du, H.; Li, J.; Zou, D.; Wu, Y.; Wu, Y. *Tetrahedron Lett.* **2018**, *59*, 1352.
- [24] (a) Menet, C. J. M.; Blanc, J.; Hodges, A. J.; Burli, R. W.; Breccia, P.; Blackaby, W. P.; Van Rompaey, L. J. C.; Fletcher, S. R. *WO 2010/010184*, **2010**.
- (b) Liu, B.; Huang, J.; Zheng, C.; Zhang, Y.; Ouyang, L.; Mao, H.; Nie, B.; Xu, J.; Chen, H. *CN 2015/10402418*, **2015**.
- [25] Chen, Y.; Liu, S.; Cui, P.; Zhang, J.; Liu, Q.; Zhou, H. *Tetrahedron Lett.* **2019**, *60*, 327.
- [26] Christopher, M. V.; Liliya, G. N.; David, W. N.; Heather, A. S.; Andreas, D.; Mark, O. B.; Felix, J. B.; Stephen, A. W. *Can. J. Chem.* **2001**, *79*, 1115.
- [27] Wheaton, S. L.; Humanayun Kabir, S. M.; Zhang, H.; Vogelsl, C. M.; Decken, A.; Westcott, S. *Cent. Eur. J. Chem.* **2010**, *8*, 725.

(Cheng, F.)