有机化学

Chinese Journal of Organic Chemistry

金属有机化学专辑

第 39 卷 第 6 期 Vol. 39 No. 6 2019

ISSN 0253-2786

中国化学会主办
中国科学院上海有机化学研究所

Strem Chemicals, Inc., established in 1964, is a privately held manufacturer & marketer of high purity specialty chemicals. Strem's products are used for R&D and commercial scale applications, especially in the pharmaceutical, microelectronics, chemicals & petrochemicals industries. Some key products include catalysts, ligands, organometallics, CVD/ALD precursors and nanomaterials. For more information, visit strem.com. Strem is an ISO 9001:2015 certified company.

STREM'S KEY PRODUCT LINES

- METAL CATALYSTS
- LIGANDS
- INORGANICS
- ORGANOMETALLICS
- PHOTOCATALYSTS
- SCREENING KITS

11-0575

27-0400

78-1350

77-0425

15-8021

15-1053

15-1060

46-2150

29-6720

15-0067

77-0410

15-0126

中国高端 均相催化剂领军者

- 10年研发经验
- 国际市场认可
- 无水无氧封装存储

贵金属催化剂：

Pd Rh Ru Ir Au Cu Ni Zr

有机配体：

P N O

MIT专利授权：

BUCHWALD LIGAND & PRECATALYST

手性膦配体：

BINAP SEGHOS PHOSPHORAMIDITE
JOSIPHOS PHOX SPIRO

手性氮配体：

BOX BINAM DACH TROST DPEN SALEN

手性催化剂：

PHOSPHORIC ACIDS NHC THIOUREA
CINCHONA ALKALOID

光催化剂：

Ir Ru ACRI. SALTS

OTHER: RAFT SILANE PORPHYRIN MOF
ALD PRECURSOR

媲美进口品质的催化剂尽在

WWW.LAAJOO.COM

TEL: 021-60702684

QQ: 3004302861

MAIL: REAGENT@SINOCOMPOUND.COM

理想的实验室伴侣！

中国人都用得起的隔膜真空泵！

旋转蒸发仪/隔膜真空泵

® 上海豫康科教仪器设备有限公司

Shanghai Yukang Science teaching Instrument equipment Co.,Ltd

地址：上海市和桥路470号

电话：021-36338262 56470425
56470426

传真：021-56470425

手机：13701684908

邮编：200435

网址：www.sh-yukang.com

Tel: 021-36338262 56470425
56470426

Fax: 021-56470425

Mobil: 13701684908

Zip Code: 200435

Web: www.sh-yukang.com

钴催化芳香族烯烃的脱氢硅化反应

程彪 陆鹏 赵家金 陆展*

(浙江大学化学系 杭州 310058)

摘要 描述了一种钴催化的高化学、区域、立体选择性的芳香族烯烃的脱氢硅化反应。亚胺吡啶咪唑啉的钴络合物可以有效促进该反应,并能显著提高该反应的化学选择性。该方法利用地球丰产过渡金属、廉价易得的烯烃和硅烷来构建具有高附加值的烯基硅烷。文中同时描述了该反应可放大到克级规模制备及可能的反应机理。

关键词 钴催化; 脱氢硅化; 烯烃; 烯基硅; 区域和立体选择性

Cobalt-Catalyzed Dehydrogenative Silylation of Vinylarenes

Cheng, Biao Lu, Peng Zhao, Jiajin Lu, Zhan*

(Department of Chemistry, Zhejiang University, Hangzhou 310058)

Abstract A highly chemo-, regio-, and stereo-selective cobalt-catalyzed dehydrogenative silylation of vinylarenes was described. The imidazoline iminopyridine cobalt complex could promote this reaction effectively and improve the chemoselectivity dramatically. This protocol used earth-abundant transition metal, readily available alkenes and hydrosilanes to construct valuable vinylsilanes. The reaction could be carried out in gram scale and the proposed mechanism was also described.

Keywords cobalt-catalyzed; dehydrogenative silylation; alkenes; vinylsilanes; regio- and stereo-selectivity

1 Introduction

Vinylsilanes are important reagents which have great applications in organic synthesis and materials science.^[1] There are various methods to prepare vinylsilanes, such as the silyl-Heck reaction,^[2] the hydrosilylation of alkynes,^[3] the cross-metathesis of vinylsilanes with olefins,^[4] and others.^[5] The dehydrogenative silylation of alkenes, which used to be viewed as an undesired side reaction in the process of alkene hydrosilylation,^[6] has attracted more and more interests from chemists as an alternative method to construct vinylsilanes (Scheme 1). The usage of transition metals, such as ruthenium,^[7] rhodium,^[8] palladium,^[9] iridium,^[10] platinum,^[11] rhenium,^[12] titanium,^[13] leads to rapid growth of the dehydrogenative silylation of alkenes.

In recent years, base metals, such as iron, cobalt, and nickel, as earth-abundant, nontoxic, lower cost, and bio-compatible transition metals, have been great attractive in alkene hydrosilylation.^[14] In 1962, Nesmeyanov and co-

workers^[15] reported a $\text{Fe}(\text{CO})_5$ -catalyzed dehydrogenative silylation of ethylene with Et_3SiH , which could obtain the vinylsilane in high chemoselectivity. After that, several iron-catalyzed dehydrogenative silylation of alkenes had been reported, however, the substrate scopes were limited or not well explored.^[16] Marciniak and co-workers^[17] reported that nickel complexes were effective for the dehydrogenative silylation of alkenes, however, the transformation should be performed at 120 °C and the chemoselectivities were moderate. As for cobalt, its application in dehydrogenative silylation was not been well explored. In 1987, Murai group^[18] reported a $\text{Co}_2(\text{CO})_8$ -catalyzed dehydrogenative silylation of α,β -unsaturated esters. In 2014, Chirik and co-workers^[19] developed a bis(imino)pyridine cobalt-catalyzed dehydrogenative silylation of aliphatic alkenes. This transformation could selectively obtained allylsilanes in most cases. As for the substrates lacking allylic hydrogens, vinylsilanes were obtained as the major products. For 3,3-dimethylbutene, the *E* isomer of the vi-

* Corresponding author. E-mail: luzhan@zju.edu.cn

Received March 10, 2019; revised April 3, 2019; published online April 16, 2019.

Project supported by the National Natural Science Foundation of China (No. 21772171), the National Basic Research Program of China (973 Program, No. 2015CB856600), the Zhejiang Provincial Natural Science Foundation (No. LR19B020001), the K. P. Chao's High Technology Development Foundation of Zhejiang University and the Fundamental Research Funds for the Central Universities.

国家自然科学基金(No. 21772171)、国家重点基础研究发展计划(No. 2015CB856600)、浙江省自然科学基金(No. LR19B020001)、浙江大学曹光彪高科技发展基金和浙江大学基本科研业务费资助项目。

nylsilane was obtained. However, the reaction of vinylarenes was not described. In 2016, Pawluć group^[20] used a cobalt complex of a tridentate Schiff-base ligand as catalyst to develop a dehydrogenative silylation of styrene with (HSiMe₂)₂O, which obtained the vinylsilanes in 98% yield with 42% of single and 63% of double addition products. Besides, dehydrogenative silylation products used to be observed as side-products in hydrosilylation of alkenes.^[21] To the best of our knowledge, highly chemo-, regio-, and stereo-selective cobalt-catalyzed dehydrogenative silylation of vinylarenes to obtain vinylsilanes has not been well developed so far.

Scheme 1 Transition metal-catalyzed hydrosilylation and dehydrogenative silylation of alkenes

2 Results and discussion

Our group is interested in earth-abundant transition metal-catalyzed asymmetric hydrofunctionalization of alkenes and alkynes.^[22–25] In the course of our research on hydrosilylation of alkenes and alkynes,^[24] an undesired side-product, vinylsilane, was observed, when using cobalt complex as the catalyst for the hydrosilylation of styrene with diphenylsilane. Then, we began to optimize the conditions for vinylsilane (Table 1). At the beginning of screening, readily available styrene and Ph₂SiH₂ were chosen as model substrates. Imidazoline iminopyridine (**IIP**) ligand cobalt complex^[23g] and NaBHET₃ were used as a precatalyst and activator, respectively. The reaction was conducted using **1a** (0.75 mmol), **2a** (0.5 mmol), **La**·CoCl₂ (5 mol%) and NaBHET₃ (15 mol%) in neat condition at room temperature for 1 h to afford a mixture of dehydrogenative silylation product **3a**, hydrosilylation product **4a** and **5a**, as well as the hydrogenation product generated from the alkenes which served as sacrificial hydrogen acceptors. When changing the steric hindrance of R³ group, we discovered that ⁱPr (**Lc**) was the most suitable group for the selectivity for **3a** (Entries 1–3). The use of more sterically hindered 2,6-diisopropyl imine (**Le**) led to further improvement of the selectivity for **3a** (Entries 3–6). Using other protected groups on nitrogen or the oxazoline iminopyridine (**OIP**)^[23,24,26] ligand led no further improvement (Entries 6–8). Adding solvents to the reaction, such as tetrahydrofuran (THF), toluene, and Et₂O, had slight influence on the results (Entries 9–11). Increasing the amount of styrene to 2 equiv. could increase the yield up to 77% (Entry 12). The reaction underwent smoothly to afford **3a** in 77% yield using 3 mol% catalyst loading within 10 minutes (Entry 13). The standard conditions

were identified as using alkene (2.0 mmol), Ph₂SiH₂ (1.0 mmol), **Le**·CoCl₂ (3 mol%) and NaBHET₃ (9 mol%) in neat condition at room temperature for 10 min.

Table 1 Optimizations for dehydrogenative silylation^a

Entry	L ·CoCl ₂ (mol%)	Solvent	Yield ^b / % of 3a/4a/5a
1	La (5)	Neat	38/19/11
2	Lb (5)	Neat	19/3.8/58
3	Lc (5)	Neat	49/5.9/10
4	Ld (5)	Neat	58/4.8/0
5	Le (5)	Neat	58/2.5/0
6	Lf (5)	Neat	56/3.5/0
7	Lg (5)	Neat	50/7.1/0
8	Lh (5)	Neat	46/17/0
9 ^c	Le (5)	THF	52/1.7/0
10 ^c	Le (5)	Toluene	57/2.1/0
11 ^c	Le (5)	Et ₂ O	56/1.9/0
12 ^d	Le (5)	Neat	77/2.0/0
13 ^{d,e}	Le (3)	Neat	77/1.1/0

^a The reaction was conducted using styrene (0.75 mmol), Ph₂SiH₂ (0.5 mmol), cobalt complex and NaBHET₃ (3 equiv. to cobalt precatalyst) at room temperature for 1 h. ^b Determined by ¹H NMR using mesitylene as an internal standard. ^c Solvent (1 mol/L) ^d **1a/2a** = 2/1. ^e 10 min.

Under the optimized conditions, the substrate scope was explored in Table 2. The substituents at *ortho*-, *meta*-, and *para*-position on phenyl group of styrene were well tolerated, which gave the corresponding vinylsilanes in 81%~86% yields (**3b**~**3d**). The rate of this reaction was influenced by the electronic effect of the substituents on phenyl group of styrene. Electron-donating functional group was well tolerated (**3f**) while the reactions of electron-deficient styrenes should extend the reaction time or increase the catalyst loading (**3g**~**3k**). Ester could be tolerated to afford the corresponding product (**3l**) in 47% yield for there were not some starting materials been consumed up. 1-Vinylnaphthalene could be converted to the corresponding product (**3m**) in 76% yield. Not only secondary silanes, but also tertiary silanes, such as Ph₂MeSiH (**2n**), PhMe₂SiH (**2o**), (EtO)₃SiH (**2p**) were well tolerated to afford the corresponding products in 42%~88% yields.

The gram-scale reaction using 2.4 equiv. of styrene

Table 2 Scope of vinylsilanes^a

$$\text{Ar}-\text{CH}=\text{CH}_2 + [\text{Si}]-\text{H} \xrightarrow[\text{neat, r.t., 10 min} \sim 3 \text{ h}]{\text{Le}\cdot\text{CoCl}_2 (3 \sim 5 \text{ mol\%}) \text{ NaHBET}_3 (9 \sim 15 \text{ mol\%})} \text{Ar}-\text{CH}=\text{CH}-[\text{Si}]$$

2 equiv.

3b, 86% yield

3c, 81% yield

3d, 86% yield

3e, 80% yield

3f, 88% yield

3g, 77% yield

3h, 75% yield^b

3i, 72% yield^b

3j, 76% yield^b

3k, 66% yield^{b,c}

3l, 47% yield^d

3m, 76% yield^{c,e}

3n, 85% yield^d

3o, 88% yield^d

3p, 42% yield^d

^a Standard conditions: alkenes (2.0 mmol), silanes (1.0 mmol), $\text{Le}\cdot\text{CoCl}_2$ (0.03 mmol), NaBHET_3 (0.09 mmol), neat, r.t., 10 min. Isolated yield. ^b 1 h. ^c $\text{Le}\cdot\text{CoCl}_2$ (5 mol%). ^d Alkenes (1.0 mmol), silanes (0.5 mmol), $\text{Le}\cdot\text{CoCl}_2$ (0.025 mmol), NaBHET_3 (0.075 mmol), 3 h. ^e 2 h.

could be smoothly performed at room temperature for 26 h to afford **3p** in 91% yield (Eq. 1), which is a very useful synthetic intermediate.^[27]

Inspired by the previously reported mechanism of dehydrogenative silylation of alkenes,^[19] the proposed mechanism was shown in Scheme 2. The cobalt-silicon species (**A**) obtained from reducing $\text{IIP}\cdot\text{CoCl}_2$ by NaBHET_3 and hydrosilane could undergo alkene insertion to generate cobalt alkyl species (**B**). The cobalt alkyl species (**B**) underwent β -hydrogen elimination to generate the dehydrogenative silylation product and the cobalt-hydrogen species (**C**). The cobalt-hydrogen species (**C**) underwent alkene insertion to generate cobalt alkyl species (**D**), which reacted with hydrosilane to regenerate cobalt-silicon species (**A**) and the hydrogenation byproduct.

3 Conclusions

In summary, we developed a highly chemo-, regio-, and stereo-selective cobalt-catalyzed dehydrogenative silylation of vinylarenes using imidazoline iminopyridine cobalt complex to deliver vinylsilanes. This protocol used earth-abundant transition metal and readily available materials to construct valuable vinylsilanes, which were useful synthetic intermediates. The reaction could be easily carried out in gram scale and the proposed mechanism was also described. The development of more non-precious metal-catalyzed reactions is continuously desirable in our laboratory.

Scheme 2 Proposed mechanism

4 Experimental section

4.1 General information

THF, ether and toluene were distilled from sodium benzophenone ketyl prior to use. Ph_2SiH_2 (97%) was purchased from TCI. NaBHET_3 (1.0 mol/L in THF) was purchased from Aldrich, CoCl_2 (99.7%) was purchased from Aladdin, $\text{Pd}(\text{OAc})_2$ (98%) was purchased from Zhejiang Metallurgical Research Institute and PPh_3 (98%), CuI (98%), DBU (98%) were purchased from Energy. All the commercially available chemicals were used as received. Alkenes were prepared according to Wittig Reaction.

NMR spectra were recorded on a Bruker-400 instrument or a BIXI-I-400 instrument. ^1H NMR chemical shifts were referenced to tetramethylsilane signal (δ 0), and ^{13}C NMR

chemical shifts were referenced to the solvent resonance (δ 77.00, CDCl_3). High-resolution mass spectra (HRMS) were recorded on a Water XEVOG2-S TOF or GCT Premier. IR spectra were recorded on a Perkin-Elmer Spectrum One FTIR spectrometer with diamond ATR accessory.

4.2 Procedures for preparation of ligands and metal complexes

A 50 mL Schlenk flask equipped with a magnetic stirrer and a flanging rubber plug was dried with flame under vacuum. When cooled to ambient temperature, it was vacuumed and flushed with N_2 . This degassed procedure was repeated for three times. Then $\text{Pd}(\text{OAc})_2$ (5 mol%), PPh_3 (10 mol%), CuI (1.0 equiv.), DBU (1.0 equiv.), imidazoline (1.2 equiv.), imidopyridine (1.0 equiv.), and DMF (0.3 mol/L) were added sequentially. The mixture was placed in an oil bath and stirred at 110 °C overnight. Then the mixture was cooled to room temperature, diluted with diethyl ether, washed with 10% aqueous ammonia and brine, dried over Na_2SO_4 anhydrous, and concentrated *in vacuo*. The crude mixture was purified by flash column chromatography to give the corresponding product.

(*S,E*)-*N*-(1-(6-(4-Benzyl-1-phenyl-4,5-dihydro-1*H*-imidazol-2-yl)pyridin-2-yl)ethylidene)-2-(*tert*-butyl)aniline (**La**): Yellow oil (0.9305 g, 63% yield). $[\alpha]_{\text{D}}^{20} +29.0$ (*c* 1.07, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 8.27~8.20 (m, 1H), 8.08~8.00 (m, 1H), 7.81 (dd, $J=8.0, 7.6$ Hz, 1H), 7.38~7.32 (m, 1H), 7.32~7.23 (m, 4H), 7.21~7.16 (m, 1H), 7.13~6.96 (m, 4H), 6.94~6.86 (m, 1H), 6.66 (d, $J=7.6$ Hz, 2H), 6.52 (dd, $J=1.2, 7.6$ Hz, 1H), 4.69~4.56 (m, 1H), 4.08 (t, $J=9.6$ Hz, 1H), 3.74 (dd, $J=7.6, 9.2$ Hz, 1H), 3.27 (dd, $J=4.8, 14.0$ Hz, 1H), 2.91 (dd, $J=8.4, 13.6$ Hz, 1H), 1.66 (s, 3H), 1.27 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ : 165.2, 160.2, 155.3, 149.4, 148.4, 143.2, 139.4, 138.0, 136.8, 129.3, 128.19, 128.16, 126.2, 126.15, 126.06, 124.9, 123.52, 123.47, 123.1, 121.8, 119.4, 65.5, 58.2, 42.0, 34.9, 29.4, 15.9; IR (neat) ν : 3059, 2921, 2856, 1675, 1497 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{33}\text{H}_{35}\text{N}_4$ [$\text{M}+\text{H}$] $^+$ 487.2862, found 487.2877.

(*S,E*)-2-(*tert*-Butyl)-*N*-(1-(6-(4-(*tert*-butyl)-1-phenyl-4,5-dihydro-1*H*-imidazol-2-yl)pyridin-2-yl)ethylidene)-aniline (**Lb**): Yellow oil (2.9426 g, 81% yield). $[\alpha]_{\text{D}}^{20} +18.7$ (*c* 1.10, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 8.25~8.19 (m, 1H), 8.08 (dd, $J=0.8, 3.6$ Hz, 1H), 7.84 (dd, $J=8.0, 7.6$ Hz, 1H), 7.36 (dd, $J=1.2, 8.0$ Hz, 1H), 7.16~7.07 (m, 3H), 7.05~6.98 (m, 1H), 6.98~6.91 (m, 1H), 6.85~6.79 (m, 2H), 6.38 (dd, $J=1.6, 8.0$ Hz, 1H), 4.20 (dd, $J=9.6, 10.8$ Hz, 1H), 4.07 (dd, $J=8.0, 11.2$ Hz, 1H), 3.72 (dd, $J=8.0, 8.8$ Hz, 1H), 1.66 (s, 3H), 1.27 (s, 9H), 1.03 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ : 165.4, 160.1, 155.4, 149.5, 148.8, 143.9, 139.6, 137.0, 128.4, 126.3, 126.2, 125.0, 123.6, 123.5, 123.3, 121.8, 119.6, 74.2, 55.4, 35.0, 34.3, 29.5, 26.0, 16.1; IR (neat) ν : 2954, 2925, 2868, 1639, 1592, 1480, 1390 cm^{-1} . HRMS (EI) calcd for $\text{C}_{30}\text{H}_{36}\text{N}_4$ [M] $^+$ 452.2940, found 452.2935.

(*S,E*)-2-(*tert*-Butyl)-*N*-(1-(6-(4-isopropyl-1-phenyl-4,5-

dihydro-(1*H*)-imidazol-2-yl)pyridin-2-yl)ethylidene)aniline (**Lc**): Yellow oil (3.3660 g, 74% yield). $[\alpha]_{\text{D}}^{20} -18.7$ (*c* 1.13, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 8.23 (dd, $J=8.0, 0.8$ Hz, 1H), 8.05 (dd, $J=8.0, 1.2$ Hz, 1H), 7.83 (dd, $J=8.0, 7.6$ Hz, 1H), 7.36 (dd, $J=7.6, 1.2$ Hz, 1H), 7.17~7.08 (m, 3H), 7.06~6.98 (m, 1H), 6.94 (dd, $J=7.6, 7.2$ Hz, 1H), 6.86~6.78 (m, 2H), 6.38 (dd, $J=7.6, 1.2$ Hz, 1H), 4.22~4.09 (m, 2H), 3.78~3.65 (m, 1H), 2.03~1.92 (m, 1H), 1.67 (s, 3H), 1.27 (s, 9H), 1.10 (d, $J=6.8$ Hz, 3H), 1.00 (d, $J=6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 165.4, 160.0, 155.4, 149.5, 148.8, 143.8, 139.6, 136.9, 128.4, 126.3, 126.2, 125.0, 123.6, 123.4, 123.1, 121.8, 119.6, 70.6, 56.6, 35.0, 33.2, 29.5, 19.0, 1 IR (neat) ν : 2957, 1639, 1592, 1479, 1390. 8.1, 16.1 cm^{-1} ; HRMS (EI) calcd for $\text{C}_{29}\text{H}_{34}\text{N}_4$ [M] $^+$ 438.2783, found 438.2784.

(*S,E*)-*N*-(1-(6-(1,4-Diisopropyl-4,5-dihydro-(1*H*)-imidazol-2-yl)pyridin-2-yl)ethylidene)-2,6-diisopropylaniline (**Lf**): Yellow solid (1.6296 g, 74% yield). $[\alpha]_{\text{D}}^{20} -39.1$ (*c* 1.05, CHCl_3). m.p. 103~105 °C. ^1H NMR (CDCl_3 , 400 MHz) δ : 8.37 (dd, $J=8.0, 0.8$ Hz, 1H), 7.94 (dd, $J=7.6, 1.2$ Hz, 1H), 7.84 (dd, $J=8.0, 7.6$ Hz, 1H), 7.20~7.13 (m, 2H), 7.09 (dd, $J=8.6, 6.6$ Hz, 1H), 4.60~4.48 (m, 1H), 3.99~3.87 (m, 1H), 3.55 (dd, $J=10.8, 9.2$ Hz, 1H), 3.24 (dd, $J=9.6, 9.2$ Hz, 1H), 2.82~2.66 (m, 2H), 2.19 (s, 3H), 1.96~1.83 (m, 1H), 1.22 (d, $J=6.8$ Hz, 3H), 1.19~1.09 (m, 15H), 1.07 (d, $J=6.8$ Hz, 3H), 0.97 (d, $J=6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 166.6, 162.5, 155.1, 150.3, 146.3, 137.0, 135.7, 135.6, 125.7, 123.5, 122.9, 122.8, 121.5, 69.6, 46.2, 45.8, 33.4, 28.2, 28.1, 23.1, 22.8, 20.9, 19.7, 19.0, 18.0, 17.1; IR (neat) ν : 2961, 1644, 1565, 1462, 1364 cm^{-1} . HRMS (ESI) calcd for $\text{C}_{28}\text{H}_{41}\text{N}_4$ [$\text{M}+\text{H}$] $^+$ 433.3331, found 433.3331.

(*S,E*)-*N*-(1-(6-(1-(*tert*-Butyl)-4-isopropyl-4,5-dihydro-(1*H*)-imidazol-2-yl)pyridin-2-yl)ethylidene)-2,6-diisopropylaniline (**Lg**): Yellow solid (0.4684 g, 22% yield). $[\alpha]_{\text{D}}^{20} -3.9$ (*c* 0.48, CHCl_3). m.p. 118~120 °C. ^1H NMR: (400 MHz, CDCl_3) δ : 8.31 (dd, $J=7.8, 0.6$ Hz, 1H), 7.81 (dd, $J=7.8, 7.6$ Hz, 1H), 7.51 (dd, $J=7.6, 1.2$ Hz, 1H), 7.20~7.12 (m, 2H), 7.12~7.04 (m, 1H), 3.92~3.81 (m, 1H), 3.68 (dd, $J=10.6, 9.4$ Hz, 1H), 3.36 (dd, $J=9.4, 9.2$ Hz, 1H), 2.80~2.64 (m, 2H), 2.18 (s, 3H), 1.93~1.81 (m, 1H), 1.19~1.10 (m, 21H), 1.06 (d, $J=6.8$ Hz, 3H), 0.99 (d, $J=6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 166.9, 163.9, 155.4, 153.5, 146.3, 136.7, 135.73, 135.68, 125.3, 123.5, 122.9, 121.0, 69.0, 54.1, 50.8, 33.4, 29.4, 28.20, 28.18, 23.11, 23.08, 22.7, 18.9, 18.2, 17.4; IR (neat) ν : 2961, 1645, 1568, 1463, 1364 cm^{-1} . HRMS (EI) calcd for $\text{C}_{29}\text{H}_{42}\text{N}_4$ [M] $^+$ 446.3409, found 446.3401.

La• CoCl_2 ~ **Lg**• CoCl_2 ^[23g] and **Lh**• CoCl_2 ^[28] were prepared according to the previously reported procedures.

4.3 Dehydrogenative silylation of vinylarenes

A 25 mL Schlenk flask equipped with a magnetic stirrer and a flanging rubber plug was dried with flame under vacuum. When cooled to ambient temperature, it was vacuumed and flushed with N_2 . This degassed procedure was repeated for three times. Then **Le**• CoCl_2 complex (3~5

mol%), alkene (2.0 equiv.), hydrosilane (1.0 equiv.), NaBHET_3 (3 equiv. to cobalt precatalyst) were added sequentially. The reaction was run for 10 min~3 h at room temperature. Then the resulting solution was quenched with 10 mL of ether and filtered through a pad of silica gel, washed with ether (10 mL $\times$ 3). The combined filtrate was concentrated and purified by flash column chromatography to give the corresponding product.

(*E*)-Diphenyl(styryl)silane (**3a**)^[29]: Colorless oil (0.2210 g, 77% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.68~7.59 (m, 4H), 7.50~7.25 (m, 11H), 7.09 (d, J =19.0 Hz, 1H), 6.72 (dd, J =3.2, 19.0 Hz, 1H), 5.25 (d, J =3.2 Hz, 1H).

(*E*)-(2-Methylstyryl)diphenylsilane (**3b**): Colorless oil (0.2593 g, 86% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.68~7.51 (m, 5H), 7.47~7.31 (m, 7H), 7.23~7.08 (m, 3H), 6.62 (dd, J =3.2, 18.8 Hz, 1H), 5.26 (d, J =3.2 Hz, 1H), 2.31 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 147.1, 137.2, 135.7, 135.5, 133.7, 130.4, 129.7, 128.3, 128.0, 126.1, 125.5, 123.2, 19.5; IR (neat) ν : 3066, 3016, 2120, 1595, 1481, 1428 cm^{-1} . HRMS (EI) calcd for $\text{C}_{21}\text{H}_{20}\text{Si}$ $[\text{M}]^+$ 300.1334, found 300.1332.

(*E*)-(3-Methylstyryl)diphenylsilane (**3c**): Colorless oil (0.2438 g, 81% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.69~7.55 (m, 4H), 7.49~7.33 (m, 6H), 7.32~7.19 (m, 3H), 7.13~7.01 (m, 2H), 6.70 (dd, J =3.2, 18.8 Hz, 1H), 5.24 (d, J =3.2 Hz, 1H), 2.34 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 149.3, 138.1, 137.8, 135.5, 133.6, 129.7, 129.4, 128.5, 128.0, 127.4, 123.9, 121.2, 21.3; IR (neat) ν : 3049, 3016, 2120, 1581, 1483, 1428 cm^{-1} . HRMS (EI) calcd for $\text{C}_{21}\text{H}_{20}\text{Si}$ $[\text{M}]^+$ 300.1334, found 300.1335.

(*E*)-(4-Methylstyryl)diphenylsilane (**3d**): Colorless oil (0.2586 g, 86% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.66~7.59 (m, 4H), 7.47~7.33 (m, 8H), 7.14 (d, J =8.0 Hz, 2H), 7.06 (d, J =19.0 Hz, 1H), 6.64 (dd, J =3.2, 19.0 Hz, 1H), 5.23 (d, J =3.2 Hz, 1H), 2.34 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 149.1, 138.6, 135.5, 135.2, 133.7, 129.7, 129.3, 128.0, 126.7, 120.0, 21.3; IR (neat) ν : 3048, 2921, 2121, 1606, 1510, 1429 cm^{-1} . HRMS (EI) calcd for $\text{C}_{21}\text{H}_{20}\text{Si}$ $[\text{M}]^+$ 300.1334, found 300.1330.

(*E*)-(4-*tert*-Butylstyryl)diphenylsilane (**3e**): Colorless oil (0.2726 g, 80% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.65~7.57 (m, 4H), 7.45~7.34 (m, 10H), 7.07 (d, J =18.8 Hz, 1H), 6.66 (dd, J =3.2, 18.8 Hz, 1H), 5.24 (d, J =3.2 Hz, 1H), 1.32 (s, 9H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 151.8, 149.0, 135.5, 135.1, 133.7, 129.7, 128.0, 126.5, 125.5, 120.3, 34.6, 31.2; IR (neat) ν : 2959, 2923, 2121, 1604, 1429, 1366 cm^{-1} . HRMS (EI) calcd for $\text{C}_{24}\text{H}_{26}\text{Si}$ $[\text{M}]^+$ 342.1804, found 342.1806.

(*E*)-(4-Methoxystyryl)diphenylsilane (**3f**): Colorless oil (0.2787 g, 88% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.66~7.58 (m, 4H), 7.46~7.34 (m, 8H), 7.03 (d, J =18.8 Hz, 1H), 6.86 (m, 2H), 6.53 (dd, J =3.6, 19.2 Hz, 1H), 5.23 (d, J =3.2 Hz, 1H), 3.81 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 160.0, 148.6, 135.5, 133.8, 130.8, 129.7, 128.0, 118.4, 113.9, 55.2; IR (neat) ν : 3067, 2927, 2120, 1605, 1510, 1427 cm^{-1} . HRMS (EI) calcd for $\text{C}_{21}\text{H}_{20}\text{OSi}$ $[\text{M}]^+$ 316.1283, found 316.1284.

(*E*)-(4-Fluorostyryl)diphenylsilane (**3g**): Colorless oil (0.2330 g, 77% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.65~7.58 (m, 4H), 7.48~7.35 (m, 8H), 7.07~6.98 (m, 3H), 6.62 (dd, J =3.2, 18.8 Hz, 1H), 5.23 (d, J =3.2 Hz, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 162.9 (d, J =247.1 Hz, 1C), 147.7, 135.5, 134.1, 133.4, 129.8, 128.3 (d, J =8.1 Hz, 1C), 128.1, 121.2, 115.5 (d, J =21.2 Hz, 1C); ^{19}F NMR (376.5 MHz, CDCl_3) δ : -112.9; IR (neat) ν : 3068, 3016, 2124, 1602, 1506, 1429 cm^{-1} . HRMS (EI) calcd for $\text{C}_{20}\text{H}_{17}\text{FSi}$ $[\text{M}]^+$ 304.1084, found 304.1081.

(*E*)-(2-Fluorostyryl)diphenylsilane (**3h**): Colorless oil (0.2288 g, 75% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.66~7.55 (m, 5H), 7.47~7.35 (m, 6H), 7.35~7.28 (m, 1H), 7.28~7.21 (m, 1H), 7.15~7.09 (m, 1H), 7.07~6.99 (m, 1H), 6.79 (dd, J =3.2, 19.2 Hz, 1H), 5.25 (d, J =3.2 Hz, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 160.3 (d, J =249.4, 1C), 140.9, 135.5, 133.3, 129.8, 128.1, 127.2, 125.7 (d, J =11.7, 1C), 124.8, 124.0, 115.7 (d, J =21.9, 1C); ^{19}F NMR (376.5 MHz, CDCl_3) δ : -118.7; IR (neat) ν : 2923, 2854, 2125, 1605, 1483, 1456, 1429 cm^{-1} . HRMS (EI) calcd for $\text{C}_{20}\text{H}_{17}\text{FSi}$ $[\text{M}]^+$ 304.1084, found 304.1080.

(*E*)-(4-Chlorostyryl)diphenylsilane (**3i**): Colorless oil (0.2300 g, 72% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.65~7.58 (m, 4H), 7.47~7.36 (m, 8H), 7.33~7.27 (m, 2H), 7.02 (d, J =19.2 Hz, 1H), 6.69 (dd, J =3.2, 19.2 Hz, 1H), 5.23 (d, J =3.2 Hz, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 147.6, 136.2, 135.5, 134.2, 133.2, 129.8, 128.7, 128.1, 127.9, 122.5; IR (neat) ν : 3067, 3017, 2124, 1602, 1489, 1428 cm^{-1} . HRMS (EI) calcd for $\text{C}_{20}\text{H}_{17}\text{ClSi}$ $[\text{M}]^+$ 320.0788, found 320.0792.

(*E*)-(2-Chlorostyryl)diphenylsilane (**3j**): Colorless oil (0.2438 g, 76% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.69~7.60 (m, 5H), 7.53 (d, J =19.2 Hz, 1H), 7.47~7.32 (m, 7H), 7.28~7.18 (m, 2H), 6.72 (dd, J =3.4, 18.8 Hz, 1H), 5.27 (d, J =3.4 Hz, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 144.8, 135.9, 135.5, 133.4, 133.2, 129.8, 129.7, 129.3, 128.1, 126.9, 126.8, 125.3; IR (neat) ν : 3066, 3016, 2124, 1593, 1466, 1430 cm^{-1} . HRMS (EI) calcd for $\text{C}_{20}\text{H}_{17}\text{ClSi}$ $[\text{M}]^+$ 320.0788, found 320.0784.

(*E*)-(3-Bromostyryl)diphenylsilane (**3k**): Colorless oil (0.2395 g, 66% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.65~7.57 (m, 5H), 7.47~7.33 (m, 8H), 7.20 (dd, J =7.6, 8.0 Hz, 1H), 6.99 (d, J =19.0 Hz, 1H), 6.73 (dd, J =3.0, 19.0 Hz, 1H), 5.24 (d, J =3.0 Hz, 1H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 147.3, 139.8, 135.5, 133.1, 131.3, 130.0, 129.9, 129.5, 128.1, 125.3, 123.7, 122.8; IR (neat) ν : 3067, 2923, 2125, 1592, 1560, 1470, 1428 cm^{-1} . HRMS (EI) calcd for $\text{C}_{20}\text{H}_{17}\text{BrSi}$ $[\text{M}]^+$ 364.0283, found 364.0282.

(*E*)-4-(2-(Diphenylsilyl)vinyl)phenyl acetate (**3l**): Colorless oil (0.0815 g, 47% yield). ^1H NMR (CDCl_3 , 400 MHz) δ : 7.67~7.55 (m, 4H), 7.50~7.34 (m, 8H), 7.10~7.01 (m, 3H), 6.66 (dd, J =3.2, 19.2 Hz, 1H), 5.24 (d, J =3.2 Hz, 1H), 2.29 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 169.3, 150.8, 147.9, 135.6, 135.5, 133.4, 129.8, 128.1, 127.7, 121.9, 121.7, 21.1; IR (neat) ν : 3067, 2123, 1766, 1604, 1504, 1429 cm^{-1} . HRMS (EI) calcd for $\text{C}_{22}\text{H}_{20}\text{O}_2\text{Si}$ $[\text{M}]^+$ 344.1233, found 344.1231.

(*E*)-(2-(Naphthalen-1-yl)vinyl)diphenylsilane (**3m**): Colorless oil (0.2554 g, 76% yield). ¹H NMR (CDCl₃, 400 MHz) δ : 8.09~8.02 (m, 1H), 7.94~7.72 (m, 4H), 7.72~7.65 (m, 4H), 7.53~7.37 (m, 9H), 6.81 (dd, *J*=4.0, 18.8 Hz, 1H), 5.34 (d, 1H); ¹³C NMR (CDCl₃, 100 MHz) δ : 146.5, 135.9, 135.5, 135.1, 133.6, 130.9, 129.8, 128.7, 128.5, 128.1, 126.2, 125.8, 125.6, 125.5, 123.9, 123.4; IR (neat) ν : 3064, 3013, 2856, 2122, 1594, 1428 cm⁻¹. HRMS (EI) calcd for C₂₄H₂₀Si [M]⁺ 336.1334, found 336.1335.

(*E*)-Methyldiphenyl(styryl)silane (**3n**):^[30] Colorless oil (0.1274 g, 85% yield). ¹H NMR (CDCl₃, 400 MHz) δ : 7.61~7.55 (m, 4H), 7.49~7.43 (m, 2H), 7.43~7.25 (m, 9H), 6.96 (d, *J*=19.2 Hz, 1H), 6.76 (d, *J*=19.2 Hz, 1H), 0.71 (s, 3H).

(*E*)-Dimethyl(phenyl)(styryl)silane (**3o**):^[30] Colorless oil (0.1054 g, 88% yield). ¹H NMR (CDCl₃, 400 MHz) δ : 7.61~7.53 (m, 2H), 7.47~7.41 (m, 2H), 7.40~7.29 (m, 5H), 7.28~7.22 (m, 1H), 6.94 (d, *J*=19.2 Hz, 1H), 6.58 (d, *J*=19.2 Hz, 1H), 0.43 (s, 6H).

(*E*)-Triethoxy(styryl)silane (**3p**):^[31] Colorless oil (0.0563 g, 42% yield). ¹H NMR (CDCl₃, 400 MHz) δ : 7.50~7.45 (m, 2H), 7.38~7.27 (m, 3H), 7.22 (d, *J*=19.6 Hz, 1H), 6.18 (d, *J*=19.6 Hz, 1H), 3.89 (q, *J*=7.0 Hz, 6H), 1.27 (t, *J*=7.0 Hz, 9H).

4.4 Gram scale reaction

The reaction was conducted according to the general procedure B using **Le**•CoCl₂ complex (0.2981 g, 0.50 mmol, 5 mol %), styrene (2.75 mL, 0.91 g/mL, 24 mmol, 2.4 equiv.), (EtO)₃SiH (1.85 mL, 0.89 g/mL, 10 mmol, 1.0 equiv.), NaBHET₃ (1.5 mL, 1.5 mmol, 1 mol/L in THF, 15 mol%). The reaction was run for 26 h at room temperature. The crude product was purified by flash column chromatography using PE/EtOAc (50/1) as the eluent to give 2.4207 g (0.91 mmol, 91% yield) of the product (**3p**).

Supporting Information ¹H NMR, ¹³C NMR spectra for the synthesized compounds. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- (a) Chan, T. H.; Fleming, I. *Synthesis* **1979**, 761.
(b) Blumenkopf, T. A.; Overman, L. E. *Chem. Rev.* **1986**, 86, 857.
(c) Langkopf, E.; Schinzer, D. *Chem. Rev.* **1995**, 95, 1375.
(d) Fleming, I.; Barbero, A.; Water, D. *Chem. Rev.* **1997**, 97, 2063.
- (a) McAtee, J. R.; Martin, S. E. S.; Ahneman, D. T.; Johnson, K. A.; Watson, D. A. *Angew. Chem., Int. Ed.* **2012**, 51, 3663.
(b) Martin, S. E. S.; Watson, D. A. *J. Am. Chem. Soc.* **2013**, 135, 13330.
- (a) Na, Y.; Chang, S. *Org. Lett.* **2000**, 2, 1887.
(b) Trost, B. M.; Ball, Z. T. *J. Am. Chem. Soc.* **2001**, 123, 12726.
(c) Kawanami, Y.; Sonoda, Y.; Mori, T.; Yamamoto, K. *Org. Lett.* **2002**, 4, 2825.
(d) Denmark, S. E.; Pan, W. *Org. Lett.* **2002**, 4, 4163.
(e) Brunner, H. *Angew. Chem., Int. Ed.* **2004**, 43, 2749.
(f) Aricó, C. S.; Cox, L. R. *Org. Biomol. Chem.* **2004**, 2, 2558.
(g) Menozzi, C.; Dalko, P. I.; Cossy, J. *J. Org. Chem.* **2005**, 70, 10717.
(h) Trost, B. M.; Ball, Z. T. *J. Am. Chem. Soc.* **2005**, 127, 17644.
- (i) Maifeld, S. V.; Tran, M. N.; Lee, D. *Tetrahedron Lett.* **2005**, 46, 105.
(j) Berthon-Gelloz, G.; Schumers, J.-M.; De Bo, G.; Markó, I. E. *J. Org. Chem.* **2008**, 73, 4190.
(k) Sore, H. F.; Blackwell, D. T.; MacDonald, S. J. F.; Spring, D. R. *Org. Lett.* **2010**, 12, 2806.
- (l) Ding, S.-T.; Song, L.-J.; Chung, L. W.; Zhang, X.-H.; Sun, J.-W.; Wu, Y.-D. *J. Am. Chem. Soc.* **2013**, 135, 13835.
(m) Guo, J.; Lu, Z. *Angew. Chem., Int. Ed.* **2016**, 55, 10835.
- (a) Pietraszuk, C.; Fischer, H.; Kujawa, M.; Marciniak, B. *Tetrahedron Lett.* **2001**, 42, 1175.
(b) Denmark, S. E.; Yang, S.-M. *Org. Lett.* **2001**, 3, 1749.
- (a) Takai, K.; Kataoka, Y.; Okazoe, T.; Utimoto, K. *Tetrahedron Lett.* **1987**, 28, 1443.
(b) Murthi, K. K.; Salomon, R. G. *Tetrahedron Lett.* **1994**, 35, 517.
(c) Hodgson, D. M.; Comina, P. J. *Tetrahedron Lett.* **1994**, 35, 9469.
(d) Itami, K.; Nokami, T.; Yoshida, J.-I. *Org. Lett.* **2000**, 2, 1299.
(e) McNulty, J.; Das, P. *Chem. Commun.* **2008**, 1244.
(f) Mu, Q.-C.; Wang, X.-B.; Ye, F.; Sun, Y.-L.; Bai, X.-F.; Chen, J.; Xia, C.-G.; Xu, L.-W. *Chem. Commun.* **2018**, 54, 12994.
- (a) Chen, C.; Hecht, M. B.; Kavara, A.; Brennessel, W. W.; Mercado, B. Q.; Weix, D. J.; Holland, P. L. *J. Am. Chem. Soc.* **2015**, 137, 13244.
(b) Ibrahim, A. D.; Entsminger, S. W.; Zhu, L.-Y.; Fout, A. R. *ACS Catal.* **2016**, 6, 3589.
(c) Jakobsson, K.; Chu, T.; Nikonov, G. I. *ACS Catal.* **2016**, 6, 7350.
(d) Raya, B.; Jing, S.; Balasanthiran, V.; RajanBabu, T. V. *ACS Catal.* **2017**, 7, 2275.
(e) Chen, Y.-J.; Ji, S.-F.; Sun, W.-M.; Chen, W.-X.; Dong, J.-C.; Wen, J.-F.; Zhang, J.; Li, Z.; Zheng, L.-R.; Chen, C.; Peng, Q.; Wang, D.-S.; Li, Y.-D. *J. Am. Chem. Soc.* **2018**, 140, 7407.
(f) Liu, J.-X.; Chen, W.-F.; Li, J.-F.; Cui, C.-M. *ACS Catal.* **2018**, 8, 2230.
- (a) Hori, Y.; Mrrsdo, T.; Watanabe, Y. *Bull. Chem. Soc. Jpn.* **1988**, 61, 3011.
(b) Christ, M. L.; Sabo-Etienne, S.; Chaudret, B. *Organometallics* **1995**, 14, 1082.
(c) Bokka, A.; Jeon, J. *Org. Lett.* **2016**, 18, 5324.
- (a) Millan, A.; Towns, E.; Maitlis, P. M. *J. Chem. Soc., Chem. Commun.* **1981**, 673.
(b) Millan, A.; Fernandez, M.-J.; Bentz, P.; Maitlis, P. M. *J. Mol. Catal.* **1984**, 26, 89.
(c) Doyle, M. P.; Devora, G. A.; Nefedov, A. O.; High, K. G. *Organometallics* **1992**, 11, 549.
(d) Takeuchi, R.; Yasue, H. *Organometallics* **1996**, 15, 2098.
(e) Truscott, B. J.; Slawin, A. M. Z.; Nolan, S. P. *Dalton Trans.* **2013**, 42, 270.
- LaPointe, A. M.; Rix, F. C.; Brookhart, M. *J. Am. Chem. Soc.* **1997**, 119, 906.
- (a) Fernández, M. J.; Esteruelas, M. A.; Jlménez, M. S.; Oro, L. A. *Organometallics* **1986**, 5, 1519.
(b) Oro, L. A.; Fernandez, M. J.; Esteruelas, M. A.; Jimenez, M. S. *J. Mol. Catal.* **1986**, 37, 151.
(c) Tanke, R. S.; Crabtree, R. H. *Organometallics* **1991**, 10, 415.
(d) Lu, B.; Falck, J. R. *J. Org. Chem.* **2010**, 75, 1701.
(e) Cheng, C.; Simmons, E. M.; Hartwig, J. F. *Angew. Chem., Int. Ed.* **2013**, 52, 8984.
- Sprengers, J. W.; de Greef, M.; Duin, M. A.; Elsevier, C. J. *Eur. J. Inorg. Chem.* **2003**, 3811.
- Jiang, Y.; Blacque, O.; Fox, T.; Frech, C. M.; Berke, H. *Chem.-Eur. J.* **2009**, 15, 2121.
- Nakamura, S.; Yonehara, M.; Uchiyama, M. *Chem.-Eur. J.* **2008**, 14, 1068.
- (a) Sun, J.; Deng, L. *ACS Catal.* **2016**, 6, 290.
(b) Du, X.-Y.; Huang, Z. *ACS Catal.* **2017**, 7, 1227.
(c) Zhang, J.-H.; Hao, X.-Q.; Wang, Z.-L.; Ren, C.-J.; Niu, J.-L.;

- Song, M.-P. *Chin. J. Org. Chem.* **2017**, *37*, 1237 (in Chinese). (张家恒, 郝新奇, 王正龙, 任常久, 牛俊龙, 宋毛平, 有机化学, **2017**, *37*, 1237.)
- (d) Gu, Z.-Y.; Ji, S.-J. *Acta Chim. Sinica* **2018**, *76*, 347 (in Chinese).
- (顾正洋, 纪顺俊, 化学学报, **2018**, *76*, 347.)
- [15] Nesmeyanov, A. N.; Freidlina, R. K.; Chukovskaya, E. C.; Petrova, R. G.; Belyavsky, A. B. *Tetrahedron* **1962**, *17*, 61.
- [16] (a) Kakiuchi, F.; Tanaka, Y.; Chatani, N.; Murai, S. *J. Organomet. Chem.* **1993**, *456*, 45.
(b) Naumov, R. N.; Itazaki, M.; Kamitani, M.; Nakazawa, H. *J. Am. Chem. Soc.* **2012**, *134*, 804.
(c) Marciniak, B.; Kownacka, A.; Kownacki, I.; Taylor, R. *Appl. Catal. A: General* **2014**, *486*, 230.
(d) Sunada, Y.; Noda, D.; Soejima, H.; Tsutsumi, H.; Nagashima, H. *Organometallics* **2015**, *34*, 2896.
(e) Du, X.-Y.; Zhang, Y.-L.; Peng, D.-J.; Huang, Z. *Angew. Chem., Int. Ed.* **2016**, *55*, 6671.
- [17] (a) Marciniak, B.; Maciejewski, H.; Kownacki, I. *J. Mol. Catal. A: Chemical* **1998**, *135*, 223.
(b) Maciejewski, H.; Marciniak, B.; Kownacki, I. *J. Organomet. Chem.* **2000**, *597*, 175.
- [18] Takeshita, K.; Seki, Y.; Kawamoto, K.; Murai, S.; Sonoda, N. *J. Org. Chem.* **1987**, *52*, 4864.
- [19] Atienza, C. C. H.; Diao, T.; Weller, K. J.; Nye, S. A.; Lewis, K. M.; Delis, J. G. P.; Boyer, J. L.; Roy, A. K.; Chirik, P. J. *J. Am. Chem. Soc.* **2014**, *136*, 12108.
- [20] Gorczyński, A.; Zaranek, M.; Witomska, S.; Bocian, A.; Stefankiewicz, A. R.; Kubicki, M.; Patroniak, V.; Pawluć, P. *Catal. Commun.* **2016**, *78*, 71.
- [21] (a) Wang, C.; Teo, W. J.; Ge, S.-Z. *ACS Catal.* **2017**, *7*, 855.
(b) Liu, Y.; Deng, L. *J. Am. Chem. Soc.* **2017**, *139*, 1798.
(c) Gao, Y.-F.; Wang, L.-J.; Deng, L. *ACS Catal.* **2018**, *8*, 9637.
(d) Basu, D.; Gilbert-Wilson, R.; Gray, D. L.; Rauchfuss, T. B. *Organometallics* **2018**, *37*, 2760.
(e) Sanagawa, A.; Nagashima, H. *Organometallics* **2018**, *37*, 2859.
- [22] For selected reviews on earth-abundant transition metal-catalyzed asymmetric hydrofunctionalization of alkenes and alkynes, see:
(a) Chen, J.-H.; Lu, Z. *Org. Chem. Front.* **2018**, *5*, 260.
(b) Chen, J.-H.; Guo, J.; Lu, Z. *Chin. J. Chem.* **2018**, *36*, 1075.
- [23] For hydroboration of alkenes and alkynes, see: (a) Chen, J.-H.; Xi, T.; Lu, Z. *Org. Lett.* **2014**, *16*, 6452.
(b) Chen, J.-H.; Xi, T.; Ren, X.; Cheng, B.; Guo, J.; Lu, Z. *Org. Chem. Front.* **2014**, *1*, 1306.
(c) Zhang, H.-Y.; Lu, Z. *ACS Catal.* **2016**, *6*, 6596.
(d) Xi, T.; Lu, Z. *ACS Catal.* **2017**, *7*, 1181.
(e) Chen, X.; Cheng, Z.-Y.; Lu, Z. *Org. Lett.* **2017**, *19*, 969.
(f) Chen, C.-H.; Shen, X.-Z.; Chen, J.-H.; Hong, X.; Lu, Z. *Org. Lett.* **2017**, *19*, 5422.
(g) Guo, J.; Cheng, B.; Shen, X.-Z.; Lu, Z. *J. Am. Chem. Soc.* **2017**, *139*, 15316.
(h) Chen, X.; Cheng, Z.-Y.; Guo, J.; Lu, Z. *Nat. Commun.* **2018**, *9*, 3939.
- [24] For hydrosilylation of alkenes and alkynes, see:
(a) Chen, J.-H.; Cheng, B.; Cao, M.-Y.; Lu, Z. *Angew. Chem., Int. Ed.* **2015**, *54*, 4661.
(b) Guo, J.; Lu, Z. *Angew. Chem., Int. Ed.* **2016**, *55*, 10835.
(c) Xi, T.; Lu, Z. *J. Org. Chem.* **2016**, *81*, 8858.
(d) Guo, J.; Shen, X.-Z.; Lu, Z. *Angew. Chem., Int. Ed.* **2017**, *56*, 615.
(e) Cheng, B.; Lu, P.; Zhang, H.-Y.; Cheng, X.-P.; Lu, Z. *J. Am. Chem. Soc.* **2017**, *139*, 9439.
(f) Cheng, B.; Liu, W.-B.; Lu, Z. *J. Am. Chem. Soc.* **2018**, *140*, 5014.
(g) Cheng, Z.-Y.; Xing, S.-P.; Guo, J.; Cheng, B.; Hu, L.-F.; Zhang, X.-H.; Lu, Z. *Chin. J. Chem.* **2019**, *37*, 457.
(h) Guo, J.; Wang, H.-L.; Xing, S.-P.; Hong, X.; Lu, Z. *Chem* **2019**, *5*, 881.
- [25] For asymmetric hydrogenation of alkenes, see: Chen, J.-H.; Chen, C.-H.; Ji, C.-L.; Lu, Z. *Org. Lett.* **2016**, *18*, 1594.
- [26] Zhang, L.; Zuo, Z.-Q.; Wan, X.-L.; Huang, Z. *J. Am. Chem. Soc.* **2014**, *136*, 15501.
- [27] Komiyama, T.; Minami, Y.; Hiyama, T. *ACS Catal.* **2017**, *7*, 631.
- [28] Guo, J.; Chen, J.-H.; Lu, Z. *Chem. Commun.* **2015**, *51*, 5725.
- [29] JoongLee, S.; KyeuPark, M.; HeeHan, B. *Silicon Chemistry* **2002**, *1*, 41.
- [30] Ramirez-Oliva, E.; Hernández, A.; Martínez-Rosales, J. M.; Aguilar-Elguezabal, A.; Herrera-Pérez, G.; Cervantes, J. *ARKIVOC* **2006**, 126.
- [31] Karabelas, K.; Hallberg, A. *J. Org. Chem.* **1986**, *51*, 5286.

(Zhao, X.)