1,4-二氮杂双环[2.2.2]辛烷催化双环邻-氨基腈衍生物的合成

颜世强^{a,b} 董道青^a 解春文^b 王文笙^b 王祖利^{*,a}^(a) 青岛农业大学化学与药学院 青岛 266109^(b) 山东达因海洋生物制药股份有限公司 威海 264300

摘要 报道了 1,4-二氮杂双环[2.2.2]辛烷(DABCO)催化下的三组分一锅法合成双环邻-氨基腈衍生物, 在该催化体系下, 21 种双环邻-氨基腈衍生物以 68%~96%的收率制备得到. 该方法于室温下即可顺利完成, 反应时间短, 目标化合物产率高. 重要的是目标产物通过简单的过滤和乙醇洗涤即可获得

关键词 1,4-二氮杂双环[2.2.2]辛烷(DABCO); 多组分反应; 邻-氨基腈衍生物; 环境友好; 乙醇

Synthesis of Bicyclic *ortho*-Aminocarbonitrile Derivatives Catalyzed by 1,4-Diazabicyclo[2.2.2]octaneYan, Shiqiang^{a,b} Dong, Daoqing^a Xie, Chunwen^b Wang, Wensheng^b Wang, Zuli^{*,a}^(a) Qingdao Agricultural University, College of Chemistry and Pharmaceutical Sciences, Qingdao, 266109^(b) Shandong Dyne Marine Biopharmaceutical Co., Ltd., Weihai, 264300

Abstract One-pot three-component reactions catalyzed by 1,4-diazabicyclo[2.2.2]octane (DABCO) for the synthesis of bicyclic *ortho*-aminocarbonitrile derivatives (21 examples, 68%~96%) have been developed. The reactions proceeded smoothly at room temperature and generated the corresponding products in short reaction time with high to excellent yields. Importantly, the desired products could be easily collected through simple filtration and washing with ethanol.

Keywords 1,4-diazabicyclo[2.2.2]octane (DABCO); multicomponent reaction; *ortho*-aminocarbonitrile derivatives; environmentally benign; ethanol

1 Introduction

One powerful trend in green and sustainable chemistry is the replacement of traditional organic reactions, which use harmful reagents that generate a large amount of chemical waste, with clean catalytic alternatives that exhibit high atom-efficiency. Multicomponent reactions (MCRs) are preferred over the stepwise synthetic methods for the following reasons: decreased usage of hazardous chemical compounds, operational simplicity and reduced reaction time, easy isolation of targeted products that avoids tedious purification and reduced costs.^[1] Most synthetic reactions are performed in harmful organic solvents, which not only cause environmental problem but also consume vast limited fossil resources. Moreover, the generated products are usually mixed with un-wanted side-products, making it neces-

sary to perform purification steps such as column chromatography. In order to resolve these problems, considerable efforts have been spent on the development of group-assistant-purification organic transformation, especially in environmentally benign solvents.^[2]

The synthesis of *ortho*-aminocarbonitriles derivatives is an important task to the research community due to their importance as core structures in organic synthesis,^[3] so many methods for their synthesis have been reported. For examples, ionic liquids were good catalyst for the synthesis of these compounds, and many different kinds of ionic liquids mediated reactions for the synthesis of *ortho*-aminocarbonitriles derivatives have been described.^[4] Organic bases, such as pyrrolidine,^[5a] morpholine,^[5b] imidazole,^[5c] ethanediamine^[5d] and NEt₃,^[5e,5f] as efficient catalysts for the synthesis of *ortho*-aminocarbonitriles de-

* Corresponding author. E-mail: wangzulichem@163.com

Received January 17, 2019; revised April 20, 2019; published online April 26, 2019.

Project supported by the National Natural Science Foundation of China (Nos. 81573340, 21402103, 21772107), the Special Projects for Independent Innovation and Achievement Transformation of Shandong Province (No. 2015ZDXX0302A02) and the Research Fund of Qingdao Agricultural University Highlevel Person (No. 631303).

国家自然科学基金(Nos. 81573340, 21402103, 21772107)、山东省自主创新及成果转化专项(No. 2015ZDXX0302A02)及青岛农业大学高层次人才基金(No. 631303)资助项目.

rivatives have also been reported. In addition, zinc titanate nanopowder,^[6] glucose-containing imidazolium salt,^[7] saccharine,^[8] triethylamine acetate (TEAA),^[8] benzyltriethylammonium chloride (TEBAC),^[9] Borax,^[10] deep eutectic solvent^[11] and cinchona alkaloid-derived bifunctional thiourea^[12] have been used to promote the formation of *ortho*-aminocarbonitriles derivatives. Despite these developments, many of these methods still have limitations such as the use of commercially unavailable reactants or catalysts, involve relatively harsh conditions, go through multistep procedures, require specific solvents or workup precautions.

Because of the basic property of 1,4-diazabicyclo[2.2.2]octane (DABCO), it is usually used as catalyst in organic reactions. At present, various of reactions have been realized in the presence of DABCO.^[13] For example, reactions of 2-(2-chloroaryl)methyleneimidazolidines with allenic esters catalyzed by DABCO have been realized and afforded imidazo(pyrido)[1,2-*a*]pyridines with good yields.^[14] In 2012, functionalized chromen-5-ones and pyrano[3,2-*c*]chromen-5-ones were prepared from the reaction of α -oxoketene-*N,S*-arylaminoacetals, aromatic aldehydes, and dimedone/4-hydroxycoumarin in the presence of DABCO.^[15] Recently, a [4 + 2] cyclo-addition reaction catalyzed by DABCO was reported by Meng and co-workers.^[16]

In our preparation of this manuscript, Chinchkar and co-workers^[18] described the multi-component reaction of cyclic ketone, aromatic aldehyde and two moles of malononitrile for the synthesis of functionalized condensed bicyclic compounds viz tetrahydroindenes, tetrahydronaphthalenes, hexahydrobenzo^[17] annulenes and hexahydrobenzo[8]-annulenes with a mixed solvent ($V_{\text{EtOH}} : V_{\text{H}_2\text{O}} = 7 : 3$) in the presence of 20 mol% DABCO at reflux condition.^[17] In continuation of our interests in organic synthesis,

herein, we describe one-pot three-component reactions catalyzed by DABCO for the construction of *ortho*-aminocarbonitriles derivatives (Scheme 1). The desired *ortho*-aminocarbonitriles crystallized directly from EtOH and were isolated through simple filtration and washing with ethanol.

Scheme 1 Synthesis of *ortho*-aminocarbonitriles

2 Results and discussion

In our initial study, cyclohexanone (**1a**), 4-methoxybenzaldehyde (**2j**) and malononitrile were chosen as model substrates to survey the reaction parameters. It was found that the solvent plays an important role, as shown in Table 1, when methanol was used as the solvent, the reaction generated the corresponding products with 87% (Table 1, Entry 1). To our pleasure, 96% yield of the product was isolated when ethanol was used as solvent (Table 1, Entry 2). Other solvents such as CH_3CN , THF and CH_2Cl_2 were also effective, but the yields were moderate (Table 1, Entries 3~5). To our disappointment, water was not suitable for this reaction and the product was only obtained in 31% yield (Table 1, Entry 6), even if the reaction time was prolonged to 30 min. We also tried the reaction in solvent-free conditions, but the yield was not high (Table 1, Entry 7). When the loading of the catalyst was increased to 0.1 equiv., almost identical yield was obtained (Table 1, Entry 8). But

Table 1 Screening of optimal reaction conditions^a

Entry	Catalyst (equiv.)	Solvent	Temp./°C	Time/min	Isolated yield/%
1	DABCO (0.05)	Methanol	r.t.	5	87
2	DABCO (0.05)	Ethanol	r.t.	5	96
3	DABCO (0.05)	Acetonitrile	r.t.	5	78
4	DABCO (0.05)	Tetrahydrofuran	r.t.	5	56
5	DABCO (0.05)	Dichloromethane	r.t.	15	42
6	DABCO (0.05)	Water	r.t.	30	31
7	DABCO (0.05)	Solvent-free	r.t.	10	55
8	DABCO (0.10)	Ethanol	r.t.	5	94
9	DABCO (0.02)	Ethanol	r.t.	5	81
10	K_2CO_3 (0.10)	Ethanol	r.t.	60	34
11	Na_2CO_3 (0.10)	Ethanol	r.t.	60	26
12	Urea (0.10)	Ethanol	80	360	51

^a Unless otherwise stated, all reactions were carried out with **1a** (10 mmol), **2i** (10 mmol), **3** (20 mmol) and solvent (10 mL). The yield of isolated product is reported.

the yield of the desired product decreased obviously when the loading of the catalyst was reduced to 0.02 equiv. (Table 1, Entry 9). Besides DABCO, other catalysts such as K_2CO_3 and Na_2CO_3 were also tested, and the target compound was obtained in moderate yields (Table 1, Entries 10, 11). When reaction was carried out in 80 °C, urea did not show good activity and 51% yield of the product was isolated (Table 1, Entry 12).

With the optimized reaction conditions in hand, the scope of the reaction was expanded to other ketone and aldehyde as shown in Table 2. The reaction proceeded smoothly to afford the desired product with good to excellent yield in most of the cases. For the mono-substituted R groups, both electron-poor and electron-rich groups were well compatible with this reaction conditions, and a slightly higher yield was observed when electron-rich groups were attached on R group (**4a**, **4e**, **4h**, **4j**). But the steric effect of the substituent in the aromatic aldehydes has no regular influence on the yields (**4b**, **4c**, **4d**, **4e**, **4f**, **4g**). Disubstituted aromatic aldehydes were also tested, a high yield was obtained when 4-hydroxy-3-methoxybenzaldehyde was subjected to this system (**4j**). Besides substituted benzaldehyde, aromatic heterocyclic aldehydes were also suitable for this reaction, affording the corresponding products in high yields (**4k**, **4l**, **4m**). But only moderate yields were obtained when aliphatic aldehydes were used as substrates (**4n**, **4o**). it is interesting to note that changing from cyclohexanone to cyclopentanone was also found to be suitable for this reaction and the desired products can be formed in moderate to high yields (**4p**~**4v**).

The same excellent results were also obtained when this method was performed at a gram scale using cyclohexanone, benzaldehyde and malononitrile as substrates (Scheme 2). This example clearly demonstrates the preparative utility of this newly developed method.

The efficiency of our method with earlier reported methods is compared by the mount of catalyst, reaction conditions, yield and time. It is clear from the table (Table 3, Entry 1), that our protocol promotes the reactions effectively as far the amount of catalyst and reaction time are

Table 2 Synthesis of **4** under optimum conditions^a

Entry	R	n	Product	Time/min	Yield ^b /%
1	C ₆ H ₅	1	4a	10	87
2	2-ClC ₆ H ₄	1	4b	5	90
3	2-FC ₆ H ₄	1	4c	3	87
4	3-FC ₆ H ₄	1	4d	5	84
5	4-FC ₆ H ₄	1	4e	5	88
6	2-O ₂ NC ₆ H ₄	1	4f	5	91
7	3-O ₂ NC ₆ H ₄	1	4g	5	88
8	4-MeC ₆ H ₄	1	4h	5	94
9	4-MeOC ₆ H ₄	1	4i	5	96
10	4-OH-3-MeOC ₆ H ₃	1	4j	10	85
11	1-Naphthyl	1	4k	3	88
12	2-Furyl	1	4l	5	80
13	2-Thenyl	1	4m	5	87
14	CH ₃ CH ₂	1	4n	15	68 ^c
15	(CH ₃) ₂ CH	1	4o	15	71 ^c
16	C ₆ H ₅	0	4p	15	78
17	4-MeOC ₆ H ₄	0	4q	10	85
18	2-ClC ₆ H ₄	0	4r	10	80
19	4-FC ₆ H ₄	0	4s	10	76
20	1-Naphthyl	0	4t	5	88
21	2-Thenyl	0	4u	10	73

^a Reaction conditions: **1** (1 mmol), **2** (1 mmol), **3** (2 mmol), ethanol (1 mL), DABCO (0.05 mmol) at room temperature. ^b Isolated yields. ^c Heated at 60 °C.

Scheme 2 Synthesis of **4a** on gram scale

Table 3 Comparison of catalytic activity of DABCO with various catalysts for the synthesis of bicyclic *ortho*-aminocarbonitrile

Entry	Catalyst	Reaction condition	Time/min	Yield/%	Ref.
1	DABCO (0.05 equiv.)	Ethanol, r.t.	5	96	This work
2	DABCO (0.20 equiv.)	V(Ethanol) : V(water)=7 : 3, reflux	60	92	[17]
3	DES (1.0 mL/mmol)	Neat, 50 °C	80	78	[11]
4	OBS (0.25 equiv.)	Ethanol, reflux	300	92	[8]
5	TEAA (0.1 mL/mmol)	Neat, 80 °C	240	90	[8]
6	Borax (0.10 equiv.)	Ethanol, r.t.	60	87	[10]
7	[BPy]BF ₄ (2.0 mL/mmol)	Neat, 60 °C	240	89	[4a]

concerned.

According to the literature^[4-12] and our experiment, the possible mechanism was proposed in Scheme 3. **A** was formed from the Knoevenagel condensation reaction of cyclohexanone and malononitrile. Then the Michael addition reaction of **A** and **B** which was formed from the Knoevenagel condensation of benzaldehyde and malononitrile occurred to form **C** in the presence of DABCO. After that, **C** undergo intramolecular addition reaction to form **D**. Finally, the isomerization of **D** provide the desired products **E**.

Scheme 3 Probable reaction mechanism

3 Conclusions

In summary, a direct method for the synthesis of *ortho*-aminocarbonitriles derivatives via DABCO-catalyzed reaction of aromatic aldehyde, cyclohexanone, and 2 equiv. of malononitrile is presented here. The reactions generated the corresponding products with good to excellent yields in most cases under simple and mild reaction conditions in a short time. Further developments of this synthetic utility of this method are currently underway in our lab.

4 Experimental

4.1 General information

All major chemicals and solvents were obtained from commercial sources and used without further purification. NMR spectra were recorded at 400 MHz for protons on JOEL JNM-ECA 400 spectrometers. ¹H NMR chemical shifts (δ) are relative to TMS ($\delta=0.0$). Chemical shifts for ¹³C NMR spectra are reported in parts per million (ppm) from tetramethylsilane with the solvent as the internal standard.

4.2 General procedure for the synthesis of bicyclic *ortho*-aminocarbonitrile derivatives

A mixture of cyclic ketone **1** (1 mmol), aldehyde **2** (1 mmol), malononitrile **3** (2 mmol), DABCO (0.05 mmol), and ethanol (1 mL) was stirred at room temperature. The progress of the reaction was monitored by thin layer chro-

matography (TLC). On completion of the reaction, the solid product was collected by simple filtration, washed with ethanol and dried. The solid was further recrystallized with 95% EtOH/DMF (*V/V*=1/4) to provide the pure product **4**.

2-Amino-4-phenyl-4a,5,6,7-tetrahydronaphthalene-1,3,3(4*H*)-tricarbonitrile (**4a**): White solid, yield 87%. m.p. 254~255 °C (lit.^[3] 255~257 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.61~7.59 (m, 1H), 7.50~7.43 (m, 4H), 7.36 (s, 2H), 5.75~5.71 (m, 1H), 3.54 (d, *J*=12.5 Hz, 1H), 2.81 (t, *J*=11.5 Hz, 1H), 2.19 (d, *J*=18.9 Hz, 1H), 2.11~1.99 (m, 1H), 1.68 (dd, *J*=11.3, 3.8 Hz, 1H), 1.52~1.37 (m, 2H), 0.85 (dt, *J*=13.5, 6.9 Hz, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ : 144.04, 135.11, 132.86, 129.40, 129.33, 129.07, 127.41, 120.80, 116.64, 113.03, 112.82, 82.01, 51.00, 43.34, 34.31, 27.47, 25.33, 21.46; HRMS (ESI) calcd for C₁₉H₁₇N₄ [M+H]⁺ 301.1448, found 301.1453.

2-Amino-4-(2-chlorophenyl)-4a,5,6,7-tetrahydronaphthalene-1,3,3(4*H*)-tricarbonitrile (**4b**): White solid, yield 90%. m.p. 270~272 °C (lit.^[3] 271~272 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.77 (dd, *J*=7.8, 1.4 Hz, 1H), 7.62 (dd, *J*=7.9, 1.3 Hz, 1H), 7.55~7.52 (m, 2H), 7.43 (s, 2H), 5.77~5.74 (m, 1H), 3.88 (d, *J*=12.4 Hz, 1H), 2.86 (t, *J*=12.1 Hz, 1H), 2.23~2.13 (m, 1H), 2.13~2.04 (m, 1H), 1.72~1.62 (m, 1H), 1.44 (ddd, *J*=13.4, 10.6, 6.8 Hz, 1H), 1.40~1.32 (m, 1H), 0.81 (td, *J*=13.4, 2.5 Hz, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ : 143.93, 135.58, 132.28, 131.14, 130.65, 129.70, 128.87, 128.38, 121.35, 116.45, 112.72, 112.04, 81.93, 46.94, 42.11, 34.96, 27.21, 25.22, 21.27; HRMS (ESI) calcd for C₁₉H₁₅ClN₄Na [M+Na]⁺ 357.0877, found 357.0869.

2-Amino-4-(2-fluorophenyl)-4a,5,6,7-tetrahydronaphthalene-1,3,3(4*H*)-tricarbonitrile (**4c**): White solid, yield 87%. m.p. 264~267 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.82~7.68 (m, 1H), 7.57~7.26 (m, 5H), 5.72~5.67 (m, 1H), 3.72 (d, *J*=12.6 Hz, 1H), 2.89 (dd, *J*=38.3, 27.5 Hz, 1H), 2.25~2.01 (m, 2H), 1.80~1.63 (m, 1H), 1.59~1.36 (m, 2H), 1.02~0.79 (m, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ : 162.31, 160.34, 143.66, 131.58, 129.63, 128.84, 125.40, 121.80, 121.32, 116.49, 112.65, 82.15, 43.25, 42.53, 34.16, 27.24, 25.25, 21.32; HRMS (ESI) calcd for C₁₉H₁₅FN₄Na [M+Na]⁺ 341.1173, found 341.1167.

2-Amino-4-(3-fluorophenyl)-4a,5,6,7-tetrahydronaphthalene-1,3,3(4*H*)-tricarbonitrile (**4d**): White solid, yield 84%. m.p. 262~265 °C (lit.^[3] 260~262 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ : 7.60~7.43 (m, 2H), 7.39 (s, 2H), 7.35~7.25 (m, 2H), 5.75~5.72 (m, 1H), 3.65 (t, *J*=13.3 Hz, 1H), 2.83 (d, *J*=8.8 Hz, 1H), 2.23~2.14 (m, 1H), 2.11~1.98 (m, 1H), 1.68 (d, *J*=7.5 Hz, 1H), 1.47 (m, 2H), 0.90~0.87 (m, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ : 143.87, 137.85, 131.19, 129.08, 128.90, 123.67, 120.98, 119.61, 116.57, 116.24, 114.49, 112.87, 112.68, 82.06, 50.38, 43.05, 34.15, 27.37, 25.31, 21.37; HRMS (ESI) calcd for C₁₉H₁₅FN₄Na [M+Na]⁺ 341.1173, found 341.1177.

2-Amino-4-(4-fluorophenyl)-4a,5,6,7-tetrahydronaph-

thalene-1,3,3(4*H*)-tricarbonitrile (**4e**): White solid, yield 88%. m.p. 264~265 °C (lit.^[3] 265~267 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.64 (s, 1H), 7.49 (s, 1H), 7.40~7.32 (m, 3H), 7.28 (t, *J*=8.0 Hz, 1H), 5.76~5.69 (m, 1H), 3.62 (d, *J*=12.5 Hz, 1H), 2.80 (t, *J*=11.4 Hz, 1H), 2.24~2.13 (m, 1H), 2.11~1.99 (m, 1H), 1.72~1.63 (m, 1H), 1.52~1.40 (m, 2H), 0.85 (q, *J*=11.3 Hz, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 163.78, 161.83, 143.88, 134.91, 131.35, 129.59, 129.22, 120.86, 116.60, 116.27, 116.10, 115.92, 115.75, 112.86, 82.04, 50.11, 43.34, 34.31, 27.43, 25.32, 21.43; HRMS (ESI) calcd for C₁₉H₁₅FN₄Na [M+Na]⁺ 341.1173, found 341.1174.

2-Amino-4-(2-nitrophenyl)-4a,5,6,7-tetrahydronaphthalene-1,3,3(4*H*)-tricarbonitrile (**4f**): White solid, yield 91%. m.p. 158~161 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 8.11~8.07 (m, 1H), 8.02 (d, *J*=7.8 Hz, 1H), 7.92 (t, *J*=7.6 Hz, 1H), 7.75 (t, *J*=7.8 Hz, 1H), 7.43 (s, 2H), 5.84~5.72 (m, 1H), 4.07 (d, *J*=12.2 Hz, 1H), 3.01 (t, *J*=11.1 Hz, 1H), 2.25~2.06 (m, 2H), 1.70 (d, *J*=8.4 Hz, 1H), 1.54~1.42 (m, 2H), 1.08~0.98 (m, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 151.37, 143.61, 134.06, 131.11, 130.15, 128.59, 128.25, 125.92, 121.67, 116.38, 112.47, 112.44, 81.94, 45.38, 42.55, 34.67, 27.07, 25.24, 21.27; HRMS (ESI) calcd for C₁₉H₁₅O₂N₅Na [M+Na]⁺ 368.1118, found 368.1117.

2-Amino-4-(3-nitrophenyl)-4a,5,6,7-tetrahydronaphthalene-1,3,3(4*H*)-tricarbonitrile (**4g**): White solid, yield 88%. m.p. 154~156 °C (lit.^[5] 156~157 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ: 8.41 (s, 1H), 8.32 (d, *J*=7.6 Hz, 1H), 8.02 (dd, *J*=9.5, 7.5 Hz, 1H), 7.81 (dt, *J*=31.0, 7.9 Hz, 1H), 7.43 (s, 2H), 5.76 (s, 1H), 3.93 (dd, *J*=24.9, 12.5 Hz, 1H), 2.92 (s, 1H), 2.20 (d, *J*=18.8 Hz, 1H), 2.07 (d, *J*=7.8 Hz, 1H), 1.67 (s, 1H), 1.45 (dd, *J*=23.4, 11.8 Hz, 2H), 0.90 (dd, *J*=24.6, 12.0 Hz, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 148.67, 148.04, 143.54, 139.32, 137.36, 134.71, 130.84, 128.85, 127.21, 124.49, 122.41, 121.15, 116.54, 112.63, 82.07, 49.94, 42.93, 33.98, 27.40, 25.28, 21.35; HRMS (ESI) calcd for C₁₉H₁₅O₂N₅Na [M+Na]⁺ 368.1118, found 368.1110.

2-Amino-4-(4-methoxyphenyl)-4a,5,6,7-tetrahydronaphthalene-1,3,3(4*H*)-tricarbonitrile (**4h**): White solid, yield 94%. m.p. 244~245 °C (lit.^[5] 244~246 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.47 (d, *J*=6.6 Hz, 1H), 7.35 (s, 2H), 7.29~7.23 (m, 3H), 5.75~5.69 (m, 1H), 3.47 (d, *J*=12.5 Hz, 1H), 2.77 (t, *J*=11.5 Hz, 1H), 2.34 (s, 3H), 2.22~2.13 (m, 1H), 2.11~2.00 (m, 1H), 1.67 (dd, *J*=9.6, 5.3 Hz, 1H), 1.51~1.39 (m, 2H), 0.89~0.78 (m, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 144.07, 138.79, 132.74, 132.06, 129.74, 129.56, 129.40, 127.22, 120.70, 116.65, 113.07, 112.89, 82.00, 50.78, 43.46, 34.37, 27.48, 25.34, 21.48, 21.21; HRMS (ESI) calcd for C₂₀H₁₉N₄ [M+H]⁺ 315.1604, found 315.1600.

2-Amino-4-(4-methoxyphenyl)-4a,5,6,7-tetrahydronaphthalene-1,3,3(4*H*)-tricarbonitrile (**4i**): White solid, yield 96%. m.p. 264~265 °C (lit.^[3] 261~262 °C); ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.50 (d, *J*=8.0 Hz, 1H), 7.34 (s, 3H), 7.02 (dd, *J*=40.6, 7.7 Hz, 2H), 5.75~5.68 (m, 1H),

3.79 (s, 3H), 3.47 (d, *J*=12.5 Hz, 1H), 2.76 (t, *J*=11.5 Hz, 1H), 2.18 (d, *J*=18.7 Hz, 1H), 2.12~1.99 (m, 1H), 1.68 (dd, *J*=9.8, 5.2 Hz, 1H), 1.54~1.39 (m, 2H), 0.90~0.79 (m, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 160.01, 144.08, 134.06, 129.45, 128.53, 126.84, 120.68, 116.67, 114.80, 114.06, 113.12, 112.97, 81.99, 55.60, 50.47, 43.63, 34.49, 27.50, 25.35, 21.50; HRMS (ESI) calcd for C₂₀H₁₈ON₄Na [M+Na]⁺ 353.1373, found 353.1370.

2-Amino-4-(3-methoxy-4-hydroxyphenyl)-4a,5,6,7-tetrahydronaphthalene-1,3,3(4*H*)-tricarbonitrile (**4j**): White solid, yield 85%. m.p. 268~272 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ: 9.23 (s, 1H), 7.32 (s, 2H), 7.15~6.71 (m, 3H), 5.71 (s, 1H), 3.77 (d, *J*=22.6 Hz, 3H), 3.41~3.35 (m, 1H), 2.74 (d, *J*=47.4 Hz, 1H), 2.18 (d, *J*=18.6 Hz, 1H), 2.05 (ddd, *J*=15.5, 7.0, 3.4 Hz, 1H), 1.69 (d, *J*=8.9 Hz, 1H), 1.60~1.38 (m, 2H), 0.87 (d, *J*=11.8 Hz, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 148.21, 147.63, 147.38, 144.16, 129.52, 125.70, 125.42, 125.42, 120.66, 119.89, 117.36, 116.70, 116.26, 115.57, 113.49, 113.12, 111.25, 81.96, 56.23, 51.17, 43.82, 34.61, 27.48, 25.37, 21.53; HRMS (ESI) calcd for C₂₀H₁₉O₂N₄ [M+H]⁺ 347.1503, found 347.1499.

2-Amino-4-(1-naphthyl)-4a,5,6,7-tetrahydronaphthalene-1,3,3(4*H*)-tricarbonitrile (**4k**): White solid, yield 88%. ¹H NMR (400 MHz, DMSO-*d*₆) δ: 8.56~8.48 (m, 1H), 8.05~7.96 (m, 2H), 7.88 (d, *J*=7.2 Hz, 1H), 7.66 (t, *J*=7.8 Hz, 1H), 7.59~7.50 (m, 2H), 7.32 (s, 2H), 5.79~5.74 (m, 1H), 4.57 (d, *J*=12.2 Hz, 1H), 2.95 (t, *J*=11.4 Hz, 1H), 2.19 (d, *J*=18.9 Hz, 1H), 2.12~1.99 (m, 1H), 1.64~1.55 (m, 1H), 1.50~1.34 (m, 2H), 0.78 (td, *J*=13.2, 2.4 Hz, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 144.48, 134.05, 133.23, 131.76, 129.71, 129.70, 129.14, 126.96, 126.49, 126.35, 125.65, 124.30, 120.66, 116.89, 113.49, 112.71, 82.30, 43.75, 43.18, 35.85, 26.96, 25.36, 21.48; HRMS (ESI) calcd for C₂₃H₁₈N₄Na [M+Na]⁺ 373.1424, found 373.1419.

2-Amino-4-(2-furyl)-4a,5,6,7-tetrahydronaphthalene-1,3,3(4*H*)-tricarbonitrile (**4l**): white solid, yield 80%. ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.80 (d, *J*=1.2 Hz, 1H), 7.38 (s, 2H), 6.63~6.51 (m, 2H), 5.73~5.65 (m, 1H), 3.82 (d, *J*=12.4 Hz, 1H), 2.69 (dd, *J*=15.1, 8.1 Hz, 1H), 2.24~2.15 (m, 1H), 2.14~1.99 (m, 1H), 1.77~1.67 (m, 1H), 1.43 (dddd, *J*=28.7, 12.5, 6.9, 3.4 Hz, 2H), 1.03 (td, *J*=13.5, 2.5 Hz, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 148.94, 144.73, 143.74, 128.75, 120.69, 116.52, 112.82, 112.71, 111.93, 111.20, 81.85, 45.91, 41.57, 34.79, 26.97, 25.20, 21.38; HRMS (ESI) calcd for C₁₇H₁₄ON₄Na [M+Na]⁺ 313.1060, found 313.1054.

2-Amino-4-(2-thenyl)-4a,5,6,7-tetrahydronaphthalene-1,3,3(4*H*)-tricarbonitrile (**4m**): White solid, yield 87%. ¹H NMR (400 MHz, DMSO-*d*₆) δ: 7.66 (d, *J*=5.0 Hz, 1H), 7.37 (s, 2H), 7.29 (d, *J*=3.3 Hz, 1H), 7.14 (dd, *J*=5.0, 3.6 Hz, 1H), 5.74~5.63 (m, 1H), 4.02 (d, *J*=12.3 Hz, 1H), 2.61 (t, *J*=11.7 Hz, 1H), 2.24~2.14 (m, 1H), 2.13~2.01 (m, 1H), 1.75~1.65 (m, 1H), 1.54~1.37 (m, 2H), 1.02~0.92 (m, 1H); ¹³C NMR (100 MHz, DMSO-*d*₆) δ: 143.83, 137.32, 129.52, 128.98, 127.76, 127.52, 120.88, 116.57,

113.25, 112.73, 81.95, 47.42, 43.81, 36.78, 27.34, 25.31, 21.45; HRMS (ESI) calcd for $C_{17}H_{14}N_4Na$ $[M+Na]^+$ 329.0831, found 329.0824.

2-Amino-4-ethyl-4a,5,6,7-tetrahydronaphthalene-1,3,3-(4*H*)-tricarbonitrile (**4n**): White solid, yield 68%. 1H NMR (400 MHz, DMSO- d_6) δ : 7.32 (s, 2H), 5.65~5.63 (m, 1H), 2.20~1.95 (m, 5H), 1.94~1.81 (m, 2H), 1.69 (ddd, J =14.9, 7.6, 3.6 Hz, 1H), 1.49~1.39 (m, 1H), 1.20 (dt, J =13.1, 6.4 Hz, 1H), 1.11 (t, J =7.5 Hz, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 143.67, 129.33, 120.41, 116.61, 114.04, 113.05, 82.09, 47.18, 41.71, 36.58, 26.78, 25.20, 24.49, 21.80, 13.48; HRMS (ESI) calcd for $C_{15}H_{16}N_4Na$ $[M+Na]^+$ 275.1267, found 275.1264.

2-Amino-4-isopropyl-4a,5,6,7-tetrahydronaphthalene-1,3,3-(4*H*)-tricarbonitrile (**4o**): White solid, yield 71%. 1H NMR (400 MHz, DMSO- d_6) δ : 7.25 (s, 2H), 5.75~5.59 (m, 1H), 2.35~2.25 (m, 2H), 2.21~2.04 (m, 4H), 1.87~1.78 (m, 1H), 1.47 (qd, J =11.4, 6.6 Hz, 1H), 1.29~1.20 (m, 4H), 1.18 (d, J =7.3 Hz, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 143.95, 129.52, 120.89, 116.66, 114.27, 113.59, 82.14, 49.67, 40.67, 34.25, 30.41, 28.09, 25.16, 23.41, 21.85, 17.12; HRMS (ESI) calcd for $C_{16}H_{18}N_4Na$ $[M+Na]^+$ 289.1424, found 289.1418.

5-Amino-1,2,7,7a-tetrahydro-7-phenylindene-4,6,6-tricarbonitrile (**4p**): White solid, yield 78%. 1H NMR (400 MHz, DMSO- d_6) δ : 7.69 (s, 2H), 7.55 (d, J =6.6 Hz, 2H), 7.51~7.40 (m, 3H), 5.50 (d, J =2.1 Hz, 1H), 3.64 (d, J =12.6 Hz, 1H), 3.40~3.27 (m, 1H), 2.43~2.26 (m, 2H), 1.93~1.82 (m, 1H), 1.18 (dq, J =20.1, 9.9 Hz, 1H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 146.59, 135.67, 135.04, 130.10, 129.53, 129.03, 119.87, 116.40, 112.81, 112.68, 77.59, 51.37, 44.21, 42.96, 31.45, 29.78; HRMS (ESI) calcd for $C_{18}H_{14}N_4Na$ $[M+Na]^+$ 309.1111, found 309.1107.

5-Amino-1,2,7,7a-tetrahydro-7-(4-methoxyphenyl)indene-4,6,6-tricarbonitrile (**4q**): White solid, yield 85%. 1H NMR (400 MHz, DMSO- d_6) δ : 7.66 (s, 2H), 7.45 (d, J =8.6 Hz, 2H), 7.01 (d, J =8.6 Hz, 2H), 5.48 (d, J =1.6 Hz, 1H), 3.78 (s, 3H), 3.56 (d, J =12.6 Hz, 1H), 3.31~3.22 (m, 1H), 2.43~2.22 (m, 2H), 1.91~1.83 (m, 1H), 1.22~1.10 (m, 1H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 160.20, 146.64, 135.79, 131.27, 126.81, 119.76, 116.43, 114.43, 112.96, 112.78, 77.57, 55.62, 50.90, 44.50, 43.13, 31.43, 29.81; HRMS (ESI) calcd for $C_{19}H_{16}ON_4Na$ $[M+Na]^+$ 339.1216, found 339.1207.

5-Amino-1,2,7,7a-tetrahydro-7-(2-chlorophenyl)indene-4,6,6-tricarbonitrile (**4r**): white solid, yield 80%. 1H NMR (400 MHz, DMSO- d_6) δ : 7.78 (d, J =6.8 Hz, 1H), 7.74 (s, 2H), 7.65~7.59 (m, 1H), 7.57~7.45 (m, 2H), 5.53 (d, J =2.1 Hz, 1H), 3.98 (d, J =12.5 Hz, 1H), 3.45~3.35 (m, 1H), 2.40~2.27 (m, 2H), 1.85~1.75 (m, 1H), 1.17~1.04 (m, 1H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 146.49, 135.48, 135.39, 132.33, 131.24, 130.56, 129.99, 128.40, 120.44, 116.23, 112.48, 111.98, 77.54, 47.51, 43.93, 42.91, 31.37, 29.80; HRMS (ESI) calcd for $C_{18}H_{13}ClN_4Na$ $[M+Na]^+$ 343.0721, found 343.0717.

5-Amino-1,2,7,7a-tetrahydro-7-(4-fluorophenyl)indene-4,6,6-tricarbonitrile (**4s**): white solid, yield 76%. 1H NMR

(400 MHz, DMSO- d_6) δ : 7.69 (s, 2H), 7.60 (dd, J =8.4, 5.5 Hz, 2H), 7.32 (t, J =8.8 Hz, 2H), 5.49 (d, J =1.7 Hz, 1H), 3.72 (d, J =12.6 Hz, 1H), 3.46~3.22 (m, 1H), 2.43~2.24 (m, 2H), 1.94~1.81 (m, 1H), 1.23~1.10 (m, 1H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 163.94, 161.99, 160.67, 146.44, 135.57, 134.04, 132.27, 131.31, 119.92, 117.51, 117.33, 116.36, 116.07, 115.90, 112.77, 112.60, 77.62, 50.50, 44.20, 43.03, 31.45, 29.70; HRMS (ESI) calcd for $C_{18}H_{13}FN_4Na$ $[M+Na]^+$ 327.1016, found 327.1007.

5-Amino-1,2,7,7a-tetrahydro-7-(1-naphthyl)indene-4,6,6-tricarbonitrile (**4t**): white solid, yield 88%. 1H NMR (400 MHz, DMSO- d_6) δ : 8.62~8.57 (m, 1H), 8.06~7.96 (m, 2H), 7.88 (d, J =7.2 Hz, 1H), 7.70~7.60 (m, 3H), 7.60~7.50 (m, 2H), 5.52 (d, J =2.1 Hz, 1H), 4.69 (d, J =12.2 Hz, 1H), 3.49 (dd, J =17.9, 9.7 Hz, 1H), 2.41~2.19 (m, 2H), 1.87~1.73 (m, 1H), 1.07 (dq, J =20.1, 9.9 Hz, 1H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 147.16, 136.21, 133.97, 133.15, 131.75, 129.84, 129.03, 126.83, 126.75, 126.50, 125.71, 124.57, 119.64, 116.66, 113.14, 112.70, 77.81, 44.59, 44.42, 44.02, 31.28, 29.78; HRMS (ESI) calcd for $C_{22}H_{16}N_4Na$ $[M+Na]^+$ 359.1267, found 359.1257.

5-Amino-1,2,7,7a-tetrahydro-7-(2-thenyl)indene-4,6,6-tricarbonitrile (**4u**): white solid, yield 73%. 1H NMR (400 MHz, DMSO- d_6) δ : 7.75~7.62 (m, 3H), 7.31 (d, J =3.2 Hz, 1H), 7.14 (dd, J =5.0, 3.6 Hz, 1H), 5.50 (d, J =2.1 Hz, 1H), 4.08 (d, J =12.4 Hz, 1H), 3.15 (dd, J =17.8, 10.1 Hz, 1H), 2.43~2.24 (m, 2H), 2.01~1.89 (m, 1H), 1.30 (dq, J =20.2, 9.9 Hz, 1H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 146.23, 137.30, 135.29, 129.42, 127.92, 127.51, 120.33, 116.34, 112.90, 112.81, 77.57, 47.84, 45.10, 44.78, 31.32, 30.03; HRMS (ESI) calcd for $C_{16}H_{12}N_4Na$ $[M+Na]^+$ 315.0675, found 315.0667.

Supporting Information 1H NMR and ^{13}C NMR spectra of compounds **4a**~**4u** are available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Tang, M.; Wu, Y.; Liu, Y.; Cai, M.; Xia, F.; Liu, S.; Hu, W.; Wu, Y. *Acta Chim. Sinica* **2016**, *74*, 54 (in Chinese). (唐敏, 吴永, 刘源, 蔡茂强, 夏飞, 刘顺英, 胡文浩, 化学学报, **2016**, *74*, 54.) (b) Yu, S.; Fu, X.; Liu, G.; Qiu, H.; Hu, W. *Acta Chim. Sinica* **2018**, *76*, 895 (in Chinese). (余思凡, 傅祥, 刘耿鑫, 邱晃, 胡文浩, 化学学报, **2018**, *76*, 895.) (c) Lu, L.-H.; Zhou, S.-J.; Sun, M.; Chen, J.-L.; Xia, W.; Yu, X.; Xu, X.; He, W.-M. *ACS Sustainable Chem. Eng.* **2019**, *7*, 1574. (d) Wu, C.; Xiao, H.-J.; Wang, S.-W.; Tang, M.-S.; Tang, Z.-L.; Xia, W.; Li, W.-F.; Cao, Z.; He, W.-M. *ACS Sustainable Chem. Eng.* **2019**, *7*, 2169. (e) Wu, C.; Hong, L.; Shu, H.; Zhou, Q.-H.; Wang, Y.; Su, N.; Jiang, S.; Cao, Z.; He, W.-M. *ACS Sustainable Chem. Eng.* **2019**, *7*, 8798. (f) Wang, Z.; Yang, L.; Liu, H.-L.; Bao, W.-H.; Tan, Y.-Z.; Wang, M.; Tang, Z.; He, W.-M. *Chin. J. Org. Chem.* **2018**, *38*, 2639 (in Chinese). (王峥, 杨柳, 刘慧兰, 谭英芝, 包文虎, 汪明, 唐子龙, 何卫民, 有机化学, **2018**, *38*, 2639.)

- (g) Liu, Q.; Wang, L.; Yue, H.; Li, J.-S.; Luo, Z.; Wei, W. *Green Chem.* **2019**, *21*, 1609.
- (h) Wei, W.; Cui, H.; Yue, H.; Yang, D. *Green Chem.* **2018**, *20*, 3197.
- (i) Wang, L.; Bao, P.; Liu, W.; Liu, S.; Hu, C.; Yue, H.; Yang, D.; Wei, W. *Chin. J. Org. Chem.* **2018**, *38*, 3189 (in Chinese).
- (王雷雷, 鲍鹏丽, 刘维伟, 刘思彤, 胡昌松, 岳会兰, 杨道山, 魏伟, 有机化学, **2018**, *38*, 3189.)
- (j) Yue, H.; Bo, P.; Wang, B. L.; Lv, X.; Yang, D.; Wei, W. *Chin. J. Org. Chem.* **2019**, *39*, 463 (in Chinese).
- (岳会兰, 鲍鹏丽, 王雷雷, 吕晓霞, 杨道山, 王桦, 魏伟, 有机化学, **2019**, *39*, 463.)
- (k) Xie, L.-Y.; Peng, S.; Fan, T.-G.; Liu, Y.-F.; Sun, M.; Jiang, L.-L.; Wang, X.-X.; Cao, Z.; He, W.-M. *Sci. China Chem.* **2019**, *62*, 460.
- (l) Li, G.; Gan, Z.; Kong, K.; Dou, X.; Yang, D. *Adv. Synth. Catal.* **2019**, *361*, 1808.
- (m) Zong, Y.; Lang, Y.; Yang, M.; Li, X.; Fan, X.; Wu, J. *Org. Lett.* **2019**, *21*, 1935.
- [2] (a) Tan, J.; Guo, Y.; Zeng, F.; Chen, G.; Xie, L.; He, W. *Chin. J. Org. Chem.* **2018**, *38*, 1740 (in Chinese).
- (谭家希, 郭也, 曾飞, 陈观荣, 谢龙勇, 何卫民, 有机化学, **2018**, *38*, 1740.)
- (b) Sha, H.-K.; Liu, F.; Lu, J.; Liu, Z.-Q.; Hao, W.-J.; Tang, J.-L.; Tu, S.-J.; Jiang, B. *Green Chem.* **2018**, *20*, 3476.
- (c) Fan, W.; Chen, K.-Y.; Chen, Q.-P.; Li, G.; Jiang, B. *Org. Biomol. Chem.* **2017**, *15*, 6493.
- (d) Xie, L.-Y.; Peng, S.; Tan, J.-X.; Sun, R.-X.; Yu, X.; Dai, N.-N.; Tang, Z.-L.; Xu, X.; He, W.-M. *ACS Sustainable Chem. Eng.* **2018**, *6*, 16976.
- (e) Xie, L.-Y.; Peng, S.; Liu, F.; Liu, Y.-F.; Sun, M.; Tang, Z.-L.; Jiang, S.; Cao, Z.; He, W.-M. *ACS Sustainable Chem. Eng.* **2019**, *7*, 7193.
- (g) Liu, K.-J.; Fu, Y.-L.; Xie, L.-Y.; Wu, C.; He, W.-B.; Peng, S.; Wang, Z.; Bao, W.-H.; Cao, Z.; Xu, X.; He, W.-M. *ACS Sustainable Chem. Eng.* **2018**, *6*, 4916.
- [3] (a) Enders, D.; Huettl, M. R. M.; Grondal, C.; Raabe, G. *Nature* **2006**, *441*, 861.
- (b) Padwa, A. *Chem. Soc. Rev.* **2009**, *38*, 3072.
- (c) Takumi, M.; Noriaki, O.; Takatoshi, I.; Toshiyuki, M. *Tetrahedron Lett.* **2000**, *41*, 1051.
- (d) Takumi, M.; Yoshio, I. *Tetrahedron* **2002**, *58*, 3155.
- (e) Saddiqa, A.; Bukhari, I. H.; Tahir, M. N.; Ahmad, M.; Shafiq, M.; Arshad, M. N. *Asian J. Chem.* **2015**, *27*, 969.
- [4] (a) Wan, Y.; Zhang, X.; Zhao, L.; Wang, C.; Chen, L.; Liu, G.; Huang, S.; Yue, S.; Zhang, W.; Wu, H. *J. Heterocycl. Chem.* **2015**, *52*, 623.
- (b) Lohar, T.; Kumbhar, A.; Barge, M.; Salunkhe, R. *J. Mol. Liq.* **2016**, *224*, 1102.
- (c) Gaikwad, D. S.; Undale, K. A.; Patil, D. B.; Patravale, A. A.; Kamble, A. A. *J. Iran. Chem. Soc.* **2018**, *15*, 1175.
- (d) Wang, X. S.; Wu, J. R.; Zhou, J.; Zhang, M. M. *J. Heterocycl. Chem.* **2011**, *48*, 1056.
- (e) Zhang, M. M.; Wu, J. R.; Zhou, J.; Wang, X. S. *Synth. Commun.* **2012**, *42*, 599.
- [5] (a) Mojtahedi, M. M.; Pourabdi, L.; Abaee, M. S.; Jami, H.; Dini, M.; Halvagar, M. R. *Tetrahedron* **2016**, *72*, 1699.
- (b) Elinson, M. N.; Ilovaisky, A. I.; Merkulova, V. M.; Barba, F.; Batanero, B. *Tetrahedron* **2013**, *69*, 7125.
- (c) Khan, M. N.; Pal, S.; Karamthulla, S.; Choudhury, L. H. *RSC Adv.* **2014**, *4*, 3732.
- (d) Wang, J.; Li, Q.; Qi, C.; Liu, Y.; Ge, Z.; Li, R. *Org. Biomol. Chem.* **2010**, *8*, 4240.
- (e) Babu, T. H.; Joseph, A. A.; Muralidharan, D.; Perumal, P. T. *Tetrahedron Lett.* **2010**, *51*, 994.
- (f) Elinson, M. N.; Vereshchagin, A. N.; Nasybullin, R. F.; Bobrovsky, S. I.; Ilovaisky, A. I.; Merkulova, V. M.; Bushmarinov, I. S.; Egorov, M. P. *RSC Adv.* **2015**, *5*, 50421.
- [6] Das, P.; Butcher, R. J.; Mukhopadhyay, C. *Green Chem.* **2012**, *14*, 1376.
- [7] Zhang, L.; Wan, Y.; Zhang, X.; Cui, H.; Zou, H.; Zhou, Q. Wu, H. *Tetrahedron Lett.* **2015**, *56*, 4934.
- [8] Maleki, B.; Rooky, R.; Rezaei-Seresht, E.; Tayeb R. *Org. Prep. Proced. Int.* **2017**, *49*, 557.
- [9] (a) Wang, X.; Zhang, M.; Li, Q.; Yao, C.; Tu, S. *Tetrahedron* **2007**, *63*, 5265.
- (b) Wang, X. S.; Zhou, J.; Yang, K.; Zhang, M. M. *Synth. Commun.* **2010**, *40*, 1065.
- [10] Molla, A.; Hussain, S. *RSC Adv.* **2014**, *4*, 29750.
- [11] Azizi, N.; Ahoioie, T. S.; Hashemi, M. M.; Azizi, N.; Ahoioie, T. S.; Hashemi, M. M. *J. Mol. Liq.* **2017**, *246*, 221.
- [12] Huang, X. F.; Zhang, Y. F.; Qi, Z. H.; Li, N. K.; Geng, Z. C.; Li, K.; Wang, X. W. *Org. Biomol. Chem.* **2014**, *12*, 4372.
- [13] For selected reactions catalyzed by DABCO see: (a) Maher, D.; Connon, S. *Tetrahedron Lett.* **2004**, *45*, 1301.
- (b) Luca, C.; Francesco, D. S.; Fabrizio, M. *Eur. J. Org. Chem.* **2006**, 4852.
- (c) Baidya, M.; Kobayashi, S.; Brotzel, F.; Schmidhammer, U.; Riedle, E.; Mayr, H. *Angew. Chem., Int. Ed.* **2007**, *46*, 6176.
- (d) Chandrasekhar, S.; Narsihmulu, C.; Saritha, B.; Sultana, S. *Tetrahedron Lett.* **2004**, *45*, 5865.
- (e) Murthy, S. N.; Madhav, B.; Nageswar, Y. V. D. *Tetrahedron Lett.* **2010**, *51*, 5252.
- (f) Zhang, W.; Xu, H. D.; Xu, H.; Tang, W. P. *J. Am. Chem. Soc.* **2009**, *131*, 3832.
- (g) Meshram, H. M.; Kumar, G. S.; Ramesh, P.; Reddy, B. C. *Tetrahedron Lett.* **2010**, *51*, 2580.
- (h) Fan, M.; Li, G.; Liang, Y. *Tetrahedron* **2006**, *62*, 6782.
- (i) Wu, J.; Sun, X. Y.; Li, Y. Z. *Eur. J. Org. Chem.* **2005**, 4271.
- [14] Li, M.; Zhou, Z.; Wen, L.; Qiu, Z. *J. Org. Chem.* **2011**, *76*, 3054.
- [15] Singh, M. S.; Nandi, G. C.; Samai, S. *Green Chem.* **2012**, *14*, 447.
- [16] Liu, Y. F.; Du, Y. L.; Yu, A. M.; Mu, H. F.; Meng, X. T. *Org. Biomol. Chem.* **2016**, *14*, 1226.
- [17] Chinchkar, S. M.; Patil, J. D.; Korade, S. N.; Gokavi, G. S.; Shejwal, R. V.; Pore, D. M. *Lett. Org. Chem.* **2017**, *14*, 403.
- [18] (a) Dong, D.-Q.; Chen, W.-J.; Chen, D.-M.; Li, L.-X.; Li, G.-H.; Wang, Z.-L.; Deng, Q.; Long, S. *Chin. J. Org. Chem.* **2019**, *39*, DOI: 10.6023/cjoc201904070 (in Chinese).
- (董道青, 陈文静, 陈德茂, 李丽霞, 李光辉, 王祖利, 邓企, 龙妹, 有机化学, **2019**, *39*, DOI: 10.6023/cjoc201904070.)
- (b) Dong, D.-Q.; Li, L.-X.; Li, G.-H.; Deng, Q.; Wang, Z.-L.; Long, S. *Chin. J. Catal.* **2019**, 1494.
- (c) Xu, X.-M.; Chen, D.-M.; Wang, Z.-L. *Chin. J. Org. Chem.* **2019**, *39*, DOI: 10.6023/cjoc201904068 (in Chinese).
- (徐鑫明, 陈德茂, 王祖利, 有机化学, **2019**, *39*, DOI: 10.6023/cjoc201904068.)
- (d) Hao, S. H.; Zhang, X. Y.; Dong, D. Q.; Wang, Z. L. *Chin. Chem. Lett.* **2015**, *26*, 599.
- (e) Zhang, H.; Dong, D.-Q.; Hao, S.-H.; Wang, Z. L. *RSC Adv.* **2016**, *6*, 8465.
- (f) Yan, S. Q.; Ding, N.; Zhang, W.; Wang, P.; Li, Y. X.; Li, M. *Carbohydr. Res.* **2012**, *354*, 6.
- (g) Chun, Y. X.; Yan, S. Q.; Li, X. P.; Ding, N.; Zhang, W.; Wang, P.; Li, M.; Li, Y. X. *Tetrahedron Lett.* **2011**, *52*, 6196.
- (h) Yan, S. Q.; Ding, N.; Zhang, W.; Wang, P.; Li, Y. X.; Li, M. *J. Carbohydr. Chem.* **2012**, *31*, 571.
- (i) Hao, S.; Li, L.; Dong, D. Q.; Wang, Z. L.; Yu, X. *Tetrahedron Lett.* **2018**, *59*, 4073.
- (j) Li, G. H.; Dong, D. Q.; Yang, Y.; Yu, X. Y.; Wang, Z. L. *Adv. Synth. Catal.* **2019**, *361*, 832.
- (k) Li, G.-H.; Dong, D.-Q.; Yu, X.-Y.; Wang, Z.-L. *New J. Chem.* **2019**, *43*, 1667.
- (l) Dong, D.-Q.; Chen, W.-J.; Yang, Y.; Gao, X.; Wang, Z.-L. *ChemistrySelect* **2019**, *4*, 2480.

(Lu, Y.)