3-去羟基 Phomonol 的合成

叶青青^a 张梦帆^a 刘耀宗^b 杨震^{*a}^(a) 绍兴文理学院 浙江省精细化工与传统工艺绿色替代技术重点实验室 绍兴 312000^(b) 甘肃银光聚银化工有限公司 白银 730900

摘要 以商业购买的(*R*)-3-羟基-4-氯丁酸乙酯或者(*S*)-1,2-环氧戊烷为原料, 经过 6 步反应, 完成了天然产物类似物 3-去羟基 phomonol 的合成, 关键步骤包括格氏反应, Prins 环化反应和钯催化分子内烷氧羰基化反应等, 无保护基合成策略的应用缩短了合成步骤, 提高了合成的效率。

关键词 合成; 3-去羟基 phomonol; Prins 环化反应; 钯催化分子内烷氧羰基化反应

Synthesis of 3-Dehydroxyphomonol

Ye, Qingqing^a Zhang, Mengfan^a Liu, Yaozong^b Yang, Zhen^{*a}^(a) Zhejiang Key Laboratory of Alternative Technologies for Fine Chemicals Process, Shaoxing University, Shaoxing 312000^(b) Gansu Yinguang Juyin Chemical Co., Ltd, Baiyin, Gansu 730900

Abstract A concise synthesis of 3-dehydroxyphomonol has been accomplished in six steps from commercially available (*R*)-4-chloro-3-hydroxy-butyric acid ethyl ester or (*S*)-2-propyl-oxirane. The key steps involved Grignard reaction, Prins cyclization and palladium-catalyzed intramolecular alkoxy carbonylation to install the tetrahydropyran ring. The synthesis demonstrated an application of protecting-group-free strategy.

Keywords synthesis; 3-dehydroxyphomonol; Prins cyclization; palladium-catalyzed intramolecular alkoxy carbonylation

1 Introduction

Functionalized tetrahydropyran rings are very important building blocks available in natural products, such as neopeltolide,^[1] cyanolide A,^[2] aspergillide A^[3] and mandelalide A.^[4] During the last several decades, many efforts and contributions had been made for the construction of tetrahydropyrans in total synthesis of natural products, including Prins cyclization,^[5] oxo-Michael addition,^[6] radical cyclization,^[7] transition metal catalyzed cyclization^[8] and hetero-Diels-Alder reaction.^[9]

Phomonol (**1**), a tetrasubstituted tetrahydropyran natural product, was isolated from the endophytic fungal strain *Phomopsis* sp. A123 (Figure 1).^[10] This strain was isolated from leaves of the mangrove species, *Kandelia candel*, collected in the Fugong Mangrove Conservation Area, Fujian, China. Its highly substituted tetrahydropyran ring has attracted many research groups.^[11] We have revealed the protecting-group-free synthesis of 4-*epi*-phomonol in our previous work.^[11f] Herein, the synthesis of

3-dehydroxyphomonol (**2**), which might helpful for further biological studies of phomonol and analogues, was reported.

Figure 1 Structures of phomonol (**1**) and 3-dehydroxyphomonol (**2**)

2 Results and discussion

Our retrosynthetic analysis of 3-dehydroxyphomonol is depicted in Scheme 1. The methyl ketone could be established by Grignard reaction from Weinreb amide **3**, while the tetrahydropyran ring was constructed by Prins cyclization from alcohol **4**, which could be easily derived from commercially available (*R*)-4-chloro-3-hydroxy-butyric acid ethyl ester **5**.

* Corresponding author. E-mail: yangzhen09@usx.edu.cn

Received March 8, 2019; revised April 18, 2019; published online April 26, 2019.

Project supported by the National Natural Science Foundation of China (No. 21302129) and the Natural Science Foundation of Zhejiang Province (No. LQ13B020002).

国家自然科学基金(No. 21302129)和浙江省自然科学基金(No. LQ13B020002)资助项目。

Scheme 1 Retrosynthetic analysis of 3-dehydroxyphomol

According to the retrosynthetic analysis, we began our synthesis of 3-dehydroxyphomol from commercially available (*R*)-4-chloro-3-hydroxy-butyric acid ethyl ester **5** (Scheme 2). Iodination of **5** afforded ethyl ester **6** under the condition of sodium iodide in acetone.^[12] Epoxidation of ester **6** was then occurred in the presence of silver(I) oxide, and the corresponding epoxy-ester **7** was obtained in 80% yield.^[13] Treating **7** with vinylmagnesium bromide obtained alcohol-ester **8**, which was converted to Weinreb amide **4** in the presence of *N*-methoxymethyl amine. Subjecting **4** to the Prins cyclization, followed by trifluoroacetyl deprotection, afforded the desired tetrahydropyran **3**. Treatment of **3** with methylmagnesium iodide obtained 3-dehydroxyphomol (**2**) in 85% yield.

Scheme 2 Synthesis of 3-dehydroxyphomol (**2**)

The synthesis strategy aforementioned was elegant and useful, but also showed obvious shortcomings, such as the use of expensive silver oxide with equivalent amount. Therefore, we developed an alternative route for the synthesis of intermediate **3** from commercially available (*S*)-2-propyl-oxirane. The retrosynthetic analysis of **3** was outlined in Scheme 3. The Weinreb amide **3** was achieved

from diol **9**, exploiting the palladium-catalyzed intramolecular alkoxy carbonylation. Diol **9** could be easily prepared from (*S*)-2-propyl-oxirane.

Scheme 3 Retrosynthetic analysis of intermediate **3**

Synthesis of intermediate **3** was begun with (*S*)-2-propyl-oxirane (Scheme 4). A three-step sequence including ring opening of **10**, ozonolysis of **11**, followed by Barbier-type reaction^[14] smoothly generated diol **9** in 53% yield with moderate stereoselectivity (*anti/cis*=4/1), and the desired isomer was separated by column chromatography. To construct the tetrahydropyran ring, a palladium-catalyzed intramolecular alkoxy carbonylation was applied to the diol **9**, providing the desired methyl ester **13** in good yield.^[14] Transformation of methyl ester **13** to Weinreb **3** ended the alternative synthesis route.

Scheme 4 An alternative synthesis of intermediate **3**

3 Conclusions

In conclusion, a concise synthesis of 3-dehydroxyphomol was achieved in 6 steps from commercially available materials. The synthesis of tetrahydropyran ring by Prins cyclization and palladium-catalyzed cyclization was highlighted. The synthesis might be helpful for the further biological studies toward phomol and analogues.

4 Experimental section

4.1 Instruments and reagents

Oxygen- and moisture-sensitive reactions were carried out under argon atmosphere. Solvents were purified and

dried by standard methods prior to use. All commercially available reagents were used without further purification unless otherwise noted. Column chromatography was performed on silica gel (200~300 mesh). Optical rotations were measured on a precision automated polarimeter. Infrared spectra were recorded on a 670 FT-IR spectrometer. ^1H NMR and ^{13}C NMR spectra were recorded on a 400 MHz spectrometers. Chemical shifts are reported as δ values relative to internal chloroform (δ 7.27 for ^1H NMR and 77.0 for ^{13}C NMR).

4.2 Experimental method

4.2.1 Synthesis of (*R*)-3-hydroxy-4-iodo-butyric acid ethyl ester (**6**)

To a solution of **5** (2.0 g, 12.0 mmol) in acetone (30 mL) was added sodium iodide (7.3 g, 48.2 mmol) in one portion. The reaction was proceeded in a sealed tube under 100 °C for 6 h. The mixture was cooled down to room temperature and solvents were evaporated, then water (30 mL) was added and the mixture was extracted with ethyl acetate twice (50 mL $\times$ 2). The combined organic phase was washed with saturated aq. Na_2SO_3 , and dried over anhydrous Na_2SO_4 . The solvent was removed, and the residue was purified by flash column chromatography with EtOAc/hexane ($V:V=1:20$) to give **6** (2.49 g, 89%) as a pale yellow oil. $[\alpha]_{\text{D}}^{20} +12$ (c 0.5, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 4.16 (dd, $J=12.0, 8.0$ Hz, 2H), 3.99~3.98 (m, 1H), 3.35~3.25 (m, 3H), 2.68~2.55 (m, 2H), 1.26 (t, $J=8.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 171.7, 67.4, 61.0, 40.7, 14.1, 12.0; IR (KBr) ν_{max} : 3434, 2981, 2925, 1726, 1184, 1024 cm^{-1} ; HRMS (ESIMS) calcd for $\text{C}_{16}\text{H}_{12}\text{ClO}_3$ $[\text{M} + \text{H}]^+$ 167.0469, found 167.0465.

4.2.2 Synthesis of (*R*)-oxiranyl-acetic acid ethyl ester (**7**)

To a solution of **6** (2.3 g, 8.91 mmol) in CH_3CN (30 mL) was added silver oxide (2.47 g, 10.7 mmol) in one portion. The mixture was stirred at room temperature for 16 h in dark. The suspension was filtered through a celite pad and the filtrate was evaporated. The residue was purified by flash column chromatography with EtOAc/hexane ($V:V=1:10$) to give **7** (1.57 g, 82.2%) as a colorless oil. $[\alpha]_{\text{D}}^{20} +9$ (c 0.1, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 4.20~4.13 (m, 2H), 3.26 (d, $J=4.0$ Hz, 1H), 2.83~2.81 (m, 1H), 2.62~2.54 (m, 3H), 1.25 (t, $J=8.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 170.0, 60.5, 47.6, 46.3, 37.7, 13.7; IR (KBr) ν_{max} : 2926, 1735, 1266, 1185, 1030 cm^{-1} ; HRMS (ESIMS) calcd for $\text{C}_6\text{H}_{11}\text{O}_3$ $[\text{M} + \text{H}]^+$ 131.0703, found 131.0696.

4.2.3 Synthesis of (*S*)-3-hydroxy-hex-5-enoic acid ethyl ester (**8**)

To a suspension of copper iodide (**I**) (82 mg, 0.43 mmol) and **7** (0.5 g, 3.9 mmol) in THF (20 mL) was slowly added vinylmagnesium bromide (1.0 mol $\cdot\text{L}^{-1}$, 6.8 mL, 6.8 mmol) at -40 °C under nitrogen. The reaction was stirred for 3 h at the temperature and then quenched by

saturated NH_4Cl . The mixture was extracted with ethyl acetate twice (50 mL $\times$ 2). The combined organic phase was washed with brine and dried over anhydrous Na_2SO_4 . The solvent was removed, and the residue was purified by flash column chromatography with EtOAc/hexane ($V:V=1:10$) to give **8** (0.55 g, 90%) as a colorless oil. $[\alpha]_{\text{D}}^{20} +35$ (c 0.28, CHCl_3); ^1H NMR (400MHz, CDCl_3) δ : 5.81~5.74 (m, 1H), 5.10~5.06 (m, 2H), 4.12 (dd, $J=16.0, 8.0$ Hz, 2H), 4.05~4.03 (m, 1H), 2.99 (br, 1H), 2.49~2.34 (m, 2H), 2.26~2.21 (m, 2H), 1.22 (t, $J=8.0, 3\text{H}$); ^{13}C NMR (100 MHz, CDCl_3) δ : 173.4, 134.5, 118.7, 67.9, 61.3, 41.5, 41.2, 14.7; IR (KBr) ν_{max} : 3438, 2926, 1727, 1642, 1374, 1265, 1175, 1029 cm^{-1} ; HRMS (ESIMS) calcd for $\text{C}_8\text{H}_{15}\text{O}_3$ $[\text{M} + \text{H}]^+$ 159.1016, found 159.1013.

4.2.4 Synthesis of (*S*)-*N*-methoxy-*N*-methyl-3-hydroxyhex-5-enoic acid amide (**4**)

To a suspension of *N,O*-dimethylhydroxylamine hydrochloride (0.7 g, 7.2 mmol) and **8** (0.28 g, 1.8 mmol) in THF (20 mL) was slowly added $i\text{-PrMgCl}$ (2.0 mol $\cdot\text{L}^{-1}$, 7.2 mL, 14.4 mmol) at -30 °C under nitrogen. The reaction was stirred for 3 h, and then quenched by saturated NH_4Cl . The mixture was extracted with ethyl acetate twice (30 mL $\times$ 2). The combined organic phase was washed with brine and dried over anhydrous Na_2SO_4 . The solvent was removed, and the residue was purified by flash column chromatography with EtOAc/hexane ($V:V=1:10$) to give **4** (0.27 g, 86%) as a colorless oil. $[\alpha]_{\text{D}}^{20} +70$ (c 0.5, CHCl_3); ^1H NMR (400MHz, CDCl_3) δ : 5.87~5.81 (m, 1H), 5.14~5.08 (m, 2H), 4.10~4.07 (m, 2H), 3.71 (br, 1H), 3.67 (s, 3H), 3.17 (s, 3H), 2.68~2.43 (m, 2H), 2.36~2.22 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ : 173.6, 134.4, 117.5, 67.3, 61.2, 40.8, 37.4, 31.8; IR (KBr) ν_{max} : 3433, 2921, 1636, 1420, 1388, 1175, 1058 cm^{-1} ; HRMS (ESIMS) calcd for $\text{C}_8\text{H}_{16}\text{NO}_3$ $[\text{M} + \text{H}]^+$ 174.1125, found 174.1119.

4.2.5 Synthesis of (2*S*,4*R*,6*S*)-2-(4-hydroxy-6-propyl-tetrahydropyran-2-yl)-*N*-methoxy-*N*-methyl-acetamide (**3**)

To a solution of **4** (0.35 g, 2.0 mmol) and *n*-butanal (0.22 g, 3.0 mmol) in dichloromethane (20 mL) was slowly added $\text{CF}_3\text{CO}_2\text{H}$ (1.48 mL, 20.0 mmol) at room temperature under nitrogen. The mixture was stirred for 3 h, and then K_2CO_3 (3.45 g, 25.0 mmol) in 10 mL of MeOH was added. The mixture stirred for another 2 h, and then extracted with ethyl acetate twice (30 mL $\times$ 2). The combined organic phase was washed with brine and dried over anhydrous Na_2SO_4 . The solvent was removed, and the residue was purified by flash column chromatography with EtOAc/hexane ($V:V=1:3$) to give **3** (0.39 g, 80%) as a colorless oil. $[\alpha]_{\text{D}}^{20} +17$ (c 0.35, CHCl_3); ^1H NMR (400MHz, CDCl_3) δ : 3.81~3.71 (m, 2H), 3.66 (s, 3H), 3.31~3.28 (m, 1H), 3.15 (s, 3H), 2.86~2.80 (m, 1H), 2.41 (dd, $J=16.0, 4.0$ Hz, 2H), 2.04~2.00 (m, 1H), 1.88~1.84 (m, 1H), 1.47~1.41 (m, 4H), 0.85 (t, $J=6.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 171.9, 75.5, 72.2, 68.0, 61.4, 41.2, 40.9, 38.2, 32.1, 29.7, 18.8, 14.1; IR (KBr) ν_{max} :

3433, 2921, 1636, 1420, 1388, 1175, 1058 cm^{-1} ; HRMS (ESIMS) calcd for $\text{C}_{12}\text{H}_{24}\text{NO}_4$ $[\text{M}+\text{H}]^+$ 246.1700, found 246.1691.

4.2.6 Synthesis of (4*S*,6*S*)-non-1-ene-4,6-diol (**9**)

To a suspension of copper iodide (**1**) (0.38 g, 2.0 mmol) and **10** (1.72 g, 20.0 mmol) in THF (20 mL) was slowly added vinylmagnesium bromide ($1.0 \text{ mol} \cdot \text{L}^{-1}$, 25 mL, 25.0 mmol) at -30°C under nitrogen. The reaction was stirred for 6 h and quenched by saturated NH_4Cl . The mixture was stirred for another 3 h and extracted with ether (50 mL $\times$ 2). The combined organic phase was washed with saturated aq. NaCl and dried over anhydrous Na_2SO_4 . The solvent was removed at 10°C to give alcohol **11**. The residue was used for next step without further purification.

The residue **11** was dissolved in dichloromethane (90 mL) and the solution was bubbled with ozone at -78°C until a deep blue colour persisted. PPh_3 (7.8 g, 30.0 mmol) was then slowly added, the mixture was stirred for 4 h at room temperature, and the solvent was removed. The residue was dissolved in Et_2O (50 mL), vigorously stirred for 1 h. The suspension was filtered and the filtrate was evaporated to give **12**, which was used for next step without further purification.

The residue of **12** and allylic bromide (4.8 g, 40.0 mmol) were dissolved in THF (30 mL), followed by addition of NH_4Cl (30 mL). The resulting mixture was cooled to -20°C , and activated zinc dust (2.0 g, 60.0 mmol) was slowly added over 30 min. The mixture was then stirred vigorously at room temperature for 4 h, filtered through a pad of Celite. The filtrate was extracted with diethyl ether. The combined organic fractions were washed with brine, dried over anhydrous Na_2SO_4 , filtered, and the solvent was removed. Flash chromatography (EtOAc /hexane, $V:V=1:5$) afforded diol **9** (1.67 g, 53% for 3 steps) as a colorless oil. $[\alpha]_{\text{D}}^{20} +18$ (c 0.5, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 5.88~5.78 (m, 1H), 5.17~5.13 (m, 2H), 4.02~3.96 (m, 2H), 2.65 (br, 2H), 2.28 (dd, $J=12.0$, 4.0 Hz, 2H), 1.64~1.62 (m, 2H), 1.45~1.25 (m, 4H), 0.94 (t, $J=8.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 134.7, 118.2, 69.0, 68.2, 42.0, 41.8, 39.6, 19.0, 14.1; IR (KBr) ν_{max} 3365, 2926, 1638, 1458, 973 cm^{-1} ; HRMS (ESIMS) calcd for $\text{C}_9\text{H}_{19}\text{O}_2$ $[\text{M}+\text{H}]^+$ 159.1380, found 159.1387.

4.2.7 Synthesis of (2*S*,4*R*,6*S*)-(4-Hydroxy-6-propyl-tetrahydropyran-2-yl)acetic acid methyl ester (**13**)

To a solution of PdCl_2 (40 mg, 0.23 mmol) and CuCl_2 (0.94 g, 7 mmol) in CH_3CN (10 mL) under an atmosphere of CO (balloon) was added a solution of diol **9** (190 mg, 1.2 mmol) in MeOH (10 mL). The reaction mixture was stirred at room temperature under an atmosphere of CO (balloon) for 6 h. The mixture was concentrated. Flash column chromatography (EtOAc /hexanes, $V:V=1:3$) afforded alcohol **13** (207 mg, 80%) as a colorless oil. $[\alpha]_{\text{D}}^{20} -25$ (c 1.0, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 3.83~3.73 (m, 2H), 3.68 (s, 3H), 3.31~3.28 (m, 1H), 2.59 (dd, $J=16.0$, 8.0 Hz, 1H), 2.43 (dd, $J=16.0$, 4.0 Hz, 1H), 2.01~1.91 (m, 2H), 1.37~1.12 (m, 4H), 0.88 (t,

$J=6.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 171.7, 75.5, 72.1, 68.0, 51.7, 41.0, 40.9, 40.7, 37.9, 18.7, 13.9; IR (KBr) ν_{max} : 3435, 2925, 1725, 1418, 1168, 1055 cm^{-1} ; HRMS (ESIMS) calcd for $\text{C}_{11}\text{H}_{21}\text{O}_4$ $[\text{M}+\text{H}]^+$ 217.1434, found 217.1441.

4.2.8 Synthesis of **3** from **13**

To a suspension of *N,O*-dimethylhydroxylamine hydrochloride (0.19 g, 2.0 mmol) and **13** (108 mg, 0.5 mmol) in THF (10 mL) was slowly added $^i\text{PrMgCl}$ ($2.0 \text{ mol} \cdot \text{L}^{-1}$, 2.0 mL, 4.0 mmol) at -30°C under nitrogen. The reaction was stirred for 3 h, and then quenched by saturated NH_4Cl . The mixture was extracted with ethyl acetate twice (10 mL $\times$ 2). The combined organic phase was washed with brine, dried over anhydrous Na_2SO_4 , and the solvent was removed. The residue was purified by flash column chromatography with EtOAc /hexane ($V:V=1:3$) to give **3** (0.1 g, 83%) as a colorless oil.

4.2.9 Synthesis of 3-dehydroxyphomonol (**2**)

To a solution of **3** (0.13 g, 0.53 mmol) in THF (10 mL) was slowly added methylmagnesium iodide ($1.0 \text{ mol} \cdot \text{L}^{-1}$, 2.0 mL, 2.0 mmol) at -30°C under nitrogen. The reaction was stirred for 3 h at the temperature and then quenched by saturated NH_4Cl , extracted with ethyl acetate twice (50 mL $\times$ 2), washed with brine, dried over anhydrous Na_2SO_4 . The solvent was removed, and the residue was purified by flash column chromatography with EtOAc /hexane ($V:V=1:5$) to give 3-dehydroxyphomonol **2** (89 mg, 84%) as a colorless oil. $[\alpha]_{\text{D}}^{20} +2$ (c 0.2, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ : 3.82~3.71 (m, 2H), 3.29~3.25 (m, 1H), 2.70 (dd, $J=16.0$, 8.0 Hz, 1H), 2.41 (dd, $J=16.0$, 8.0 Hz, 1H), 2.15 (s, 3H), 1.95~1.88 (m, 2H), 1.56 (s, 1H), 1.47~1.25 (m, 4H), 1.25~1.11 (m, 2H), 0.85 (t, $J=6.0$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 207.1, 75.4, 71.9, 68.0, 49.7, 41.0, 41.0, 38.1, 31.0, 18.8, 14.0; IR (KBr) ν_{max} : 3411, 2928, 1712, 1365, 1261, 1140, 1175, 1115 cm^{-1} ; HRMS (ESIMS) calcd for $\text{C}_{11}\text{H}_{21}\text{O}_3$ $[\text{M}+\text{H}]^+$ 201.1485, found 201.1477.

Supporting Information ^1H NMR and ^{13}C NMR spectra of compounds. This material is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Wright, A. E.; Botelho, J. C.; Guzman, E.; Harmody, D.; Linley, P.; McCarthy, P. J.; Pitts, T. P.; Pomponi, S. A.; Reed, J. K. *J. Nat. Prod.* **2007**, 70, 412.
(b) Yu, J. F.; Feng, R. K.; Yang, Z. *Chin. J. Org. Chem.* **2017**, 37, 2526 (in Chinese).
- [2] Pereira, A. R.; McCue, C. F.; Gerwick, W. H. *J. Nat. Prod.* **2010**, 73, .
- [3] Kito, K.; Ookura, R.; Yoshida, S.; Namikoshi, M.; Ooi, T.; Kusumi, T. *Org. Lett.* **2008**, 10, 225.
- [4] Sikorska, J.; Hau, A. M.; Anklin, C.; Parker-Nance, S.; Davies-C, M. T.; Ishmael, J. E.; McPhail, K. L. *J. Org. Chem.* **2012**, 77, 6066.
- [5] Han, X.; Peh, G. R.; Floreancig, P. E. *Eur. J. Org. Chem.* **2013**, 1193.
- [6] Hua, J. L.; Bian, M.; Ding, H. F. *Tetrahedron Lett.* **2016**, 57, 5519.
- [7] Romero, K. J.; Galliher, M. S.; Pratt, D. A.; Stephenson-Corey, R. J.

- Chem. Soc. Rev.* **2018**, 47, 7851.
- [8] Bai, Y.; Davis, D. C.; Dai, M. J. *J. Org. Chem.* **2017**, 82, 2319.
- [9] Heravi, M. M.; Ahmadi, T.; Ghavidel, M.; Heidari, B.; Hamidi, H.; *RSC Adv.* **2015**, 5, 101999.
- [10] Li Y. Y.; Wang, M. Z.; Huang, Y. J.; Shen, Y. M. *Mycology* **2010**, 254.
- [11] (a) Krishna, P. R.; Prabhakar, S. *Tetrahedron Lett.* **2013**, 54, 3788.
(b) Reddy, B-V. S.; Srinivas, L. P.; Reddy, S.; Reddy, B. P.; Prasad, A. R.; Yadav, J. S. *Helv. Chim. Acta* **2014**, 97, 1326.
(c) Bhaskar. K.; Jangilib, P.; Anireddy, J. S. *J. Pharm. Res.* **2015**, 9, 550.
(d) Garrido, F.; Santalla, H.; Góme, G.; Fall, Y. *Tetrahedron Lett.* **2016**, 57, 2631.
(e) Gahalawat, S.; Pandey, S. K. *Tetrahedron Lett.* **2017**, 58, 2898.
(f) Wang, X.; Feng, R. K.; Qi, C. Z.; Yang, Z. *Tetrahedron Lett.* **2017**, 58, 3588.
- [12] Časar, Z. *Synlett* **2008**, 2036.
- [13] LarchevGque, M.; Henrot, S. *Tetrahedron* **1990**, 46, 4277.
- [14] Yang, Z.; Xie, X.; Jing, P.; Zhao, G.; Zheng, J.; Zhao, C.; She, X. *Org. Biomol. Chem.* **2011**, 9, 984.

(Li, L.; Fan, Y.)