有机化学

Chinese Journal of Organic Chemistry

金属有机化学专辑

第 39 卷 第 6 期 Vol. 39 No. 6 2019

ISSN 0253-2786

中国化学会主办
中国科学院上海有机化学研究所

Strem Chemicals, Inc., established in 1964, is a privately held manufacturer & marketer of high purity specialty chemicals. Strem's products are used for R&D and commercial scale applications, especially in the pharmaceutical, microelectronics, chemicals & petrochemicals industries. Some key products include catalysts, ligands, organometallics, CVD/ALD precursors and nanomaterials. For more information, visit strem.com. Strem is an ISO 9001:2015 certified company.

STREM'S KEY PRODUCT LINES

- METAL CATALYSTS
- LIGANDS
- INORGANICS
- ORGANOMETALLICS
- PHOTOCATALYSTS
- SCREENING KITS

11-0575

27-0400

78-1350

77-0425

15-8021

15-1053

15-1060

46-2150

29-6720

15-0067

77-0410

15-0126

中国高端 均相催化剂领军者

- 10年研发经验
- 国际市场认可
- 无水无氧封装存储

贵金属催化剂：

Pd Rh Ru Ir Au Cu Ni Zr

有机配体：

P N O

MIT专利授权：

BUCHWALD LIGAND & PRECATALYST

手性膦配体：

BINAP SEGPPOS PHOSPHORAMIDITE
JOSIPHOS PHOX SPIRO

手性氮配体：

BOX BINAM DACH TROST DPEN SALEN

手性催化剂：

PHOSPHORIC ACIDS NHC THIOUREA
CINCHONA ALKALOID

光催化剂：

Ir Ru ACRI. SALTS

OTHER: RAFT SILANE PORPHYRIN MOF
ALD PRECURSOR

媲美进口品质的催化剂尽在

WWW.LAAJOO.COM

TEL: 021-60702684

QQ: 3004302861

MAIL: REAGENT@SINOCOMPOUND.COM

理想的实验室伴侣！

中国人都用得起的隔膜真空泵！

旋转蒸发仪/隔膜真空泵

® 上海豫康科教仪器设备有限公司

Shanghai Yukang Science teaching Instrument equipment Co.,Ltd

地址：上海市和桥路470号

电话：021-36338262 56470425
56470426

传真：021-56470425

手机：13701684908

邮编：200435

网址：www.sh-yukang.com

Tel: 021-36338262 56470425
56470426

Fax: 021-56470425

Mobil: 13701684908

Zip Code: 200435

Web: www.sh-yukang.com

蝴蝶型双核铜配合物催化氧化合成亚胺的研究

张玲娟* 党宇娇 张献明*

(山西师范大学 教育部磁性分子与磁信息材料重点实验室 临汾 041004)

摘要 无论是在实验室还是工业生产中,通过胺的选择性氧化制备亚胺类化合物都具有重要的意义。秉承原子经济性、环境友好的理念,以蝴蝶型络合物 $\text{Cu}_2(\text{o phen})_2$ 为催化剂,实现了需氧型的胺的选择性氧化。值得注意的是,该催化体系对醇与胺的氧化交叉偶联、一级胺的氧化偶联和二级胺的氧化脱氢反应都是有效的。反应产率高达 93%,亚胺的选择性高达 99%。在反应体系中,避免了昂贵的氮氧自由基类化合物的使用,使得该体系更具有潜在的实际应用价值。

关键词 需氧氧化;双核铜配合物;无碱;无氮-氧自由基;高选择性

Diversified Synthesis of Imines via Aerobic Oxidation Catalyzed by Dinuclear Butterfly-Like Cu(I) Complex

Zhang, Lingjuan* Dang, Yujiao Zhang, Xianming*

(Key Laboratory of Magnetic Molecules & Magnetic Information Materials, Ministry of Education, School of Chemical and Material Science, Shanxi Normal University, Linfen 041004)

Abstract The selective catalytic oxidation of amines for the synthesis of imines is important both in laboratory and industrial production. From atom-efficient, economic and environmental view of points, dioxygen selective oxidation of amines was achieved by using wings-opened butterfly-like complex $\text{Cu}_2(\text{o phen})_2$ as catalyst. It was worth noting that the catalytic system was efficient to the cross-coupling of alcohols with amines, homocoupling of primary amines and oxidative dehydrogenation of secondary amines. The yield is up to 93% and the selectivity of imines is as high as 99%. Avoiding the use of expensive nitroxyl derivatives and base was suitable for practical application.

Keywords aerobic oxidation; dinuclear Cu-complex; base-free; N—O radical free; high selectivity

1 Introduction

Imines are widely used as practical synthons in the process of constructing biologically or pharmaceutically active compounds and heterocycles.^[1] In addition to the classic condensation of carbonyl compounds with amines, three attractive methods have been extensively applied in synthesizing imines with green air or dioxygen as terminal oxidant: (a) dehydrogenated cross-coupling of alcohols with amines,^[2] (b) homocoupling of primary amines, in which the major challenge is to avoid the formation of nitrile or azocompound,^[2h,3] (c) oxidative dehydrogenation of secondary amines.^[4] Various catalyst systems like photo catalyzed, bioinspired, metal (Ru, Os, Pd, Pt, Au, Co, Mn, Cu, etc.) and metal-free catalyzed have been widely reported.^[1e] Due to low-cost, well-sourced and high-active for dioxygen activation, copper catalysts have been re-

ceived widespread attention for aerobic imine formation among various available systems.^[1a,5] Nevertheless, many of these “Cu/O₂” catalytic systems shared two problems: first, extra base^[5c,5g,6] or expensive nitroxyl derivatives^[5f,7] as co-catalysts were usually required to expand the substrate scope, decrease the reaction temperature or increase the selectivity of product (Scheme 1). Second, there was hardly a catalytic system that can effectively promote the synthesis of imines simultaneously through the above three ways so far. From sustainable development point of view, searching high efficient and universal catalyst for the synthesis of imines (C=N) is ever-appealing in the catalyst research field.

Recently, our group reported a 2,2,6,6-tetramethylpiperidin-1-oxyl (TEMPO)/base-free catalytic system for selective oxidation of alcohols using a planar wings opened butterfly-like complex $\text{Cu}_2(\text{o phen})_2$ as catalyst.^[8] Moreo-

* Corresponding authors. E-mail: zhanglj@sxnu.edu.cn; zhangxm@dns.sxnu.edu.cn

Received March 15, 2019; revised April 3, 2019; published online May 15, 2019.

Project supported by the National Natural Science Foundation of China (No. 21402112), and the 1331 Project of Shanxi Province.

国家自然科学基金(No.21402112)、山西省 1331 工程资助项目。

Scheme 1 Various “Cu/O₂” catalytic systems for the synthesis of imines

ver, density functional theory (DFT) calculations revealed that molecular oxygen was effectively activated via wings-flapping of Cu₂(ophen)₂. It is generally known that amine and alcohol have similar structure and physicochemical property. Hence, we may anticipate aerobic oxidation of amines to imines catalyzed by dinuclear Cu₂(ophen)₂.

Herein, a novel, convenient approach for the synthesis of imines using butterfly-like complex Cu₂(ophen)₂ as catalyst under O₂/air atmosphere is present. Interestingly, the catalytic system is effective both to promote oxidative dehydrogenation of secondary amines, cross-coupling of alcohols with amines and to homocoupling of primary amines in the absence of 2,2,6,6-tetramethylpiperidinoxy (TEMPO) and base. It is worth noting that oxidative coupling reaction of alcohols and amines can be carried out successfully at room temperature.^[2a,2c,5b,5c,6a]

2 Results and discussion

2.1 Oxidative cross-coupling of alcohols with amines

Firstly, the dehydrogenative cross-coupling of alcohol and amine to construct imines was chosen in which the structures of products are diversified and water was the only byproduct. Under dioxygen atmosphere (101 kPa), *N*-benzylidene-1-phenylmethanamine (**3aa**) could be obtained in 92% yield from the reaction of benzyl alcohol (**1a**) with benzylamine (**2a**) in *N,N*-dimethylformamide (DMF) in the presence of Cu₂(ophen)₂ (2.5 mol%) at room temperature (Table 1, Entry 6). Compared with the oxidation of alcohol catalyzed by Cu₂(ophen)₂, the reducing of cross-coupling reaction temperature showed that the generation of imine effectively promoted the oxidative transformation of alcohol. Likewise, the process of cross-coupling was no need for any base and nitroxyl co-catalyst.

With the optimal conditions for the preparation of imine **3aa** in hand, the scope of this cross-coupling reaction was explored. The electronic properties and position of substituents in benzene ring didn't have obvious impact on the reaction when using aromatic methyl alcohols and amines as raw materials (Table 2, Entries 1~8, 11~13). In line with the report of literatures, the synthesis of ketimine was ineffective by employing secondary aromatic alcohol (**1i**) as a substrate.^[5g] Unfortunately, aniline was less active to the conversion even increasing the temperature to 60 °C (Table 2, Entry 14).

Table 1 Screening and optimization of the reaction conditions cross-coupling of alcohols with amines.^[a]

Entry	Oxid.	Solvent	Cat.	Conv. ^c /%	Yield ^d /%
1	Air	CH ₃ CN	Cu ₂ (ophen) ₂	59	7
2	O ₂	CH ₃ CN	Cu ₂ (ophen) ₂	89	88
3	N ₂	CH ₃ CN	Cu ₂ (ophen) ₂	0	0
4	O ₂	THF	Cu ₂ (ophen) ₂	90	88
5	O ₂	Toluene	Cu ₂ (ophen) ₂	62	60
6	O ₂	DMF	Cu ₂ (ophen) ₂	92	91
7 ^b	O ₂	DMF	CuCl ₂ +phen	42	39
8 ^b	O ₂	DMF	CuCl+phen	69	65
9	O ₂	DMF	Cu powder+phen	26	22

^a Reaction conditions unless otherwise stated, benzyl alcohol (0.5 mmol), benzylamine (0.6 mmol), Cu₂(ophen)₂ (2.5 mol%), solvent (2 mL), O₂ balloon, room temperature, 12 h; ^b [CuX] (5 mol%), phen (5 mol%); ^c determined by GC using nitrobenzene as an internal standard; ^d determined by ¹H NMR.

2.2 Oxidative homocoupling of amines and oxidative dehydrogenation of secondary amine

The direct oxidative homocoupling of amines was another attractive strategy for the construction of imines.^[3] It should be noted that a major challenge of the homocoupling reaction is avoiding the over oxidation to nitrile.^[9] Next, the dinuclear Cu₂L₂-catalyzed aerobic homocoupling reactions of amines were investigated. It was fortunately observed that high amines conversion and excellent imines selectivity were obtained in the presence of Cu₂(ophen)₂ (2.5 mol%) at 100 °C under air atmosphere, only trace nitrile was observed. A series of aromatic methylamines **2** could be readily oxidized to produce imines **4** in yields up to 94% (Table 3). The oxidative coupling reactions of aliphatic amine were relatively less efficient but with good selectivities for imines (Table 3, **4b**). Unfortunately, the oxidative coupling reactions of secondary amine and the cross-coupling of benzylamine with aniline were unsuccessful (Table 3, **4g**, **4h**). Interestingly, under the same condition, the oxidative dehydrogenation of symmetric dibenzylamine could be achieved to imine in medium conversions (Scheme 2).

We found that the oxidative homocoupling of amine is more difficult than cross-coupling of alcohols with amines (temperature from r.t. to 100 °C). In addition, no or only trace aldehyde was observed by GC during the reaction process, the condensation of aldehyde from the hydrolysis of imine intermediate **7** with another amine to give **4** was unlikely in our system.^[10] The product yield decreased obviously to 3% in the presence of superoxide scavenger (*p*-benzoquinone, BQ, Table 3, **3aa**)^[3f] and the characteristic peaks DMPO-O₂^{•−} were observed by the electron spin resonance (ESR) spectra spin-trap technique,^[3c] indicating that the formation of a superoxide radical L₂Cu₂-O₂^{•−} is an

Table 2 Oxidative cross-coupling of alcohols with amines^a

Entry	Compd. 1	R ¹	R ²	Compd. 2	R ³	Compd. 3	Yield ^b /%
1	1a	C ₆ H ₅	H	2a	C ₆ H ₅ CH ₂	3aa	90
2	1b	4-MeOC ₆ H ₄	H	2a		3ba	89
3	1c	2-MeOC ₆ H ₄	H	2a		3ca	93
4	1d	4-ClC ₆ H ₄	H	2a		3da	87
5	1e	4-BrC ₆ H ₄	H	2a		3ea	85
6	1f	C ₆ H ₅ CH=CH	H	2a		3fa	93
7	1g	2-Pyridyl	H	2a		3ga	92
8	1h	2-Furyl	H	2a		3ha	75
9	1i	C ₆ H ₅	CH ₃	2a		3ia	0 (16) ^c
10	1j	Cyclohexyl	H	2a		3ja	91
11	1a			2b	CH ₃ (CH ₂) ₅	3ab	93
12	1a			2c	Cyclohexyl	3ac	89
13	1a			2d	4-MeOC ₆ H ₄ CH ₂	3ad	83
14	1a			2e	C ₆ H ₅	3ae	25 (46) ^c

^a Conditions: alcohol (0.5 mmol), amine (0.6 mmol), Cu₂(ophen)₂ (2.5 mol%), DMF (2 mL), O₂ balloon, room temperature, 12 h. ^b Determined by GC-MS on the amount of alcohol using nitrobenzene as an internal standard. ^c The reaction temperature rased up to 60°C.

Table 3 Oxidative dehydrogenation homocoupling of amines^a **3aa** 94% (> 99%), trace (adding BQ) **4b**: 20% (100%), 24 h^b **4c** 91% (> 99%), 24 h **4d** 94% (98%) **4e** 60% (>99%) **4f** 71% (> 99%), 24 h **4g** trace **4h** trace

^a Conditions: amine (1 mmol), Cu₂(ophen)₂ (2.5 mol%), DMF (2 mL), open air, 100 °C, 12 h. ^btoluene as solvent (when using DMF as solvent, N-hexylformamide was obtained in 95% conversion and trace imine was observed).

essential step in homocoupling of amines (Scheme 3). Based on the above results, a favourable mechanism is proposed: dinuclear Cu₂(ophen)₂ activates dioxygen molecule to give L₂Cu₂-O₂^{••}, amine is oxidized by activated O₂ to give a reactive imine intermediate **7**. Then, the target molecule was formed through the direct condensation of imine **7** with another amine with the release of NH₃. The demand for high reaction temperature may be due to the weak elec-

Scheme 2 Oxidative dehydrogenation of secondary amine **Scheme 3** Proposed mechanism for Cu₂(ophen)₂ catalyzed homocoupling of amines

trophilicity of imine than aldehyde.

3 Conclusions

In this work, a simple, facile method of preparing imines by the aerobic oxidation of alcohols and amines was developed. Employing dinuclear Cu₂(ophen)₂ as catalyst, diversified imines were obtained with moderate to high yields under air or oxygen atmosphere, avoiding the use of excesses of bases and nitroxyl derivatives. It's worth mentioning that the efficient catalytic system was suitable for the dehydrogenated cross-coupling of alcohols with

amines, homocoupling of primary amines and oxidative dehydrogenation of secondary amines. Importantly, high selectivity of imines also can be achieved in the homocoupling of primary amines.

4 Experimental

[Cu₂(ophen)₂] was synthesized according to literature methods.^[11] Other chemicals were analytically pure from commercial sources and used without further purification. The catalyst was dried at room temperature before the catalytic reactions. Gas chromatography (GC) analyses were obtained by an Agilent 7820A (FID from AGILENT 7820) equipped with a cross-linked (95%)-dimethyl-(5%)-diphenylpolysil-oxane column (HP-5, 30 m × 0.32 mm × 0.25 μm), helium, injector temperature 250 °C, detector temperature 300 °C, and oven temperature program 45 °C (3 min)-20 °C/min-280 °C (2 min). GC-MS spectra were recorded on a Thermo Fisher ITQ-900. Electron paramagnetic resonance (EPR) measurement was conducted on a A300-10/12 (Bruker). ¹H NMR were determined on Bruker Ascend™-600.

4.1 General procedure for the oxidative cross-coupling of alcohols with amines

To a 5 mL round bottom flask, the mixture of benzyl alcohol **1a** (0.5 mmol, 0.052 mL), benzylamine **2a** (0.5 mmol, 0.066 mL) and Cu₂(ophen)₂ (0.025 mmol, 13 mg) in dimethylformamide (DMF) (2 mL) was stirred at room temperature under oxygen atmosphere (oxygen ball). The reaction was monitored and conversions were determined by GC using nitrobenzene as an internal standard. At the end of the reaction, the mixture was extracted with ethyl acetate. The organic phase was dried over anhydrous Na₂SO₄. The crude residue was obtained after evaporation of the solvent in vacuum, and the residue was purified by flash chromatography with petroleum and EtOAc as the eluent to give the pure product.

4.2 General procedure for the oxidative homocoupling of amines

To a 5 mL round bottom flask, the mixture of benzylamine **2a** (1.0 mmol, 0.132 mL) and Cu₂(ophen)₂ (0.025 mmol, 13 mg) in DMF (2 mL) was stirred at 100 °C under air. The reaction was monitored and conversions were determined by GC using nitrobenzene as an internal standard. At the end of the reaction, the reaction mixture was cooled to room temperature and extracted with ethyl acetate. The organic phase was dried over anhydrous Na₂SO₄. The crude residue was obtained after evaporation of the solvent in vacuum, and the residue was purified by flash chromatography with petroleum and EtOAc as the eluent to give the pure product.

4.3 Spectral data of products

N-Benzylidene-*p*-benzylamine (**3aa**):^[2m] ¹H NMR (600 MHz, CDCl₃) δ: 8.26 (s, 1H), 7.75~7.70 (m, 2H), 7.28 (d, *J*=2.2 Hz, 3H), 7.25 (d, *J*=4.4 Hz, 4H), 7.24 (s, 1H), 4.75 (d, *J*=1.3 Hz, 2H).

N-(4-Methoxybenzylidene)-1-phenylmethan-amine (**3ba**):^[12] ¹H NMR (600 MHz, CDCl₃) δ: 8.28 (s, 1H), 7.69 (d, *J*=8.7 Hz, 2H), 7.28~7.35 (m, 5H), 6.88 (d, *J*=8.7 Hz, 2H), 4.78 (s, 2H), 3.78 (s, 3H).

N-(2-Methoxybenzylidene)-1-phenylmethan-amine (**3ca**):^[2m] ¹H NMR (600 MHz, CDCl₃) δ: 8.81 (s, 1H), 7.37~6.95 (m, 9H), 4.75 (s, 2H), 3.25 (s, 3H).

N-(4-Chlorobenzylidene)-1-phenylmethanamine (**3da**):^[2m] ¹H NMR (600 MHz, CDCl₃) δ: 8.27 (s, 1H), 7.69~7.64 (m, 1H), 7.34~7.29 (m, 3H), 7.36~7.23 (m, 3H), 7.28~7.14 (m, 1H), 4.73 (s, 2H).

N-(4-Bromobenzylidene)-1-phenylmethanamine (**3ea**):^[2m] ¹H NMR (600 MHz, CDCl₃) δ: 8.24 (s, 1H), 7.55~7.54 (d, *J*=8.0 Hz, 2H), 7.45 (d, *J*=8.2 Hz, 2H), 7.28 (m, 4H), 7.22 (m, 1H), 4.71 (s, 2H).

N-(3-Phenylallylidene)-1-phenylmethanamine (**3fa**):^[2m] ¹H NMR (600 MHz, CDCl₃) δ: 8.12 (s, 1H), 7.44 (d, *J*=8.5 Hz, 2H), 7.34~7.26 (m, 7H), 7.25~7.21 (m, 1H), 6.94 (d, *J*=3 Hz, 2H), 4.70 (s, 2H).

N-(Pyridin-2-ylmethylene)-1-phenylmethanamine (**3ga**):^[2m] ¹H NMR (600 MHz, CDCl₃) δ: 8.42 (d, *J*=4 Hz, 1H), 8.37 (s, 1H), 8.14 (d, *J*=8.5 Hz, 1H), 7.74 (d, *J*=7.5 Hz, 1H), 7.34~7.30 (m, 1H), 7.29~7.21 (m, 4H), 7.20~7.14 (m, 1H), 4.54 (d, *J*=6 Hz, 2H).

N-(Furan-2-ylmethylene)-1-phenylmethanamine (**3ha**):^[2m] ¹H NMR (600 MHz, CDCl₃) δ: 8.30 (s, 1H), 8.01 (s, 1H), 7.69 (d, *J*=6 Hz, 2H), 7.41 (s, 1H), 7.31 (d, *J*=6.5 Hz, 2H), 7.24~7.20 (m, 4H), 7.17~7.13 (m, 1H), 6.66 (d, *J*=4 Hz, 1H), 6.39~6.35 (m, 1H), 4.70 (s, 2H).

N-Benzylidene-1-hexanamine (**3ab**):^[12] ¹H NMR (600 MHz, CDCl₃) δ: 8.17 (s, 1H), 7.62 (m, 2H), 7.30 (m, 3H), 3.51 (t, *J*=7.0 Hz, 2H), 1.64 (m, 2H), 1.26~1.32 (m, 6H), 0.81 (m, 3H).

N-Benzylidene-1-cyclohexanamine (**3ac**):^[2m] ¹H NMR (600 MHz, CDCl₃) δ: 8.29 (s, 1H), 7.67~7.72 (m, 2H), 7.37~7.32 (m, 3H), 3.17 (m, 1H), 1.89~1.22 (m, 10H).

N-Benzylideneaniline (**3ae**):^[2m] ¹H NMR (600 MHz, CDCl₃) δ: 8.44 (s, 1H), 7.91 (m, 2H), 7.46 (m, 3H), 7.40 (m, 2H), 7.22 (m, 3H).

N-Hexylidenehexan-1-amine (**4b**):^[13] ¹H NMR (600 MHz, CDCl₃) δ: 7.55 (t, *J*=4.8 Hz, 1H), 3.30 (t, *J*=6.9 Hz, 2H), 2.25~2.11 (m, 2H), 1.61~1.49 (m, 4H), 1.32~1.27 (m, 10H), 0.88~0.82 (m, 6H) ppm;

N-(4-Fluorobenzylidene)-*p*-fluorobenzylamine (**4c**):^[3f] ¹H NMR (600 MHz, CDCl₃) δ: 8.31 (s, 1H), 7.75~7.71 (m, 2H), 7.29~7.18 (m, 2H), 7.10~7.02 (m, 2H), 7.03~6.96 (m, 2H), 4.75 (s, 1H).

N-(4-Methoxybenzylidene)-*p*-methoxybenzylamine (**4d**):^[3f] ¹H NMR (600 MHz, CDCl₃) δ: 8.21 (s, 1H), 7.69~7.62 (m, 2H), 7.20 (d, *J*=8.8 Hz, 2H), 6.90~6.80 (m, 4H), 4.70 (s, 2H), 3.72 (s, 3H), 3.70 (s, 3H).

1-(Thiophen-2-yl)-*N*-(thiophen-2-ylmethyl)methanimine (**4f**):^[3f] ¹H NMR (600 MHz, CDCl₃) δ: 8.32 (d, *J*=1.2 Hz, 1H), 7.32 (dt, *J*=5.0, 1.0 Hz, 1H), 7.24 (dd, *J*=3.6, 1.1 Hz, 1H), 7.19 (dd, *J*=4.9, 1.5 Hz, 1H), 7.01 (dd, *J*=5.0, 3.6 Hz, 1H), 6.95~6.89 (m, 2H), 4.89 (s, 2H).

Supporting Information The structure of the catalyst, the GC spectra, mass spectra of products and the ESR spectrum are available. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Largeron, M. *Eur. J. Org. Chem.* **2013**, 2013, 5225.
(b) Kobayashi, S.; Mori, Y.; Fossey, J. S.; Salter, M. M. *Chem. Rev.* **2011**, 111 (4), 2626.
(c) Chen, B.; Wang, L.; Gao, S. *ACS Catal.* **2015**, 5, 5851.
(d) Zhang, Z. G.; Li, J. L.; Huang, G. Q.; Sun, K. Zhang, G. S.; Ma, N. N.; Liu, Q. F.; Liu, T. X. *Chin. J. Chem.* **2016**, 34, 1309.
(e) Patil, R. D.; Adimurthy, S. *Asian J. Org. Chem.* **2013**, 2, 726.
- [2] (a) Mukherjee, A.; Nerush, A.; Leitun, G.; Shimon, L. J. W.; Ben David, Y.; Espinosa Jalapa, N. A.; Milstein, D. *J. Am. Chem. Soc.* **2016**, 138, 4298.
(b) Chen, B.; Li, J.; Dai, W.; Wang, L.; Gao, S. *Green Chem.* **2014**, 16, 332.
(c) Zhang, G.; Hanson, S. K. *Org. Lett.* **2013**, 15, 650.
(d) Zhang, E.; Tian, H.; Xu, S.; Yu, X.; Xu, Q. *Org. Lett.* **2013**, 15 (11), 2704.
(e) Montag, M.; Zhang, J.; Milstein, D. *J. Am. Chem. Soc.* **2012**, 134 (25), 10325.
(f) Maggi, A.; Madsen, R. *Organometallics* **2012**, 31, 451.
(g) He, W.; Wang, L.; Sun, C.; Wu, K.; He, S.; Chen, J.; Wu, P.; Yu, Z. *Chem.-Eur. J.* **2011**, 17, 13308.
(h) Kegnaes, S.; Mielby, J.; Mentzel, U. V.; Christensen, C. H.; Riisager, A. *Green Chem.* **2010**, 12, 1437.
(i) Jaiswal, G.; Landge, V. G.; Jagadeesan, D.; Balaraman, E. *Green Chem.* **2016**, 18, 3232.
(j) Tamura, M.; Tomishige, K. *Angew. Chem., Int. Ed.* **2015**, 54, 864.
(k) Saranya, S.; Ramesh, R.; Grzegorz Malecki, J. *Eur. J. Org. Chem.* **2017**, 2017, 6726.
(l) Chandra, M. D.; Sadhukha, A.; Maayan, G. *J. Catal.* **2017**, 355, 139.
(m) Han, L.; Xing, P.; Jiang, B. *Org. Lett.* **2014**, 16, 3428.
- [3] (a) Wang, Z.; Lang, X. *Appl. Catal. B-Environ.* **2018**, 224, 404.
(b) Rodríguez-Lugo, R. E.; Chacón-Terán, M. A.; De León, S.; Vogt, M.; Rosenthal, A. J.; Landaeta, V. R. *Dalton Trans.* **2018**, 47, 2061.
(c) Jin, J.; Yang, C.; Zhang, B.; Deng, K. *J. Catal.* **2018**, 361, 33.
(d) Xu, B.; Hartigan, E. M.; Feula, G.; Huang, Z.; Lumb, J.-P.; Arndtsen, B. A. *Angew. Chem., Int. Ed.* **2016**, 55, 15802.
(e) Su, C.; Tandiana, R.; Tian, B.; Sengupta, A.; Tang, W.; Su, J.; Loh, K. P. *ACS Catal.* **2016**, 3594.
(f) Raza, F.; Park, J. H.; Lee, H.-R.; Kim, H.-I.; Jeon, S.-J.; Kim, J.-H. *ACS Catal.* **2016**, 2754.
(g) Sun, D.; Ye, L.; Li, Z. *Appl. Catal. B- Environ.* **2015**, 164, 428.
(h) Johnson, J. A.; Luo, J.; Zhang, X.; Chen, Y.-S.; Morton, M. D.; Echeverría, E.; Torres, F. E.; Zhang, J. *ACS Catal.* **2015**, 5, 5283.
(i) Wu, X.-F.; Petrosyan, A.; Ghochikyan, T. V.; Saghyian, A. S.; Langer, P. *Tetrahedron Lett.* **2013**, 54, 3158.
(j) Furukawa, S.; Ohno, Y.; Shishido, T.; Teramura, K.; Tanaka, T. *ACS Catal.* **2011**, 1, 1150.
(k) Varyani, M.; Khatri, P. K.; Jain, S. L. *Tetrahedron Lett.* **2016**, 57, 723.
- [4] Nikbakht, F.; Heydari, A. *CR Chim.* **2015**, 18, 132.
- [5] (a) Liu, Y.; Chen, Y.; Zhou, W.; Jiang, B.; Zhang, X.; Tian, G. *Catal. Sci. Technol.* **2018**, 8, 5535.
(b) Dutta, I.; De, S.; Yadav, S.; Mondol, R.; Bera, J. K. *J. Organomet. Chem.* **2017**, 849-850, 117.
(c) Bai, L.; Dang, Z. *RSC Adv.* **2015**, 5, 10341.
(d) Wang, J.; Lu, S.; Cao, X.; Gu, H. *Chem. Comm.* **2014**, 50, 5637.
(e) Ryland, B. L.; Stahl, S. S. *Angew. Chem., Int. Ed.* **2014**, 53, 8824.
(f) Tian, H.; Yu, X.; Li, Q.; Wang, J.; Xu, Q. *Adv. Synth. Catal.* **2012**, 354, 2671.
(g) Kang, Q.; Zhang, Y. *Green Chem.* **2012**, 14, 1016.
- [6] (a) Lan, Y.-S.; Liao, B.-S.; Liu, Y.-H.; Peng, S.-M.; Liu, S.-T. *Eur. J. Org. Chem.* **2013**, 2013, 5160.
(b) Pérez, J. M.; Cano, R.; Yus, M.; Ramón, D. J. *Eur. J. Org. Chem.* **2012**, 2012, 4548.
- [7] (a) Huang, B.; Tian, H.; Lin, S.; Xie, M.; Yu, X.; Xu, Q. *Tetrahedron Lett.* **2013**, 54, 2861.
(b) Sonobe, T.; Oisaki, K.; Kanai, M. *Chem. Sci.* **2012**, 3, 3249.
- [8] Zhang, L. J.; Liu, J.; Zhang, F. Q.; Zhang, X.-M. *J. Catal.* **2017**, 354, 78.
- [9] (a) Shen, Y.; Zhou, Y.; Jiang, L.; Ding, G.; Luo, L.; Zhang, Z.; Xie, X. *Tetrahedron* **2018**, 74, 4266.
(b) Tao, C.; Wang, B.; Sun, L.; Liu, Z.; Zhai, Y.; Zhang, X.; Wang, J. *OrgBioChem* **2017**, 15, 328.
(c) Ray, R.; Chandra, S.; Yadav, V.; Mondal, P.; Maiti, D.; Lahiri, G. K. *Chem. Commun.* **2017**, 53, 4006.
(d) Liu, G.; Sun, L.; Liu, J.; Wang, F.; Guild, C. J. *Mol. Catal.* **2017**, 440, 148.
(e) Pavel, O. D.; Goodrich, P.; Cristian, L.; Coman, S. M.; Părvulescu, V. I.; Hardacre, C. *Catal. Sci. Technol.* **2015**, 5, 2696.
- [10] Patil, R. D.; Adimurthy, S. *RSC Adv.* **2012**, 2, 5119.
- [11] Zhang, X.-M.; Tong, M.-L.; Gong, M. L.; Lee, H.-K.; Luo, L.; Li, K. F.; Tong, Y.-X.; Chen, X.-M. *Chem.-Eur. J.* **2002**, 8, 3187.
- [12] Zhang, G. Q.; Hanson, S. K. *Org. Lett.* **2013**, 15, 3650.
- [13] Zhang, Y.; Lu, F.; Huang, R.; Zhang, H.; Zhao, J. *Catal. Commun.* **2016**, 81, 10.

(Lu, Y.)