膦催化水杨醛亚胺与马来酰亚胺的[4+1]环化反应及螺[苯并呋喃-2,3'-吡咯烷]衍生物的合成

杨 梅^a 曹仕轩^a 贺峥杰^{*,a,b}

(^a南开大学元素有机化学国家重点实验室 天津 300071)

(^b天津化学化工协同创新中心 天津 300071)

摘要 报道了一个膦催化水杨醛亚胺与马来酰亚胺的[4+1]环化反应, 以44%~99%收率生成螺[苯并呋喃-2,3'-吡咯烷]衍生物。产物包含一对可经硅胶柱层析分离的 *syn* 和 *anti* 异构体(*dr* 1.6 : 1~5 : 1), 由此提供了合成螺[苯并呋喃-2,3'-吡咯烷]简单有效的方法。该反应起始于 PPh₃ 与马来酰亚胺原位形成非烯丙基磷叶立德活性中间体, 经亲核加成/分子内取代的串联过程完成, 代表了新一例经此类磷叶立德完成的膦催化[4+1]环化反应。

关键词 膦; [4+1]环化反应; 亚胺; 马来酰亚胺; 合成

Phosphine-Catalyzed [4+1] Annulation of Salicyl Imines with Maleimides and Synthesis of Spiro[benzofuran-2,3'-pyrrolidine] Derivatives

Yang, Mei^a Cao, Shixuan^a He, Zhengjie^{*,a,b}

(^a State Key Laboratory of Elemento-organic Chemistry, College of Chemistry, Nankai University, Tianjin 300071)

(^b Collaborative Innovation Center of Chemical Science and Engineering (Tianjin), Tianjin 300071)

Abstract In this work, a phosphine-catalyzed [4+1] annulation between salicyl imines and maleimides has been successfully developed, which readily produces spiro[benzofuran-2,3'-pyrrolidine] derivatives in 44%~99% yields. The annulation products were obtained as a pair of *syn*- and *anti*-isomers with *dr* 1.6 : 1~5 : 1. The *syn*- and *anti*-isomers can be readily separated by column chromatography on silica gel. Thus, this reaction constitutes a simple and efficient method for the synthesis of spiro[benzofuran-2,3'-pyrrolidines]. Presumably, the reaction is initiated by *in situ* generated non-allylic P-ylide from maleimide and PPh₃, and proceeds through a cascade sequence of nucleophilic addition/intramolecular S_N2-like substitution. It accordingly represents a new example of the phosphine-catalyzed [4+1] annulation via *in situ* generated non-allylic P-ylides.

Keywords phosphine; [4+1] annulation reaction; imine; maleimide; synthesis

1 Introduction

The 2,3-dihydrobenzofuran ring is the core structure of a number of bioactive natural and artificial molecules, and its effective synthesis has therefore attracted much interest from chemists.^[1] As a special subset of 2,3-dihydrobenzofurans, the spirocyclic benzofuran skeletons have also been found in many biologically interesting compounds (Figure 1).^[2] For instance, the telomerase inhibitor griseorhodin A (I), probably the most heavily oxidized bacterial polyketide, features a unique spirobenzofuran moiety which is

crucial for its activity.^[2a,2b] Griseofulvin (II), first isolated from filamentous fungi, contains a spirobenzofuranone core, and has been used to treat ringworm infections of the skin and nails in both humans and animals for decades.^[2c~2g] The spiro[benzofuran-2,3'-pyrrolidine] derivative (III) was used as a key precursor in the synthesis of a KA receptor antagonist.^[2h~2j] Since a spirocycle contains two perpendicular rings commonly connected through one quaternary carbon, this structural feature imposes more challenges in its synthesis.^[3] Thus, synthesis of the spirobenzofuran skeleton has become an attractive research

* Corresponding author. E-mail: zhengjiehe@nankai.edu.cn

Received April 16, 2019; revised May 7, 2019; published online May 15, 2019.

Dedicated to the 100th anniversary of the birth of Professor Ruyu Chen.

Project supported by the National Natural Science Foundation of China (Nos. 21472096, 21272119).

国家自然科学基金(Nos. 21472096, 21272119)资助项目。

topic. In the reported effective syntheses, the spirobenzofurans were generally constructed through multistep synthetic routes.^[2] The catalytic cascade synthetic protocols are relatively rare.^[2f,4] Accordingly, new and efficient synthetic strategies for spirocyclic benzofurans remain highly desirable.

Figure 1 Representative examples of bioactive spirocyclic benzofuran derivatives

During the past two decades, the phosphine catalytic annulation strategy has made great progress and has provided a powerful toolbox for constructing diverse cycles.^[5] Among the annulation reactions, the allylic phosphorus ylide-initiated annulations are proven to be prevailing.^[5e,5g] As the key reactive intermediates, allylic P-ylides are *in situ* generated from popular substrates such as electron-deficient allenes and Morita-Baylis-Hillman (MBH) derivatives with phosphines. The allylic P-ylide-mediated annulations generally proceed through a tandem sequence of Michael addition/intramolecular S_N2' -like substitution.^[6] By this mode, the ring-closure step can effectively circumvent the steric hindrance arising from the bulky phosphonium moiety.^[6] On the other hand, the non-allylic P-ylide-initiated annulations were seldom reported through a similar mode of Michael addition/intramolecular S_N2 -like substitution.^[7] Recently our group and Shi's group independently developed novel phosphine-catalyzed [4 + 1] annulation reactions of electron-deficient 4π -conjugated systems such as 1,3-dienes or 1,3-azadienes with maleimides, providing a convenient and highly efficient method to construct important azaspiro[4.4]nonane skeletons (Scheme 1).^[7] By our results, the annulation reaction is initiated with the *in situ* formed non-allylic P-ylide from maleimide and PPh_3 , and proceeds through a sequence of the Michael addition/intramolecular S_N2 -like substitution. Accordingly this reaction also represents the first example of phosphine-catalyzed [4 + 1] annulation reaction via non-allylic phosphorus ylides (Scheme 1).^[7b] Encouraged by the feasibility of this annulation reaction in the construction of spirocycles, recently we further extended this [4 + 1] annulation strategy to dual functional salicyl imines

1 and successfully developed a new and convenient method to synthesize spiro[benzofuran-2,3'-pyrrolidine]-2',5'-diones (Scheme 1). Herein we report the relevant results.

Previous work from Shi and He groups

This work

Scheme 1 Phosphine-catalyzed [4 + 1] annulations via non-allylic P-ylides

2 Results and discussion

We initiated the investigation with the stoichiometric reaction of salicyl *N*-tosylimine **1a** and phosphorus ylide **2aa**^[8] in CH_2Cl_2 at room temperature. To our delight, an expected [4 + 1] annulation product **3a** was obtained in 88% combined yield after column chromatographic isolation (Scheme 2). This result revealed that a [4 + 1] annulation reaction between salicyl *N*-tosylimine **1a** and phosphorus ylide **2aa** could be readily achieved.

Scheme 2 Reaction of salicyl imine **1a** and phosphorus ylide **2aa**

Considering that phosphorus ylide **2aa** can be *in situ* generated from the corresponding *N*-phenyl maleimide and PPh_3 in the presence of benzoic acid,^[7b] we then investigated a phosphine catalytic version of the [4 + 1] annulation by using salicyl *N*-tosylimine **1a** and *N*-phenyl maleimide **2a** as the model substrates (Table 1). In the presence of PPh_3 (20 mol%, relative to **2a**) and benzoic acid (20 mol%), a mixture of **1a** (0.2 mmol) and **2a** (0.24 mmol) in CH_2Cl_2 (2.0 mL) was stirred at room temperature for 48 h, and the expected spirocyclic benzofuran **3a** was obtained but in only 23% isolated yield with *dr* 2.5 : 1 (Entry 1). Considering the fact that salicyl *N*-tosylimine **1a** bearing an acidic phenolic hydroxyl may act as a proton shuttle instead of benzoic acid, this reaction was repeated in the absence of benzoic acid and the product **3a** was obtained in 34% yield with 2.9 : 1 *dr* (Entry 2). Without catalyst PPh_3 ,

the annulation reaction could not occur at all (Entry 3). With the catalyst PPh_3 (20 mol%) employed, the substrate ratio of **1a/2a** was surveyed. Gratifyingly, the total yields of **3a** were significantly increased as the ratio of **1a/2a** was adjusted from 1 : 1.2 to 3 : 1 (Entries 2 and 4~7). With the ratio of **1a/2a** 2 : 1 selected, several triarylphosphines were screened (Entries 8~12). The results unveiled that 4-fluoro, 4-chloro, 4-methyl, and 4-methoxy substituted triarylphosphines were all effective for the reaction (Entries 8~11); but $(4\text{-CF}_3\text{C}_6\text{H}_4)_3\text{P}$ was ineffective (Entry 12). With PPh_3 (20 mol%) employed as the catalyst, a couple of common solvents including chloroform, toluene, acetonitrile and methanol were then examined (Entries 13~16). It was found that the model reaction readily gave the product **3a** in moderate yields in nonpolar or polar aprotic solvents (Entries 13~15). Protic solvent methanol was detrimental to the reaction and none of **3a** was obtained (Entry 16). Finally, the loading of the catalyst PPh_3 was also surveyed. As the loading of PPh_3 was increased to 100 mol%, the total yield of **3a** was improved to 99% (Entries 17~19). Thus, the optimal conditions were established as listed in Entry 19, Table 1. With regard to the ratio of *syn*-**3a** versus *anti*-**3a**, in all cases, the product **3a** was isolated with *dr* varying from 1.7 : 1 to 6.7 : 1 (Table 1, Entries 1~19).

With the optimized conditions in hand, the substrate scope of this [4+1] annulation reaction was then explored, and the results are summarized in Table 2. With salicyl

N-tosylimine **1a** used as a representative partner, a series of *N*-substituted maleimides **2** were firstly surveyed (Entries 1~9). Except *N*-(*tert*-butoxycarbonyl) maleimide **2i**, *N*-aryl, *N*-benzyl and *N*-butyl maleimides **2** were all effective substrates in the [4+1] annulations with **1a**, readily affording the corresponding products **3** in modest to excellent yields. With *N*-phenyl maleimide **2a** chosen as one reactant, a series of different salicyl *N*-tosylimines **1** were further examined (Entries 10~18). Both electron-donating and electron-withdrawing substituents R^1 on the benzene ring of salicyl *N*-tosylimines were well tolerated, yielding the corresponding products in modest to excellent yields (Entries 10~16). Salicyl *N*-benzenesulfonylimine **1i** also uneventfully delivered the normal product **3p** in 99% yield (Table 2, Entry 17). Furthermore, the bulky substrate (2-hydroxynaphthalen-1-yl) *N*-tosylimine **1j** was also effective in the reaction with maleimide **2a**, readily giving the corresponding tricyclic product **3q** in good yield (Entry 18). In all cases, the annulation products **3** were obtained as a pair of separable diastereomeric isomers *syn*-**3** and *anti*-**3** in the variable ratio of 1.6 : 1~5.0 : 1 (Table 2). In most of cases, products **3** were isolated as pure diastereomers. Thus, the results indicate that the phosphine-catalyzed [4+1] annulation reaction between salicyl imines **1** and maleimides **2** has a flexible substrate scope and accordingly provides a convenient and efficient method to construct spiro[benzofuran-2,3'-pyrrolidine]-2',5'-diones.

Table 1 Survey of conditions for the phosphine-catalyzed [4+1] annulation reaction of **1a** and **2a**^a

Entry	Catalyst (mol%)	Ratio of 1a/2a	Solvent	Time/h	Yield ^b /%	<i>dr</i> ^c
1 ^d	PPh_3 (20)	1/1.2	CH_2Cl_2	48	23	2.5 : 1
2	PPh_3 (20)	1/1.2	CH_2Cl_2	48	34	2.9 : 1
3	None	1/1.2	CH_2Cl_2	48	0	—
4	PPh_3 (20)	1.2/1	CH_2Cl_2	24	63	2.2 : 1
5	PPh_3 (20)	1.5/1	CH_2Cl_2	24	56	2.3 : 1
6	PPh_3 (20)	2/1	CH_2Cl_2	24	74	1.7 : 1
7	PPh_3 (20)	3/1	CH_2Cl_2	24	78	2.8 : 1
8	$(4\text{-FC}_6\text{H}_4)_3\text{P}$ (20)	2/1	CH_2Cl_2	48	61	2.4 : 1
9	$(4\text{-ClC}_6\text{H}_4)_3\text{P}$ (20)	2/1	CH_2Cl_2	48	41	3.6 : 1
10	$(4\text{-MeC}_6\text{H}_4)_3\text{P}$ (20)	2/1	CH_2Cl_2	24	70	2.7 : 1
11	$(4\text{-MeOC}_6\text{H}_4)_3\text{P}$ (20)	2/1	CH_2Cl_2	24	49	3 : 1
12	$(4\text{-CF}_3\text{C}_6\text{H}_4)_3\text{P}$ (20)	2/1	CH_2Cl_2	48	Trace	—
13	PPh_3 (20)	2/1	CHCl_3	36	50	3 : 1
14	PPh_3 (20)	2/1	Toluene	36	59	3.4 : 1
15	PPh_3 (20)	2/1	CH_3CN	24	66	6.7 : 1
16	PPh_3 (20)	2/1	CH_3OH	24	0	—
17	PPh_3 (50)	2/1	CH_2Cl_2	24	84	3 : 1
18	PPh_3 (70)	2/1	CH_2Cl_2	24	86	2.9 : 1
19	PPh_3 (100)	2/1	CH_2Cl_2	24	99	2.4 : 1

^a Typical conditions: a mixture of **1a**, **2a** and phosphine in solvent (2.0 mL) was stirred at room temperature for a specified time. ^b Isolated yields. ^c Determined by ^1H NMR assay of the crude product. ^d Benzoic acid was added.

Table 2 Synthesis of spiro[benzofuran-2,3'-pyrrolidine]-2',5'-diones **3**^a

Entry	R ¹	R ² (1)	R ³ (2)	Time/h	3a	Yield ^b / % of 3	dr ^c
1	H	Ts (1a)	Ph (2a)	24	3a	99	2.4 : 1
2	H	Ts (1a)	4-MeOC ₆ H ₄ (2b)	24	3b	96	3.9 : 1
3	H	Ts (1a)	4-MeC ₆ H ₄ (2c)	24	3c	44	2.2 : 1
4	H	Ts (1a)	4-ClC ₆ H ₄ (2d)	24	3d	81	2.0 : 1
5	H	Ts (1a)	4-FC ₆ H ₄ (2e)	24	3e	71	2.3 : 1
6	H	Ts (1a)	4-NO ₂ C ₆ H ₄ (2f)	12	3f	80	3.4 : 1
7	H	Ts (1a)	Bn (2g)	12	3g	77	3.2 : 1
8	H	Ts (1a)	<i>n</i> -Bu (2h)	24	3h	65	3.4 : 1
9	H	Ts (1a)	Boc (2i)	96		Trace	—
10	3,5-Cl ₂	Ts (1b)	Ph (2a)	24	3i	92	2.1 : 1
11	3,5-Br ₂	Ts (1c)	Ph (2a)	24	3j	66	3.2 : 1
12	5-Me	Ts (1d)	Ph (2a)	24	3k	86	2.7 : 1
13	5-MeO	Ts (1e)	Ph (2a)	24	3l	91	2.4 : 1
14	5-NO ₂	Ts (1f)	Ph (2a)	96	3m	73	1.6 : 1
15	4-MeO	Ts (1g)	Ph (2a)	24	3n	64	2.7 : 1
16	3-MeO	Ts (1h)	Ph (2a)	96	3o	92	5.0 : 1
17	H	Bs (1i)	Ph (2a)	24	3p	99	3.6 : 1
18	5,6-Benzo	Ts (1j)	Ph (2a)	96	3q	74	3.7 : 1

^a Typical conditions: under N₂ atmosphere, a mixture of salicyl imines **1** (1.0 mmol), maleimide **2** (0.5 mmol) and PPh₃ (0.5 mmol) in CH₂Cl₂ (4.0 mL) was stirred at room temperature for a specified time. ^b Isolated yields based on **2**. ^c Determined by ¹H NMR assay of the crude product.

The structures of products **3** were well identified by ¹H and ¹³C NMR and HRMS. Representative compounds *syn*-**3a** (CCDC 1906851) and *anti*-**3a** (CCDC 1906852) were further confirmed by X-ray crystallographic analyses.

According to the stoichiometric reaction showed in Scheme 2 and our previous work,^[7b,9] a plausible mechanism is exemplified in Scheme 3 to rationalize the formation of **3**. A phosphorus ylide **B** is *in situ* generated via the nucleophilic attack of phosphine at maleimide **2a** followed by proton transfer. The ylide **B** then undergoes a nucleophilic addition to salicyl imine like **1a** to generate intermediate **C**, which subsequently transforms into intermediate **D** by proton transfer. Finally, cyclization occurs by an intramolecular S_N2-like substitution to furnish spirocyclic benzofuran like **3a** and regenerate the phosphine catalyst (Scheme 3). In light of our previous report,^[7b] the *in situ* formation of P-ylide **B** is aided with a protic acid additive such as benzoic acid (20 mol%). In this reaction, the substrate salicyl imines like **1a** bearing a phenolic hydroxyl presumably also acts as a protic acid additive and assists formation of P-ylide **B** with the proton transfer process (Scheme 3). Since the substrate salicyl imine **1a** is in excess as the protic additive, the excessive salicyl imine **1a** may inhibit the [4+1] annulation reaction by protonating the reactive P-ylide **B**. This inhibition may be compensated with the increased loading of the catalyst PPh₃. That is why a relatively higher (up to 100 mol%) catalyst loading is required to attain a good yield (Table 1, Entry 19).

Scheme 3 A plausible mechanism for the [4+1] annulation

3 Conclusions

In conclusion, we have successfully developed a novel phosphine-catalyzed [4+1] annulation reaction between dual functional salicyl imines and maleimides. This work definitely expands the substrate scope of the non-allylic P-ylide-initiated [4+1] annulation strategy. Also, this reaction provides a simple and efficient method for constructing spirobenzofurans from readily available materials and under mild conditions. Mechanistically, this reaction

proceeds through a cascade sequence of nucleophilic addition/intramolecular S_N2 -like substitution involving reactive P-ylide intermediates. Accordingly, it represents a new mode of the non-allylic P-ylide-mediated [4+1] annulation reaction. Further expanding its scope and developing its asymmetric version are currently under investigation in our laboratory.

4 Experimental section

4.1 General methods

Unless otherwise noted, all reactions were carried out in a nitrogen atmosphere under anhydrous conditions. All the solvents were purified according to the standard procedures. ^1H NMR and ^{13}C NMR spectra were recorded on a Bruker AV 400 in CDCl_3 with tetramethylsilane (TMS) as the internal standard. Melting points were measured on a RY-I apparatus and uncorrected. High resolution ESI mass spectra were acquired with a Varian 7.0 T FTMS instrument. X-ray crystal diffraction data were collected on a Nonius Kappa CCD diffractometer with Mo $K\alpha$ radiation ($\lambda=0.7107\text{ \AA}$) at room temperature. Column chromatography was performed on silica gel (200~300 mesh) using a mixture of petroleum ether/ethyl acetate as eluent. Substrates **1** were prepared according to literature methods.^[10] Maleimides **2** were prepared through the reported method from maleic anhydride and the corresponding amines.^[11]

4.2 General procedure for PPh_3 -catalyzed [4+1] annulation reaction between **1** and **2**

Under a N_2 atmosphere, a mixture of **1** (1 mmol), **2** (0.5 mmol), and PPh_3 (0.5 mmol) in CH_2Cl_2 (4.0 mL) was stirred at room temperature for the specified hours (monitored by TLC). After the reaction was completed, the mixture was concentrated on a rotary evaporator under reduced pressure and the residue was subjected to purification by column chromatography on silica gel (gradient elution with a mixture of petroleum ether/ethyl acetate from 5 : 1 to 3 : 1) to afford the [4+1] annulation products **3**.

N-(2',5'-Dioxo-1'-phenyl-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)-4-methylbenzenesulfonamide (**3a**): 99% yield, 2.4 : 1 *dr*. For *syn*-**3a**, white solid, m.p. 198~200 °C; ^1H NMR (CDCl_3 , 400 MHz) δ : 7.80 (d, $J=8.3$ Hz, 2H), 7.54~7.32 (m, 7H), 7.23 (t, $J=7.9$ Hz, 1H), 6.93~6.79 (m, 2H), 6.72 (d, $J=7.5$ Hz, 1H), 5.53 (d, $J=7.2$ Hz, 1H), 5.30 (d, $J=7.2$ Hz, 1H), 3.79 (d, $J=18.6$ Hz, 1H), 2.96 (d, $J=18.6$ Hz, 1H), 2.44 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 174.6, 172.4, 157.5, 144.7, 135.8, 131.5, 131.2, 130.3, 129.3, 129.0, 127.3, 126.7, 124.2, 123.1, 122.4, 110.7, 88.6, 59.3, 37.0, 21.6. For *anti*-**3a**, white solid, m.p. 191~194 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.01 (d, $J=5.7$ Hz, 1H), 7.89 (d, $J=8.2$ Hz, 2H), 7.57~7.38 (m, 7H), 7.22 (t, $J=7.8$ Hz, 1H), 6.89 (d, $J=8.0$ Hz, 1H), 6.84 (t, $J=7.4$ Hz, 1H), 6.40 (d, $J=7.5$ Hz, 1H), 5.52 (d, $J=5.7$ Hz, 1H), 3.51 (s, 2H), 2.43 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 173.6, 172.4, 158.9, 143.7, 138.3, 132.5, 130.7, 130.3, 129.5, 129.2, 127.7, 127.2, 125.2, 124.4, 121.9, 110.1, 89.8, 64.4, 41.4,

21.5; HRMS-ESI calcd for $\text{C}_{24}\text{H}_{21}\text{N}_2\text{O}_5\text{S}$ [$\text{M} + \text{H}$] $^+$ 449.1166, found 449.1167.

N-(1'-(4-Methoxyphenyl)-2',5'-dioxo-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)-4-methylbenzenesulfonamide (**3b**): 96% yield, 3.9 : 1 *dr*. For *syn*-**3b**, white solid, m.p. 206~208 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.14 (d, $J=8.1$ Hz, 1H), 7.83 (d, $J=8.2$ Hz, 2H), 7.49 (d, $J=8.1$ Hz, 2H), 7.31~7.25 (m, 3H), 7.10 (d, $J=9.0$ Hz, 2H), 6.96~6.89 (m, 2H), 6.67 (d, $J=7.5$ Hz, 1H), 5.17 (d, $J=8.1$ Hz, 1H), 3.82 (s, 3H), 3.50 (d, $J=18.2$ Hz, 1H), 3.21 (d, $J=18.2$ Hz, 1H), 2.42 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 175.0, 172.1, 159.3, 157.3, 143.6, 137.4, 130.6, 130.1, 128.2, 126.6, 124.8, 124.2, 123.5, 122.0, 114.3, 110.0, 87.8, 59.7, 55.4, 36.4, 21.0; HRMS-ESI calcd for $\text{C}_{25}\text{H}_{23}\text{N}_2\text{O}_6\text{S}$ [$\text{M} + \text{H}$] $^+$ 479.1272, found 479.1284.

N-(2',5'-Dioxo-1'-(*p*-tolyl)-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)-4-methylbenzenesulfonamide (**3c**): 44% yield, 2.2 : 1 *dr*. For *syn*-**3c**, white solid, m.p. 200~202 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.15 (s, 1H), 7.83 (d, $J=8.2$ Hz, 2H), 7.48 (d, $J=8.1$ Hz, 2H), 7.35 (d, $J=8.2$ Hz, 2H), 7.30~7.22 (m, 3H), 6.97~6.88 (m, 2H), 6.67 (d, $J=7.5$ Hz, 1H), 5.17 (s, 1H), 3.51 (d, $J=18.2$ Hz, 1H), 3.23 (d, $J=18.2$ Hz, 1H), 2.41 (s, 3H), 2.37 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 175.4, 172.4, 157.8, 144.1, 138.9, 137.9, 131.1, 130.6, 130.0, 129.6, 127.3, 127.1, 125.4, 124.0, 122.5, 110.5, 88.3, 60.2, 36.9, 21.5, 21.3. For *anti*-**3c**, white solid, m.p. 188~190 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.02 (d, $J=6.9$ Hz, 1H), 7.90 (d, $J=8.2$ Hz, 2H), 7.48 (d, $J=8.1$ Hz, 2H), 7.34 (d, $J=8.3$ Hz, 2H), 7.29 (d, $J=8.4$ Hz, 2H), 7.22 (t, $J=7.8$ Hz, 1H), 6.90 (d, $J=8.0$ Hz, 1H), 6.84 (t, $J=7.5$ Hz, 1H), 6.40 (d, $J=7.5$ Hz, 1H), 5.51 (d, $J=6.9$ Hz, 1H), 3.50 (s, 2H), 2.43 (s, 3H), 2.37 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 173.7, 172.5, 158.9, 143.7, 138.8, 138.3, 130.7, 130.3, 129.9, 129.9, 127.5, 127.2, 125.1, 124.4, 121.9, 110.1, 89.8, 64.3, 41.3, 21.5, 21.2; HRMS-ESI calcd for $\text{C}_{25}\text{H}_{23}\text{N}_2\text{O}_5\text{S}$ [$\text{M} + \text{H}$] $^+$ 463.1322, found 463.1334.

N-(1'-(4-Chlorophenyl)-2',5'-dioxo-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)-4-methylbenzenesulfonamide (**3d**): 81% yield, 2.0 : 1 *dr*. For *syn*-**3d**, white solid, m.p. 192~193 °C; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ : 9.16 (d, $J=8.1$ Hz, 1H), 7.82 (d, $J=8.2$ Hz, 2H), 7.64 (d, $J=8.7$ Hz, 2H), 7.47 (d, $J=8.1$ Hz, 2H), 7.41 (d, $J=8.7$ Hz, 2H), 7.27 (t, $J=7.7$ Hz, 1H), 6.96~6.88 (m, 2H), 6.65 (d, $J=7.5$ Hz, 1H), 5.19 (d, $J=8.1$ Hz, 1H), 3.52 (d, $J=18.1$ Hz, 1H), 3.24 (d, $J=18.1$ Hz, 1H), 2.40 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 174.6, 171.6, 157.3, 143.6, 137.4, 133.4, 130.6, 130.5, 130.1, 129.2, 128.8, 126.6, 124.8, 123.5, 122.0, 110.1, 87.9, 59.9, 36.5, 21.0. For *anti*-**3d**, white solid, m.p. 182~184 °C; ^1H NMR (CDCl_3 , 400 MHz) δ : 7.81 (d, $J=8.3$ Hz, 2H), 7.45~7.40 (m, 2H), 7.33 (dt, $J=3.9, 2.2$ Hz, 4H), 7.22 (t, $J=7.7$ Hz, 1H), 6.85 (t, $J=7.6$ Hz, 2H), 6.71~6.66 (m, 1H), 5.68 (d, $J=9.5$ Hz, 1H), 5.26 (d, $J=9.5$ Hz, 1H), 3.33 (s, 2H), 2.45 (s, 3H); ^{13}C NMR (100 MHz, $\text{DMSO}-d_6$) δ : 172.5, 171.7, 158.4, 144.5, 136.6, 134.8, 131.1, 130.1, 129.6, 129.5, 128.0, 127.2, 124.2, 123.5, 122.4, 110.5, 88.1, 64.0, 41.4, 21.6;

HRMS-ESI calcd for $C_{24}H_{23}ClN_3O_5S$ $[M + NH_4]^+$ 500.1041, found 500.1048.

N-(1'-(4-Fluorophenyl)-2',5'-dioxo-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)-4-methylbenzenesulfonamide (**3e**): 71% yield, 2.3 : 1 *dr*. For *syn*-**3e**, white solid, m.p. 216~218 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 9.16 (d, J =8.1 Hz, 1H), 7.83 (d, J =8.2 Hz, 2H), 7.48 (d, J =8.1 Hz, 2H), 7.45~7.37 (m, 4H), 7.27 (t, J =7.7 Hz, 1H), 6.97~6.87 (m, 2H), 6.66 (d, J =7.5 Hz, 1H), 5.19 (d, J =8.1 Hz, 1H), 3.52 (d, J =18.1 Hz, 1H), 3.24 (d, J =18.1 Hz, 1H), 2.41 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 175.3, 172.3, 162.2 (d, J_{CF} =245.7 Hz), 157.8, 144.1, 137.9, 131.1, 130.6, 129.8 (d, J_{CF} =9.0 Hz), 128.4 (d, J_{CF} =2.9 Hz), 127.1, 125.3, 124.0, 122.5, 116.6 (d, J_{CF} =23.0 Hz), 110.6, 88.3, 60.4, 37.0, 21.5. For *anti*-**3e**, white solid, m.p. 174~176 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 9.02 (d, J =7.0 Hz, 1H), 7.89 (d, J =8.2 Hz, 2H), 7.52~7.39 (m, 6H), 7.23 (t, J =7.8 Hz, 1H), 6.90 (d, J =8.1 Hz, 1H), 6.85 (t, J =7.5 Hz, 1H), 6.42 (d, J =7.5 Hz, 1H), 5.52 (d, J =7.0 Hz, 1H), 3.51 (s, 2H), 2.43 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 173.5, 172.3, 162.2 (d, J =245.5 Hz), 158.9, 143.7, 138.3, 130.8, 130.3, 129.9 (d, J =8.9 Hz), 128.7 (d, J =2.8 Hz), 127.2, 125.2, 124.3, 121.9, 116.5 (d, J =23.0 Hz), 110.1, 89.9, 64.4, 41.3, 21.5; HRMS-ESI calcd for $C_{24}H_{23}FN_3O_5S$ $[M + NH_4]^+$ 484.1337, found 484.1343.

4-Methyl-*N*-(1'-(4-nitrophenyl)-2',5'-dioxo-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)benzenesulfonamide (**3f**): 80% yield, 3.4 : 1 *dr*. For *syn*-**3f**, white solid, m.p. 193~195 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 9.18 (d, J =8.1 Hz, 1H), 8.44 (d, J =8.9 Hz, 2H), 7.83 (d, J =8.1 Hz, 2H), 7.71 (d, J =8.9 Hz, 2H), 7.47 (d, J =8.0 Hz, 2H), 7.28 (t, J =7.7 Hz, 1H), 6.93 (dd, J =13.8, 7.6 Hz, 2H), 6.66 (d, J =7.4 Hz, 1H), 5.23 (d, J =8.1 Hz, 1H), 3.55 (d, J =18.1 Hz, 1H), 3.30 (d, J =18.1 Hz, 1H), 2.41 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 174.8, 171.7, 157.8, 147.5, 144.1, 137.9, 137.6, 131.1, 130.5, 128.5, 127.1, 125.4, 124.9, 123.9, 122.6, 110.6, 88.4, 60.4, 37.1, 21.5; HRMS-ESI calcd for $C_{24}H_{23}N_4O_7S$ $[M + NH_4]^+$ 511.1282, found 511.1286.

N-(1'-Benzyl-2',5'-dioxo-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)-4-methylbenzenesulfonamide (**3g**): 77% yield, 3.2 : 1 *dr*. For *syn*-**3g**, white solid, m.p. 157~159 °C; 1H NMR (CDCl₃, 400 MHz) δ : 7.67 (d, J =8.2 Hz, 2H), 7.42~7.29 (m, 7H), 7.20 (t, J =7.8 Hz, 1H), 6.84 (t, J =7.5 Hz, 1H), 6.78 (d, J =8.1 Hz, 1H), 6.64 (d, J =7.5 Hz, 1H), 5.24 (d, J =7.6 Hz, 1H), 5.07 (d, J =7.6 Hz, 1H), 4.74 (d, J =14.3 Hz, 1H), 4.67 (d, J =14.3 Hz, 1H), 3.59 (d, J =18.6 Hz, 1H), 2.83 (d, J =18.6 Hz, 1H), 2.45 (s, 3H); ^{13}C NMR (CDCl₃, 100 MHz) δ : 175.0, 172.7, 157.7, 144.6, 136.0, 135.1, 131.2, 130.2, 128.7(2C), 128.0, 127.2, 124.4, 123.5, 122.4, 110.5, 88.3, 77.4, 77.2, 77.0, 76.7, 58.8, 42.7, 36.8, 21.6. For *anti*-**3g**, white solid, m.p. 187~189 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 8.93 (d, J =6.8 Hz, 1H), 7.87 (d, J =8.2 Hz, 2H), 7.47 (d, J =8.1 Hz, 2H), 7.33 (q, J =3.3 Hz, 4H), 7.31~7.26 (m, 1H), 7.19 (t, J =7.8 Hz, 1H), 6.87~6.78 (m, 2H), 6.36 (d, J =7.5 Hz, 1H), 5.46 (d, J =6.8 Hz, 1H), 4.70 (d, J =15.3 Hz, 1H), 4.60 (d,

J =15.3 Hz, 1H), 3.41 (s, 2H), 2.43 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 174.0, 173.5, 158.7, 143.7, 138.1, 135.9, 130.6, 130.2, 129.0, 127.8, 127.7, 127.4, 125.0, 124.8, 121.8, 110.1, 89.6, 63.5, 42.2, 40.8, 21.5; HRMS-ESI calcd for $C_{25}H_{23}N_2O_5S$ $[M + H]^+$ 463.1322, found 463.1323.

N-(1'-Butyl-2',5'-dioxo-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)-4-methylbenzenesulfonamide (**3h**): 65% yield, 3.4 : 1 *dr*. For *syn*-**3h**, white solid, m.p. 127~129 °C; 1H NMR (CDCl₃, 400 MHz) δ : 7.78 (d, J =8.2 Hz, 2H), 7.38 (d, J =8.1 Hz, 2H), 7.20 (t, J =7.8 Hz, 1H), 6.85 (t, J =7.5 Hz, 1H), 6.79 (d, J =8.1 Hz, 1H), 6.67 (d, J =7.5 Hz, 1H), 5.34 (d, J =7.6 Hz, 1H), 5.17 (d, J =7.6 Hz, 1H), 3.56 (q, J =8.6 Hz, 2H), 3.55 (d, J =18.5 Hz, 1H), 2.79 (d, J =18.5 Hz, 1H), 2.47 (s, 3H), 1.66~1.57 (m, 2H), 1.41~1.31 (m, 2H), 0.95 (t, J =7.3 Hz, 3H); ^{13}C NMR (CDCl₃, 100 MHz) δ : 175.3, 173.3, 157.6, 144.6, 136.2, 131.1, 130.2, 127.3, 124.3, 123.6, 122.3, 110.6, 88.4, 58.8, 39.1, 36.8, 29.5, 21.6, 20.0, 13.6. For *anti*-**3h**, white solid, m.p. 161~163 °C; 1H NMR (CDCl₃, 400 MHz) δ : 7.83 (d, J =8.3 Hz, 2H), 7.38 (d, J =8.0 Hz, 2H), 7.25 (t, J =7.7 Hz, 1H), 6.92~6.88 (m, 1H), 6.83 (d, J =8.1 Hz, 1H), 6.73 (d, J =7.5 Hz, 1H), 5.54 (d, J =10.0 Hz, 1H), 5.29 (d, J =10.0 Hz, 1H), 3.55~3.49 (m, 2H), 3.17 (d, J =18.4 Hz, 1H), 3.09 (d, J =18.4 Hz, 1H), 2.49 (s, 3H), 1.66~1.58 (m, 2H), 1.37 (dq, J =14.7, 7.3 Hz, 2H), 0.96 (t, J =7.3 Hz, 3H); ^{13}C NMR (CDCl₃, 101 MHz) δ : 173.6, 173.0, 158.3, 144.4, 137.1, 130.8, 130.1, 127.1, 124.2, 124.1, 122.3, 110.4, 87.6, 63.1, 40.9, 39.0, 29.5, 21.6, 20.0, 13.6; HRMS-ESI calcd for $C_{22}H_{28}N_3O_5S$ $[M + NH_4]^+$ 446.1744, found 446.1747.

N-(5,7-Dichloro-2',5'-dioxo-1'-phenyl-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)-4-methylbenzenesulfonamide (**3i**): 92% yield, 2.1 : 1 *dr*. For *syn*-**3i**, white solid, m.p. 209~211 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 9.13 (d, J =8.5 Hz, 1H), 7.84 (d, J =8.1 Hz, 2H), 7.59~7.53 (m, 3H), 7.52~7.47 (m, 3H), 7.38 (d, J =7.7 Hz, 2H), 6.28 (s, 1H), 5.36 (d, J =8.5 Hz, 1H), 3.52 (d, J =18.1 Hz, 1H), 3.41 (d, J =18.1 Hz, 1H), 2.43 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 173.6, 171.3, 152.6, 143.9, 137.6, 131.5, 130.1, 129.9, 129.0, 128.8, 127.8, 127.0, 126.7, 125.9, 123.6, 115.2, 89.3, 59.7, 36.1, 21.0. For *anti*-**3i**, white solid, m.p. 199~201 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 9.13 (d, J =7.4 Hz, 1H), 7.89 (d, J =8.2 Hz, 2H), 7.58~7.46 (m, 6H), 7.44~7.35 (m, 2H), 6.05 (dd, J =1.9, 1.1 Hz, 1H), 5.71 (d, J =7.4 Hz, 1H), 3.65 (d, J =17.7 Hz, 1H), 3.50 (d, J =17.7 Hz, 1H), 2.45 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 172.7, 171.8, 153.9, 144.1, 138.0, 132.3, 130.4, 130.0, 129.5, 129.3, 128.4, 127.7, 127.2, 125.9, 123.7, 115.2, 91.1, 64.2, 40.8, 21.5; HRMS-ESI calcd for $C_{24}H_{22}Cl_2N_3O_5S$ $[M + NH_4]^+$ 534.0652, found 534.0656.

N-(5,7-Dibromo-2',5'-dioxo-1'-phenyl-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)-4-methylbenzenesulfonamide (**3j**): 66% yield, 3.2 : 1 *dr*. For *syn*-**3j**, white solid, m.p. 221~224 °C; 1H NMR (400 MHz, DMSO- d_6) δ : 9.10 (d, J =8.4 Hz, 1H), 7.82 (d, J =7.7 Hz, 2H), 7.74 (s, 1H), 7.55

(t, $J=7.0$ Hz, 2H), 7.48 (d, $J=7.8$ Hz, 3H), 7.37 (d, $J=7.4$ Hz, 2H), 6.31 (d, $J=8.3$ Hz, 1H), 5.37 (d, $J=8.3$ Hz, 1H), 3.50 (d, $J=18.0$ Hz, 1H), 3.36 (d, $J=18.0$ Hz, 1H), 2.43 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 174.1, 171.7, 155.0, 144.4, 138.1, 135.4, 132.0, 130.6, 129.5, 129.4, 128.3, 127.5, 127.4, 127.2, 113.8, 103.9, 89.5, 60.3, 36.6, 21.6; HRMS-ESI calcd for $\text{C}_{24}\text{H}_{22}\text{Br}_2\text{N}_3\text{O}_5\text{S}$ [$\text{M}+\text{NH}_4$] $^+$ 621.9641, found 621.9647.

4-Methyl-*N*-(5-methyl-2',5'-dioxo-1'-phenyl-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)benzenesulfonamide (**3k**): Isolated as a diastereomeric mixture, 86% yield, *dr* 2.7 : 1; white solid. For the major *syn*-**3k**, ^1H NMR (400 MHz, DMSO- d_6) δ : 9.05 (d, $J=8.2$ Hz, 1H), 7.81 (d, $J=8.1$ Hz, 2H), 7.58~7.46 (m, 5H), 7.35 (d, $J=7.6$ Hz, 2H), 7.05 (d, $J=8.2$ Hz, 1H), 6.80 (d, $J=8.2$ Hz, 1H), 6.26 (s, 1H), 5.14 (d, $J=8.1$ Hz, 1H), 3.50 (d, $J=18.1$ Hz, 1H), 3.20 (d, $J=18.1$ Hz, 1H), 2.42 (s, 3H), 2.11 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 174.8, 171.9, 155.3, 143.6, 137.6, 131.7, 130.8, 130.5, 130.1, 129.1, 128.8, 127.0, 126.7, 125.1, 123.6, 109.6, 87.9, 59.7, 36.5, 21.0, 20.2. For the minor *anti*-**3k**, ^1H NMR (400 MHz, DMSO- d_6) δ : 8.89 (d, $J=7.2$ Hz, 1H), 7.89 (d, $J=8.1$ Hz, 2H), 7.58~7.46 (m, 5H), 7.39 (d, $J=7.5$ Hz, 2H), 7.01 (d, $J=8.4$ Hz, 1H), 6.77 (d, $J=8.2$ Hz, 1H), 6.05 (s, 1H), 5.45 (d, $J=7.1$ Hz, 1H), 3.48 (s, 1H), 3.34 (s, 1H), 2.44 (s, 3H), 2.08 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 174.5, 173.0, 156.5, 143.2, 137.9, 132.0, 130.8, 130.5, 129.8, 129.0, 128.7, 127.2, 126.8, 125.0, 123.9, 109.2, 89.3, 63.9, 40.9, 21.0, 20.3; HRMS-ESI calcd for $\text{C}_{25}\text{H}_{23}\text{N}_2\text{O}_5\text{S}$ [$\text{M}+\text{H}$] $^+$ 463.1322, found 463.1331.

N-(5-Methoxy-2',5'-dioxo-1'-phenyl-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)-4-methylbenzenesulfonamide (**3l**): Isolated as a diastereomeric mixture, 91% yield, *dr* 2.4 : 1; white solid. For the major *syn*-**3l**, ^1H NMR (400 MHz, DMSO- d_6) δ : 9.09 (d, $J=8.3$ Hz, 1H), 7.84 (d, $J=8.3$ Hz, 2H), 7.58~7.46 (m, 5H), 7.42~7.34 (m, 2H), 6.80 (tt, $J=8.8$, 5.6 Hz, 2H), 5.95 (d, $J=2.3$ Hz, 1H), 5.16 (d, $J=8.2$ Hz, 1H), 3.53 (s, 3H), 3.50 (d, $J=18.0$ Hz, 1H), 3.21 (d, $J=18.0$ Hz, 1H), 2.41 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 175.2, 172.3, 155.0, 151.7, 144.2, 138.3, 132.2, 130.6, 129.6, 129.3, 127.5, 127.2, 124.8, 116.7, 111.0, 110.0, 88.6, 60.5, 56.0, 37.0, 21.5. For the minor *anti*-**3l**, ^1H NMR (400 MHz, DMSO- d_6) δ : 8.91 (d, $J=7.4$ Hz, 1H), 7.91 (d, $J=8.2$ Hz, 2H), 7.58~7.46 (m, 5H), 7.42~7.34 (m, 2H), 6.80 (tt, $J=8.8$, 5.6 Hz, 2H), 5.73 (d, $J=2.3$ Hz, 1H), 5.48 (d, $J=7.4$ Hz, 1H), 3.50 (s, 5H), 2.41 (s, 3H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 173.6, 172.3, 154.5, 152.8, 143.8, 138.1, 132.5, 130.3, 129.5, 129.2, 127.7, 127.3, 125.2, 116.0, 110.4, 110.3, 89.9, 64.6, 55.9, 41.3, 21.5; HRMS-ESI calcd for $\text{C}_{25}\text{H}_{23}\text{N}_2\text{O}_6\text{S}$ [$\text{M}+\text{H}$] $^+$ 479.1272, found 479.1279.

4-Methyl-*N*-(5-nitro-2',5'-dioxo-1'-phenyl-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)benzenesulfonamide (**3m**): 73% yield, 1.6 : 1 *dr*. For *anti*-**3m**, white solid, m.p. 175~177 $^{\circ}\text{C}$; ^1H NMR (CDCl_3 , 400 MHz) δ : 8.18 (dd, $J=8.9$, 2.1 Hz, 1H), 7.87 (d, $J=8.0$ Hz, 2H), 7.51 (t, $J=7.7$ Hz, 2H), 7.44 (dd, $J=10.8$, 7.6 Hz, 3H), 7.36 (d, $J=8.0$

Hz, 2H), 7.33 (s, 1H), 6.92 (d, $J=8.9$ Hz, 1H), 5.58 (d, $J=10.3$ Hz, 1H), 5.40 (d, $J=10.3$ Hz, 1H), 3.43 (s, 2H), 2.51 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 171.8, 171.2, 163.0, 145.5, 143.3, 136.3, 130.8, 130.5, 129.4, 129.4, 127.9, 127.1, 126.6, 125.8, 120.9, 110.8, 89.9, 62.3, 40.5, 21.7; HRMS-ESI calcd for $\text{C}_{24}\text{H}_{23}\text{N}_4\text{O}_7\text{S}$ [$\text{M}+\text{NH}_4$] $^+$ 511.1282, found 511.1289.

N-(6-Methoxy-2',5'-dioxo-1'-phenyl-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)-4-methylbenzenesulfonamide (**3n**): 64% yield, *dr* 2.7 : 1. For *syn*-**3n**, white solid; m.p. 210~211 $^{\circ}\text{C}$; ^1H NMR (CDCl_3 , 400 MHz) δ : 7.81 (d, $J=7.9$ Hz, 2H), 7.44 (d, $J=7.9$ Hz, 2H), 7.38~7.30 (m, 4H), 7.26 (s, 1H), 6.55 (d, $J=8.3$ Hz, 1H), 6.50~6.26 (m, 2H), 5.22 (d, $J=7.2$ Hz, 1H), 5.07 (d, $J=7.2$ Hz, 1H), 3.82 (d, $J=18.6$ Hz, 1H), 3.75 (s, 3H), 2.99 (d, $J=18.6$ Hz, 1H), 2.46 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 174.3, 172.2, 162.7, 159.1, 144.7, 136.1, 131.5, 130.3, 129.3, 129.0, 127.3, 126.6, 124.4, 115.0, 109.0, 96.6, 89.3, 59.0, 55.6, 37.0, 21.6. For *anti*-**3n**, white solid; m.p. 182~184 $^{\circ}\text{C}$; ^1H NMR (CDCl_3 , 400 MHz) δ : 7.83 (d, $J=8.3$ Hz, 2H), 7.52~7.46 (m, 2H), 7.45~7.33 (m, 5H), 6.60 (d, $J=8.2$ Hz, 1H), 6.42 (dt, $J=6.4$, 2.1 Hz, 2H), 5.47 (d, $J=9.8$ Hz, 1H), 5.23 (d, $J=9.9$ Hz, 1H), 3.75 (s, 3H), 3.34 (d, $J=18.3$ Hz, 1H), 3.28 (d, $J=18.3$ Hz, 1H), 2.46 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 172.7, 171.9, 162.5, 159.9, 144.5, 137.0, 131.2, 130.1, 129.2, 129.0, 127.2, 126.7, 124.6, 115.4, 108.7, 96.6, 88.7, 63.6, 55.6, 41.5, 21.6; HRMS-ESI calcd for $\text{C}_{25}\text{H}_{26}\text{N}_3\text{O}_6\text{S}$ [$\text{M}+\text{NH}_4$] $^+$ 496.1537, found 496.1538.

N-(7-Methoxy-2',5'-dioxo-1'-phenyl-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)-4-methylbenzenesulfonamide (**3o**): 92% yield, *dr* 5 : 1. For *syn*-**3o**, white solid; m.p. 168~170 $^{\circ}\text{C}$; ^1H NMR (CDCl_3 , 400 MHz) δ : 7.80 (d, $J=8.2$ Hz, 2H), 7.50 (t, $J=7.4$ Hz, 2H), 7.44 (d, $J=7.3$ Hz, 1H), 7.40~7.31 (m, 4H), 6.88~6.80 (m, 2H), 6.33 (d, $J=7.0$ Hz, 1H), 5.46 (d, $J=7.2$ Hz, 1H), 5.33 (d, $J=7.2$ Hz, 1H), 3.86 (s, 3H), 3.81 (d, $J=18.7$ Hz, 1H), 3.03 (d, $J=18.7$ Hz, 1H), 2.44 (s, 3H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 174.2, 172.3, 146.0, 144.7, 135.8, 131.5, 130.3, 129.2, 129.0, 127.3, 126.7, 124.3, 123.4, 115.6, 113.5, 89.0, 59.8, 56.1, 37.0, 21.6. For *anti*-**3o**, white solid; m.p. 200~202 $^{\circ}\text{C}$; ^1H NMR (CDCl_3 , 400 MHz) δ : 7.84 (d, $J=8.1$ Hz, 2H), 7.53~7.32 (m, 7H), 6.90~6.79 (m, 2H), 6.37 (d, $J=6.9$ Hz, 1H), 5.55 (d, $J=9.9$ Hz, 1H), 5.34 (d, $J=9.9$ Hz, 1H), 3.87 (s, 3H), 3.44 (d, $J=18.5$ Hz, 1H), 3.31 (d, $J=18.5$ Hz, 1H), 2.45 (s, 3H); ^{13}C NMR (CDCl_3 , 101 MHz) δ : 172.4, 171.9, 146.8, 144.5, 136.8, 131.2, 130.2, 129.2, 129.0, 127.2, 126.7, 124.9, 123.3, 115.8, 113.5, 88.4, 64.1, 56.1, 41.4, 21.6; HRMS-ESI calcd for $\text{C}_{25}\text{H}_{26}\text{N}_3\text{O}_6\text{S}$ [$\text{M}+\text{NH}_4$] $^+$ 496.1537, found 496.1545.

4-Bromo-*N*-(2',5'-dioxo-1'-phenyl-3*H*-spiro[benzofuran-2,3'-pyrrolidin]-3-yl)benzenesulfonamide (**3p**): Isolated as a diastereomeric mixture, 99% yield, *dr* 3.6 : 1; white solid. For the major *syn*-**3p**, ^1H NMR (400 MHz, DMSO- d_6) δ : 9.22 (d, $J=7.8$ Hz, 1H), 7.99~7.90 (m, 2H), 7.77~7.72 (m, 1H), 7.69 (d, $J=7.7$ Hz, 2H), 7.59~7.53 (m, 2H), 7.50~7.46 (m, 1H), 7.40~7.35 (m, 2H), 7.27 (t, $J=$

7.7 Hz, 1H), 6.99~6.84 (m, 2H), 6.58 (d, $J=7.6$ Hz, 1H), 5.22 (d, $J=7.6$ Hz, 1H), 3.52 (d, $J=18.1$ Hz, 1H), 3.26 (d, $J=18.1$ Hz, 1H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 175.3, 172.3, 157.9, 140.9, 133.7, 132.2, 131.1, 130.2, 129.6, 129.3, 127.5, 127.0, 125.3, 124.0, 122.5, 110.6, 88.4, 60.3, 37.0. For the minor *anti*-**3p**, ^1H NMR (400 MHz, DMSO- d_6) δ : 9.10 (d, $J=7.1$ Hz, 1H), 8.06~7.99 (m, 2H), 7.77~7.72 (m, 1H), 7.66 (d, $J=1.2$ Hz, 2H), 7.59~7.53 (m, 2H), 7.50~7.46 (m, 1H), 7.44~7.40 (m, 2H), 7.21 (d, $J=8.6$ Hz, 1H), 6.99~6.84 (m, 1H), 6.82 (t, $J=7.4$ Hz, 1H), 6.33 (d, $J=7.5$ Hz, 1H), 5.56 (d, $J=7.1$ Hz, 1H), 3.53 (s, 2H); ^{13}C NMR (100 MHz, DMSO- d_6) δ : 173.6, 172.4, 158.9, 141.1, 133.4, 132.5, 130.8, 129.9, 129.5, 129.2, 127.7, 127.2, 125.1, 124.3, 121.9, 110.2, 89.8, 64.3, 41.3; HRMS-ESI calcd for $\text{C}_{23}\text{H}_{22}\text{N}_3\text{O}_5\text{S} [\text{M}+\text{NH}_4]^+$ 452.1274, found 452.1282.

N-(2',5'-Dioxo-1'-phenyl-1*H*-spiro[naphtho[2,1-*b*]furan-2,3'-pyrrolidin]-1-yl)-4-methylbenzenesulfonamide (**3q**): 74% yield, 3.7 : 1 *dr*. For *syn*-**3q**, white solid, m.p. 196~198 °C; ^1H NMR (CDCl_3 , 400 MHz) δ : 7.83 (d, $J=8.2$ Hz, 2H), 7.73 (t, $J=8.2$ Hz, 2H), 7.56~7.30 (m, 7H), 7.25 (t, $J=7.3$ Hz, 1H), 7.13~7.02 (m, 2H), 6.55 (d, $J=8.3$ Hz, 1H), 5.64 (d, $J=7.4$ Hz, 1H), 5.12 (d, $J=7.4$ Hz, 1H), 3.87 (d, $J=18.7$ Hz, 1H), 3.08 (d, $J=18.7$ Hz, 1H), 2.50 (s, 3H); ^{13}C NMR (CDCl_3 , 100 MHz) δ : 174.3, 172.1, 156.6, 144.8, 136.2, 132.8, 131.5, 130.3, 130.0, 129.5, 129.2, 129.2, 128.9, 127.6, 127.4, 126.6, 123.9, 120.9, 113.5, 112.0, 89.4, 59.5, 37.4, 21.7; HRMS-ESI calcd for $\text{C}_{28}\text{H}_{26}\text{N}_3\text{O}_5\text{S} [\text{M}+\text{NH}_4]^+$ 516.1587, found 516.1591.

Supporting Information Copies of ^1H NMR and ^{13}C NMR spectra for **3**, X-ray crystallographic data, and ORTEP drawings for *syn*-**3a** and *anti*-**3a**. This material is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Hemelaere, R.; Carreaux, F.; Carboni, B. *Eur. J. Org. Chem.* **2015**, 11, 2470.
(b) Wu, B.; Chen, M.-W.; Ye, Z.-S.; Yu, C.-B.; Zhou, Y.-G. *Adv. Synth. Catal.* **2014**, 356, 383.
(c) Luo, L.; Liu, C.; Hou, Z.; Wang, Y.; Dai, L. *RSC Adv.* **2014**, 4, 29527.
(d) Liu, Y.; Lu, A.; Hu, K.; Wang, Y.; Song, H.; Zhou, Z.; Tang, C. *Eur. J. Org. Chem.* **2013**, 22, 4836.
(e) Coy B, E. D.; Jovanovic, L.; Sefkow, M. *Org. Lett.* **2010**, 12, 1976.
(f) Xie, P.; Huang, Y.; Chen, R. *Org. Lett.* **2010**, 12, 3768.
(g) Leticia, J.-G.; Sergio, G.-M.; Miriam, A.-C.; Manuel, M.-D.; Ignacio, R.-G. *Chem. Eur. J.* **2007**, 13, 557.
(h) Barton, D. H. R.; Nakanishi, K.; MethCohn, O.; Kelly, J. W. *Comprehensive Natural Products Chemistry*, Pergamon Press, Oxford, UK, **1999**.
(i) Qin, Z.; Liu, W.; Wang, D.; He, Z. *J. Org. Chem.* **2016**, 81, 4690.
(j) Cheng, Y.; Fang, Z.; Li, W.; Li, P. *Org. Chem. Front.* **2018**, 5, 2728.
- [2] (a) Li, A.; Piel, J. *Chem. Biol.* **2002**, 9, 1017.
(b) Tsang, K. Y.; Brimble, M. A. *Tetrahedron* **2007**, 63, 6015.
(c) Pirrung, M. C.; Brown, W. L.; Rage, S.; Laughton, P. *J. Am. Chem. Soc.* **1991**, 113, 8561.
(d) Usegilo, M.; Castellano, P. M.; Operto, M. A.; Torres, R.; Kaufman, T. S. *Bioorg. Med. Chem. Lett.* **2006**, 16, 5097.
(e) Padmanaban, M.; Bijou, A. T.; Glorius, F. *Org. Lett.* **2011**, 13, 5624.
(f) Dong, N.; Li, X.; Wang, F.; Cheng, J.-P. *Org. Lett.* **2013**, 15, 4896.
(g) Petersen, A. B.; Ronnest, M. H.; Larsen, T. O.; Clausen, M. H. *Chem. Rev.* **2014**, 114, 12088.
(h) Effland, R. C.; Gardner, B. A.; Strupczewski, J. J. *Heterocycl. Chem.* **1981**, 18, 811.
(i) Kyei, A. S.; Tchabanenko, K.; Baldwin, J. E.; Adlington, R. M. *Tetrahedron Lett.* **2004**, 45, 8931.
(j) Cohen, J. L.; Limon, A.; Miledi, R.; Chamberlin, A. R. *Bioorg. Med. Chem. Lett.* **2006**, 16, 2189.
- [3] (a) Han, B.; Huang, W.; Ren, W.; He, G.; Wang, J.; Peng, C. *Adv. Synth. Catal.* **2015**, 357, 561.
(b) Wen, J.; Wei, W.; Xue, S.; Yang, D.; Lou, Y.; Gao, C.; Wang, H. *J. Org. Chem.* **2015**, 80, 4966.
(c) Yang, C.; Li, J.; Zhou, R.; Chen, X.; Gao, Y.; He, Z. *Org. Biomol. Chem.* **2015**, 13, 4869.
(d) Liang, L.; Li, E.; Xie, P.; Huang, Y. *Chem. Asian J.* **2014**, 9, 1270.
(e) Huang, X.-F.; Zhang, Y.-F.; Qi, Z.-H.; Li, N.-K.; Geng, Z.-C.; Li, K.; Wang, X.-W. *Org. Biomol. Chem.* **2014**, 12, 4372.
(f) Sannigrahi, M. *Tetrahedron*. **1999**, 55, 9007.
(g) Heathcock, C. H.; Graham, S. L.; Pirrung, M. C.; Plavac, F.; White, C. T. In *The Total Synthesis of Natural Products*, Vol. 5, Ed.: Simon, J., John Wiley and Sons, New York, **1983**, p. 264.
- [4] Karthikeyan, K.; Perumal, P. T. *Synlett* **2009**, 2366.
- [5] For selected reviews on phosphine-catalyzed annulations, see:
(a) Lu, X.; Zhang, C.; Xu, Z. *Acc. Chem. Res.* **2001**, 34, 535.
(b) Ye, L.-W.; Zhou, J.; Tang, Y. *Chem. Soc. Rev.* **2008**, 37, 1140.
(c) Marinetti, A.; Voituriez, A. *Synlett* **2010**, 174.
(d) Zhao, Q.-Y.; Lian, Z.; Wei, Y.; Shi, M. *Chem. Commun.* **2012**, 48, 1724.
(e) Zhou, R.; Liu, R.; Li, R.; He, Z. *Chin. J. Org. Chem.* **2014**, 34, 2385 (in Chinese).
(周荣, 刘蓉芳, 李瑞丰, 贺峥杰, 有机化学, **2014**, 34, 2385.)
(f) Xie, P.; Huang, Y. *Org. Biomol. Chem.* **2015**, 13, 8578.
(g) Zhou, R.; He, Z. *Eur. J. Org. Chem.* **2016**, 1937.
(h) Wang, T.; Han, X.; Zhong, F.; Yao, W.; Lu, Y. *Acc. Chem. Soc.* **2016**, 49, 1369.
- [6] Zhou, R.; Duan, C.; Yang, C.; He, Z. *Chem. Asian J.* **2014**, 9, 1183.
- [7] (a) Zhang, X.-N.; Chen, G.-Q.; Tang, X.-Y.; Wei, Y.; Shi, M. *Angew. Chem. Int. Ed.* **2014**, 53, 10768.
(b) Yang, M.; Wang, T.; Cao, S.; He, Z. *Chem. Commun.* **2014**, 50, 13506.
- [8] The P-ylide **2a** can be readily prepared from Ph_3P and maleimide **2a** in the presence of an acid, see: Hedaya, E.; Theodoropoulos, S. *Tetrahedron* **1968**, 24, 2241.
- [9] Zhou, R.; Wang, J.; Yu, J.; He, Z. *J. Org. Chem.* **2013**, 78, 10596.
- [10] Wynne, J. H.; Price, S. E.; Rorer, J. R.; Stalick, W. M. *Synth. Commun.* **2009**, 33, 341.
- [11] Sortino, M.; Garibotto, F.; Filho, V. C.; Gupta, M.; Enriz, R.; Zaccchino, S. *Bioorg. Med. Chem.*, **2011**, 19, 2823.

(Li, L.; Fan, Y.)