铜催化 1-芳基-1-环醇和 1,3-二羰基化合物的环化反应: 合成环烷烃稠合的二氢呋喃衍生物

杨胜彪^a 李燕^{*,a} 张前^{*,a,b}

(^a 东北师范大学化学学院 长春 130024)

(^b 中国科学院上海有机化学研究所 金属有机化学国家重点实验室 上海 200032)

摘要 报道了一种铜催化的 1-芳基-1-环醇和 1,3-二羰基化合物的环化反应,以中等到优秀的产率合成了一系列环烷烃稠合的二氢呋喃衍生物。该反应条件温和,官能团兼容性好。机理研究表明此环化反应可能经历了自由基中间体。

关键词 环化反应; 环醇; 1,3-二羰基化合物; 二氢呋喃

Copper-Catalyzed Cyclization of 1-Aryl-1-cycloalcohols and 1,3-Dicarbonyl Compounds: Synthesis of Cycloalkane-Fused Dihydrofuran Derivatives

Yang, Shengbiao^a Li, Yan^{*,a} Zhang, Qian^{*,a,b}

(^a Department of Chemistry, Northeast Normal University, Changchun 130024)

(^b State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032)

Abstract A novel method of copper-catalyzed cyclization of 1-aryl-1-cycloalcohols and 1,3-dicarbonyl compounds has been developed, affording various cycloalkane-fused dihydrofurans in moderate to excellent yields. This transformation proceeds under mild conditions and with a good tolerance of functional groups. The mechanistic investigation revealed that the reaction involved radical intermediates.

Keywords cyclization; cycloalcohol; 1,3-dicarbonyl compound; dihydrofuran

1 Introduction

Cycloalkane-fused dihydrofurans are widely found in many natural products as well as therapeutic agents, such as melicodin C,^[1] clerodin,^[2] and hugonianene A,^[3] whose anti-proliferative, human immunodeficiency virus (HIV) protease inhibitors,^[4] or anti-feedant activity have attracted much attention in related biological studies (Figure 1). Thus, various synthetic methods for such compounds have been developed,^[5] mainly including cyclization of 1,3-cyclohexanedione and alkenes^[6] or 1,3-dicarbonyl compounds and cycloalkenes,^[7] resulting the formation of cyclopentane- or cyclohexane-fused hydrofuran derivatives (Schemes 1a, 1b). Nevertheless, these methods usually could not access to cyclobutane-fused dihydrofurans might

due to the strong strain of cyclobutenes. Recently, Yu and co-workers^[8] synthesized cyclobutane-fused dihydrofurans through TfOH-catalyzed cyclopropane ring enlargement/C—C formation/etherification of alkylcyclopropanes and 1,3-dicarbonyl compounds (Scheme 1c). While the substrates alkylcyclopropanes need to be pre-prepared. Therefore, the development of general methods for various cycloalkane-fused dihydrofuran derivatives starting from easily available materials is highly desirable. Herein, we present copper-catalyzed cyclization of 1-aryl-1-cycloalcohols and 1,3-dicarbonyl compounds, providing a series of cycloalkane-fused dihydrofurans in excellent yields starting from readily available materials (Scheme 1d).

* Corresponding authors. E-mail: liy078@nenu.edu.cn, zhangq651@nenu.edu.cn

Received March 21, 2019; revised May 16, 2019; published online May 21, 2019.

Dedicated to the 100th anniversary of the birth of Professor Ruyi Chen.

Project supported by the National Natural Science Foundation of China (No. 21831002) and Ten Thousand Talents Program.

国家自然科学基金(No. 21831002)和万人计划资助项目。

Figure 1 Some examples of cycloalkane-fused dihydrofurans skeleton in natural products and therapeutic agents

Previous work:

(a) Cyclization of 1,3-cyclohexanedione and alkenes

(b) 1,3-Dicarbonyl compounds and cycloalkenes

(c) Yu and co-workers' work

(d) This work:

Scheme 1 Synthesis of cycloalkane-fused dihydrofurans

2 Results and discussion

Initially, 1-phenyl-1-cyclopentanol (**1aa**) and acetylacetone (**2a**) were chosen as the model substrates. After the reaction of **1aa** (0.2 mmol) and **2a** (3 equiv.) was performed in the presence of CuCl (10 mol%) and K₂S₂O₈ (2 equiv.) in CH₃CN (2 mL) at 50 °C under nitrogen atmosphere for 17 h, the desired cyclopentane-fused dihydrofuran **3a** was obtained in 14% yield, along with the formation of cyclopentene **4a** (44%) (Table 1, Entry 1). Other copper salts, such as CuBr, CuI, Cu(CH₃CN)₄PF₆, CuTc, CuCl₂, Cu(OAc)₂ could also promote the reaction (Entries 2~8). Cu(CH₃CN)₄PF₆ gave the best result, affording **3a** in 81% yield and no **4a** being detected (Entry 4). Other metal salts, such as FeCl₃ and ZnCl₂ could not improve the yield of **3a** (Entries 9 and 10). Without the metal salt, **4a** became the main product (58%), along with a small amount of **3a** (10% yield) (Entry 11). Without the oxidant, **3a** could not

be formed (Entry 12). When oxidant K₂S₂O₈ was replaced by PhI(OAc)₂, *tert*-butyl hydroperoxid (TBHP), or 3-chloroperbenzoic acid (*m*-CPBA), only **4a** was formed (Entries 13~15). Upon increasing or decreasing the reaction temperature, the yield of **3a** decreased to 72% and 64%, respectively (Entries 16 and 17). Using tetrahydrofuran (THF), toluene, dichloroethane (DCE), or dichloromethane (DCM) as solvents, no better results were observed (Entries 18~21).

Table 1 Optimization of reaction conditions^a

Entry	Catalyst	Oxidant	Temp./°C	Solvent	Yield ^b /%	
					3a	4a
1	CuCl	K ₂ S ₂ O ₈	50	CH ₃ CN	14	44
2	CuBr	K ₂ S ₂ O ₈	50	CH ₃ CN	44	13
3	CuI	K ₂ S ₂ O ₈	50	CH ₃ CN	23	37
4	Cu(CH ₃ CN) ₄ PF ₆	K ₂ S ₂ O ₈	50	CH ₃ CN	81	0
5	CuTc	K ₂ S ₂ O ₈	50	CH ₃ CN	19	48
7	CuCl ₂	K ₂ S ₂ O ₈	50	CH ₃ CN	30	12
8	Cu(OAc) ₂	K ₂ S ₂ O ₈	50	CH ₃ CN	18	41
9	FeCl ₃	K ₂ S ₂ O ₈	50	CH ₃ CN	18	25
10	ZnCl ₂	K ₂ S ₂ O ₈	50	CH ₃ CN	18	51
11	—	K ₂ S ₂ O ₈	50	CH ₃ CN	10	58
12	Cu(CH ₃ CN) ₄ PF ₆	—	50	CH ₃ CN	0	90
13	Cu(CH ₃ CN) ₄ PF ₆	PhI(OAc) ₂	50	CH ₃ CN	0	86
14	Cu(CH ₃ CN) ₄ PF ₆	TBHP	50	CH ₃ CN	0	57
15	Cu(CH ₃ CN) ₄ PF ₆	<i>m</i> -CPBA	50	CH ₃ CN	0	47
16	Cu(CH ₃ CN) ₄ PF ₆	K ₂ S ₂ O ₈	30	CH ₃ CN	64	11
17	Cu(CH ₃ CN) ₄ PF ₆	K ₂ S ₂ O ₈	70	CH ₃ CN	72	0
18	Cu(CH ₃ CN) ₄ PF ₆	K ₂ S ₂ O ₈	50	THF	3	47
19	Cu(CH ₃ CN) ₄ PF ₆	K ₂ S ₂ O ₈	50	Toluene	0	89
20	Cu(CH ₃ CN) ₄ PF ₆	K ₂ S ₂ O ₈	50	DCE	6	83
21	Cu(CH ₃ CN) ₄ PF ₆	K ₂ S ₂ O ₈	50	DCM	18	24

^a General reaction conditions: **1aa** (0.2 mmol), **2a** (3 equiv., 0.6 mmol), catalyst (10 mol%), oxidant (2 equiv.), solvent (2 mL), 17 h, N₂ atmosphere.

^b Isolated yields.

With these established conditions, the scope of 1-aryl-1-cyclopentanol (**1a**) was examined. As shown in Table 2, various aryls containing both electron-donating and withdrawing groups, such as alkyl, alkoxy, halide, CF₃, and OCF₃, were tolerant, providing the desired cyclopentane-fused dihydrofurans **3a**~**3l** in good to excellent yields. It should be noted that halogen substituents such as Cl (**3h** and **3j**) were well tolerated, which provides the possibility for further transformation. And, the fluorine and trifluoromethyl groups were intact in this novel reaction, which makes the dihydrofurans **3g**, **3i** and **3k** much potential in biological and pharmacological activity study. The naphthyl substituted cyclopentanol (**1am**) was also suitable substrate, affording the corresponding **3m** in excellent yield. Additionally, hetero-aromatic substituent such as thiophenyl (**1an**) was also suitable for this reaction,

Table 2 Variety of 1-aryl-1-cyclopentanols **1a**^a

^a General reaction conditions: **1a** (0.2 mmol), **2a** (3 equiv., 0.6 mmol), catalyst (10 mol%), oxidant (2 equiv.), solvent (2 mL), 17 h, N₂ atmosphere. Isolated yields.

albeit the desired product **3n** was obtained in a relatively low yield (47%). Unfortunately, furanyl substituted substrate **1ao** could not undergo this reaction.

Next, the ring size of 1-aryl-1-cycloalcohols **1** was explored under the optimal conditions. The results are summarized in Table 3. Starting from 1-aryl-1-cyclohexanols **1b**, the oxidative cyclization reaction could proceed smoothly and give the corresponding cyclohexane-fused dihydrofurans **5a**~**5d** in medium to excellent yields (73%~98%). To our delight, various 1-aryl-1-cyclobutanols **1c** were founded to be well suitable substrates, affording the desired cyclobutane-fused dihydrofurans **6a**~**6h** in good to excellent yields (61%~99%). The structure of **6e** was further confirmed by X-ray analysis.^[9] Notably, the medium-sized cycloalcohols could be well-tolerated. Cycloheptanols **1d** could efficiently undergo this reaction, and a series of cycloheptane-fused dihydrofurans **7a**~**7e** were obtained in 64%~96% yields. Moreover, using 1-phenylcyclododecanol (**1e**) as the substrate, the expected product **8** was smoothly formed with an excellent yield (98%). 1-Phenylcyclopropane (**1f**) was also tested, but no desired product was detected. In all reaction, no obvious electronic effect was observed. Additionally, cyclic 1,3-diketones **2b**~**2d** were also examined under standard conditions (Scheme 2a). We were pleased to find that 1,3-cyclopentanedione (**2b**), 1,3-cyclohexanedione (**2c**), dimedone (**2d**) and cyclopentanol (**1ak**) underwent the oxidative cyclization reaction, forming tricycle compounds **9a**~**9c** in good to excellent yields (79%~99%). Furthermore, unsymmetrical 1,3-dicarbonyl compounds, such as 1-phenylbutan-1,3-dione (**2e**) and hexane-2,4-dione (**2f**) could afford the desired products **10a/10a'** and **10b/10b'** in good yields with a relatively low

Table 3 Variety of ring-sizes of alicyclic alcohols **1b**~**1e**^a

^a General reaction conditions: 1-aryl-1-cycloalcohols **1b**~**1e** (0.2 mmol), **2a** (0.6 mmol), Cu(CH₃CN)₄PF₆ (10 mol%), K₂S₂O₈ (0.4 mmol), 50 °C for 17 h, CH₃CN (2 mL), N₂ atmosphere.

(a) Reaction of **1ak** and cyclic 1,3-diketones **2b ~ 2d** (b) Reaction of **1ak** and unsymmetrical 1,3-diketones **2e, 2f** Scheme 2 Reaction of **1ak** and 1,3-diketones **2b ~ 2f**

regioselectivity (Scheme 2b).

In order to probe the reaction mechanism, some control experiments were conducted (Scheme 3). When adding 1.5 equiv. of 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) or 1.5 equiv. of 2,6-di-*tert*-butyl-4-methylphenol (BHT) to the reaction of **1ak** and **2a**, the reaction was completely inhibited. These results suggested that the cyclization reaction might involve radical species. Additionally, starting

Scheme 3 Control experiments

from 1-phenyl-cyclopentene (**4a**), **3a** could be formed in 52% yield under standard conditions. This indicated that the cycloalkene might be intermediate.

Based on the above experiments and previous works,^[10] a proposed mechanism is shown in Scheme 4. Initially, 1-phenyl-1-cyclopentanol (**1aa**) underwent dehydration to form 1-phenyl-1-cyclopentene (**4a**). Meanwhile, the depro-

Scheme 4 Plausible mechanism

tonation of 1,3-dicarbonyl compound **2a** formed radical **A** in the presence of $K_2S_2O_8$, which added to **4a** to form benzylic radical **B**. **B** was oxidized to give benzylic cation **C**, which could tautomerize to enol benzylic cation **D**. Finally, **D** underwent the intramolecular nucleophilic attack and the loss of proton to produce **3a**. For unsymmetrical 1,3-dicarbonyl compounds **2e** and **2f**, the low regioselectivity of oxa-cyclization might result from the low selectivity of ketone-enol tautomerization.

3 Conclusions

In summary, a novel copper-catalyzed cyclization reaction of 1,3-dicarbonyl compounds and 1-aryl-1-cycloalcohols, facially constructing a wide range of cycloalkane-fused dihydrofurans derivatives in good to excellent yields under mild conditions was developed. This reaction would provide a new way toward cycloalkane-fused dihydrofurans.

4 Experimental section

4.1 General information

All commercially available chemicals were used without further purification. 1H NMR spectra were recorded at 25 °C on a Varian 600 or 500 MHz, ^{13}C NMR spectra were recorded at 25 °C on a Varian 150 or 125 MHz spectrometer with TMS as internal standard. High resolution mass spectra were recorded on a Bruker microtof spectrometer. Flash column chromatography was performed on silica gel 60 (particle size 300~400 mesh ASTM, purchased from Taizhou, China).

4.2 Preparation of 1-substituted cycloalcohols

All glassware were flame-dried before use. Cyclopentanone (13.4 mmol) was dissolved in 25 mL of diethyl ether and purged with N_2 . The temperature was reduced to 0 °C. The appropriate Grignard reagent (14.7 mmol) was added dropwise with stirring. The reaction was allowed to stir for an additional 3 h. Water was added slowly to quench the reaction. The organic layer was removed, and the aqueous layer was extracted three times with diethyl ether (25 mL $\times$ 3). The organic layers were combined, dried with Na_2SO_4 , filtered and the solvent was evaporated under vacuum. The residue was purified by column chromatography (petroleum ether/ethyl acetate, $V:V=25:1$) to give the corresponding product 1-phenyl-1-cyclopentanol (**1aa**).

4.3 General procedure for cycloalkane-fused dihydrofurans (3 and 5~10)

$Cu(CH_3CN)_4PF_6$ (0.02 mmol, 7.5 mg), $K_2S_2O_8$ (0.4 mmol, 108 mg), **1aa** (0.2 mmol) and CH_3CN (2 mL) were added into a screw-cap test tube under N_2 atmosphere in glove-box. The test tube was then stirred at room temperature for 5 min. **2a** (60 μ L, 0.6 mmol) was added into reaction mixture and the test tube was then sealed off with a screw-cap and the reaction mixture was stirred at 50 °C for 17 h. After the reaction finished, the solvent was evap-

orated under vacuum. The residue was purified by column chromatography (petroleum ether/ethyl ether, $V:V=20:1$) to give the corresponding product **3a**.

1-(2-Methyl-6a-phenyl-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)ethanone (**3a**): light yellow oil, 81% yield (39.3 mg). 1H NMR (600 MHz, $CDCl_3$) δ : 7.38~7.33 (m, 4H), 7.28~7.24 (m, 1H), 3.62 (d, $J=8.4$ Hz, 1H), 2.32 (s, 3H), 2.32~2.27 (m, 1H), 2.19 (s, 3H), 2.13~1.98 (m, 2H), 1.95~1.82 (m, 2H), 1.75~1.60 (m, 1H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 194.3, 167.5, 144.2, 128.3, 127.3, 124.2, 116.1, 99.3, 55.3, 42.9, 34.9, 29.1, 24.7, 15.1; HRMS (ESI-TOF) calcd for $C_{16}H_{18}O_2Na$ $[M+Na]^+$ 265.1199, found 265.1208.

1-(2-Methyl-6a-(*o*-tolyl)-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)ethanone (**3b**): light yellow oil, 83% yield (42.6 mg). 1H NMR (600 MHz, $CDCl_3$) δ : 7.49 (q, $J=3.0$ Hz, 1H), 7.19~7.18 (m, 3H), 3.86 (d, $J=8.4$ Hz, 1H), 2.46 (s, 3H), 2.31 (s, 3H), 2.25~2.11 (m, 3H), 2.21 (s, 3H), 1.87 (q, $J=6.0$ Hz, 2H), 1.77~1.68 (m, 1H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 194.2, 166.8, 141.0, 134.7, 132.5, 127.4, 125.8, 125.0, 116.9, 100.5, 53.1, 41.5, 34.3, 29.2, 24.8, 21.3, 15.2; HRMS (ESI-TOF) calcd for $C_{17}H_{20}O_2Na$ $[M+Na]^+$ 279.1356, found 279.1358.

1-(2-Methyl-6a-(*m*-tolyl)-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)ethanone (**3c**): yellow oil, 87% yield (44.6 mg). 1H NMR (600 MHz, $CDCl_3$) δ : 7.27~7.22 (m, 1H), 7.22~7.13 (m, 2H), 7.08 (d, $J=7.8$ Hz, 1H), 3.62 (d, $J=8.4$ Hz, 1H), 2.36 (s, 3H), 2.32 (s, 3H), 2.29 (dd, $J=13.8, 6.0$ Hz, 1H), 2.20 (s, 3H), 2.15~1.97 (m, 2H), 1.95~1.79 (m, 2H), 1.74~1.67 (m, 1H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 194.4, 167.6, 144.2, 138.0, 128.3, 128.1, 124.9, 121.3, 116.1, 99.4, 55.2, 43.0, 34.9, 29.1, 24.7, 21.5, 15.1; HRMS (ESI-TOF) calcd for $C_{17}H_{21}O_2$ $[M+H]^+$ 257.1536, found 257.1536.

1-(2-Methyl-6a-(*p*-tolyl)-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)ethanone (**3d**): light yellow oil, 83% yield (42.6 mg). 1H NMR (600 MHz, $CDCl_3$) δ : 7.26 (d, $J=7.8$ Hz, 2H), 7.16 (d, $J=7.8$ Hz, 2H), 3.61 (d, $J=7.8$ Hz, 1H), 2.34 (s, 3H), 2.32 (s, 3H), 2.31~2.27 (m, 1H), 2.20 (s, 3H), 2.11~1.97 (m, 2H), 1.89~1.30 (m, 2H), 1.74~1.65 (m, 1H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 194.8, 168.1, 141.1, 137.2, 129.1, 124.3, 116.1, 99.5, 55.1, 42.7, 34.9, 29.0, 24.7, 21.0, 15.2; HRMS (ESI-TOF) calcd for $C_{17}H_{20}O_2Na$ $[M+Na]^+$ 279.1356, found 279.1357.

1-(6a-(4-Methoxyphenyl)-2-methyl-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)ethanone (**3e**): light yellow oil, 96% yield (52.3 mg). 1H NMR (600 MHz, $CDCl_3$) δ : 7.30 (d, $J=8.4$ Hz, 2H), 6.88 (d, $J=9.0$ Hz, 2H), 3.80 (s, 3H), 3.60 (d, $J=8.4$ Hz, 1H), 2.35~2.25 (m, 4H), 2.20 (s, 3H), 2.07~1.94 (m, 2H), 1.90~1.76 (m, 2H), 1.75~1.61 (m, 1H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 194.5, 167.6, 158.9, 136.2, 125.7, 116.2, 113.7, 99.3, 55.3, 54.9, 42.4, 34.9, 29.2, 24.6, 15.2; HRMS (ESI-TOF) calcd for $C_{17}H_{21}O_3$ $[M+H]^+$ 273.1485, found 273.1485.

1-(2-Methyl-6a-(4-(trifluoromethoxy)phenyl)-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)ethanone (**3f**): colorless oil, yield 74% (48.3 mg). 1H NMR (600 MHz,

CDCl_3) δ : 7.43~7.36 (m, 2H), 7.20 (d, $J=8.4$ Hz, 2H), 3.60 (d, $J=8.4$ Hz, 1H), 2.37~2.26 (m, 1H), 2.32 (s, 3H), 2.21 (s, 3H), 2.10~2.07 (m, 1H), 2.04~1.96 (m, 1H), 1.91~1.85 (m, 2H), 1.76~1.68 (m, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.3, 167.2, 148.4, 143.0, 125.8, 120.9, 120.4 (q, $J=255.3$ Hz) 116.2, 98.7, 55.5, 43.1, 34.9, 29.2, 24.7, 15.1; HRMS (ESI-TOF) calcd for $\text{C}_{17}\text{H}_{18}\text{F}_3\text{O}_3$ $[\text{M}+\text{H}]^+$ 327.1203, found 327.1205.

1-(6a-(4-Fluorophenyl)-2-methyl-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)ethanone (**3g**): colorless oil, yield 92% (47.9 mg). ^1H NMR (600 MHz, CDCl_3) δ : 7.39~7.29 (m, 2H), 7.09~6.98 (m, 2H), 3.59 (d, $J=8.4$ Hz, 1H), 2.32 (s, 3H), 2.31~2.26 (m, 1H), 2.21 (s, 3H), 2.13~1.94 (m, 2H), 1.93~1.81 (m, 2H), 1.76~1.63 (m, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.3, 167.3, 162.0 (d, $J=7.8$, 244.5 Hz), 140.0 (d, $J=3.2$ Hz), 126.1 (d, $J=8.1$ Hz), 116.2, 115.1 (d, $J=21.2$ Hz), 98.9, 55.3, 42.9, 34.9, 29.2, 24.7, 15.1; HRMS (ESI-TOF) calcd for $\text{C}_{16}\text{H}_{17}\text{FO}_2\text{Na}$ $[\text{M}+\text{Na}]^+$ 283.1105, found 283.1104.

1-(6a-(4-Chlorophenyl)-2-methyl-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)ethanone (**3h**): colorless oil, yield 93% (51.5 mg). ^1H NMR (600 MHz, CDCl_3) δ : 7.34~7.28 (m, 4H), 3.58 (d, $J=8.4$ Hz, 1H), 2.31 (s, 3H), 2.31~2.27 (m, 1H), 2.20 (s, 3H), 2.09~2.02 (m, 1H), 2.01~1.92 (m, 1H), 1.93~1.82 (m, 2H), 1.76~1.60 (m, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.3, 167.3, 142.8, 133.1, 128.5, 125.8, 116.1, 98.7, 55.4, 43.0, 34.9, 29.2, 24.7, 15.1; HRMS (ESI-TOF) calcd for $\text{C}_{16}\text{H}_{18}\text{ClO}_2$ $[\text{M}+\text{H}]^+$ 277.0990, found 277.0997.

1-(6a-(3-Fluorophenyl)-2-methyl-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)ethanone (**3i**): light yellow oil, 97% yield (50.5 mg). ^1H NMR (600 MHz, CDCl_3) δ : 7.32 (td, $J=7.8$, 6.0 Hz, 1H), 7.13 (d, $J=7.8$ Hz, 1H), 7.08 (dt, $J=10.2$, 2.4 Hz, 1H), 6.96 (td, $J=7.8$, 2.4 Hz, 1H), 3.60 (d, $J=8.4$ Hz, 1H), 2.33 (s, 3H), 2.29 (m, 1H), 2.20 (s, 3H), 2.12~2.03 (m, 1H), 2.03~1.95 (m, 1H), 1.93~1.83 (m, 2H), 1.76~1.66 (m, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.3, 167.3, 162.8 (d, $J=244.2$ Hz), 147.1 (d, $J=244.2$ Hz), 130.0 (d, $J=8.2$ Hz), 119.8 (d, $J=2.7$ Hz), 116.1, 114.0, 111.6 (d, $J=22.7$ Hz), 98.7, 55.6, 43.1, 34.9, 29.2, 24.7, 15.1; HRMS (ESI-TOF) calcd for $\text{C}_{16}\text{H}_{18}\text{FO}_2$ $[\text{M}+\text{H}]^+$ 261.1285, found 261.1283.

1-(6a-(3-Chlorophenyl)-2-methyl-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)ethanone (**3j**): colorless oil, 92% yield (50.9 mg). ^1H NMR (600 MHz, CDCl_3) δ : 7.36 (t, $J=1.8$ Hz, 1H), 7.32~7.26 (m, 1H), 7.26~7.19 (m, 2H), 3.60 (d, $J=8.4$ Hz, 1H), 2.33 (s, 3H), 2.33~2.27 (m, 1H), 2.21 (s, 3H), 2.12~2.03 (m, 1H), 2.02~1.94 (m, 1H), 1.93~1.80 (m, 2H), 1.80~1.62 (m, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.2, 167.2, 146.5, 134.4, 129.7, 127.4, 124.6, 122.5, 116.1, 98.6, 55.5, 43.2, 34.9, 29.2, 24.7, 15.1; HRMS (ESI-TOF) calcd for $\text{C}_{17}\text{H}_{18}\text{ClO}_2$ $[\text{M}+\text{H}]^+$ 277.0990, found 277.0995.

1-(2-Methyl-6a-(4-(trifluoromethyl)phenyl)-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)ethanone (**3k**): brown oil, 96% yield (59.6 mg). ^1H NMR (600 MHz, CDCl_3) δ : 7.61 (d, $J=7.8$ Hz, 2H), 7.49 (d, $J=8.4$ Hz,

2H), 3.62 (d, $J=8.4$ Hz, 1H), 2.36~2.28 (m, 4H), 2.21 (s, 3H), 2.18~2.05 (m, 1H), 2.05~1.97 (m, 1H), 1.96~1.87 (m, 2H), 1.78~1.69 (m, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.4, 167.4, 148.4, 129.6 (q, $J=32.3$ Hz), 125.4 (q, $J=3.8$ Hz), 124.7, 124.0 (q, $J=270.3$ Hz), 116.1, 98.8, 55.7, 43.4, 35.0, 29.1, 24.8, 15.1; HRMS (ESI-TOF) calcd for $\text{C}_{17}\text{H}_{17}\text{F}_3\text{O}_2\text{Na}$ $[\text{M}+\text{Na}]^+$ 333.1073, found 333.1077.

1-(6a-(3,5-Dimethylphenyl)-2-methyl-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)ethanone (**3l**): colorless oil, 81% yield (43.8 mg). ^1H NMR (600 MHz, CDCl_3) δ : 6.98 (s, 2H), 6.91 (s, 1H), 3.62 (d, $J=7.8$ Hz, 1H), 2.34~2.25 (m, 10H), 2.20 (s, 3H), 2.10~1.95 (m, 2H), 1.91~1.81 (m, 2H), 1.73~1.64 (m, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.5, 167.6, 144.2, 138.0, 129.0, 122.1, 116.1, 99.4, 55.1, 43.0, 35.0, 29.2, 24.7, 21.4, 15.2; HRMS (ESI-TOF) calcd for $\text{C}_{18}\text{H}_{23}\text{O}_2$ $[\text{M}+\text{H}]^+$ 271.1693, found 271.1691.

1-(2-Methyl-6a-(naphthalen-1-yl)-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)ethanone (**3m**): yellow oil, 98% yield (57.3 mg). ^1H NMR (600 MHz, CDCl_3) δ : 8.09 (d, $J=8.4$ Hz, 1H), 7.88 (d, $J=7.2$ Hz, 1H), 7.78 (d, $J=7.8$ Hz, 1H), 7.74 (d, $J=7.2$ Hz, 1H), 7.52~7.46 (m, 2H), 7.43 (t, $J=7.8$ Hz, 1H), 4.12 (d, $J=8.4$ Hz, 1H), 2.53~2.45 (m, 2H), 2.37 (s, 3H), 2.35~2.32 (m, 1H), 2.01~1.97 (m, 2H), 1.91~1.83 (m, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.3, 166.5, 138.4, 134.7, 129.7, 129.4, 128.7, 125.7, 125.2, 125.0, 124.7, 122.5, 117.1, 100.6, 53.3, 42.4, 34.5, 29.2, 25.0, 15.2; HRMS (ESI-TOF) calcd for $\text{C}_{20}\text{H}_{20}\text{O}_2\text{Na}$ $[\text{M}+\text{Na}]^+$ 315.1356, found 315.1352.

1-(2-Methyl-6a-(thiophen-2-yl)-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)ethanone (**3n**): yellow oil, 47% yield (23.3 mg). ^1H NMR (600 MHz, CDCl_3) δ : 7.24 (dd, $J=4.8$, 1.2 Hz, 1H), 7.02~6.95 (m, 2H), 3.68 (d, $J=9.0$ Hz, 1H), 2.43~2.39 (m, 1H), 2.28 (s, 3H), 2.23 (s, 3H), 2.16~2.01 (m, 2H), 1.88~1.79 (m, 2H), 1.74~1.66 (m, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.4, 167.1, 147.8, 126.9, 124.6, 122.7, 116.0, 97.0, 55.5, 42.4, 34.4, 29.3, 24.5, 15.2; HRMS (ESI-TOF) calcd for $\text{C}_{14}\text{H}_{17}\text{O}_2\text{S}$ $[\text{M}+\text{H}]^+$ 249.0944, found 249.0949.

1-(2-Methyl-7a-phenyl-3a,4,5,6,7,7a-hexahydrobenzofuran-3-yl)ethanone (**5a**): light yellow oil, 87% yield (44.6 mg). ^1H NMR (500 MHz, CDCl_3) δ : 7.37~7.30 (m, 4H), 7.25 (t, $J=7.0$ Hz, 1H), 3.31 (t, $J=6.5$ Hz, 1H), 2.33 (s, 3H), 2.16 (s, 3H), 2.15~2.01 (m, 2H), 1.90~1.77 (m, 1H), 1.77~1.52 (m, 4H), 1.53~1.34 (m, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ : 194.3, 167.0, 147.1, 128.2, 127.1, 124.2, 119.2, 90.3, 47.0, 34.7, 29.2, 27.6, 19.7, 19.5, 15.6; HRMS (ESI-TOF) calcd for $\text{C}_{17}\text{H}_{20}\text{O}_2\text{Na}$ $[\text{M}+\text{Na}]^+$ 279.1356, found 279.1356.

1-(2-Methyl-7a-(*p*-tolyl)-3a,4,5,6,7,7a-hexahydrobenzofuran-3-yl)ethanone (**5b**): yellow oil, 98% yield (53.0 mg). ^1H NMR (600 MHz, CDCl_3) δ : 7.25 (d, $J=7.8$ Hz, 2H), 7.13 (d, $J=8.4$ Hz, 2H), 3.29 (t, $J=6.6$ Hz, 1H), 2.32 (s, 3H), 2.32 (s, 3H), 2.16 (s, 3H), 2.14~2.00 (m, 2H), 1.84~1.75 (m, 1H), 1.67~1.50 (m, 3H), 1.45~1.36 (m, 2H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.3, 166.9, 144.1, 136.7,

128.8, 124.1, 119.2, 90.3, 46.9, 34.7, 29.1, 27.6, 20.9, 19.7, 19.5, 15.5; HRMS (ESI-TOF) calcd for $C_{18}H_{22}O_2Na$ $[M+Na]^+$ 293.1512, found 293.1522.

1-(7a-(4-Fluorophenyl)-2-methyl-3a,4,5,6,7,7a-hexahydrobenzofuran-3-yl)ethanone (**5c**): colorless oil, 93% yield (51.0 mg). 1H NMR (600 MHz, $CDCl_3$) δ : 7.36~7.29 (m, 2H), 7.04~6.96 (m, 2H), 3.27 (t, $J=6.6$ Hz, 1H), 2.32 (s, 3H), 2.17 (s, 3H), 2.15~2.01 (m, 2H), 1.82~1.75 (m, 1H), 1.68~1.51 (m, 3H), 1.50~1.44 (m, 1H), 1.43~1.36 (m, 1H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 194.2, 166.7, 161.3 (d, $J=244.1$ Hz), 142.8 (d, $J=3.0$ Hz), 125.9 (d, $J=6.8$ Hz), 119.3, 114.9 (d, $J=21.2$ Hz), 89.9, 47.0, 34.8, 32.9, 32.8, 29.2, 27.6, 19.6, 19.4, 15.5; HRMS (ESI-TOF) calcd for $C_{17}H_{19}FO_2Na$ $[M+Na]^+$ 297.1261, found 297.1265.

1-(2-Nethyl-7a-(4-(trifluoromethyl)phenyl)-3a,4,5,6,7,7a-hexahydrobenzofuran-3-yl)ethanone (**5d**): colorless oil, 73% yield (47.4 mg). 1H NMR (600 MHz, $CDCl_3$) δ : 7.58 (d, $J=8.4$ Hz, 2H), 7.48 (d, $J=7.8$ Hz, 2H), 3.28 (t, $J=6.6$ Hz, 1H), 2.34 (s, 3H), 2.19~2.10 (m, 4H), 2.10~2.05 (m, 1H), 1.83~1.75 (m, 1H), 1.69~1.62 (m, 2H), 1.60~1.52 (m, 1H), 1.50~1.38 (m, 2H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 194.1, 166.6, 151.2, 129.4 (q, $J=2.4$ Hz), 126.8, 125.2 (q, $J=3.8$ Hz), 123.2 (d, $J=270.2$ Hz, 2H), 119.5, 89.8, 46.9, 34.7, 29.2, 27.7, 19.6, 19.5, 15.5; HRMS (ESI-TOF) calcd for $C_{18}H_{19}O_2Na$ $[M+Na]^+$ 347.1229, found 347.1223.

1-(3-Methyl-1-phenyl-2-oxabicyclo[3.2.0]hept-3-en-4-yl)ethanone (**6a**): colorless oil, 75% yield (34.2 mg). 1H NMR (500 MHz, $CDCl_3$) δ : 7.47 (d, $J=7.5$ Hz, 2H), 7.42 (t, $J=7.5$ Hz, 2H), 7.33 (t, $J=7.5$ Hz, 1H), 3.83 (t, $J=3.5$, 1H), 2.88~2.67 (m, 2H), 2.57~2.41 (m, 1H), 2.36 (s, 3H), 2.17 (s, 3H), 2.13~1.99 (m, 1H); ^{13}C NMR (125 MHz, $CDCl_3$) δ : 194.9, 169.1, 140.8, 128.6, 127.9, 124.7, 117.5, 90.9, 49.3, 35.3, 29.5, 24.6, 15.3; HRMS (ESI-TOF) calcd for $C_{15}H_{17}O_2$ $[M+H]^+$ 229.1223, found 229.1232.

1-(3-Methyl-1-(*o*-tolyl)-2-oxabicyclo[3.2.0]hept-3-en-4-yl)ethanone (**6b**): colorless oil, 97% yield (47.0 mg). 1H NMR (600 MHz, $CDCl_3$) δ : 7.40 (d, $J=7.2$ Hz, 1H), 7.30~7.20 (m, 3H), 4.17 (d, $J=7.2$ Hz, 1H), 2.92~2.80 (m, 1H), 2.77~2.67 (m, 1H), 2.40~2.30 (m, 4H), 2.28 (s, 3H), 2.23 (s, 3H), 2.06~1.97 (m, 1H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 194.8, 168.7, 137.7, 136.4, 131.8, 128.7, 125.6, 125.6, 117.5, 92.4, 46.2, 35.4, 29.4, 24.2, 20.1, 15.3; HRMS (ESI-TOF) calcd for $C_{16}H_{18}O_2Na$ $[M+Na]^+$ 265.1199, found 265.1197.

1-(3-Methyl-1-(*m*-tolyl)-2-oxabicyclo[3.2.0]hept-3-en-4-yl)ethanone (**6c**): yellow oil, 63% yield (30.5 mg). 1H NMR (600 MHz, $CDCl_3$) δ : 7.30 (t, $J=7.2$ Hz, 1H), 7.28~7.24 (m, 2H), 7.15 (d, $J=7.2$ Hz, 1H), 3.82 (d, $J=7.2$ Hz, 1H), 2.84~2.68 (m, 2H), 2.53~2.41 (m, 1H), 2.39 (s, 3H), 2.35 (s, 3H), 2.17 (s, 3H), 2.11~2.02 (m, 1H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 195.0, 169.1, 140.8, 138.4, 128.8, 128.6, 125.5, 121.8, 117.5, 91.0, 49.3, 35.3, 29.5, 24.7, 21.5, 15.4; HRMS (ESI-TOF) calcd for $C_{16}H_{18}O_2Na$ $[M+Na]^+$ 265.1199, found 265.1198.

1-(3-Methyl-1-(*p*-tolyl)-2-oxabicyclo[3.2.0]hept-3-en-4-

yl)ethanone (**6d**): yellow oil, 61% yield (29.6 mg). 1H NMR (600 MHz, $CDCl_3$) δ : 7.35 (d, $J=7.8$ Hz, 2H), 7.22 (d, $J=7.8$ Hz, 2H), 3.80 (d, $J=7.2$ Hz, 1H), 2.84~2.67 (m, 2H), 2.51~2.39 (m, 1H), 2.37 (s, 3H), 2.34 (s, 3H), 2.16 (s, 3H), 2.10~2.02 (m, 1H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 195.0, 169.1, 137.8, 137.8, 129.2, 124.8, 117.5, 90.9, 49.2, 35.2, 29.4, 24.5, 21.1, 15.3; HRMS (ESI-TOF) calcd for $C_{16}H_{18}O_2Na$ $[M+Na]^+$ 265.1199, found 265.1201.

1-(1-(2,4-Dimethylphenyl)-3-methyl-2-oxabicyclo[3.2.0]hept-3-en-4-yl)ethanone (**6e**): white solid, 62% yield (31.8 mg). m.p. 95~96 °C; 1H NMR (600 MHz, $CDCl_3$) δ : 7.29 (d, $J=7.8$ Hz, 1H), 7.07 (s, 1H), 7.03 (d, $J=7.8$ Hz, 1H), 4.13 (t, $J=1.8$ Hz, 1H), 2.92~2.78 (m, 1H), 2.75~2.67 (m, 1H), 2.40~2.29 (m, 7H), 2.27 (s, 3H), 2.22 (s, 3H), 2.04~1.96 (m, 1H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 194.9, 168.9, 138.6, 137.6, 133.6, 132.7, 126.1, 125.7, 117.5, 92.3, 46.3, 35.4, 29.4, 24.3, 20.9, 20.0, 15.4; HRMS (ESI-TOF) calcd for $C_{17}H_{20}O_2Na$ $[M+Na]^+$ 279.1356, found 279.1357.

1-(1-(4-Fluorophenyl)-3-methyl-2-oxabicyclo[3.2.0]hept-3-en-4-yl)ethanone (**6f**): light yellow oil, 96% yield (47.3 mg). 1H NMR (600 MHz, $CDCl_3$) δ : 7.45~7.41 (m, 2H), 7.09 (t, $J=8.4$ Hz, 2H), 3.80 (d, $J=7.8$ Hz, 1H), 2.83~2.77 (m, 1H), 2.75~2.65 (m, 1H), 2.52~2.41 (m, 1H), 2.35 (s, 3H), 2.17 (s, 3H), 2.11~2.03 (m, 1H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 194.8, 168.9, 162.4 (d, $J=244.5$ Hz), 136.8, 126.7 (d, $J=9$ Hz), 117.5, 115.5 (d, $J=21.0$ Hz), 90.4, 49.4, 35.3, 29.5, 24.5, 15.3; HRMS (ESI-TOF) calcd for $C_{15}H_{16}FO_2$ $[M+H]^+$ 247.1129, found 247.1128.

1-(1-(4-Chlorophenyl)-3-methyl-2-oxabicyclo[3.2.0]hept-3-en-4-yl)ethanone (**6g**): yellow oil, yield 98% (51.5 mg). 1H NMR (600 MHz, $CDCl_3$) δ : 7.40~7.36 (m, 4H), 3.78 (d, $J=7.8$ Hz, 1H), 2.82~2.77 (m, 1H), 2.70~2.66 (m, 1H), 2.46~2.43 (m, 1H), 2.35 (s, 3H), 2.17 (s, 3H), 2.10~2.05 (m, 1H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 194.9, 169.0, 139.5, 133.8, 128.7, 126.2, 117.6, 90.3, 49.5, 35.4, 29.4, 24.6, 15.3; HRMS (ESI-TOF) calcd for $C_{15}H_{15}ClO_2Na$ $[M+Na]^+$ 285.0653, found 285.0653.

1-(3-Methyl-1-(naphthalen-1-yl)-2-oxabicyclo[3.2.0]hept-3-en-4-yl)ethanone (**6h**): yellow solid, 99% yield (55.1 mg). m.p. 45~46 °C; 1H NMR (600 MHz, $CDCl_3$) δ : 7.99~7.97 (m, 1H), 7.91~7.89 (m, 1H), 7.87 (d, $J=8.4$ Hz, 1H), 7.60 (d, $J=7.2$ Hz, 1H), 7.54~7.47 (m, 3H), 4.36 (d, $J=6.6$ Hz, 1H), 3.12~3.07 (m, 1H), 2.85~2.80 (m, 1H), 2.40~2.34 (m, 1H), 2.32 (s, 1H), 2.28 (s, 3H), 2.13~2.08 (m, 1H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 195.0, 168.9, 134.5, 134.3, 130.8, 129.7, 128.9, 126.2, 125.9, 125.6, 124.7, 124.0, 117.5, 92.2, 46.6, 36.4, 29.5, 24.7, 15.5; HRMS (ESI-TOF) calcd for $C_{19}H_{18}O_2Na$ $[M+Na]^+$ 301.1199, found 301.1199.

1-(2-Methyl-8a-phenyl-4,5,6,7,8,8a-hexahydro-3aH-cyclohepta[b]furan-3-yl)ethanone (**7a**): colorless oil, 96% yield (51.9 mg). 1H NMR (600 MHz, $CDCl_3$) δ : 7.39 (d, $J=7.2$ Hz, 2H), 7.33 (t, $J=7.8$ Hz, 2H), 7.27~7.21 (m, 1H), 3.59 (d, $J=5.4$ Hz, 1H), 2.34 (s, 3H), 2.19 (s, 3H), 2.14~1.97 (m, 3H), 1.88~1.79 (m, 1H), 1.65~1.58 (m,

2H), 1.56~1.41 (m, 4H); ^{13}C NMR (150 MHz, CDCl_3) δ : 193.9, 165.6, 148.3, 128.2, 127.0, 123.8, 116.7, 94.0, 53.3, 40.3, 31.0, 29.8, 29.3, 26.3, 23.9, 15.5; HRMS (ESI-TOF) calcd for $\text{C}_{18}\text{H}_{22}\text{O}_2\text{Na}$ $[\text{M} + \text{Na}]^+$ 293.1512, found 293.1506.

1-(2-Methyl-8a-(*o*-tolyl)-4,5,6,7,8,8a-hexahydro-3aH-cyclohepta[*b*]furan-3-yl)ethanone (**7b**): light yellow oil, 83% yield (47.2 mg). ^1H NMR (600 MHz, CDCl_3) δ : 7.43~7.34 (m, 1H), 7.18~7.07 (m, 3H), 3.64 (s, 1H), 2.45 (s, 3H), 2.40 (s, 3H), 2.30~2.21 (m, 2H), 2.21 (s, 3H), 2.02~1.95 (m, 1H), 1.88~1.73 (m, 2H), 1.69~1.57 (m, 3H), 1.36~1.17 (m, 2H); ^{13}C NMR (150 MHz, CDCl_3) δ : 193.7, 165.1, 146.2, 133.6, 132.6, 126.9, 125.7, 123.5, 116.4, 94.8, 52.0, 37.7, 31.3, 29.7, 29.4, 24.8, 24.5, 21.4, 15.3; HRMS (ESI-TOF) calcd for $\text{C}_{19}\text{H}_{25}\text{O}_2$ $[\text{M} + \text{H}]^+$ 285.1849, found 285.1851.

1-(2-Methyl-8a-(*p*-tolyl)-4,5,6,7,8,8a-hexahydro-3aH-cyclohepta[*b*]furan-3-yl)ethanone (**7c**): colorless oil, 71% yield (40.4 mg). ^1H NMR (600 MHz, CDCl_3) δ : 7.29 (d, $J=8.4$ Hz, 2H), 7.14 (d, $J=8.4$ Hz, 2H), 3.57 (d, $J=5.4$ Hz, 1H), 2.33 (s, 3H), 2.32 (s, 3H), 2.19 (s, 3H), 2.13~1.91 (m, 3H), 1.91~1.78 (m, 1H), 1.64~1.56 (m, 2H), 1.55~1.43 (m, 4H); ^{13}C NMR (150 MHz, CDCl_3) δ : 193.9, 165.7, 145.2, 136.7, 128.9, 123.9, 116.8, 94.1, 53.3, 40.3, 31.0, 29.8, 29.3, 26.5, 23.8, 20.9, 15.5; HRMS (ESI-TOF) calcd for $\text{C}_{19}\text{H}_{25}\text{O}_2$ $[\text{M} + \text{H}]^+$ 285.1849, found 285.1850.

1-(8a-(4-Fluorophenyl)-2-methyl-4,5,6,7,8,8a-hexahydro-3aH-cyclohepta[*b*]furan-3-yl)ethanone (**7d**): colorless oil, 64% yield (36.9 mg). ^1H NMR (600 MHz, CDCl_3) δ : 7.36 (dd, $J=8.4$, 5.4 Hz, 2H), 7.01 (t, $J=8.4$ Hz, 2H), 3.54 (d, $J=5.4$ Hz, 1H), 2.33 (s, 3H), 2.21 (s, 3H), 2.09~1.98 (m, 3H), 1.88~1.80 (m, 1H), 1.68~1.54 (m, 2H), 1.55~1.41 (m, 4H); ^{13}C NMR (150 MHz, CDCl_3) δ : 193.8, 165.4, 161.7 (d, $J=244.1$ Hz), 144.0, 125.6 (d, $J=7.8$ Hz), 116.8, 115.0 (d, $J=21.2$ Hz), 93.7, 53.4, 40.4, 31.0, 29.7, 29.3, 26.4, 23.8, 15.5; HRMS (ESI-TOF) calcd for $\text{C}_{18}\text{H}_{22}\text{FO}_2$ $[\text{M} + \text{H}]^+$ 289.1598, found 289.1597.

1-(8a-(4-methoxyphenyl)-2-methyl-4,5,6,7,8,8a-hexahydro-3aH-cyclohepta[*b*]furan-3-yl)ethanone (**7e**): yellow oil, yield 93% (55.9 mg). ^1H NMR (600 MHz, CDCl_3) δ : 7.33 (d, $J=8.4$ Hz, 2H), 6.86 (d, $J=8.8$ Hz, 2H), 3.79 (s, 3H), 3.56 (d, $J=8.4$ Hz, 1H), 2.31 (s, 3H), 2.20 (s, 3H), 2.10~1.90 (m, 3H), 1.86~1.82 (m, 1H), 1.63~1.44 (m, 6H); ^{13}C NMR (150 MHz, CDCl_3) δ : 193.9, 165.6, 158.6, 140.1, 125.3, 116.9, 113.5, 94.0, 55.2, 53.2, 40.3, 31.0, 29.8, 29.3, 26.7, 23.8, 15.6; HRMS (ESI-TOF) calcd for $\text{C}_{19}\text{H}_{24}\text{O}_3\text{Na}$ $[\text{M} + \text{Na}]^+$ 323.1618, found 323.1626.

1-(2-Methyl-13a-phenyl-3a,4,5,6,7,8,9,10,11,12,13,13a-dodecahydrocyclohepta[*b*]furan-3-yl)ethanone (**8**): light yellow oil, 98% yield (66.7 mg). ^1H NMR (600 MHz, CDCl_3) δ : 7.35~7.28 (m, 4H), 7.25~7.21 (m, 1H), 3.27 (d, $J=7.8$ Hz, 1H), 2.28 (s, 3H), 2.14 (s, 3H), 2.11~2.01 (m, 1H), 2.01~1.91 (m, 1H), 1.91~1.81 (m, 1H), 1.68~1.35 (m, 14H), 1.22~1.14 (m, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.0, 165.9, 145.8, 127.8, 127.0, 124.8, 119.2, 93.5, 51.1, 35.6, 29.1, 28.4, 27.1, 26.9, 26.6, 25.5, 24.8,

23.6, 23.5, 22.2, 15.5; HRMS (ESI-TOF) calcd for $\text{C}_{23}\text{H}_{33}\text{O}_2$ $[\text{M} + \text{H}]^+$ 341.2475, found 341.2476.

4a-(4-(Trifluoromethyl)phenyl)-2,3,5,6,7,7a-hexahydro-dicyclopenta[*b,d*]furan-1(4aH)-one (**9a**): black oil, yield 99% (61.0 mg). ^1H NMR (600 MHz, CDCl_3) δ : 7.64 (d, $J=7.8$ Hz, 2H), 7.50 (d, $J=8.4$ Hz, 2H), 3.54 (d, $J=7.8$ Hz, 1H), 2.91~2.73 (m, 2H), 2.72~2.56 (m, 2H), 2.51~2.46 (m, 1H), 2.07~1.91 (m, 4H), 1.83~1.73 (m, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 196.4, 194.2, 147.1, 130.0 (q, $J=2.4$ Hz), 125.6 (q, $J=3.8$ Hz), 124.7, 123.8 (q, $J=157.4$ Hz), 113.9, 50.4, 43.0, 40.6, 32.5, 25.3, 22.2; HRMS (ESI-TOF) calcd for $\text{C}_{17}\text{H}_{16}\text{F}_3\text{O}_2$ $[\text{M} + \text{H}]^+$ 309.1097, found 309.1090.

3a-(4-(Trifluoromethyl)phenyl)-3,3a,5,6,7,8b-hexahydro-1H-cyclopenta[*b*]benzofuran-8(2H)-one (**9b**): colorless oil, 99% yield (63.8 mg). ^1H NMR (600 MHz, CDCl_3) δ : 7.61 (d, $J=8.4$ Hz, 2H), 7.49 (d, $J=8.4$ Hz, 2H), 3.65 (d, $J=7.8$ Hz, 1H), 2.50~2.54 (m, 1H), 2.53~2.46 (m, 1H), 2.44~2.26 (m, 3H), 2.12~1.94 (m, 5H), 1.92~1.88 (m, 1H), 1.72~1.63 (m, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 195.3, 176.3, 147.9, 129.7 (q, $J=2.3$ Hz), 125.4 (q, $J=3.8$ Hz), 124.8, 124.5 (q, $J=270.3$ Hz), 115.9, 102.5, 52.0, 43.5, 36.6, 33.3, 24.9, 23.8, 21.8; HRMS (ESI-TOF) calcd for $\text{C}_{18}\text{H}_{18}\text{F}_3\text{O}_2$ $[\text{M} + \text{H}]^+$ 323.1253, found 323.1271.

6,6-Dimethyl-3a-(4-(trifluoromethyl)phenyl)-3,3a,5,6,7,8b-hexahydro-1H-cyclopenta[*b*]benzofuran-8(2H)-one (**9c**): yellow solid, 79% yield (55.4 mg). m.p. 117~118 °C; NMR Spectroscopy: ^1H NMR (600 MHz, CDCl_3) δ : 7.61 (d, $J=8.4$ Hz, 2H), 7.48 (d, $J=7.8$ Hz, 2H), 3.63 (d, $J=7.8$ Hz, 1H), 2.42~2.34 (m, 3H), 2.27~2.17 (m, 2H), 2.10~1.87 (m, 4H), 1.73~1.62 (m, 1H), 1.13 (s, 3H), 1.11 (s, 3H); ^{13}C NR (150 MHz, CDCl_3) δ : 194.6, 175.3, 147.9, 129.7 (q, $J=2.3$ Hz), 125.5 (q, $J=3.8$ Hz), 124.5 (q, $J=270.3$ Hz), 124.8, 114.4, 102.7, 52.0, 51.1, 43.3, 37.7, 34.1, 33.3, 29.1, 28.0, 25.0; HRMS (ESI-TOF) calcd for $\text{C}_{20}\text{H}_{22}\text{F}_3\text{O}_2$ $[\text{M} + \text{H}]^+$ 351.1566, found 351.1567.

(2-Methyl-6a-(4-(trifluoromethyl)phenyl)-4,5,6,6a-tetrahydro-3aH-cyclopenta[*b*]furan-3-yl)(phenyl)methanone (**10a**): colorless oil, 34% yield (25.3 mg). ^1H NMR (600 MHz, CDCl_3) δ : 7.63 (d, $J=8.4$ Hz, 2H), 7.53 (d, $J=8.4$ Hz, 2H), 7.51~7.48 (m, 2H), 7.46 (d, $J=7.2$ Hz, 1H), 7.40 (t, $J=7.2$ Hz, 2H), 3.90 (d, $J=8.4$ Hz, 1H), 2.36 (dd, $J=13.8$, 6.0 Hz, 1H), 2.11~1.96 (m, 3H), 1.96~1.88 (m, 1H), 1.87~1.74 (m, 4H); ^{13}C NMR (150 MHz, CDCl_3) δ : 193.3, 167.5, 148.6, 141.2, 131.0, 129.5 (q, $J=32.6$ Hz), 128.3, 127.7, 125.5 (q, $J=3.6$ Hz), 124.8, 124.1 (q, $J=270.8$ Hz), 116.3, 98.7, 56.3, 43.7, 34.7, 25.0, 15.5; HRMS (ESI-TOF) calcd for $\text{C}_{22}\text{H}_{19}\text{F}_3\text{O}_2\text{Na}$ $[\text{M} + \text{Na}]^+$ 395.1229, found 395.1217.

1-(2-Phenyl-6a-(4-(trifluoromethyl)phenyl)-4,5,6,6a-tetrahydro-3aH-cyclopenta[*b*]furan-3-yl)ethanone (**10a'**): white solid, yield 34% (25.3 mg). m.p. 119~120 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.64 (d, $J=8.4$ Hz, 2H), 7.62~7.54 (m, 4H), 7.54~7.46 (m, 3H), 3.85 (d, $J=8.4$ Hz, 1H), 2.42 (dd, $J=13.8$, 6.0 Hz, 1H), 2.24~2.13 (m, 1H), 2.13~2.00 (m, 2H), 2.00~1.93 (m, 1H), 1.92~1.81 (m, 4H); ^{13}C NMR (150 MHz, CDCl_3) δ : 194.3, 166.1,

148.5, 131.0, 130.7, 129.7 (q, $J=32.4$ Hz), 129.1, 128.5, 125.5 (q, $J=3.6$ Hz), 124.8, 124.1 (q, $J=270.8$ Hz), 118.5, 98.7, 56.3, 44.0, 35.1, 29.0, 25.1; HRMS (ESI-TOF) calcd for $C_{22}H_{19}F_3O_2Na$ $[M+Na]^+$ 395.1229, found 395.1229.

1-(2-Methyl-6a-(4-(trifluoromethyl)phenyl)-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)propan-1-one (**10b**): colorless oil, 56% yield (36.3 mg). 1H NMR (600 MHz, $CDCl_3$) δ : 7.61 (d, $J=8.4$ Hz, 2H), 7.50 (d, $J=8.4$ Hz, 2H), 3.62 (d, $J=8.4$ Hz, 1H), 2.48 (q, $J=7.2$ Hz, 2H), 2.35 (s, 3H), 2.32 (dd, $J=13.8, 6.0$ Hz, 1H), 2.16~2.06 (m, 1H), 2.05~1.97 (m, 1H), 1.94~1.87 (m, 2H), 1.80~1.69 (m, 1H), 1.07 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 197.4, 166.7, 148.5, 129.5 (q, $J=32.1$ Hz), 125.3 (q, $J=3.6$ Hz), 124.0 (q, $J=270.3$ Hz), 124.7, 115.3, 98.6, 55.6, 43.2, 35.2, 34.0, 24.9, 15.0, 7.9; HRMS (ESI-TOF) calcd for $C_{18}H_{19}F_3O_2Na$ $[M+Na]^+$ 347.1229, found 347.1227.

1-(2-Ethyl-6a-(4-(trifluoromethyl)phenyl)-4,5,6,6a-tetrahydro-3aH-cyclopenta[b]furan-3-yl)ethanone (**10b'**): colorless oil, 42% yield (27.2 mg). 1H NMR (600 MHz, $CDCl_3$) δ : 7.54 (d, $J=8.4$ Hz, 2H), 7.42 (d, $J=7.8$ Hz, 2H), 3.53 (d, $J=8.4$ Hz, 1H), 2.78~2.59 (m, 2H), 2.24 (dd, $J=13.8, 6.0$ Hz, 1H), 2.12 (s, 3H), 2.08~1.98 (m, 1H), 1.95~1.89 (m, 1H), 1.87~1.77 (m, 2H), 2.78~2.59 (m, 1H), 1.17 (t, $J=7.6$ Hz, 3H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 194.1, 172.1, 148.6, 129.5 (q, $J=32.4$ Hz), 125.3 (q, $J=3.6$ Hz), 124.6, 129.5 (q, $J=124.01$ Hz), 114.9, 98.3, 55.9, 43.5, 35.1, 29.1, 24.7, 22.0, 11.5; HRMS (ESI-TOF) calcd for $C_{18}H_{19}F_3O_2Na$ $[M+Na]^+$ 347.1229, found 347.1233.

Supporting Information Crystal structures of **6e**, and 1H NMR and ^{13}C NMR spectra of new compounds **3**, **5**~**10**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] Nakashima, K.; Masayoshi Oyama, M.; Ito, T.; Akao, Y.; Witono, J. R.; Darnaedi, D.; Tanaka, T.; Murata, J.; Iinuma, M. *Tetrahedron* **2012**, *68*, 2421.
- [2] Lallemand, J.-Y.; Six, Y.; Ricard, L. *Eur. J. Org. Chem.* **2002**, 503.
- [3] Baraza, L. D.; Joseph, C. C.; Nkunya, M. H. *Nat. Prod. Res.* **2007**, *21*, 1027.
- [4] Ghosh, A. K.; Parham, G. L.; Martyr, C. D.; Nyalapatla, P. R.; Osswald, H. L.; Agniswamy, J.; Wang, Y.-F.; Amamo, M.; Weber, I. T.; Mitsuya, H. *J. Med. Chem.* **2013**, *56*, 6792.
- [5] (a) Iqbal, J.; Bhatia, B.; Nayyar, N. K. *Chem. Rev.* **1994**, *94*, 519.
(b) Lee, Y. R.; Kim, B. S.; Kim, D. H. *Tetrahedron* **2000**, *56*, 8845.
(c) Pan, X.-Q.; Zou, J.-P.; Zhang, W. *Mol. Diversity* **2009**, *13*, 421.
(d) Liu, X.; Chen, X.; Mohr, J. T. *Chem.-Eur. J.* **2016**, *22*, 2274.
(e) Calter, M. A.; Phillips, R. M.; Flaschenriem, C. J. *Am. Chem. Soc.* **2005**, *127*, 14566.
(f) Chennapuram, M.; Emmadi, N. R.; Bingi, C.; Nanubolu, J. B.; Atmakur, K. *Green Chem.* **2014**, *16*, 3237.
(g) Chawla, R.; Singh, A. K.; Yadav, L. D. S. *Tetrahedron Lett.* **2012**, *53*, 3382.
- [6] (a) Mekonnen, A.; Carlson, R. *Synthesis* **2006**, 1657.
(b) Zhong, C.; Liao, T.; Tuguldur, O.; Shi, X. *Org. Lett.* **2010**, *12*, 2064.
(c) Heiba, E.-A. I.; Dessau, R. M. *J. Org. Chem.* **1974**, *39*, 3456.
(d) Riveira, M. J.; Mischne, M. P. *J. Org. Chem.* **2014**, *79*, 8244.
(e) Zhang, Z.; Li, C.; Wang, S. H.; Zhang, F. M.; Han, X.; Tu, Y. Q.; Zhang, X. M. *Org. Biomol. Chem.* **2017**, *15*, 3239.
- [7] (a) Karade, N. N.; Shiroadkar, S. G.; Patil, M. N.; Potrekar, R. A.; Karade, H. N. *Tetrahedron Lett.* **2003**, *44*, 6729.
(b) Nishino, H.; Fujino, R. *Synthesis* **2005**, 731.
(c) Yilmaz, M.; Pekel, A. T. *J. Fluorine Chem.* **2011**, *132*, 628.
(d) Dengiz, C.; Caliskan, R.; Balci, M. *Tetrahedron Lett.* **2012**, *53*, 550.
(e) Yirsaw, A. M.; Carlson, R. E. *ARKIVOC* **2017**, iv, 74.
(f) Lal, S.; Chowdhury, A.; Namboothiri, I. N. N. *Tetrahedron* **2017**, *73*, 1297.
- [8] Ye, S.; Yu, Z.-X. *Chem. Commun.* **2011**, 47, 794.
- [9] CCDC 1589697 (**6e**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.
- [10] (a) Wang, H.; Guo, L.-N.; Duan, X.-H. *Org. Lett.* **2013**, *15*, 5254.
(b) Wang, G.-W.; Dong, Y.-W.; Wu, P.; Yuan, T.-T.; Shen, Y.-B. *J. Org. Chem.* **2008**, *73*, 7088.
(c) Li, P.; Zhao, J.; Xia, C.; Li, F. *Org. Lett.* **2014**, *16*, 5992.
(d) Casey, B. M.; Eakin, C. A.; Jiao, J.; Sadasivam, D. V.; Flowers, R. A. *Tetrahedron* **2009**, *65*, 10762.
(e) Yi, H.; Liao, Z.; Zhang, G.; Zhang, G.; Fan, C.; Zhang, X.; Bunel, E. E.; Pao, C. W.; Lee, J. F.; Lei, A. *Chem.-Eur. J.* **2015**, *21*, 18925.

(Zhao, C.)