有机化学

Chinese Journal of Organic Chemistry

金属有机化学专辑

第 39 卷 第 6 期 Vol. 39 No. 6 2019

ISSN 0253-2786

中国化学会主办
中国科学院上海有机化学研究所

Strem Chemicals, Inc., established in 1964, is a privately held manufacturer & marketer of high purity specialty chemicals. Strem's products are used for R&D and commercial scale applications, especially in the pharmaceutical, microelectronics, chemicals & petrochemicals industries. Some key products include catalysts, ligands, organometallics, CVD/ALD precursors and nanomaterials. For more information, visit strem.com. Strem is an ISO 9001:2015 certified company.

STREM'S KEY PRODUCT LINES

- METAL CATALYSTS
- LIGANDS
- INORGANICS
- ORGANOMETALLICS
- PHOTOCATALYSTS
- SCREENING KITS

11-0575

27-0400

78-1350

77-0425

15-8021

15-1053

15-1060

46-2150

29-6720

15-0067

77-0410

15-0126

中国高端 均相催化剂领军者

- 10年研发经验
- 国际市场认可
- 无水无氧封装存储

贵金属催化剂：

Pd Rh Ru Ir Au Cu Ni Zr

有机配体：

P N O

MIT专利授权：

BUCHWALD LIGAND & PRECATALYST

手性膦配体：

BINAP SEGPPOS PHOSPHORAMIDITE
JOSIPHOS PHOX SPIRO

手性氮配体：

BOX BINAM DACH TROST DPEN SALEN

手性催化剂：

PHOSPHORIC ACIDS NHC THIUREA
CINCHONA ALKALOID

光催化剂：

Ir Ru ACRI. SALTS

OTHER: RAFT SILANE PORPHYRIN MOF
ALD PRECURSOR

媲美进口品质的催化剂尽在

WWW.LAAJOO.COM

TEL: 021-60702684

QQ: 3004302861

MAIL: REAGENT@SINOCOMPOUND.COM

理想的实验室伴侣！

中国人都用得起的隔膜真空泵！

旋转蒸发仪/隔膜真空泵

® 上海豫康科教仪器设备有限公司

Shanghai Yukang Science teaching Instrument equipment Co.,Ltd

地址：上海市和桥路470号

电话：021-36338262 56470425
56470426

传真：021-56470425

手机：13701684908

邮编：200435

网址：www.sh-yukang.com

Tel: 021-36338262 56470425
56470426

Fax: 021-56470425

Mobil: 13701684908

Zip Code: 200435

Web: www.sh-yukang.com

由饱和酮与肼/醛肼的[3+2]/[2+3]环化反应区域选择性合成多取代吡唑

师晓楠 田苗苗 王慕华 张新迎 范学森*

(河南师范大学化学化工学院 河南省有机功能分子与药物创新重点实验室
绿色化学介质与反应教育部重点实验室 新乡 453007)

摘要 报道了一种经由二价铜催化饱和酮与肼或醛肼的串联反应合成 1,3-二取代吡唑或 1,3,4-三取代吡唑类化合物的简便和区域选择性新方法。从机理上看, 1,3-二取代吡唑的生成经历了烯酮中间体的原位生成及其与肼的[3+2]环化反应, 而 1,3,4-三取代吡唑的生成则经历了烯酮中间体的原位生成及其与醛肼的[2+3]环化反应。与文献方法相比, 该方法具有原料简单易得、底物适用范围广、区域选择性好、效率高、原子经济性优异等优势。

关键词 吡唑合成; 二价铜催化反应; $C(sp^3)$ -H 官能团化; [3+2]环化; [2+3]环化

Regioselective Synthesis of Substituted Pyrazoles via [3+2]/[2+3] Cyclization of Saturated Ketone with Hydrazine/Aldehyde Hydrazone

Shi, Xiaonan Tian, Miaomiao Wang, Muhua Zhang, Xinying Fan, Xuesen*

(Henan Key Laboratory of Organic Functional Molecules and Drug Innovation, Key Laboratory of Green Chemical Media and Reactions, Ministry of Education, School of Chemistry and Chemical Engineering, Henan Normal University, Xinxiang 453007)

Abstract A highly convenient and regioselective synthesis of 1,3-disubstituted pyrazoles or 1,3,4-trisubstituted pyrazoles from Cu(II)-catalyzed cascade reactions of saturated ketones with hydrazines or aldehyde hydrazones is presented. Mechanistically, the formation of 1,3-disubstituted pyrazoles involves the *in situ* generation of an enone intermediate followed by its [3+2] annulations with hydrazine. On the other hand, the formation of 1,3,4-trisubstituted pyrazoles is believed to go through a cascade process including enone formation and its subsequent [2+3] annulation with aldehyde hydrazone. Compared with literature methods, the notable features of the protocol include simple starting materials, general and broad substrate scope, high regioselectivity, good efficiency and excellent atom-economy.

Keywords pyrazole synthesis; Cu(II) catalyzed reaction; $C(sp^3)$ -H functionalization; [3+2] annulation; [2+3] annulation

1 Introduction

It is well known that substituted pyrazoles are essential building blocks of numerous pharmaceuticals, agrochemicals, dyes, pigments and functional materials.^[1,2] This could be representatively exemplified by the following clinical drugs such as rimonabant, celebrex, viagra, ceftolozane and omarigliptin (Figure 1).

Due to their importance, a number of methods to construct the pyrazole scaffold have been developed, which mainly include: (1) condensation of hydrazine with enone, 1,3-dicarbonyl compounds or 2-ethoxymethylene-3-oxo-

butanenitrile,^[3] (2) [3+2] cycloaddition of 1,3-dipoles with dipolarophiles,^[4] (3) multi-component reactions of vinyl azide with aldehydes and tosylhydrazine,^[5a] arylglyoxal with tosylhydrazine and aldehydes/ ketones,^[5b] arylhydrazines with alkynes and CO,^[5c] or terminal alkynes with aldehydes and hydrazines,^[5d] (4) transformations of hydrazone derivatives, such as oxidative cyclization of α,β -unsaturated hydrazones,^[6a] reactions of hydrazones with α -bromoketones,^[6b] oxidative cyclization of β,γ -unsaturated hydrazones,^[6c] Sonogashira coupling/cyclization of *N*-propargyl sulfonyl hydrazones,^[6d] [3+2]-cyclo-

* Corresponding author. E-mail: xuesen.fan@htu.cn

Received March 23, 2019; revised May 14, 2019; published online May 21, 2019.

Project supported by the National Natural Science Foundation of China (No. 21572047) and the Plan for Scientific Innovation Talents of Henan Province (No. 184200510012).

国家自然科学基金(No. 21572047)和河南省科技创新杰出人才(No. 184200510012)资助项目。

Figure 1 Clinical drugs containing a pyrazole scaffold

addition of ynones with CF_3CHN_2 .^[7] While these methods are generally efficient and reliable, some of them still suffer from difficulty in accessing highly functionalized substrates, poor regioselectivity and monotonous substitution pattern of products. Therefore, the development of novel methods for the regioselective preparation of diversely substituted pyrazoles starting from simple and easily obtainable substrates is still highly desirable.

In recent years, direct functionalization of the inert $\text{C}(\text{sp}^3)\text{—H}$ bonds is emerging as an attractive strategy in organic synthesis due to their advantages such as simple substrates, high sustainability and step-economy. In this regard, direct oxidative dehydrogenation of inactivated saturated ketones has been developed as a sustainable and reliable approach toward enones.^[8] More interestingly, this enone formation protocol could be conveniently combined with other transformations to form advanced molecular scaffolds from simple and economic starting materials in a one-pot manner with remarkably reduced operation steps.^[9] Inspired by these pioneering studies and as continuation of our own interest in this aspect,^[10] we have explored the reaction of saturated ketones with hydrazines under the catalysis of copper catalyst. From this study, a regioselective and sustainable synthesis of 1,3-disubstituted pyrazoles was established.

2 Results and discussion

Our study was commenced by treating propiophenone (**1a**) with $\text{Cu}(\text{OAc})_2$, 2,2,6,6-tetramethyl piperidine-*N*-oxyl (TEMPO) and 2,2'-bipyridine (bpy) in chlorobenzene (PhCl) at 120 °C under N_2 for 10 h.^[9] The resulting mixture was then treated with phenylhydrazine (**2a**) at 120 °C under air for 8 h. From this reaction, 1,3-diphenyl-1*H*-pyrazole (**3a**) was obtained in a yield of 52%. Meanwhile, the formation of 1,5-diphenyl-1*H*-pyrazole (**3a'**), the regioisomer of **3a**, was not observed (Eq. 1).

To improve the efficiency, several parameters were screened. First of all, the amount of TEMPO was increased

Scheme 1 Formation of **3a** from the one-pot reaction of **1a** with **2a**

from 1 to 2 equiv., and the yield of **3a** increased slightly (Table 1, Entry 2). When 0.5 equiv. of TEMPO was used, the reaction efficiency dropped obviously (Entry 3). In the absence of TEMPO, the formation of **3a** was not observed (Entry 4). As for the effect of different reaction media, dioxane, 1,2-dichloroethane (DCE) and CH_3CN were found to be less efficient than PhCl (Entries 5~7). Next, $\text{Cu}(\text{OAc})_2$ was replaced by CuBr_2 or CuCl_2 as the catalyst. Under this circumstance, the yield of **3a** decreased significantly (Entries 8, 9 vs. 1). Then, the second stage of this cascade reaction was carried out under oxygen instead of air. To our pleasure, the yield of **3a** was improved to 70% (Entry 10). It was also found that a period of 5 h was the most suitable time for the second stage of this cascade process (Entry 11 vs. 10, 12). Further studies showed that varying the ratio of **1a** : **2a** did not improve the yield of **3a** (Entries 13 and 14). Moreover, increasing or reducing the amount of $\text{Cu}(\text{OAc})_2$ did not result in obvious change of the reaction efficiency (Entries 15 and 16). As a summary, **3a** was formed in a yield of 71% by treating 0.6 mmol of **1a** with 0.1 mmol of $\text{Cu}(\text{OAc})_2$, 0.5 mmol of TEMPO and 0.05 mmol of bpy in PhCl at 120 °C under N_2 for 10 h, and then treating the resulting mixture with 0.5 mmol of **2a** at 120 °C under O_2 for 5 h (Entry 11).

With the optimum reaction conditions in hands, we then the scope and generality of this 1,3-disubstituted pyrazole forming reaction were explored, and the results were listed in Table 2. First, a range of ketones **1** were screened by

Table 1 Optimization studies for the formation of **3a**^a

Entry	Catalyst	Oxidant (equiv.)	Solvent	t/h	Yield ^b /%
1	Cu(OAc) ₂	TEMPO (1)	PhCl	10+8	52
2	Cu(OAc) ₂	TEMPO (2)	PhCl	10+8	54
3	Cu(OAc) ₂	TEMPO (0.5)	PhCl	10+8	30
4	Cu(OAc) ₂	TEMPO (0)	PhCl	10+8	ND
5	Cu(OAc) ₂	TEMPO (1)	Dioxane	10+8	47
6	Cu(OAc) ₂	TEMPO (1)	CH ₃ CN	10+8	45
7	Cu(OAc) ₂	TEMPO (1)	DCE	10+8	45
8	CuBr ₂	TEMPO (1)	PhCl	10+8	32
9	CuCl ₂	TEMPO (1)	PhCl	10+8	23
10	Cu(OAc) ₂	TEMPO (1)/O ₂	PhCl	10+8	70
11	Cu(OAc) ₂	TEMPO (1)/O ₂	PhCl	10+5	71
12	Cu(OAc) ₂	TEMPO (1)/O ₂	PhCl	10+3	52
13 ^c	Cu(OAc) ₂	TEMPO (1)/O ₂	PhCl	10+5	60
14 ^d	Cu(OAc) ₂	TEMPO (1)/O ₂	PhCl	10+5	71
15 ^e	Cu(OAc) ₂	TEMPO (1)/O ₂	PhCl	10+5	73
16 ^f	Cu(OAc) ₂	TEMPO (1)/O ₂	PhCl	10+5	41

^a Reaction conditions: **1a** (0.6 mmol), **2a** (0.5 mmol), catalyst (0.1 mmol), bpy (0.05 mmol), solvent (3 mL). ^b Isolated yield. ^c **1a** (0.5 mmol), **2a** (0.6 mmol).

^d **1a** (0.75 mmol). ^e Cu(OAc)₂ (0.15 mmol). ^f Cu(OAc)₂ (0.05 mmol).

using **2a** as a model substrate. Among them, propiophenones bearing various substituents on the phenyl ring reacted with **2a** smoothly to give **3a**~**3i** in moderate to good yields. Notably, a number of functional groups, from the electron-donating methoxy to the electron-withdrawing fluoro, were well compatible with the reaction conditions. Moreover, 2-propionyl thiophene as a hetero aromatic ketone took part in this cascade process smoothly to give **3j** with good efficiency. Interestingly, two aliphatic ketones, pentan-3-one and 1-cyclohexylpropan-1-one were also suitable substrates for this reaction to afford **3k** and **3l**. When 1-phenylbutan-1-one, 1-phenylpentan-1-one or 1,3-di-phenylpropan-1-one was tried, the formation of the corresponding products **3'**, **3''** or **3'''** was not observed, most likely due to steric effect. The structure of **3f** was confirmed by its X-ray single crystal diffraction analysis.^[11]

Next, the scope of hydrazines (**2**) was studied by using **1a** as a model substrate. The results included in Table 3 showed that a series of phenyl hydrazine derivatives bearing a methyl unit on the *meta*-position, or a fluoro, chloro, methyl, or methoxy group on the *para*-position of the phenyl moiety took part in this reaction smoothly to give **3m**~**3q** in moderate to good yields. In addition, the reactions of two aliphatic hydrazines, methyl hydrazine and isopropyl hydrazine, proceeded smoothly to afford **3r** and **3s** in yields of 45% and 42%, respectively.

Thus far, a novel synthesis of 1,3-disubstituted pyrazoles via Cu(II)-promoted cascade reactions of saturated ketones with hydrazines were established (Tables 2 and 3). Next, we turned our focus from the synthesis of 1,3-disubstituted pyrazoles to the preparation of 1,3,4-tri-substituted pyra-

Table 2 Scope of ketones for the synthesis of **3**^{a,b}

^a Reaction conditions: **1** (0.6 mmol), **2a** (0.5 mmol), Cu(OAc)₂ (0.1 mmol), TEMPO (0.5 mmol), bpy (0.05 mmol), PhCl (3 mL). ^b Isolated yields.

Table 3 Scope of hydrazines for the synthesis of **3**^{a,b}

^a Reaction conditions: **1a** (0.6 mmol), **2** (0.5 mmol), Cu(OAc)₂ (0.1 mmol), TEMPO (0.5 mmol), bpy (0.05 mmol), PhCl (3 mL). ^b Isolated yields.

zoles bearing a 4-acyl moiety, which are well known for their potent biological activities and rich chemical reactivity.^[1,12] In this regard, a facile synthesis of 1,3,4-trisubstituted pyrazoles via the cascade reactions of saturated ketones with ketone hydrazones was reported.^[10b] While this synthetic protocol provides an efficient and easy-to-

run approach toward pyrazole derivatives, it nevertheless has the drawback of being relatively poor in atom-economy since part of the ketone hydrazone used therein is lost during the formation of the pyrazole scaffold (Scheme 2). To solve this problem, aldehyde hydrazones **4** were used to replace ketone hydrazones as the substrates to react with **1**. Thus, **1a** was subjected to $\text{Cu}(\text{OAc})_2/\text{TEMPO}/\text{bpy}$, and then treated with 1-benzylidene-2-phenylhydrazine (**4a**). To our delight, the desired (1,3-diphenyl-1*H*-pyrazol-4-yl)(phenyl)methanone (**5a**) was obtained in 63% yield (Scheme 2). It is worth to be noted herein that although the yield of **5a** is slightly lower than that with our previously reported method (63% vs 72%),^[10b] the direct formation of **5a** from **1a** and **4a** is still synthetically useful, not only because it is more atom economy, but also because aldehyde hydrazones are much more readily available and economically sustainable than ketone hydrazones.

Our previous work, Ref. [10b]

Scheme 2 Synthesis of 1,3,4-trisubstituted pyrazoles

To explore the substrate scope of this reaction, several saturated ketones (**1**) were tried to react with **4a**. The results included in Table 4 demonstrated that propiophenones bearing various substituents on the phenyl ring reacted with **4a** smoothly to give **5a**~**5e** in moderate yields. Notably, the electronic nature of the phenyl unit did not render an obvious effect on the reaction efficiency as demonstrated in the formation of **5b**~**5e**. Moreover, 2-propionyl thiophene and 3-propionylpyridine as heteroaromatic ketone substrates took part in this cascade reaction smoothly to give **5f** and **5g** in moderate yields. Finally, 1-cyclohexyl propan-1-one as an aliphatic ketone substrate could also react with **4a** to give **5h** in a yield of 51%.

Next, the suitability of different aldehyde hydrazones **4** for the synthesis of **5** was explored, and the results were listed in Table 5. First, hydrazones derived from diversely substituted benzaldehydes were let to react with **1a**. From these reactions, **5i**~**5o** were obtained in 42%~74% yields. Notably, the reaction of 1-phenyl-2-(thiophen-2-ylmethylene)hydrazine with **1a** proceeded smoothly to give **5p**. Second, aldehyde hydrazones **4** derived from different hydrazines were found to react with **1a** readily to afford **5q**~**5y** in moderate yields. Third, diversely substituted propiophenones reacted with various aldehyde hydrazones

Table 4 Scope of ketones for the synthesis of **5** (*I*)^{a,b}

^a Reaction conditions: **1** (0.6 mmol), **4a** (0.5 mmol), $\text{Cu}(\text{OAc})_2$ (0.1 mmol), TEMPO (0.5 mmol), bpy (0.05 mmol), PhCl (3 mL). ^b Isolated yields.

to give **5z**~**5dd**, thus resulting in a highly versatile synthesis of 1,3,4-trisubstituted pyrazoles.

To get some insight into the reaction mechanism accounting for the formation of **3a**, the following control experiments were carried out. First, the pre-prepared enone **A** was treated with **2a** under standard reaction conditions. From this reaction, **3a** was obtained in a yield of 78% (Scheme 3). This result told that enone **A** should be an intermediate in the formation of **3a**. Second, **A** was treated with **2a** in ethanol at room temperature for 10 min to give **B** in a yield of 92% (Scheme 3). Third, stirring the solution of **B** in PhCl at 120 °C under O_2 for 5 h afforded **3a** in a yield of 93% (Scheme 3). These results indicated that **B** should be another key intermediate in the formation of **3a**.

Scheme 3 Control experiment

Based on the above described experimental results and previous reports,^[6,8] a plausible mechanism accounting for the formation of **3a** is proposed in Scheme 4. Initially, **1a**

Table 5 Scope of hydrazines for the synthesis of **5**^{a,b}

^a Reaction conditions: **1** (0.6 mmol), **4** (0.5 mmol), Cu(OAc)₂ (0.1 mmol), TEMPO (0.5 mmol), bpy (0.05 mmol), PhCl (3 mL). ^b Isolated yields.

is transformed into enone **A** through Cu(II)-catalyzed oxidative dehydrogenation. Next, condensation of **A** with **2a** affords an α,β -unsaturated hydrazone **II**, most likely via the formation of intermediate **I**. The following intramolecular Michael addition of **II** leads to the formation of dihydropyrazole **B**. Finally, an oxidative aromatization occurs with **B** to afford **3a**.

Scheme 4 Plausible mechanism for the formation of **3a**

As for the formation of **5a** from the reaction of **1a** with **4a**, it should also involve an initial oxidative dehydrogenation of **1a** to give enone **A**, which then undergoes a conjugate addition with **4a** to afford adduct **III**. Next, an intramolecular nucleophilic addition occurs with **III** to afford tetrahydropyrazole **IV**. Under the reaction conditions, **IV** undergoes oxidative dehydrogenation to afford **5a** as the final product (Scheme 5).

Scheme 5 Plausible mechanism for the formation of **5a**

This proposed mechanism shown in Scheme 5 is partly supported by the following control experiment. The pre-prepared enone **A** was treated with **4a** under standard reaction conditions. From this reaction, **5a** was obtained in a yield of 52% (Eq. 1). This result told that enone **A** should be an intermediate in the formation of **5a**.

Finally, to determine whether this newly developed pyrazole synthesis is suitable for large scale applications, the

preparation of **3a** was carried out in an enlarged scale of 5 mmol. From this reaction, **3a** was obtained in a yield of 60% (Scheme 6).

Scheme 6 Gram-scale synthesis of **3a**

3 Conclusions

In summary, we have developed a novel synthesis of 1,3-disubstituted pyrazoles from the reaction of saturated ketones with hydrazines, and an improved synthesis of 1,3,4-trisubstituted pyrazoles from the reaction of saturated ketones with aldehyde hydrazones. Compared with literature methods, notable features of these new protocols include simple and economical starting materials, high regioselectivity and excellent step- and atom-economy. The development of more applications of saturated ketones as enone precursors in the synthesis of heterocyclic compounds is currently under way in our laboratories.

4 Experimental section

4.1 General experimental information

Commercial reagents were used without further purification. Melting points were recorded with a micro melting point apparatus and uncorrected. The ^1H NMR spectra were recorded on a Bruker DPX spectrometer at 400 MHz or 600 MHz. The ^{13}C NMR spectra were recorded on a Bruker DPX spectrometer at 100 MHz or 150 MHz. High resolution mass spectra (HRMS) were obtained via ESI mode by using a MicrOTOF-QII mass spectrometer. The conversion of starting materials was monitored by thin layer chromatography (TLC) using silica gel plates (silica gel 60 F254 0.25 mm), and components were visualized by observation under UV light (254 and 365 nm).

4.2 Experimental procedures

4.2.1 Typical procedure for the synthesis of **3a**

To a reaction tube equipped with a stir bar were added propiophenone (**1a**, 81 mg, 0.6 mmol), PhCl (3 mL), $\text{Cu}(\text{OAc})_2$ (18 mg, 0.1 mmol), bpy (8 mg, 0.05 mmol) and TEMPO (78 mg, 0.5 mmol). The mixture was stirred at 120 °C under N_2 for 10 h. To the resulting mixture was added phenylhydrazine (**2a**, 54 mg, 0.5 mmol). It was then stirred at 120 °C under O_2 for 5 h. Upon completion, the mixture was diluted with dichloromethane (DCM) (15 mL), and washed with water (5 mL) and brine (5 mL). The organic layer was dried over anhydrous Na_2SO_4 , and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel with petroleum ether/ethyl acetate ($V:V=50:1$) as the eluent to give **3a**. **3b**~**3s** were obtained in a similar manner.

1,3-Diphenyl-1H-pyrazole (**3a**): Yellow solid (78 mg,

71%). m.p. 79~81 °C (lit.^[13] (82~85 °C)); ^1H NMR (400 MHz, CDCl_3) δ : 6.75 (d, $J=2.4$ Hz, 1H), 7.25~7.29 (m, 1H), 7.31~7.35 (m, 1H), 7.40~7.46 (m, 4H), 7.74~7.77 (m, 2H), 7.91~7.93 (m, 3H); ^{13}C NMR (100 Hz, CDCl_3) δ : 105.1, 119.1, 125.9, 126.4, 128.0, 128.1, 128.7, 129.5, 133.2, 140.3, 153.0; IR (KBr) ν : 1513, 1252, 1021, 952, 827, 745, 702 cm^{-1} ; HRMS calcd for $\text{C}_{15}\text{H}_{13}\text{N}_2$ [$\text{M}+\text{H}$] $^+$ 221.1073, found 221.1064.

3-(3-Fluorophenyl)-1-phenyl-1H-pyrazole (**3b**)^[14]: Yellow solid (68 mg, 57%). m.p. 57~59 °C; ^1H NMR (400 MHz, CDCl_3) δ : 6.76 (d, $J=2.8$ Hz, 1H), 7.00~7.05 (m, 1H), 7.29~7.33 (m, 1H), 7.36~7.41 (m, 1H), 7.45~7.49 (m, 2H), 7.62~7.69 (m, 2H), 7.75~7.78 (m, 2H), 7.96 (d, $J=2.8$ Hz, 1H); ^{13}C NMR (100 Hz, CDCl_3) δ : 105.2, 112.7 (d, $^2J_{\text{C-F}}=22.6$ Hz), 114.8 (d, $^2J_{\text{C-F}}=21.8$ Hz), 119.1, 121.5 (d, $^4J_{\text{C-F}}=2.9$ Hz), 126.6, 128.2, 129.5, 130.2 (d, $^3J_{\text{C-F}}=8.0$ Hz), 135.4 (d, $^3J_{\text{C-F}}=8.0$ Hz), 140.1, 151.8 (d, $^4J_{\text{C-F}}=2.9$ Hz), 163.2 (d, $^1J_{\text{C-F}}=243.7$ Hz); IR (KBr) ν : 1516, 1252, 1026, 953, 825, 742, 699 cm^{-1} ; HRMS calcd for $\text{C}_{15}\text{H}_{12}\text{FN}_2$ [$\text{M}+\text{H}$] $^+$ 239.0979, found 239.0975.

3-(3-Chlorophenyl)-1-phenyl-1H-pyrazole (**3c**)^[14]: Yellow solid (77 mg, 61%). m.p. 92~93 °C; ^1H NMR (400 MHz, CDCl_3) δ : 6.68 (d, $J=2.4$ Hz, 1H), 7.21~7.29 (m, 3H), 7.37~7.41 (m, 2H), 7.68~7.72 (m, 3H), 7.85~7.88 (m, 2H); ^{13}C NMR (100 Hz, CDCl_3) δ : 104.1, 118.1, 122.9, 124.9, 125.6, 126.9, 127.2, 128.4, 128.9, 133.6, 133.9, 139.1, 150.5; IR (KBr) ν : 1519, 1252, 1023, 954, 825, 745, 701 cm^{-1} ; HRMS calcd for $\text{C}_{15}\text{H}_{12}\text{ClN}_2$ [$\text{M}+\text{H}$] $^+$ 255.0684, found 255.0680.

3-(3-Methoxyphenyl)-1-phenyl-1H-pyrazole (**3d**)^[6c]: Yellow solid (70 mg, 56%). m.p. 68~69 °C; ^1H NMR (400 MHz, CDCl_3) δ : 3.81 (s, 3H), 6.68 (d, $J=2.4$ Hz, 1H), 6.81~6.83 (m, 1H), 7.21 (t, $J=7.2$ Hz, 1H), 7.26 (t, $J=8.0$ Hz, 1H), 7.36~7.43 (m, 4H), 7.68~7.70 (m, 2H), 7.86 (d, $J=2.4$ Hz, 1H); ^{13}C NMR (100 Hz, CDCl_3) δ : 55.4, 105.2, 111.0, 114.0, 118.5, 119.1, 126.4, 128.0, 129.4, 129.7, 134.5, 140.2, 152.8, 160.0; IR (KBr) ν : 1516, 1258, 1021, 954, 824, 742, 701 cm^{-1} ; HRMS calcd for $\text{C}_{16}\text{H}_{15}\text{N}_2\text{O}$ [$\text{M}+\text{H}$] $^+$ 251.1179, found 251.1173.

3-(4-Fluorophenyl)-1-phenyl-1H-pyrazole (**3e**): Yellow solid (69 mg, 58%). m.p. 84~85 °C (lit.^[14] 87~88 °C); ^1H NMR (400 MHz, CDCl_3) δ : 6.64 (d, $J=2.8$ Hz, 1H), 7.01~7.06 (m, 2H), 7.21 (t, $J=7.2$ Hz, 1H), 7.37~7.41 (m, 2H), 7.68 (d, $J=8.0$ Hz, 2H), 7.79~7.82 (m, 2H), 7.86 (d, $J=2.4$ Hz, 1H); ^{13}C NMR (100 Hz, CDCl_3) δ : 104.8, 115.6 (d, $^2J_{\text{C-F}}=21.8$ Hz), 119.0, 122.9, 126.4, 127.5 (d, $^3J_{\text{C-F}}=8.0$ Hz), 128.1, 129.5, 140.2, 152.1, 162.8 (d, $^1J_{\text{C-F}}=245.1$ Hz); IR (KBr) ν : 1513, 1252, 1023, 953, 824, 742, 698 cm^{-1} ; HRMS calcd for $\text{C}_{15}\text{H}_{12}\text{FN}_2$ [$\text{M}+\text{H}$] $^+$ 239.0979, found 239.0971.

3-(4-Chlorophenyl)-1-phenyl-1H-pyrazole (**3f**): Yellow solid (80 mg, 63%). m.p. 113~115 °C (lit.^[13] 119~120 °C); ^1H NMR (400 MHz, CDCl_3) δ : 6.66 (d, $J=2.4$ Hz, 1H), 7.22 (t, $J=7.2$ Hz, 1H), 7.31 (d, $J=8.4$ Hz, 2H), 7.39 (t, $J=7.6$ Hz, 2H), 7.67 (t, $J=7.6$ Hz, 2H), 7.77 (d, $J=8.4$ Hz, 2H), 7.86 (d, $J=2.4$ Hz, 1H); ^{13}C NMR (100 Hz, CDCl_3) δ : 103.9, 118.1, 125.5, 126.1, 127.1, 127.8,

128.4, 130.6, 132.7, 139.1, 150.8; IR (KBr) ν : 1513, 1255, 1023, 956, 825, 746, 696 cm^{-1} ; HRMS calcd for $\text{C}_{15}\text{H}_{12}\text{ClN}_2$ $[\text{M}+\text{H}]^+$ 255.0684, found 255.0672.

1-Phenyl-3-(*p*-tolyl)-1*H*-pyrazole (**3g**): Yellow solid (83 mg, 71%). m.p. 90~92 °C (lit.^[13] 92~94 °C); ^1H NMR (400 MHz, CDCl_3) δ : 2.29 (s, 3H), 6.62 (d, $J=2.0$ Hz, 1H), 7.13~7.18 (m, 3H), 7.35 (t, $J=7.6$ Hz, 2H), 7.65~7.67 (m, 2H), 7.72 (d, $J=8.0$ Hz, 2H), 7.81 (d, $J=2.4$ Hz, 1H); ^{13}C NMR (100 Hz, CDCl_3) δ : 21.4, 104.9, 119.0, 125.8, 126.2, 127.9, 129.40, 129.44, 130.4, 137.9, 140.3, 153.0; IR (KBr) ν : 1516, 1252, 1026, 956, 825, 742, 702 cm^{-1} ; HRMS calcd for $\text{C}_{16}\text{H}_{15}\text{N}_2$ $[\text{M}+\text{H}]^+$ 235.1230, found 235.1232.

3-(4-Methoxyphenyl)-1-phenyl-1*H*-pyrazole (**3h**): Yellow solid (85 mg, 68%). m.p. 100~102 °C (lit.^[13] 101~103 °C); ^1H NMR (400 MHz, CDCl_3) δ : 3.77 (s, 3H), 6.62 (d, $J=2.4$ Hz, 1H), 6.88~6.91 (m, 2H), 7.19 (d, $J=7.6$ Hz, 1H), 7.36~7.40 (m, 2H), 7.67~7.69 (m, 2H), 7.75~7.79 (m, 2H), 7.85 (d, $J=2.4$ Hz, 1H); ^{13}C NMR (100 Hz, CDCl_3) δ : 55.3, 104.6, 114.1, 119.0, 126.0, 126.2, 127.1, 127.9, 129.4, 140.3, 152.8, 159.6; IR (KBr) ν : 1516, 1258, 1023, 953, 827, 742, 699 cm^{-1} ; HRMS calcd for $\text{C}_{16}\text{H}_{15}\text{N}_2\text{O}$ $[\text{M}+\text{H}]^+$ 251.1179, found 251.1188.

3-(Benzo[*d*][1,3]dioxol-5-yl)-1-phenyl-1*H*-pyrazole (**3i**): Yellow solid (80 mg, 61%). m.p. 113~115 °C (lit.^[14] 116~117 °C); ^1H NMR (400 MHz, CDCl_3) δ : 5.98 (s, 2H), 6.66 (d, $J=2.8$ Hz, 1H), 6.86 (d, $J=8.0$ Hz, 1H), 7.27 (t, $J=7.6$ Hz, 1H), 7.38 (dd, $J_1=8.4$ Hz, $J_2=1.6$ Hz, 1H), 7.43~7.47 (m, 3H), 7.73~7.75 (m, 2H), 7.91 (d, $J=2.4$ Hz, 1H); ^{13}C NMR (150 Hz, CDCl_3) δ : 101.1, 104.7, 106.5, 108.5, 118.9, 119.6, 126.3, 127.5, 127.9, 129.4, 140.2, 147.6, 148.1, 152.7; IR (KBr) ν : 1513, 1258, 1026, 953, 824, 745, 701 cm^{-1} ; HRMS calcd for $\text{C}_{16}\text{H}_{13}\text{N}_2\text{O}_2$ $[\text{M}+\text{H}]^+$ 265.0972, found 265.0968.

1-Phenyl-3-(thiophen-2-yl)-1*H*-pyrazole (**3j**)^[14]: Yellow solid (70 mg, 62%). m.p. 86~87 °C; ^1H NMR (400 MHz, CDCl_3) δ : 6.65 (d, $J=2.8$ Hz, 1H), 7.05~7.08 (m, 1H), 7.24~7.28 (m, 2H), 7.39~7.45 (m, 3H), 7.71~7.73 (m, 2H), 7.88 (d, $J=2.4$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 105.1, 119.1, 124.2, 125.0, 126.4, 127.5, 128.0, 129.5, 136.4, 140.0, 148.3; IR (KBr) ν : 1516, 1255, 1026, 956, 824, 745, 698 cm^{-1} ; HRMS calcd for $\text{C}_{13}\text{H}_{11}\text{N}_2\text{S}$ $[\text{M}+\text{H}]^+$ 227.0632, found 227.0634.

3-Ethyl-1-phenyl-1*H*-pyrazole (**3k**)^[13]: Yellow oil (38 mg, 44%). ^1H NMR (400 MHz, CDCl_3) δ : 1.31 (t, $J=7.6$ Hz, 3H), 2.76 (q, $J=7.6$ Hz, 2H), 6.27 (d, $J=2.4$ Hz, 1H), 7.24 (d, $J=8.0$ Hz, 1H), 7.39~7.44 (m, 2H), 7.64~7.67 (m, 2H), 7.82 (d, $J=2.4$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 13.9, 21.7, 106.0, 118.9, 125.9, 127.3, 129.4, 140.3, 156.6; HRMS calcd for $\text{C}_{11}\text{H}_{13}\text{N}_2$ $[\text{M}+\text{H}]^+$ 173.1073, found 173.1053.

3-Cyclohexyl-1-phenyl-1*H*-pyrazole (**3l**)^[15]: Yellow oil (46 mg, 41%). ^1H NMR (400 MHz, CDCl_3) δ : 1.18~1.45 (m, 5H), 1.59~1.77 (m, 3H), 1.95~1.98 (m, 2H), 2.65~2.71 (m, 1H), 6.18 (d, $J=2.4$ Hz, 1H), 7.15 (t, $J=7.6$ Hz, 1H), 7.33 (t, $J=7.6$ Hz, 2H), 7.57 (d, $J=7.6$ Hz, 2H), 7.72 (d, $J=2.4$ Hz, 1H); ^{13}C NMR (150 Hz, CDCl_3) δ : 25.1,

25.3, 32.2, 36.7, 103.6, 117.9, 124.8, 125.9, 128.3, 139.3, 159.2; IR (KBr) ν : 1516, 1252, 1023, 959, 824, 745, 701 cm^{-1} ; HRMS calcd for $\text{C}_{15}\text{H}_{19}\text{N}_2$ $[\text{M}+\text{H}]^+$ 227.1543, found 227.1538.

3-Phenyl-1-(*m*-tolyl)-1*H*-pyrazole (**3m**): Yellow solid (89 mg, 76%). m.p. 70~72 °C (lit.^[13] 71~73 °C); ^1H NMR (400 MHz, CDCl_3) δ : 2.36 (s, 3H), 6.68 (d, $J=2.4$ Hz, 1H), 7.03 (d, $J=7.6$ Hz, 1H), 7.24~7.28 (m, 2H), 7.35 (t, $J=7.6$ Hz, 2H), 7.46 (d, $J=8.0$ Hz, 1H), 7.55 (s, 1H), 7.84~7.86 (m, 3H); ^{13}C NMR (100 Hz, CDCl_3) δ : 20.5, 103.9, 115.1, 118.9, 124.8, 126.1, 126.96, 127.04, 127.6, 128.2, 132.2, 138.5, 139.2, 151.8; IR (KBr) ν : 1514, 1251, 1025, 954, 826, 702 cm^{-1} ; HRMS calcd for $\text{C}_{16}\text{H}_{15}\text{N}_2$ $[\text{M}+\text{H}]^+$ 235.1230, found 235.1228.

1-(4-Fluorophenyl)-3-phenyl-1*H*-pyrazole (**3n**): Yellow solid (64 mg, 54%). m.p. 86~88 °C (lit.^[13] 90~92 °C); ^1H NMR (600 MHz, CDCl_3) δ : 6.74 (s, 1H), 7.13 (t, $J=7.8$ Hz, 2H), 7.33 (t, $J=7.2$ Hz, 1H), 7.42 (t, $J=7.2$ Hz, 2H), 7.69~7.71 (m, 2H), 7.84 (s, 1H), 7.90 (t, $J=7.2$ Hz, 2H); ^{13}C NMR (150 Hz, CDCl_3) δ : 105.1, 116.2 (d, $^2J_{\text{C-F}}=23.0$ Hz), 120.8 (d, $^3J_{\text{C-F}}=8.7$ Hz), 125.9, 128.13, 128.15, 128.7, 133.0, 136.6 (d, $^4J_{\text{C-F}}=2.1$ Hz), 153.0, 161.1 (d, $^2J_{\text{C-F}}=243.9$ Hz); IR (KBr) ν : 1516, 1252, 1044, 1018, 824, 703 cm^{-1} ; HRMS calcd for $\text{C}_{15}\text{H}_{12}\text{FN}_2$ $[\text{M}+\text{H}]^+$ 239.0979, found 239.0971.

1-(4-Chlorophenyl)-3-phenyl-1*H*-pyrazole (**3o**): Yellow solid (76 mg, 60%). m.p. 125~127 °C (lit.^[13] 129~130 °C); ^1H NMR (400 MHz, CDCl_3) δ : 6.70 (d, $J=2.8$ Hz, 1H), 7.27 (t, $J=7.2$ Hz, 1H), 7.34~7.37 (m, 4H), 7.62~7.65 (m, 2H), 7.82~7.84 (m, 3H); ^{13}C NMR (150 Hz, CDCl_3) δ : 104.4, 119.1, 124.8, 126.9, 127.2, 127.7, 128.5, 130.7, 131.8, 137.7, 152.2; IR (KBr) ν : 1516, 1252, 1026, 956, 827, 743, 702 cm^{-1} ; HRMS calcd for $\text{C}_{15}\text{H}_{12}\text{ClN}_2$ $[\text{M}+\text{H}]^+$ 255.0684, found 255.0671.

3-Phenyl-1-(*p*-tolyl)-1*H*-pyrazole (**3p**): Yellow solid (81 mg, 69%). m.p. 107~109 °C (lit.^[13] 111~112 °C); ^1H NMR (600 MHz, CDCl_3) δ : 2.28 (s, 3H), 6.64 (s, 1H), 7.15 (t, $J=7.2$ Hz, 2H), 7.24~7.25 (m, 1H), 7.33 (t, $J=6.6$ Hz, 2H), 7.54 (d, $J=7.2$ Hz, 2H), 7.79~7.83 (m, 3H); ^{13}C NMR (150 Hz, CDCl_3) δ : 21.0, 104.8, 119.1, 125.9, 128.0, 128.7, 130.0, 133.3, 136.2, 138.1, 152.7; IR (KBr) ν : 1516, 1255, 1018, 954, 825, 743, 701 cm^{-1} ; HRMS calcd for $\text{C}_{16}\text{H}_{15}\text{N}_2$ $[\text{M}+\text{H}]^+$ 235.1230, found 235.1224.

1-(4-Methoxyphenyl)-3-phenyl-1*H*-pyrazole (**3q**): Yellow solid (78 mg, 62%). m.p. 100~102 °C (lit.^[14] 103~104 °C); ^1H NMR (400 MHz, CDCl_3) δ : 3.73 (s, 3H), 6.63 (d, $J=2.4$ Hz, 1H), 6.87 (d, $J=9.2$ Hz, 2H), 7.23 (t, $J=7.2$ Hz, 1H), 7.33 (t, $J=7.2$ Hz, 2H), 7.56 (d, $J=8.8$ Hz, 2H), 7.73 (d, $J=2.4$ Hz, 1H), 7.82 (d, $J=7.6$ Hz, 2H); ^{13}C NMR (100 Hz, CDCl_3) δ : 55.6, 104.6, 114.5, 120.8, 125.8, 127.9, 128.1, 128.7, 133.3, 134.1, 152.6, 158.2; IR (KBr) ν : 1669, 1393, 1073, 758, 692 cm^{-1} ; HRMS calcd for $\text{C}_{16}\text{H}_{15}\text{N}_2\text{O}$ $[\text{M}+\text{H}]^+$ 251.1179, found 251.1170.

1-Methyl-3-phenyl-1*H*-pyrazole (**3r**)^[16]: Yellow oil (36 mg, 45%). ^1H NMR (600 MHz, CDCl_3) δ : 3.85 (s, 3H), 6.45 (s, 1H), 7.21~7.22 (m, 1H), 7.28~7.32 (m, 3H), 7.71 (d, $J=7.2$ Hz, 2H); ^{13}C NMR (100 Hz, CDCl_3) δ :

39.0, 102.8, 125.5, 127.5, 128.6, 131.4, 133.6, 151.6; IR (KBr) ν : 1651, 1593, 1528, 1293, 960, 795 cm^{-1} ; HRMS calcd for $\text{C}_{10}\text{H}_{11}\text{N}_2$ $[\text{M} + \text{H}]^+$ 159.0917, found 159.0909.

1-Isopropyl-3-phenyl-1*H*-pyrazole (**3s**)^[17]: Yellow oil (39 mg, 42%). ^1H NMR (400 MHz, CDCl_3) δ : 1.52 (d, $J=6.8$ Hz, 6H), 4.47~4.58 (m, 1H), 6.52 (d, $J=2.4$ Hz, 1H), 7.24~7.28 (m, 1H), 7.35~7.41 (m, 3H), 7.79~7.82 (m, 2H); ^{13}C NMR (150 MHz, CDCl_3) δ : 23.1, 53.9, 102.3, 125.6, 127.37, 127.40, 128.6, 134.0, 150.8; HRMS calcd for $\text{C}_{12}\text{H}_{15}\text{N}_2$ $[\text{M} + \text{H}]^+$ 187.1230, found 187.1216.

4.2.2 Typical procedure for the synthesis of **5a**

To a reaction tube equipped with a stir bar were added propiophenone (**1a**, 81 mg, 0.6 mmol), PhCl (3 mL), $\text{Cu}(\text{OAc})_2$ (18 mg, 0.1 mmol), bpy (8 mg, 0.05 mmol) and TEMPO (78 mg, 0.5 mmol). The mixture was stirred at 120 $^\circ\text{C}$ under N_2 for 10 h. To the resulting mixture was added 1-benzylidene-2-phenyl hydrazine (**4a**, 98 mg, 0.5 mmol). It was then stirred at 120 $^\circ\text{C}$ under air for 5 h. Upon completion, the mixture was diluted with DCM (15 mL), and washed with water (5 mL) and brine (5 mL). The organic layer was dried over anhydrous Na_2SO_4 , and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel with petroleum ether/ethyl acetate ($V:V=20:1$) as the eluent to give **5a**. **5b**~**5dd** were obtained in a similar manner.

(1,3-Diphenyl-1*H*-pyrazol-4-yl)(phenyl)methanone (**5a**)^[10b]: White solid (102 mg, 63%). m.p. 141~142 $^\circ\text{C}$. ^1H NMR (400 MHz, CDCl_3) δ : 7.32~7.42 (m, 6H), 7.47~7.54 (m, 3H), 7.72~7.74 (m, 2H), 7.79 (d, $J=8.0$ Hz, 2H), 7.83~7.85 (m, 2H), 8.28 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 119.6, 121.2, 127.5, 128.2, 128.4, 128.6, 128.9, 129.5, 129.7, 132.1, 132.3, 132.6, 138.9, 139.3, 154.0, 190.0; IR (KBr) ν : 1645, 1534, 1246, 1038, 799, 764, 753 cm^{-1} ; MS m/z : 325 $[\text{M} + \text{H}]^+$.

(1,3-Diphenyl-1*H*-pyrazol-4-yl)(3-methoxyphenyl)methanone (**5b**)^[10b]: White solid (103 mg, 58%), m.p. 133~134 $^\circ\text{C}$. ^1H NMR (400 MHz, CDCl_3) δ : 3.79 (s, 3H), 7.05~7.08 (m, 1H), 7.28~7.43 (m, 7H), 7.48~7.52 (m, 2H), 7.72~7.75 (m, 2H), 7.77~7.80 (m, 2H), 8.30 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 55.4, 113.7, 119.3, 119.6, 121.2, 122.3, 127.6, 128.2, 128.6, 128.9, 129.4, 129.7, 132.1, 132.3, 139.3, 140.2, 154.0, 159.6, 189.7; IR (KBr) ν : 1634, 1604, 1522, 1305, 1158, 896, 839, 752, 692 cm^{-1} ; MS m/z : 355 $[\text{M} + \text{H}]^+$.

(1,3-Diphenyl-1*H*-pyrazol-4-yl)(4-fluorophenyl)methanone (**5c**)^[10b]: White solid (118 mg, 69%), m.p. 141~143 $^\circ\text{C}$. ^1H NMR (400 MHz, CDCl_3) δ : 7.01~7.07 (m, 2H), 7.30~7.38 (m, 4H), 7.49 (t, $J=8.0$ Hz, 2H), 7.65~7.70 (m, 2H), 7.78~7.80 (m, 2H), 7.82~7.87 (m, 2H), 8.29 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 115.5 (d, $^2J_{\text{C-F}}=21.9$ Hz), 119.6, 121.2, 127.6, 128.2, 128.7, 128.9, 129.7, 132.0, 132.07, 132.14, 135.0 (d, $^3J_{\text{C-F}}=3.3$ Hz), 139.2, 153.8, 165.5 (d, $^1J_{\text{C-F}}=252.6$ Hz), 188.6; IR (KBr) ν : 1604, 1161, 750, 689 cm^{-1} ; MS m/z : 343 $[\text{M} + \text{H}]^+$.

(1,3-Diphenyl-1*H*-pyrazol-4-yl)(4-(trifluoromethyl)phenyl)methanone (**5d**): White solid (90 mg, 53%), m.p. 146~148 $^\circ\text{C}$ (lit.^[10b] 159~161 $^\circ\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 2.40 (s, 3H), 7.21 (d, $J=8.0$ Hz, 2H), 7.33~7.38 (m, 4H), 7.49 (t, $J=7.6$ Hz, 2H), 7.73~7.80 (m, 6H), 8.26 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 21.7, 119.6, 121.4, 127.5, 128.2, 128.6, 128.9, 129.1, 129.6, 129.7, 132.0, 132.2, 136.3, 139.3, 143.5, 153.9, 189.8; IR (KBr) ν : 1648, 1534, 1326, 1106, 1059, 859, 768, 712, 683 cm^{-1} ; MS m/z : 339 $[\text{M} + \text{H}]^+$.

(1,3-Diphenyl-1*H*-pyrazol-4-yl)(4-(trifluoromethyl)phenyl)methanone (**5e**): White solid (115 mg, 59%), m.p. 163~165 $^\circ\text{C}$ (lit.^[10b] 163~165 $^\circ\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 7.31~7.38 (m, 4H), 7.48 (t, $J=7.6$ Hz, 2H), 7.62 (d, $J=8.4$ Hz, 2H), 7.67~7.69 (m, 2H), 7.77~7.79 (m, 2H), 7.87 (d, $J=8.0$ Hz, 2H), 8.29 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 119.6, 120.9, 123.7 (q, $^1J_{\text{C-F}}=271.2$ Hz), 125.4 (q, $^3J_{\text{C-F}}=3.3$ Hz), 127.8, 128.2, 128.9, 129.0, 129.6, 129.7, 131.8, 132.6, 133.8 (q, $^2J_{\text{C-F}}=31.8$ Hz), 139.1, 141.9, 154.2, 188.8; IR (KBr) ν : 1640, 1516, 1223, 827, 745, 706 cm^{-1} ; MS m/z : 393 $[\text{M} + \text{H}]^+$.

(1,3-Diphenyl-1*H*-pyrazol-4-yl)(thiophen-2-yl)methanone (**5f**)^[10b]: White solid (79 mg, 48%), m.p. 150~152 $^\circ\text{C}$. ^1H NMR (400 MHz, CDCl_3) δ : 7.07~7.09 (m, 1H), 7.35~7.41 (m, 4H), 7.51 (t, $J=8.4$ Hz, 2H), 7.62~7.63 (m, 1H), 7.66 (dd, $J_1=4.8$ Hz, $J_2=0.8$ Hz, 1H), 7.78~7.82 (m, 4H), 8.40 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 119.6, 121.1, 127.6, 128.0, 128.3, 128.7, 128.8, 129.7, 131.2, 132.0, 133.9, 134.0, 139.3, 145.0, 153.4, 181.4; IR (KBr) ν : 1634, 1581, 1229, 1070, 897, 742, 701, 683 cm^{-1} ; MS m/z : 331 $[\text{M} + \text{H}]^+$.

(1,3-Diphenyl-1*H*-pyrazol-4-yl)(pyridin-3-yl)methanone (**5g**)^[10b]: White solid (93 mg, 57%). m.p. 131~133 $^\circ\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.30~7.40 (m, 5H), 7.50 (t, $J=8.0$ Hz, 2H), 7.66~7.69 (m, 2H), 7.79 (d, $J=7.6$ Hz, 2H), 8.06~8.08 (m, 1H), 8.36 (s, 1H), 8.71 (d, $J=3.2$ Hz, 1H), 9.00 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 119.7, 120.9, 123.4, 127.8, 128.3, 128.9, 129.1, 129.7, 131.8, 132.5, 134.3, 136.6, 139.1, 150.4, 152.9, 154.1, 188.1; IR (KBr) ν : 1528, 1455, 1208, 1164, 903, 752, 715, 661 cm^{-1} ; MS m/z : 326 $[\text{M} + \text{H}]^+$.

Cyclohexyl(1,3-diphenyl-1*H*-pyrazol-4-yl)methanone (**5h**): White solid (84 mg, 51%). m.p. 110~111 $^\circ\text{C}$ (lit.^[10b] 113~114 $^\circ\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 1.12~1.23 (m, 3H), 1.44~1.53 (m, 2H), 1.65~1.66 (m, 1H), 1.76~1.86 (m, 4H), 2.77~2.84 (m, 1H), 7.36 (t, $J=7.6$ Hz, 1H), 7.41~7.47 (m, 3H), 7.49 (t, $J=7.6$ Hz, 2H), 7.71~7.74 (m, 2H), 7.78 (d, $J=7.6$ Hz, 2H), 8.42 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 25.78, 25.84, 29.4, 48.9, 119.7, 121.7, 127.6, 128.1, 128.8, 129.3, 129.6, 130.9, 132.7, 139.3, 153.9, 199.0; IR (KBr) ν : 1634, 1458, 1205, 1109, 902, 762, 712, 662 cm^{-1} ; MS m/z : 331 $[\text{M} + \text{H}]^+$.

(3-(2-Fluorophenyl)-1-phenyl-1*H*-pyrazol-4-yl)(phenyl)methanone (**5i**): White solid (72 mg, 42%), m.p. 109~110 $^\circ\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 6.95~7.00 (m, 1H), 7.20 (td, $J_1=7.6$ Hz, $J_2=1.2$ Hz, 1H), 7.29~7.40 (m, 4H), 7.46~7.52 (m, 3H), 7.69 (td, $J_1=7.6$ Hz, $J_2=2.0$ Hz, 1H), 7.77~7.79 (m, 2H), 7.82~7.84 (m, 2H), 8.32 (s,

1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 115.6 (d, $^2J_{\text{C-F}}=21.9$ Hz), 119.7, 120.8 (d, $^2J_{\text{C-F}}=14.3$ Hz), 122.6, 124.2 (d, $^3J_{\text{C-F}}=3.3$ Hz), 127.7, 128.3, 129.3, 129.7, 130.6 (d, $^3J_{\text{C-F}}=8.7$ Hz), 130.8 (d, $^4J_{\text{C-F}}=2.3$ Hz), 131.4, 132.5, 138.5, 139.3, 148.8, 160.0 (d, $^1J_{\text{C-F}}=247.2$ Hz), 189.5; IR (KBr) ν : 1519, 1452, 1223, 897, 822, 739, 705, 684 cm^{-1} ; HRMS calcd for $\text{C}_{22}\text{H}_{15}\text{FN}_2\text{O}_2$ [M+Na] $^+$ 365.1061, found 365.1041.

(3-(3-Fluorophenyl)-1-phenyl-1H-pyrazol-4-yl)(phenyl)methanone (**5j**): White solid (94 mg, 55%), m.p. 132~134 $^{\circ}\text{C}$ (lit.^[10b] 129~131 $^{\circ}\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 7.01~7.06 (m, 1H), 7.27~7.33 (m, 1H), 7.38 (t, $J=7.2$ Hz, 1H), 7.43 (t, $J=7.6$ Hz, 2H), 7.48~7.57 (m, 5H), 7.78 (d, $J=7.6$ Hz, 2H), 7.85 (d, $J=7.2$ Hz, 2H), 8.28 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 115.5 (d, $^2J_{\text{C-F}}=20.7$ Hz), 115.8 (d, $^2J_{\text{C-F}}=23.0$ Hz), 119.6, 121.3, 124.7 (d, $^4J_{\text{C-F}}=2.3$ Hz), 127.7, 128.5, 129.5, 129.6, 129.7 (d, $^3J_{\text{C-F}}=4.4$ Hz), 132.5, 132.8, 134.2 (d, $^3J_{\text{C-F}}=8.7$ Hz), 138.9, 139.1, 152.7, 162.6 (d, $^1J_{\text{C-F}}=243.9$ Hz), 189.8; IR (KBr) ν : 1634, 1516, 1288, 900, 827, 756, 708, 689 cm^{-1} ; MS m/z : 343 [M+H] $^+$.

(3-(4-Fluorophenyl)-1-phenyl-1H-pyrazol-4-yl)(phenyl)methanone (**5k**): White solid (118 mg, 69%), m.p. 140~142 $^{\circ}\text{C}$ (lit.^[10b] 135~136 $^{\circ}\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 7.02~7.06 (m, 2H), 7.37 (t, $J=7.6$ Hz, 1H), 7.42 (t, $J=7.6$ Hz, 2H), 7.49 (t, $J=7.6$ Hz, 2H), 7.55 (t, $J=7.6$ Hz, 1H), 7.75~7.78 (m, 4H), 7.82~7.84 (m, 2H), 8.27 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 115.2 (d, $^2J_{\text{C-F}}=21.9$ Hz), 119.6, 121.0, 127.7, 128.3 (d, $^4J_{\text{C-F}}=3.3$ Hz), 128.5, 129.4, 129.7, 130.8 (d, $^3J_{\text{C-F}}=8.9$ Hz), 132.6, 132.7, 139.0, 139.2, 153.1, 163.1 (d, $^1J_{\text{C-F}}=246.0$ Hz), 189.9; IR (KBr) ν : 1637, 1522, 1223, 821, 742 cm^{-1} ; MS m/z : 343 [M+H] $^+$.

(3-(4-Chlorophenyl)-1-phenyl-1H-pyrazol-4-yl)(phenyl)methanone (**5l**): White solid (131 mg, 73%), m.p. 147~148 $^{\circ}\text{C}$ (lit.^[10b] 147~149 $^{\circ}\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 7.31~7.39 (m, 3H), 7.43 (t, $J=7.6$ Hz, 2H), 7.49 (t, $J=7.6$ Hz, 2H), 7.56 (t, $J=7.2$ Hz, 1H), 7.72~7.78 (m, 4H), 7.84 (d, $J=8.4$ Hz, 2H), 8.26 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 119.6, 121.1, 127.7, 128.4, 128.5, 129.4, 129.7, 130.3, 130.6, 132.6, 132.8, 134.7, 138.9, 139.1, 152.9, 189.8; IR (KBr) ν : 1643, 1528, 1452, 1223, 742, 709 cm^{-1} ; MS m/z : 359 [M+H] $^+$.

Phenyl(1-phenyl-3-(*p*-tolyl)-1H-pyrazol-4-yl)methanone (**5m**): White solid (125 mg, 74%), m.p. 142~144 $^{\circ}\text{C}$ (lit.^[10b] 146~148 $^{\circ}\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 2.35 (s, 3H), 7.15 (d, $J=8.0$ Hz, 2H), 7.35 (t, $J=7.2$ Hz, 1H), 7.41 (t, $J=7.6$ Hz, 2H), 7.46~7.55 (m, 3H), 7.64 (d, $J=8.0$ Hz, 2H), 7.76~7.79 (m, 2H), 7.83~7.86 (m, 2H), 8.25 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 21.4, 119.6, 121.1, 127.4, 128.4, 128.8, 128.9, 129.2, 129.5, 129.6, 132.3, 132.6, 138.5, 139.0, 139.3, 154.1, 190.0; IR (KBr) ν : 1645, 1311, 1123, 850, 767, 718, 692 cm^{-1} ; MS m/z : 339 [M+H] $^+$.

Phenyl(1-phenyl-3-(4-(trifluoromethyl)phenyl)-1H-pyrazol-4-yl)methanone (**5n**): White solid (122 mg, 62%), m.p. 137~139 $^{\circ}\text{C}$ (lit.^[10b] 141~143 $^{\circ}\text{C}$); ^1H NMR (400

MHz, CDCl_3) δ : 7.38 (t, $J=7.2$ Hz, 1H), 7.44 (t, $J=7.6$ Hz, 2H), 7.51 (t, $J=7.6$ Hz, 2H), 7.57 (t, $J=7.2$ Hz, 1H), 7.61 (d, $J=8.0$ Hz, 2H), 7.78 (d, $J=8.0$ Hz, 2H), 7.85 (d, $J=7.2$ Hz, 2H), 7.91 (d, $J=8.0$ Hz, 2H), 8.28 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 119.7, 121.3, 122.3 (q, $^1J_{\text{C-F}}=286.5$ Hz), 125.1 (q, $^3J_{\text{C-F}}=3.3$ Hz), 127.9, 128.6, 129.2, 129.4, 129.7, 130.5 (q, $^2J_{\text{C-F}}=32.9$ Hz), 132.7, 132.9, 135.7, 138.8, 139.1, 152.7, 189.7; IR (KBr) ν : 1643, 1522, 1311, 1123, 1067, 897, 765, 717 cm^{-1} ; MS m/z : 393 [M+H] $^+$.

(3-(4-Methoxyphenyl)-1-phenyl-1H-pyrazol-4-yl)(phenyl)methanone (**5o**): White solid (96 mg, 54%), m.p. 129~131 $^{\circ}\text{C}$ (lit.^[10b] 132~134 $^{\circ}\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 3.81 (s, 3H), 6.87~6.90 (m, 2H), 7.35 (t, $J=7.2$ Hz, 1H), 7.41 (t, $J=8.0$ Hz, 2H), 7.48 (t, $J=7.6$ Hz, 2H), 7.53 (t, $J=7.6$ Hz, 1H), 7.70~7.74 (m, 2H), 7.77 (d, $J=7.6$ Hz, 2H), 7.83~7.85 (m, 2H), 8.24 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 55.3, 113.6, 119.5, 120.9, 124.6, 127.4, 128.4, 129.5, 129.6, 130.3, 132.4, 132.6, 139.1, 139.3, 153.8, 160.0, 190.1; IR (KBr) ν : 1640, 1519, 1226, 1021, 896, 834, 746, 706 cm^{-1} ; MS m/z : 355 [M+H] $^+$.

Phenyl(1-phenyl-3-(thiophen-2-yl)-1H-pyrazol-4-yl)methanone (**5p**): White solid (102 mg, 62%), m.p. 147~148 $^{\circ}\text{C}$ (lit.^[10b] 137~139 $^{\circ}\text{C}$); ^1H NMR (600 MHz, CDCl_3) δ : 7.04~7.07 (m, 1H), 7.33~7.37 (m, 2H), 7.45~7.49 (m, 4H), 7.58 (t, $J=7.2$ Hz, 1H), 7.75 (d, $J=7.6$ Hz, 2H), 7.86~7.88 (m, 2H), 7.93~7.95 (m, 1H), 8.18 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 119.6, 120.4, 126.8, 127.5, 127.7, 128.6, 128.9, 129.4, 129.7, 132.6, 132.9, 134.1, 139.0, 139.5, 148.3, 189.5; IR (KBr) ν : 1648, 1516, 1276, 1211, 884, 844, 681 cm^{-1} ; MS m/z : 331 [M+H] $^+$.

(1-(4-Chlorophenyl)-3-phenyl-1H-pyrazol-4-yl)(phenyl)methanone (**5q**): White solid (113 mg, 63%), m.p. 163~165 $^{\circ}\text{C}$ (lit.^[10b] 165~167 $^{\circ}\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 7.31~7.35 (m, 3H), 7.39 (t, $J=7.6$ Hz, 2H), 7.44~7.47 (m, 2H), 7.52 (t, $J=7.2$ Hz, 1H), 7.69~7.75 (m, 4H), 7.81~7.83 (m, 2H), 8.25 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 120.7, 121.6, 128.2, 128.4, 128.8, 128.9, 129.5, 129.8, 131.9, 132.1, 132.7, 133.1, 137.8, 138.7, 154.2, 189.9; IR (KBr) ν : 1628, 1493, 1440, 1293, 1211, 897, 830, 730, 699 cm^{-1} ; MS m/z : 359 [M+H] $^+$.

(1-(4-Nitrophenyl)-3-phenyl-1H-pyrazol-4-yl)(phenyl)methanone (**5r**): White solid (116 mg, 63%), m.p. 143~145 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.15 (s, 1H), 7.38~7.48 (m, 3H), 7.53~7.57 (m, 2H), 7.66~7.71 (m, 3H), 7.87~7.90 (m, 2H), 8.00~8.02 (m, 2H), 8.27~8.30 (m, 2H); ^{13}C NMR (150 MHz, CDCl_3) δ : 112.4, 124.5, 125.0, 126.0, 128.9, 129.0, 129.1, 129.9, 131.4, 134.2, 136.8, 140.8, 144.8, 146.7, 152.7, 185.3; HRMS calcd for $\text{C}_{22}\text{H}_{15}\text{N}_3\text{O}_3\text{Na}$ [M+Na] $^+$ 392.1006, found 392.0988.

Phenyl(3-phenyl-1-(*p*-tolyl)-1H-pyrazol-4-yl)methanone (**5s**): White solid (122 mg, 72%), m.p. 148~149 $^{\circ}\text{C}$ (lit.^[10b] 148~150 $^{\circ}\text{C}$); ^1H NMR (400 MHz, CDCl_3) δ : 2.41 (s, 3H), 7.28~7.35 (m, 5H), 7.40 (t, $J=7.6$ Hz, 2H), 7.52 (t, $J=7.6$ Hz, 1H), 7.66 (d, $J=8.4$ Hz, 2H), 7.71~7.74 (m, 2H), 7.82~7.84 (m, 2H), 8.25 (s, 1H); ^{13}C NMR

(150 MHz, CDCl_3) δ : 21.0, 119.5, 121.0, 128.1, 128.3, 128.5, 128.9, 129.5, 130.1, 132.2, 132.3, 132.6, 137.0, 137.5, 139.0, 153.9, 190.1; IR (KBr) ν : 1634, 1522, 1220, 897, 821, 770, 736, 686 cm^{-1} ; MS m/z : 339 $[\text{M} + \text{H}]^+$.

Phenyl(3-phenyl-1-(pyridin-2-yl)-1*H*-pyrazol-4-yl)-methanone (**5t**): White solid (94 mg, 58%). m.p. 145~147 °C. ^1H NMR (400 MHz, CDCl_3) δ : 7.00 (s, 1H), 7.07~7.11 (m, 1H), 7.37~7.48 (m, 5H), 7.52~7.56 (m, 1H), 7.77~7.82 (m, 1H), 7.89~7.99 (m, 5H), 8.08~8.10 (m, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 108.4, 115.0, 122.0, 126.1, 128.5, 128.7, 128.8, 129.4, 132.1, 133.2, 137.4, 138.5, 141.4, 147.5, 151.2, 152.7, 187.6; HRMS calcd for $\text{C}_{21}\text{H}_{16}\text{N}_3\text{O}$ $[\text{M} + \text{H}]^+$ 326.1288, found 326.1284.

(1-Methyl-3-phenyl-1*H*-pyrazol-4-yl)(phenyl)methanone (**5u**): White solid (60 mg, 46%). m.p. 88~90 °C (lit.^[10b] 159~161 °C); ^1H NMR (600 MHz, CDCl_3) δ : 3.98 (s, 3H), 7.29~7.31 (m, 3H), 7.36 (t, $J=7.6$ Hz, 2H), 7.48 (t, $J=7.6$ Hz, 1H), 7.61~7.64 (m, 2H), 7.75~7.77 (m, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 39.3, 119.6, 128.1, 128.2, 128.3, 128.8, 129.4, 132.3, 132.4, 136.1, 139.2, 153.2, 189.9; IR (KBr) ν : 1640, 1522, 1446, 897, 727, 692, 671 cm^{-1} ; MS m/z : 263 $[\text{M} + \text{H}]^+$.

(1-Benzyl-3-phenyl-1*H*-pyrazol-4-yl)(phenyl)methanone (**5v**): White solid (88 mg, 52%). m.p. 100~101 °C (lit.^[10b] 102~103 °C); ^1H NMR (400 MHz, CDCl_3) δ : 5.37 (s, 2H), 7.28~7.39 (m, 10H), 7.46 (t, $J=7.2$ Hz, 1H), 7.63~7.65 (m, 2H), 7.73~7.74 (m, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 56.5, 119.9, 128.06, 128.10, 128.2, 128.3, 128.6, 128.9, 129.4, 132.36, 132.40, 135.2, 135.3, 139.0, 153.2, 190.0; IR (KBr) ν : 1645, 1522, 1437, 1290, 1158, 894, 774, 730, 693 cm^{-1} ; MS m/z : 339 $[\text{M} + \text{H}]^+$.

(1-Isopropyl-3-phenyl-1*H*-pyrazol-4-yl)(phenyl)methanone (**5w**): White solid (68 mg, 47%). m.p. 100~101 °C (lit.^[10b] 98~99 °C); ^1H NMR (400 MHz, CDCl_3) δ : 1.52 (d, $J=6.8$ Hz, 6H), 4.45~4.55 (m, 1H), 7.19~7.23 (m, 3H), 7.27~7.30 (m, 2H), 7.38~7.42 (m, 1H), 7.54~7.56 (m, 2H), 7.67~7.69 (m, 2H), 7.75 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 21.8, 53.4, 118.0, 127.0, 127.1, 127.2, 127.9, 128.3, 131.2, 131.5, 131.7, 138.2, 151.6, 189.2; IR (KBr) ν : 1634, 1513, 1452, 1296, 1076, 900, 780, 743, 717, 671 cm^{-1} ; MS m/z : 291 $[\text{M} + \text{H}]^+$.

(1-(*tert*-Butyl)-3-phenyl-1*H*-pyrazol-4-yl)(phenyl)methanone (**5x**): White solid (61 mg, 40%). m.p. 108~110 °C; ^1H NMR (600 MHz, CDCl_3) δ : 1.66 (s, 9H), 7.26~7.28 (m, 3H), 7.35 (t, $J=8.0$ Hz, 2H), 7.47 (t, $J=7.6$ Hz, 1H), 7.62~7.64 (m, 2H), 7.74~7.76 (m, 2H), 7.89 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 29.7, 59.5, 118.8, 128.0, 128.1, 128.2, 129.0, 129.4, 131.9, 132.1, 133.0, 139.3, 152.3, 190.5; HRMS calcd for $\text{C}_{20}\text{H}_{21}\text{N}_2\text{O}$ $[\text{M} + \text{H}]^+$ 305.1648, found 305.1630.

(1-Cyclohexyl-3-phenyl-1*H*-pyrazol-4-yl)(phenyl)methanone (**5y**): White solid (74 mg, 45%). m.p. 79~81 °C; ^1H NMR (600 MHz, CDCl_3) δ : 1.17~1.23 (m, 1H), 1.31~1.40 (m, 2H), 1.63~1.72 (m, 3H), 1.81~1.86 (m, 2H), 2.15~2.19 (m, 2H), 4.06~4.12 (m, 1H), 7.17~7.23 (m, 3H), 7.27 (t, $J=7.6$ Hz, 2H), 7.37~7.41 (m, 1H), 7.54~7.56 (m, 2H), 7.66~7.68 (m, 2H), 7.72 (s, 1H); ^{13}C

NMR (150 MHz, CDCl_3) δ : 25.2, 25.3, 33.4, 61.8, 118.9, 128.0, 128.1, 128.2, 128.9, 129.4, 132.2, 132.7, 139.3, 152.5, 190.2; HRMS calcd for $\text{C}_{22}\text{H}_{23}\text{N}_2\text{O}$ $[\text{M} + \text{H}]^+$ 331.1805, found 331.1826.

(3-Methoxyphenyl)(3-(3-methoxyphenyl)-1-phenyl-1*H*-pyrazol-4-yl)methanone (**5z**): White solid (117 mg, 61%). m.p. 166~167 °C; ^1H NMR (600 MHz, CDCl_3) δ : 3.78 (s, 3H), 3.79 (s, 3H), 6.88 (dd, $J_1=7.8$ Hz, $J_2=1.8$ Hz, 1H), 7.07 (dd, $J_1=8.4$ Hz, $J_2=1.8$ Hz, 1H), 7.24 (d, $J=7.2$ Hz, 1H), 7.29~7.31 (m, 3H), 7.37 (t, $J=7.2$ Hz, 2H), 7.41 (d, $J=7.8$ Hz, 1H), 7.50 (t, $J=7.8$ Hz, 2H), 7.79 (d, $J=7.8$ Hz, 2H), 8.31 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 55.3, 55.4, 113.7, 114.0, 115.0, 119.4, 119.6, 121.4, 121.5, 122.3, 127.6, 129.2, 129.4, 129.6, 132.3, 133.4, 139.2, 140.2, 153.7, 159.4, 159.6, 189.8; IR (KBr) ν : 1637, 1508, 1437, 1088, 900, 849, 752, 690 cm^{-1} ; HRMS calcd for $\text{C}_{24}\text{H}_{20}\text{N}_2\text{NaO}_3$ $[\text{M} + \text{Na}]^+$ 407.1366, found 407.1348.

(4-Fluorophenyl)(3-(4-fluorophenyl)-1-phenyl-1*H*-pyrazol-4-yl)methanone (**5aa**): White solid (117 mg, 65%). m.p. 135~136 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.01~7.10 (m, 4H), 7.37 (t, $J=7.6$ Hz, 1H), 7.50 (t, $J=7.6$ Hz, 2H), 7.70~7.74 (m, 2H), 7.77 (d, $J=7.6$ Hz, 2H), 7.84~7.87 (m, 2H), 8.26 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 115.2 (d, $^2J_{\text{C-F}}=21.8$ Hz), 115.6 (d, $^2J_{\text{C-F}}=21.9$ Hz), 119.6, 120.9, 127.7, 128.1 (d, $^4J_{\text{C-F}}=3.6$ Hz), 129.7, 130.8 (d, $^3J_{\text{C-F}}=8.7$ Hz), 132.0, 132.1 (d, $^3J_{\text{C-F}}=12.4$ Hz), 135.1 (d, $^4J_{\text{C-F}}=2.9$ Hz), 139.1, 152.9, 163.1 (d, $^1J_{\text{C-F}}=237.1$ Hz), 165.6 (d, $^1J_{\text{C-F}}=243.7$ Hz), 188.4; IR (KBr) ν : 1643, 1505, 1443, 1091, 896, 839, 755, 686 cm^{-1} ; HRMS calcd for $\text{C}_{22}\text{H}_{14}\text{F}_2\text{N}_2\text{NaO}$ $[\text{M} + \text{Na}]^+$ 383.0966, found 383.0951.

(4-Chlorophenyl)(3-(4-chlorophenyl)-1-phenyl-1*H*-pyrazol-4-yl)methanone (**5bb**): White solid (125 mg, 64%). m.p. 142~143 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.34 (d, $J=8.4$ Hz, 2H), 7.37~7.42 (m, 3H), 7.50 (t, $J=7.8$ Hz, 2H), 7.70 (d, $J=9.0$ Hz, 2H), 7.76~7.79 (m, 4H), 8.25 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 119.6, 120.8, 127.8, 128.5, 128.8, 129.7, 130.2, 130.4, 130.8, 132.4, 134.8, 137.2, 139.1, 139.2, 152.8, 188.5; IR (KBr) ν : 1640, 1584, 1505, 1446, 1091, 897, 837, 753, 686 cm^{-1} ; HRMS calcd for $\text{C}_{22}\text{H}_{14}\text{Cl}_2\text{N}_2\text{NaO}$ $[\text{M} + \text{Na}]^+$ 415.0375, found 415.0364.

(1-Phenyl-3-(*p*-tolyl)-1*H*-pyrazol-4-yl)(*p*-tolyl)methanone (**5cc**): White solid (120 mg, 68%). m.p. 137~138 °C; ^1H NMR (400 MHz, CDCl_3) δ : 2.35 (s, 3H), 2.41 (s, 3H), 7.16 (d, $J=7.6$ Hz, 2H), 7.22 (d, $J=8.0$ Hz, 2H), 7.35 (t, $J=7.2$ Hz, 1H), 7.49 (d, $J=7.6$ Hz, 2H), 7.65 (d, $J=8.0$ Hz, 2H), 7.76~7.78 (m, 4H), 8.23 (s, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 21.3, 21.7, 119.5, 121.2, 127.4, 128.7, 128.9, 129.1, 129.3, 129.6, 129.7, 132.0, 136.4, 138.5, 139.4, 143.4, 153.9, 189.8; IR (KBr) ν : 1631, 1513, 1446, 1223, 1158, 902, 825, 753, 690 cm^{-1} ; HRMS calcd for $\text{C}_{24}\text{H}_{20}\text{N}_2\text{NaO}$ $[\text{M} + \text{Na}]^+$ 375.1468, found 375.1481.

(1-Phenyl-3-(4-(trifluoromethyl)phenyl)-1*H*-pyrazol-4-yl)(4-(trifluoromethyl)phenyl)methanone (**5dd**): White solid (116 mg, 50%). m.p. 120~121 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.32 (t, $J=7.2$ Hz, 1H), 7.43 (t, $J=7.6$ Hz, 2H), 7.54 (d, $J=8.0$ Hz, 2H), 7.61 (d, $J=8.0$ Hz, 2H),

7.69 (d, $J=7.6$ Hz, 2H), 7.79~7.85 (m, 4H), 8.20 (s, 1H). ^{13}C NMR (150 MHz, CDCl_3) δ : 119.7, 120.9, 123.6 (q, $^1J_{\text{C-F}}=271.4$ Hz), 124.1 (q, $^1J_{\text{C-F}}=270.2$ Hz), 125.2 (q, $^3J_{\text{C-F}}=3.3$ Hz), 125.6 (q, $^3J_{\text{C-F}}=3.3$ Hz), 128.1, 129.3, 129.5, 129.8, 130.7 (q, $^2J_{\text{C-F}}=32.9$ Hz), 132.9, 134.1 (q, $^2J_{\text{C-F}}=32.7$ Hz), 135.4, 138.9, 141.9, 152.9, 188.4; IR (KBr) ν : 1642, 1596, 1506, 1316, 1127, 1062, 900, 842, 688 cm^{-1} ; HRMS calcd for $\text{C}_{24}\text{H}_{14}\text{F}_6\text{N}_2\text{NaO}$ $[\text{M}+\text{Na}]^+$ 483.0903, found 483.0878.

4.2.3 Control experiments

To a reaction tube equipped with a stir bar were added 1-phenylprop-2-en-1-one (**A**, 66 mg, 0.5 mmol), PhCl (3 mL), $\text{Cu}(\text{OAc})_2$ (18 mg, 0.1 mmol), bpy (8 mg, 0.05 mmol), TEMPO (78 mg, 0.5 mmol) and phenylhydrazine (**2a**, 54 mg, 0.5 mmol). The resulting mixture was then stirred at 120 $^\circ\text{C}$ under O_2 for 5 h. Upon completion, the mixture was diluted with DCM (15 mL), and washed with water (5 mL) and brine (5 mL). The organic layer was dried over anhydrous Na_2SO_4 , and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel with petroleum ether/ethyl acetate ($V:V=50:1$) as the eluent to give **3a** in a yield of 78%.

To a reaction tube equipped with a stir bar were added 1-phenylprop-2-en-1-one (**A**, 66 mg, 0.5 mmol), phenylhydrazine (**2a**, 54 mg, 0.5 mmol) and ethanol (2 mL). The resulting mixture was then stirred at room temperature for 10 min. The solid thus formed was collected by filtration to give 1,3-diphenyl-4,5-dihydro-1H-pyrazole (**B**) in a yield of 92%. White solid (102 mg, 92%). ^1H NMR (400 MHz, CDCl_3) δ : 3.25 (t, $J=10.4$ Hz, 2H), 3.89 (t, $J=10.4$ Hz, 2H), 6.85 (t, $J=7.2$ Hz, 1H), 7.14 (d, $J=8.0$ Hz, 2H), 7.28~7.34 (m, 3H), 7.37~7.41 (m, 2H), 7.72~7.75 (m, 2H); ^{13}C NMR (150 Hz, CDCl_3) δ : 32.0, 48.3, 113.0, 119.1, 125.7, 128.5, 129.1, 133.0, 145.9, 149.0; HRMS calcd for $\text{C}_{15}\text{H}_{15}\text{N}_2$ $[\text{M}+\text{H}]^+$ 223.1230, found 223.1239.

To a reaction tube equipped with a stir bar were added 1,3-diphenyl-4,5-dihydro-1H-pyrazole (**B**, 111 mg, 0.5 mmol) and PhCl (3 mL). It was then stirred at 120 $^\circ\text{C}$ under O_2 for 5 h. Upon completion, the mixture was concentrated under reduced pressure, and the residue was purified by column chromatography on silica gel with petroleum ether/ethyl acetate ($V:V=50:1$) as the eluent to give **3a** in a yield of 93%.

4.2.4 Gram-scale synthesis of **3a**

To a reaction tube equipped with a stir bar were added propiophenone (**1a**, 810 mg, 6 mmol), PhCl (25 mL), $\text{Cu}(\text{OAc})_2$ (180 mg, 1 mmol), bpy (80 mg, 0.5 mmol) and TEMPO (780 mg, 5 mmol). The mixture was stirred at 120 $^\circ\text{C}$ under N_2 for 18 h. To the resulting mixture was added phenylhydrazine (**2a**, 540 mg, 5 mmol). It was then stirred at 120 $^\circ\text{C}$ under O_2 for 8 h. Upon completion, the mixture was diluted with DCM (80 mL), and washed with water (20 mL) and brine (20 mL). The organic layer was dried over anhydrous Na_2SO_4 , and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel with petroleum ether/ethyl acetate

($V:V=50:1$) as the eluent to give **3a** (661 mg, 60%).

Supporting Information ^1H NMR and ^{13}C NMR spectra of compounds **3a**~**3s**, **5a**~**5dd** and **B**, and X-ray crystal structure and data of **3f**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Fustero, S.; Sánchez-Roselló, M.; Barrio, P.; Simón-Fuentes, A. *Chem. Rev.* **2011**, *111*, 6984.
(b) Dadiboyena, S.; Nefzi, A. *Eur. J. Med. Chem.* **2011**, *46*, 5258.
(c) Wen, J.-J.; Zhu, Y.; Zhan, Z.-P. *Asian J. Org. Chem.* **2012**, *1*, 108.
(d) Li, M.; Zhao, B.-X. *Eur. J. Med. Chem.* **2014**, *85*, 311.
(e) Küçüküzümlü, Ş. G.; Şenkardeş, S. *Eur. J. Med. Chem.* **2015**, *97*, 786.
(f) Hughes, D. L. *Org. Process Res. Dev.* **2017**, *21*, 430.
(g) Faria, J. V.; Vegi, P. F.; Miguita, A. G. C.; dos Santos, M. S.; Boechat, N.; Bernardino, A. M. R. *Bioorg. Med. Chem.* **2017**, *25*, 5891.
- [2] (a) Schmidt, A.; Dreger, A. *Curr. Org. Chem.* **2011**, *15*, 2897.
(b) Haydl, A. M.; Xu, K.; Breit, B. *Angew. Chem., Int. Ed.* **2015**, *54*, 7149.
(c) Fang, Z.; Liu, J.; Qiao, Y. *Chin. J. Org. Chem.* **2018**, *38*, 1985 (in Chinese).
(房智兴, 刘巨艳, 乔艳红, 有机化学, **2018**, *38*, 1985.)
(d) Li, Y.; Dong, C.-E. *Chin. Chem. Lett.* **2015**, *26*, 623.
(e) Yu, X.; Zhang, J. *Chem.-Eur. J.* **2012**, *18*, 12945.
- [3] (a) Zhang, X.; Kang, J.; Niu, P.; Wu, J.; Yu, W.; Chang, J. *J. Org. Chem.* **2014**, *79*, 10170.
(b) Sar, D.; Bag, R.; Yashmeen, A.; Bag, S. S.; Punniyamurthy, T. *Org. Lett.* **2015**, *17*, 5308.
(c) Sun, J.; Qiu, J.-K.; Zhu, Y.-L.; Guo, C.; Hao, W.-J.; Jiang, B.; Tu, S.-J. *J. Org. Chem.* **2015**, *80*, 8217.
(d) Ding, Y.; Zhang, T.; Chen, Q.-Y.; Zhu, C. *Org. Lett.* **2016**, *18*, 4206 and references cited therein.
(e) Muzalevskiy, V. M.; Rulev, A. Y.; Romanov, A. R.; Kondrashov, E. V.; Ushakov, I. A.; Chertkov, V. A.; Nenajdenko, V. G. *J. Org. Chem.* **2017**, *82*, 7200.
(f) Černuchová, P.; Vo-Thanh, G.; Milata, V.; Loupy, A.; Jantová, S.; Theiszová, M. *Tetrahedron* **2005**, *61*, 5379.
- [4] Chen, Z.; Zheng, Y.; Ma, J.-A. *Angew. Chem., Int. Ed.* **2017**, *56*, 4569.
- [5] (a) Zhang, G.; Ni, H.; Chen, W.; Shao, J.; Liu, H.; Chen, B.; Yu, Y. *Org. Lett.* **2013**, *15*, 5967.
(b) Shu, W.-M.; Zheng, K.-L.; Ma, J.-R.; Sun, H.-Y.; Wang, M.; Wu, A.-X. *Org. Lett.* **2015**, *17*, 1914.
(c) Tu, Y.; Zhang, Z.; Wang, T.; Ke, J.; Zhao, J. *Org. Lett.* **2017**, *19*, 3466.
(d) Harigae, R.; Moriyama, K.; Togo, H. *J. Org. Chem.* **2014**, *79*, 2049.
- [6] (a) Li, X.; He, L.; Chen, H.; Wu, W.; Jiang, H. *J. Org. Chem.* **2013**, *78*, 3636.
(b) Fan, X.-W.; Lei, T.; Zhou, C.; Meng, Q.-Y.; Chen, B.; Tung, C.-H.; Wu, L.-Z. *J. Org. Chem.* **2016**, *81*, 7127.
(c) Pünner, F.; Sohtome, Y.; Sodeoka, M. *Chem. Commun.* **2016**, *52*, 14093.
(d) Yang, Y.; Hu, Z.-L.; Li, R.-H.; Chen, Y.-H.; Zhan, Z.-P. *Org. Biomol. Chem.* **2018**, *16*, 197.
- [7] Li, F.; Wang, J.; Pei, W.; Li, H.; Zhang, H.; Song, M.; Guo, L.; Zhang, A.; Liu, L. *Tetrahedron Lett.* **2017**, *58*, 4344.
- [8] (a) Diao, T.; Pun, D.; Stahl, S. S. *J. Am. Chem. Soc.* **2013**, *135*, 8205.
(b) Pun, D.; Diao, T.; Stahl, S. S. *J. Am. Chem. Soc.* **2013**, *135*,

- 8213.
- [9] (a) Shang, Y.; Jie, X.; Zhou, H.; Hu, P.; Huang, S.; Su, W. *Angew. Chem., Int. Ed.* **2013**, 52, 1299.
(b) Gandeepan, P.; Rajamalli, P.; Cheng, C.-H. *ACS Catal.* **2014**, 4, 4485.
(c) Jie, X.; Shang, Y.; Zhang, X.; Su, W. *J. Am. Chem. Soc.* **2016**, 138, 5623.
(d) Zhan, J.-L.; Wu, M.-W.; Chen, F.; Han, B. *J. Org. Chem.* **2016**, 81, 11994.
(e) Guo, W.; Liu, D.; Liao, J.; Ji, F.; Wu, W.; Jiang, H. *Chem. Front.* **2017**, 4, 1107.
- [10] (a) Wang, Z.; Chen, G.; Zhang, X.; Fan, X. *Org. Chem. Front.* **2017**, 4, 612.
(b) Tian, M.; Shi, X.; Zhang, X.; Fan, X. *J. Org. Chem.* **2017**, 82, 7363.
(c) Chen, G.; Wang, Z.; Zhang, X.; Fan, X. *J. Org. Chem.* **2017**, 82, 11230.
- [11] CCDC 1857401 (**3f**) contains the crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.
- [12] (a) Li, H.; Liu, C.; Zhang, Y.; Sun, Y.; Wang, B.; Liu, W. *Org. Lett.* **2015**, 17, 932.
(b) Neumann, J. J.; Suri, M.; Glorius, F. *Angew. Chem., Int. Ed.* **2010**, 49, 7790.
(c) Sun, Z.; Khan, J.; Makowska-Grzyska, M.; Zhang, M.; Cho, J. H.; Suebsuwong, C.; Vo, P.; Gollapalli, D. R.; Kim, Y.; Joachimiak, A.; Hedstrom, L.; Cuny, G. D. *J. Med. Chem.* **2014**, 57, 10544.
- [13] Voronin, V. V.; Ledovskaya, M. S.; Gordeev, E. G.; Rodygin, K. S.; Ananikov, V. P. *J. Org. Chem.* **2018**, 83, 3819.
- [14] Tang, X.; Huang, L.; Yang, J.; Xu, Y.; Wu, W.; Jiang, H. *Chem. Commun.* **2014**, 50, 14793.
- [15] Shan, G.; Liu, P.; Rao, Y. *Org. Lett.* **2011**, 13, 1746.
- [16] Zhang G.; Zhao, Y.; Ge, H. *Angew. Chem., Int. Ed.* **2013**, 52, 2559.
- [17] Comas-Barcel, J.; Foster, R. S.; Fiser, B.; Gomez-Bengoa, E.; Harity, J. P. A. *Chem.-Eur. J.* **2015**, 21, 3257.

(Zhao, C.)