磷催化 α' -亚甲基环戊烯酮的形式[6+2]环加成反应

石崇慧 肖本现 杜 玮 陈应春*

(四川大学华西药学院 成都 610041)

摘要 采用三环己基膦活化 α' -亚甲基环戊烯酮底物, 通过质子转移使得反应位点迁移, 促进远端 γ -位与 2-烯基-1,3-茛二酮化合物发生分子间 Rauhut-Currier 类反应, 并进一步在 β' -位发生分子内 Michael 加成, 从而实现了形式[6+2]环加成反应, 以中等至良好收率以及优秀的非对映选择性构建了一系列多取代双环并螺 1,3-茛二酮骨架。该反应策略拓展了传统 Rauhut-Currier 反应的常规反应模式, 而该新颖的[6+2]环加成反应途径也有望在有机合成中得到更多的应用。

关键词 叔膦催化; α' -亚甲基环戊烯酮; Rauhut-Currier 反应; [6+2]环加成反应

Phosphine-Catalyzed Formal [6+2] Cycloadditions of α' -Methylene 2-Cyclopentenones

Shi, Chonghui Xiao, Benxian Du, Wei Chen, Yingchun*

(West China School of Pharmacy, Sichuan University, Chengdu 610041)

Abstract The formal [6+2] cycloaddition reaction of α' -methylene-2-cyclopentenones and 2-alkylidene-1,3-indanediones was developed under the catalysis of tricyclohexyl phosphine, proceeding in a remote γ -regioselective Rauhut-Currier-type reaction followed by a β' -regioselective intramolecular Michael addition process. An array of fused bicyclic frameworks incorporating a spiro-1,3-indanedione motif were produced in fair to good yields of 48% ~ 76% with excellent diastereoselectivity (generally > 19 : 1 *dr*). This protocol expands the reaction mode of traditional Rauhut-Currier reaction, which might find further application in organic synthesis.

Keywords phosphine catalysis; α' -methylene-2-cyclopentenone; Rauhut-Currier reaction; [6+2] cycloaddition

1 Introduction

The cycloaddition reaction represents one of the indispensable methods to construct cyclic scaffolds in organic synthesis. For decades, a variety of methods, such as conventional [2+2], [3+2], [3+3], [4+2] pericyclic or formal cycloadditions, have been extensively explored.^[1] Among them, the [6+2] cycloaddition reaction based on fulvenes to produce fused bi- or tri-cyclic skeletons has attracted special attention since the pioneering work of Houk.^[2] Recently, we also reported an amine catalyzed asymmetric [6+2] cycloaddition reaction of α' -alkylidene-2-cyclopentenones by *in situ* generating 4-aminofulvene intermediates.^[3] Nevertheless, some significant substrate limitations were observed in the above strategies.

The Rauhut-Currier (RC) reaction, as a powerful and economic protocol to construct C—C bond between activated olefins, also has aroused widespread attention,

whereas the chemoselectivity represents one of the long-standing challenges in this field.^[4] Meanwhile, according to the well-established catalytic cycle, a normal RC-type reaction can not proceed by using the activated olefins with an α -substitution, because the regeneration of Lewis base catalyst relies on an elimination process involving proton transfer (Scheme 1a, II). However, the activated olefins still can form zwitterionic enolates with suitable Lewis bases. In 2006, Schaus *et al.*^[5] reported the dimerization reaction of α -substituted 1,4-dien-3-ones to deliver a variety of bicyclo[3.2.1]octenones through cascade [4+2] cycloaddition/Wittig reaction in the presence of phosphine. In addition, we^[6] also disclosed a phosphine-mediated interrupted Morita-Baylis-Hillman reaction and some unexpected cascade processes of 3-olefinic oxindoles. However, stoichiometric phosphine substances were generally required due to the generation of phosphine oxides.

* Corresponding author. E-mail: ycchen@scu.edu.cn

Received March 27, 2019; revised April 29, 2019; published online May 21, 2019.

Dedicated to the 100th anniversary of the birth of Professor Ruyu Chen.

Project supported by the National Natural Science Foundation of China (No. 21572135).

国家自然科学基金(No. 21572135)资助项目。

In 2005, The Roush group^[7] reported a phosphine-catalyzed intramolecular aldol reaction of unsaturated 1,5-diketones. It was proposed that the coulombic interaction between the phosphonium unit and the adjacent carbonyl in intermediate **IV**, generated from the RC intermediate **III** via protonation, could increase the acidity of the α' -proton, thus the reactive site was transferred from α -site to less sterically hindered α' -site to furnish the high regioselectivity of intramolecular aldol reaction. Trifluoroethanol (TFE) was found to be crucial for the success of the reaction (Scheme 1b).^[7] Based on the above considerations, we envisaged that the α' -methylene-2-cyclopentenone would be attacked by a tertiary phosphine to generate the zwitterionic intermediate **VI**. It was hypothesized that the acidity of the γ -CH of the similarly protonated intermediate **VII** would be significantly enhanced, thus leading to the formation of the intermediate **VIII** via deprotonation. As a result, the reactive site would be shifted from α' -site to remote γ -site, and a cascade RC-type reaction/phosphine elimination/*anti*-Baldwin cyclization (5-*endo-trig*)^[8] would furnish a formal [6+2] cycloaddition process accordingly (Scheme 1c).

2 Results and discussion

The initial investigation began with the reaction of α' -methylene- β -phenylcyclopent-2-en-1-one (**1a**)^[9] and 2-benzylidene-1,3-indanedione (**2a**)^[10] in TFE using *n*Bu₃P as the catalyst. To our delight, the corresponding formal [6+2] cycloaddition product **3a** was obtained in 50% yield (Table 1, Entry 1). Then a few phosphine catalysts were investigated. Me₃P gave the product in 36% yield (Table 1,

Entry 2), and a better yield was obtained with Cy₃P (Table 1, Entry 3). Nevertheless, inferior results were generally delivered with phosphines with diverse aromatic groups (Table 1, Entries 4~8). Solvent screenings (Table 1, Entries 9~14) indicated that TFE was also crucial for the observed reaction, but the reaction did not work in hexafluoroisopropanol (HFIP, Table 1, Entry 15). These results are consistent with the Roush's work and demonstrate that TFE could facilitate the migration of proton, thus enabling the switch of reactive sites.^[7] Finally, it was found that much shorter time was required by conducting the reaction with higher loadings of Cy₃P and at higher temperature, though the same yield for product **3a** was obtained (Table 1, Entry 16). Moreover, a racemic bifunctional phosphine **C5** was found to be a superior catalyst, giving **3a** in an excellent yield (Table 1, Entry 17).

Considering the ready availability of catalyst, the generality of this formal [6+2] cycloaddition reaction with a variety of substrates was investigated under the catalysis of commercial Cy₃P. The results are summarized in Table 2. Firstly, 2-alkylidene-1,3-indanediones **2** with diversely substituted aromatic and heteroaromatic rings were investigated in the reactions with dienone **1a**, and the corresponding cycloadducts **3b**~**3k** were generally produced in fair to good yields with exclusive diastereoselectivity (Table 2, Entries 2~11). Moreover, an array of dienones **1** with various β -aryl groups were tested in the reactions with acceptor **2a**, and similar results were afforded for the products **3l**~**3s** (Table 2, Entries 12~19). Unfortunately, it was found that the dienones **1** bearing either a β -methyl or β -H substituent showed inert reactivity with **2a** under the current

Scheme 1 Diverse reaction patterns via the RC process

Table 1 Optimization conditions of the phosphine-catalyzed formal [6+2] cycloaddition^a

Entry	Cat.	Solvent	T/°C	Yield ^b /%
1	<i>n</i> Bu ₃ P	TFE	40	53
2	Me ₃ P	TFE	40	36
3	Cy ₃ P	TFE	40	56
4	PPh ₃	TFE	40	45
5	C1	TFE	40	47
6	C2	TFE	40	48
7	C3	TFE	40	49
8	C4	TFE	40	30
9	Cy ₃ P	toluene	40	trace
10	Cy ₃ P	MeCN	40	NR
11	Cy ₃ P	CHCl ₃	40	NR
12	Cy ₃ P	THF	40	NR
13	Cy ₃ P	MeOH	40	32
14	Cy ₃ P	<i>i</i> PrOH	40	NR
15	Cy ₃ P	HFIP	40	NR
16 ^c	Cy ₃ P	TFE	60	56 ^d
17 ^c	C5	TFE	60	92 ^d

^a Unless otherwise noted, the reactions were carried out with **2a** (0.05 mmol), **1a** (0.1 mmol), phosphine (20 mol%) in solvent (0.25 mL) at 40 °C for 48 h under Ar. ^b Yield of isolated product; *dr* (>19 : 1) determined by ¹H NMR analysis. ^c Catalyst (30 mol%) was used. ^d For 12 h.

catalytic conditions. Furthermore, bifunctional **C5** was also employed as the catalyst in some cases, and most of them provided higher yields (Table 2, data in parentheses).

The potential asymmetric formal [6+2] cycloaddition reaction of dienone **1a** and acceptor **2a** was also explored. As outlined in Scheme 2, the chiral phosphine *S*-**C5** failed to provide any enantiocontrol, though an excellent yield for the cycloadduct **3a** could be attained. In addition, the chiral bifunctional phosphine **C6** exhibited moderate catalytic activity in TFE, but no enantioselectivity could be induced either. Similarly, moderate to good yields without enantiocontrol were obtained in the presence of chiral phosphines **C7**~**C9**. Moreover, very poor conversions were

Table 2 Substrate scope of the phosphine-catalyzed formal [6+2] cycloaddition reaction^a

Entry	R ¹	R ²	3	Yield ^b /%
1	Ph	Ph	3a	56 ^c (92)
2	Ph	4-FC ₆ H ₄	3b	45 (64)
3	Ph	4-ClC ₆ H ₄	3c	70
4	Ph	4-BrC ₆ H ₄	3d	55 (54)
5	Ph	2-MeC ₆ H ₄	3e	50 (76)
6	Ph	3-MeC ₆ H ₄	3f	69 (66)
7	Ph	4-MeC ₆ H ₄	3g	57
8	Ph	4- ^t BuC ₆ H ₄	3h	48 (54)
9	Ph	3-MeOC ₆ H ₄	3i	44 (59)
10	Ph	4-MeOC ₆ H ₄	3j	75
11	Ph	2-Furyl	3k	60 (62)
12	4-FC ₆ H ₄	Ph	3l	76 (79)
13	3,5-(CF ₃) ₂ C ₆ H ₃	Ph	3m	48
14	3-MeC ₆ H ₄	Ph	3n	57
15	4-MeC ₆ H ₄	Ph	3o	59 (69)
16	3,5-(CH ₃) ₂ C ₆ H ₃	Ph	3p	59 (78)
17	3-MeOC ₆ H ₄	Ph	3q	62 (79)
18	4-MeOC ₆ H ₄	Ph	3r	50
19	2-Naphthyl	Ph	3s	55 (79)

^a Unless otherwise noted, the reactions were carried out with **2** (0.1 mmol), **1** (0.2 mmol), Cy₃P (30 mol%) in TFE (0.5 mL) at 60 °C for 12~48 h under Ar. ^b Yield of the isolated product [data in parentheses were obtained with **C5** (30 mol%)]; *dr* (>19 : 1) determined by ¹H NMR analysis. ^c The relative configuration of racemic **3a** was determined by X-ray analysis (CCDC 1905888). The other products were assigned by analogy.

Scheme 2 Attempts in asymmetric formal [6+2] cycloaddition reaction

still observed by replacing TFE with other solvents, such as toluene, MeCN, CHCl₃ and THF. It was proposed that

the γ -reactive site is far from the chiral catalyst, which may make the stereocontrol quite difficult. On the other hand, TFE would also seriously affect the potential concerted hydrogen bonding interaction between the chiral catalyst and substrates. As a result, the development of an asymmetric [6+2] cycloaddition reaction still remained to be explored.

In addition, more electrophiles were explored. Fortunately, the desired product **5** could be generated by using 3-olefinic-7-azaioxindole **4** as the electrophile, albeit in fair yield and diastereoselectivity (Scheme 3).

Scheme 3 More exploration of phosphine-catalyzed [6+2] cycloaddition reaction

3 Conclusions

A highly γ,β' -regioselective formal [6+2] cycloaddition reaction between α' -methylene-2-cyclopentenones and 2-alkylidene-1,3-indanediones catalyzed by a tertiary phosphine was successfully developed. This reaction relied on forming zwitterionic dienolate species via proton transfer, proceeding in a γ -regioselective Rauhut-Currier-type reaction followed by an intramolecular Michael addition process. This method provided a facile and economic protocol to construct fused bicyclic frameworks incorporating a spiro-1,3-indanedione motif in fair to good yields with exclusive diastereoselectivity, which may find further application in organic synthesis. This work also expanded the reaction mode of enone substrates bearing α -substitutions under phosphine catalysis. Further investigation in this field, especially developing asymmetric reactions, is under way in our laboratory.

4 Experimental section

4.1 General

NMR data were obtained for ^1H at 400 MHz or 600 MHz, and for ^{13}C at 100 MHz or 150 MHz. Chemical shifts were reported from tetramethylsilane with the solvent resonance as the internal standard in CDCl_3 solution. ESI HRMS was recorded on a Waters SYNAPT G2. Column chromatography was performed on silica gel (200~300 mesh) eluting with ethyl acetate and petroleum ether. TLC was performed on glass-backed silica plates. UV light, I_2 , and solution of potassium permanganate were used to visualize products or starting materials. All chemicals were used without purification as commercially available unless otherwise noted. Petroleum ether and ethyl acetate (EtOAc) were distilled. THF was freshly distilled from sodium/benzophenone before use. Experiments involving moisture and/or air sensitive components were

performed under a positive pressure of argon in oven-dried glassware equipped with a rubber septum inlet. Dried solvents and liquid reagents were transferred by oven-dried syringes. 2-Alkylidene-1,3-indanediones **2** were synthesized following the literature procedures.^[10]

4.2 General procedure for the synthesis of substrates 1

To a mixture of β -phenylcyclopent-2-en-1-one (1.87 g, 11.8 mmol) and paraformaldehyde (0.71 g, 23.6 mmol) in dry toluene (100 mL) was added $(^i\text{Pr})_2\text{H}_2\text{N}^+\text{CF}_3\text{COO}^-$ (2.54 g, 11.8 mmol) and trifluoroacetic acid (0.13 mL, 1.18 mmol). The reaction mixture was stirred at 80 °C for 2 h. The mixture became clear and was cooled down to room temperature. Then more paraformaldehyde (0.71 g, 23.6 mmol) was added. The mixture was stirred at 80 °C for another 6 h. The mixture was concentrated in vacuo and purified by flash column chromatography on silica gel eluting with EtOAc/petroleum ether ($V:V=1:10$) to give 5-methylene-3-phenylcyclopent-2-en-1-one (**1a**).^[9] yellow solid, 1.20 g, 59.0% yield. m.p. 137~139 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.72~7.62 (m, 2H), 7.52~7.4 (m, 3H), 6.79 (s, 1H), 6.15 (s, 1H), 5.51 (s, 1H), 3.65 (s, 2H); ^{13}C NMR (150 MHz, CDCl_3) δ : 195.7, 167.9, 142.3, 131.5, 129.05, 128.9, 126.9, 116.5, 34.0.

4.3 General procedure for phosphine-catalyzed [6+2] cycloaddition

5-Methylene-2-cyclopentenone **1** (0.2 mmol), 2-alkylidene-1,3-indanedione **2** (0.1 mmol), and phosphine Cy_3P (8.4 mg, 0.03 mmol; or **C5**, 13.6 mg, 0.03 mmol) were dissolved in TFE (0.5 mL). Then the mixture was stirred at 60 °C under Ar for the indicated time (12~48 h). After completion, it was concentrated and purified by flash chromatography on silica gel eluting with EtOAc/petroleum ether ($V:V=1:3\sim 1:5$) to give the product **3**.

1',6'-Diphenyl-1,3',3a',6a'-tetrahydro-4'H-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3a**): White solid, 23.0 mg, 56% yield (with **C5**, 37.2 mg, 92% yield). m.p. 155~159 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.81 (d, $J=7.8$ Hz, 1H), 7.75 (d, $J=7.2$ Hz, 1H), 7.72~7.64 (m, 2H), 7.22~7.14 (m, 3H), 7.04 (d, $J=7.6$ Hz, 2H), 7.10~6.94 (m, 5H), 6.34 (s, 1H), 4.78~4.69 (m, 1H), 3.88 (td, $J=9.8, 7.2$ Hz, 1H), 3.40 (d, $J=9.8$ Hz, 1H), 2.40 (dd, $J=13.2, 9.6$ Hz, 1H), 2.28 (dd, $J=13.2, 9.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 208.9, 203.2, 200.4, 176.5, 141.5, 136.7, 135.8, 135.6, 132.7, 130.6, 128.4, 128.3, 128.0, 128.0, 127.9, 127.4, 126.0, 123.1, 123.1, 69.3, 58.0, 52.7, 51.6, 34.3; HRMS calcd for $\text{C}_{28}\text{H}_{20}\text{O}_3\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 427.1305, found 427.1317.

1'-(4-Fluorophenyl)-6'-phenyl-1,3',3a',6a'-tetrahydro-4'H-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3b**): Pale yellow solid, 19.0 mg, 45% yield (with **C5**, 27.3 mg, 64% yield). m.p. 177~179 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.85 (dd, $J=6.4, 1.6$ Hz, 1H), 7.78 (dd, $J=6.4, 1.2$ Hz, 1H), 7.76~7.69 (m, 2H), 7.22 (t, $J=7.6$ Hz, 1H), 7.19~7.13 (m, 2H), 7.08 (t, $J=7.6$ Hz, 2H), 7.01~6.91 (m, 2H), 6.68 (t, $J=8.8$ Hz, 2H), 6.33 (d, $J=1.2$ Hz, 1H), 4.69 (t,

$J=8.0$ Hz, 1H), 3.86 (td, $J=9.6, 7.2$ Hz, 1H), 3.38 (d, $J=10.0$ Hz, 1H), 2.39 (dd, $J=13.6, 9.6$ Hz, 1H), 2.26 (dd, $J=13.6, 9.6$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 208.7, 203.0, 200.3, 176.3, 161.9 ($J_{\text{CF}}^1=245.4$ Hz), 141.5, 141.4, 136.1, 135.9, 132.7, 132.6, 132.5, 130.8, 130.0, 130.0 ($J_{\text{CF}}^3=8.1$ Hz), 127.9, 126.1, 123.2, 115.3 ($J_{\text{CF}}^2=21.5$ Hz), 69.2, 57.1, 52.9, 51.5, 34.4; HRMS calcd for $\text{C}_{28}\text{H}_{19}\text{FO}_3\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 445.1210, found 445.1212.

1'-(4-Chlorophenyl)-6'-phenyl-1',3',3a',6a'-tetrahydro-4'*H*-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3c**): Pale yellow semi-solid, 30.4 mg, 70% yield; ^1H NMR (600 MHz, CDCl_3) δ : 7.84 (d, $J=7.8$ Hz, 1H), 7.78 (d, $J=7.2$ Hz, 1H), 7.76~7.67 (m, 2H), 7.23 (t, $J=7.2$ Hz, 1H), 7.19~7.12 (m, 2H), 7.08 (d, $J=7.8$ Hz, 2H), 7.01~6.90 (m, 4H), 6.32 (s, 1H), 4.74~4.65 (m, 1H), 3.84 (td, $J=9.6, 7.2$ Hz, 1H), 3.37 (d, $J=9.6$ Hz, 1H), 2.38 (dd, $J=13.8, 9.6$ Hz, 1H), 2.25 (dd, $J=13.8, 9.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 208.6, 202.8, 200.2, 176.2, 141.5, 141.4, 136.1, 136.0, 135.5, 133.3, 132.7, 130.9, 129.8, 128.6, 128.2, 127.9, 126.2, 123.3, 123.2, 69.1, 57.0, 52.87, 51.5, 34.7; HRMS calcd for $\text{C}_{28}\text{H}_{19}\text{ClO}_3\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 461.0915 (^{35}Cl), 463.0885 (^{37}Cl), found 461.0916 (^{35}Cl), 463.0910 (^{37}Cl).

1'-(4-Bromophenyl)-6'-phenyl-1',3',3a',6a'-tetrahydro-4'*H*-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3d**): Pale yellow solid, 26.5 mg, 55% yield (with **C5**, 26.2 mg, 54% yield). m.p. 222~224 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.89 (dd, $J=6.8, 1.6$ Hz, 1H), 7.82 (dd, $J=6.8, 2.0$ Hz, 1H), 7.76~7.70 (m, 2H), 7.30~7.20 (m, 1H), 7.19~7.05 (m, 6H), 6.88 (d, $J=8.4$ Hz, 2H), 6.33 (s, 1H), 4.72~4.63 (m, 1H), 3.85 (td, $J=9.6, 7.2$ Hz, 1H), 3.36 (d, $J=10.0$ Hz, 1H), 2.39 (dd, $J=13.6, 9.6$ Hz, 1H), 2.25 (dd, $J=13.6, 9.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 208.6, 200.2, 202.8, 176.2, 141.5, 141.4, 136.2, 136.0, 132.7, 131.5, 130.9, 130.2, 128.2, 127.9, 126.2, 123.3, 123.3, 121.4, 69.0, 57.0, 52.9, 51.5, 34.8; HRMS calcd for $\text{C}_{28}\text{H}_{19}\text{BrO}_3\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 505.0410 (^{79}Br), 507.0389 (^{81}Br), found 505.0407 (^{79}Br), 507.0409 (^{81}Br).

6'-Phenyl-1'-(*o*-tolyl)-1',3',3a',6a'-tetrahydro-4'*H*-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3e**): White solid, 21.2 mg, 50% yield (with **C5**, 32.0 mg, 76% yield). m.p. 200~203 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.85 (d, $J=7.6$ Hz, 1H), 7.74 (d, $J=7.6$ Hz, 1H), 7.72~7.64 (m, 2H), 7.45 (d, $J=8.0$ Hz, 1H), 7.20~7.14 (m, 1H), 7.11~7.05 (m, 2H), 7.06~6.95 (m, 3H), 6.90~6.81 (m, 1H), 6.62 (d, $J=7.2$ Hz, 1H), 6.33 (s, 1H), 4.67~4.53 (m, 1H), 3.88 (td, $J=9.6, 7.6$ Hz, 1H), 3.76 (d, $J=9.6$ Hz, 1H), 2.40 (dd, $J=13.2, 9.6$ Hz, 1H), 2.29 (dd, $J=13.2, 9.6$ Hz, 1H), 1.58 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 209.1, 203.6, 200.6, 176.7, 141.8, 141.4, 137.1, 135.9, 135.7, 135.6, 132.6, 130.7, 130.4, 128.0, 127.8, 127.3, 127.0, 126.2, 125.7, 123.1, 123.1, 68.9, 54.7, 52.4, 51.6, 34.7, 19.2; HRMS calcd for $\text{C}_{29}\text{H}_{22}\text{O}_3\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 441.1461, found 441.1460.

6'-Phenyl-1'-(*m*-tolyl)-1',3',3a',6a'-tetrahydro-4'*H*-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3f**): Pale yellow solid, 29.0 mg, 69% yield (with **C5**, 27.6 mg, 66% yield). m.p. 162~164 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.85~7.80 (m,

1H), 7.79~7.74 (m, 1H), 7.65~7.72 (m, 2H), 7.21 (d, $J=7.4$ Hz, 1H), 7.18~7.14 (m, 2H), 7.04 (dd, $J=8.4, 7.2$ Hz, 2H), 6.91~6.80 (m, 2H), 6.75 (dd, $J=7.2, 1.6$ Hz, 1H), 6.71 (s, 1H), 6.34 (d, $J=1.2$ Hz, 1H), 4.67~4.72 (m, 1H), 3.87 (td, $J=9.6, 7.2$ Hz, 1H), 3.35 (d, $J=9.6$ Hz, 1H), 2.39 (dd, $J=13.2, 9.6$ Hz, 1H), 2.27 (dd, $J=13.2, 9.6$ Hz, 1H), 2.06 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 209.1, 203.1, 200.5, 176.6, 141.6, 141.5, 137.9, 136.7, 135.8, 135.6, 132.8, 130.7, 129.4, 128.3, 128.1, 128.0, 126.0, 125.3, 123.1, 123.0, 69.3, 58.2, 52.9, 51.7, 34.3, 21.1; HRMS calcd for $\text{C}_{29}\text{H}_{22}\text{O}_3\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 441.1461, found 441.1462.

6'-Phenyl-1'-(*p*-tolyl)-1',3',3a',6a'-tetrahydro-4'*H*-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3g**): Pale yellow semi-solid, 23.7 mg, 57% yield; ^1H NMR (400 MHz, CDCl_3) δ : 7.81 (d, $J=6.8$ Hz, 1H), 7.78~7.74 (m, 1H), 7.74~7.65 (m, 2H), 7.22~7.12 (m, 3H), 7.02 (t, $J=8.0$ Hz, 2H), 6.85 (d, $J=7.6$ Hz, 2H), 6.76 (d, $J=7.6$ Hz, 2H), 6.33 (d, $J=1.2$ Hz, 1H), 4.73~4.63 (m, 1H), 3.85 (dd, $J=9.6, 7.2$ Hz, 1H), 3.36 (d, $J=9.6$ Hz, 1H), 2.36 (dd, $J=13.2, 9.6$ Hz, 1H), 2.24 (dd, $J=13.2, 9.6$ Hz, 1H), 2.10 (s, 3H). ^{13}C NMR (150 MHz, CDCl_3) δ : 209.1, 203.1, 200.6, 176.6, 141.6, 136.9, 135.8, 135.6, 133.6, 132.7, 130.6, 129.0, 128.3, 128.0, 128.0, 126.0, 123.1, 123.1, 69.3, 57.6, 52.8, 51.6, 34.4, 20.9; HRMS calcd for $\text{C}_{29}\text{H}_{22}\text{O}_3\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 441.1461, found 441.1459.

1'-(4-(*tert*-Butyl)phenyl)-6'-phenyl-1',3',3a',6a'-tetrahydro-4'*H*-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3h**): Yellow solid, 22.0 mg, 48% yield (with **C5**, 24.8 mg, 54% yield). m.p. 242~244 $^{\circ}\text{C}$; ^1H NMR (400 MHz, CDCl_3) δ : 7.84 (dd, $J=6.4, 1.2$ Hz, 1H), 7.77 (dd, $J=6.0, 1.2$ Hz, 1H), 7.85~7.68 (m, 2H), 7.19~7.12 (m, 3H), 7.03~6.94 (m, 4H), 6.88 (d, $J=8.3$ Hz, 2H), 6.37~6.29 (m, 1H), 4.81~4.62 (m, 1H), 3.85 (td, $J=9.6, 7.2$ Hz, 1H), 3.36 (d, $J=9.6$ Hz, 1H), 2.39 (dd, $J=13.6, 9.6$ Hz, 1H), 2.37 (dd, $J=13.6, 9.6$ Hz, 1H), 1.13 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ : 209.2, 203.3, 200.5, 176.6, 150.4, 141.7, 141.6, 135.7, 135.5, 133.6, 132.7, 130.6, 128.2, 128.0, 123.0, 125.9, 125.2, 123.1, 69.3, 57.8, 53.1, 51.6, 34.4, 34.3, 31.1; HRMS calcd for $\text{C}_{32}\text{H}_{28}\text{O}_3\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 483.1931, found 483.1942.

1'-(3-Methoxyphenyl)-6'-phenyl-1',3',3a',6a'-tetrahydro-4'*H*-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3i**): Pale yellow semi-solid, 19.0 mg, 44% yield (with **C5**, 25.2 mg, 59% yield); ^1H NMR (400 MHz, CDCl_3) δ : 7.84 (dd, $J=8.0, 1.2$ Hz, 1H), 7.77 (dd, $J=6.4, 1.2$ Hz, 1H), 7.85~7.68 (m, 2H), 7.25~7.17 (m, 3H), 7.06 (dd, $J=8.4, 7.2$ Hz, 2H), 6.90 (t, $J=8.4$ Hz, 1H), 6.61 (d, $J=7.6$ Hz, 1H), 6.54~6.46 (m, 2H), 6.35 (d, $J=1.2$ Hz, 1H), 4.74~4.65 (m, 1H), 3.88 (td, $J=9.6, 7.6$ Hz, 1H), 3.58 (s, 3H), 3.36 (d, $J=9.6$ Hz, 1H), 2.39 (dd, $J=13.6, 9.6$ Hz, 1H), 2.26 (dd, $J=13.6, 9.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 208.9, 203.1, 200.4, 176.5, 159.4, 141.6, 138.4, 135.9, 135.7, 132.8, 130.7, 129.4, 128.1, 128.0, 126.0, 123.1, 123.1, 120.8, 113.9, 113.4, 69.2, 58.1, 55.1, 52.9, 51.6, 34.4; HRMS calcd for $\text{C}_{29}\text{H}_{22}\text{O}_4\text{Na}$ [$\text{M}+\text{Na}$] $^+$ 457.1410, found 457.1412.

1'-(4-Methoxyphenyl)-6'-phenyl-1',3',3a',6a'-tetrahydro-4'*H*-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3j**): Yellow

solid, 32.7 mg, 75% yield. m.p. 173~175 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.81 (dd, $J=6.8, 1.6$ Hz, 1H), 7.75 (dd, $J=6.8, 1.6$ Hz, 1H), 7.65~7.73 (m, 2H), 7.25~7.13 (m, 3H), 7.06 (d, $J=7.6$ Hz, 2H), 6.89 (d, $J=8.4$ Hz, 2H), 6.50 (d, $J=8.8$ Hz, 2H), 6.32 (d, $J=1.2$ Hz, 1H), 4.80~4.59 (m, 1H), 3.84 (td, $J=9.6, 7.6$ Hz, 1H), 3.61 (s, 3H), 3.36 (d, $J=9.6$ Hz, 1H), 2.36 (dd, $J=13.6, 9.6$ Hz, 1H), 2.24 (dd, $J=13.6, 9.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 209.0, 203.3, 200.7, 176.6, 158.7, 141.6, 141.6, 135.9, 135.7, 132.8, 130.7, 129.5, 128.8, 128.1, 128.0, 126.0, 123.1, 123.1, 113.7, 69.3, 57.3, 55.1, 52.9, 51.6, 34.4; HRMS calcd for $\text{C}_{29}\text{H}_{22}\text{O}_3\text{Na}$ $[\text{M}+\text{Na}]^+$ 457.1410, found 457.1408.

1'-(Furan-3-yl)-6'-phenyl-1',3',3a',6a'-tetrahydro-4'H-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3k**): White solid, 23.7 mg, 60% yield (with **C5**, 24.6 mg, 62% yield); ^1H NMR (400 MHz, CDCl_3) δ : 7.91 (dd, $J=6.8, 2.0$ Hz, 1H), 7.81 (dd, $J=6.4, 2.0$ Hz, 1H), 7.79~7.72 (m, 2H), 7.31~7.24 (m, 3H), 7.17 (d, $J=7.6$ Hz, 2H), 7.08 (d, $J=2.0$ Hz, 1H), 6.74 (s, 1H), 6.33~6.25 (m, 1H), 6.21 (d, $J=2.0$ Hz, 1H), 4.61~4.48 (m, 1H), 3.77 (td, $J=9.6, 7.2$ Hz, 1H), 3.30 (d, $J=9.6$ Hz, 1H), 2.34 (dd, $J=13.6, 9.6$ Hz, 1H), 2.22 (dd, $J=13.6, 9.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 208.9, 203.3, 200.4, 176.2, 143.3, 141.7, 141.6, 140.6, 136.0, 135.9, 132.9, 131.0, 128.1, 128.0, 126.1, 123.3, 123.2, 121.2, 109.5, 68.3, 52.9, 51.5, 48.0, 34.0; HRMS calcd for $\text{C}_{26}\text{H}_{18}\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 417.1097, found 417.1094.

6'-(4-Fluorophenyl)-1'-phenyl-1',3',3a',6a'-tetrahydro-4'H-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3l**): White solid, 32.4 mg, 76% yield (with **C5**, 28.4 mg, 79% yield). m.p. 211~213 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.85~7.80 (m, 1H), 7.78~7.74 (m, 1H), 7.73~7.64 (m, 2H), 7.19~7.12 (m, 2H), 7.03~6.95 (m, 5H), 6.75~6.65 (m, 2H), 6.30 (d, $J=1.2$ Hz, 1H), 4.71~4.67 (m, 1H), 3.87 (td, $J=9.6, 7.2$ Hz, 1H), 3.37 (d, $J=9.6$ Hz, 1H), 2.40 (dd, $J=13.2, 9.6$ Hz, 1H), 2.27 (dd, $J=13.2, 9.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 208.7, 203.0, 200.4, 175.1, 164.0 (d, $J_{\text{CF}}^1=251.1$ Hz), 141.6, 141.5, 136.7, 135.9, 135.7, 130.0 (d, $J_{\text{CF}}^3=8.6$ Hz), 128.9, 128.9, 128.5, 128.4, 127.6, 125.8, 123.1, 115.3 (d, $J_{\text{CF}}^2=21.6$ Hz), 69.2, 58.0, 52.9, 51.6, 34.4; HRMS calcd for $\text{C}_{28}\text{H}_{19}\text{FO}_3\text{Na}$ $[\text{M}+\text{Na}]^+$ 445.1210, found 445.1208.

6'-(3,5-Bis(trifluoromethyl)phenyl)-1'-phenyl-1',3',3a',6a'-tetrahydro-4'H-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3m**): White solid, 25.8 mg, 48% yield. m.p. 162~166 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.82 (d, $J=7.8$ Hz, 1H), 7.76 (d, $J=7.8$ Hz, 1H), 7.74~7.65 (m, 3H), 7.62 (s, 2H), 7.26 (s, 1H), 6.98 (s, 4H), 6.46 (s, 1H), 4.83~4.75 (m, 1H), 3.92 (td, $J=9.6, 7.8$ Hz, 1H), 3.37 (d, $J=9.6$ Hz, 1H), 2.41 (dd, $J=13.2, 9.6$ Hz, 1H), 2.29 (dd, $J=13.2, 9.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 208.0, 202.8, 200.0, 172.3, 141.5, 141.5, 136.0, 135.9, 135.6, 135.0, 131.8 (q, $J_{\text{CF}}^2=33.6$), 128.8, 128.8, 128.0, 127.9, 127.6, 123.9 (q, $J_{\text{CF}}^3=5.7$), 123.2, 122.6 (q, $J_{\text{CF}}^1=217.5$), 69.1, 57.9, 52.7, 51.7, 34.1; HRMS calcd for $\text{C}_{30}\text{H}_{18}\text{F}_6\text{O}_3\text{Na}$ $[\text{M}+\text{Na}]^+$ 563.1052, found 563.1058.

1'-Phenyl-6'-(*m*-tolyl)-1',3',3a',6a'-tetrahydro-4'H-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3n**): White semi-solid, 23.6 mg, 56% yield; ^1H NMR (400 MHz, CDCl_3) δ : 7.82 (dd, $J=7.6, 1.6$ Hz, 1H), 7.76 (dd, $J=7.6, 1.6$ Hz, 1H), 7.73~7.64 (m, 2H), 7.18~7.07 (m, 1H), 7.03~6.97 (m, 7H), 6.84 (s, 1H), 6.35 (d, $J=1.2$ Hz, 1H), 4.75~4.65 (m, 1H), 3.88 (td, $J=9.6, 7.2$ Hz, 1H), 3.40 (d, $J=9.6$ Hz, 1H), 2.39 (dd, $J=13.6, 9.6$ Hz, 1H), 2.26 (dd, $J=13.6, 9.6$ Hz, 1H), 1.95 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 209.0, 203.1, 200.5, 176.6, 141.6, 137.8, 137.1, 135.9, 135.7, 132.5, 131.5, 129.1, 128.5, 128.4, 128.1, 127.4, 125.7, 125.2, 123.1, 123.1, 69.3, 58.0, 52.8, 51.7, 34.5, 20.9; HRMS calcd for $\text{C}_{29}\text{H}_{22}\text{O}_3\text{Na}$ $[\text{M}+\text{Na}]^+$ 441.1461, found 441.1463.

1'-Phenyl-6'-(*p*-tolyl)-1',3',3a',6a'-tetrahydro-4'H-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3o**): White solid, 24.8 mg, 59% yield (with **C5**, 29.0 mg, 69% yield). m.p. 230~234 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.81~7.79 (m, 1H), 7.78~7.74 (m, 1H), 7.73~7.64 (m, 2H), 7.06 (dd, $J=6.4, 1.6$ Hz, 2H), 6.93~6.91 (m, 5H), 6.84 (d, $J=8.0$ Hz, 2H), 6.33 (d, $J=1.2$ Hz, 1H), 4.73~7.68 (m, 1H), 3.87 (td, $J=9.6, 7.2$ Hz, 1H), 3.41 (d, $J=9.6$ Hz, 1H), 2.39 (dd, $J=13.6, 9.6$ Hz, 1H), 2.27 (dd, $J=13.6, 9.6$ Hz, 1H), 2.22 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 208.9, 203.1, 200.6, 176.4, 141.6, 141.3, 136.9, 135.8, 135.6, 129.9, 128.9, 128.5, 128.4, 128.1, 127.4, 125.2, 123.1, 123.1, 69.4, 58.1, 52.6, 51.7, 34.4, 21.4; HRMS calcd for $\text{C}_{29}\text{H}_{22}\text{O}_3\text{Na}$ $[\text{M}+\text{Na}]^+$ 441.1461, found 441.1453.

6'-(3,5-Dimethylphenyl)-1'-phenyl-1',3',3a',6a'-tetrahydro-4'H-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3p**): White solid, 25.4 mg, 59% yield (with **C5**, 29.3 mg, 78% yield). m.p. 194~196 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.77~7.72 (m, 1H), 7.71~7.66 (m, 1H), 7.66~7.56 (m, 2H), 6.99~6.89 (m, 5H), 6.76 (s, 1H), 6.71 (s, 2H), 6.27 (d, $J=1.2$ Hz, 1H), 4.63~4.59 (m, 1H), 3.81 (td, $J=9.6, 7.6$ Hz, 1H), 3.32 (d, $J=9.6$ Hz, 1H), 2.30 (dd, $J=13.6, 9.6$ Hz, 1H), 2.17 (dd, $J=13.6, 9.6$ Hz, 1H), 1.92 (s, 6H); ^{13}C NMR (150 MHz, CDCl_3) δ : 208.9, 203.0, 200.5, 176.8, 141.5, 137.7, 137.2, 135.8, 135.6, 132.5, 132.3, 128.6, 128.3, 127.4, 126.2, 125.3, 123.1, 123.0, 69.3, 57.9, 52.89, 51.6, 34.5, 20.8; HRMS calcd for $\text{C}_{30}\text{H}_{24}\text{O}_3\text{Na}$ $[\text{M}+\text{Na}]^+$ 455.1618, found 455.1622.

6'-(3-Methoxyphenyl)-1'-phenyl-1',3',3a',6a'-tetrahydro-4'H-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3q**): White semi-solid, 27.1 mg, 62% yield (with **C5**, 34.4 mg, 79% yield); ^1H NMR (400 MHz, CDCl_3) δ : 7.81~7.78 (m, 1H), 7.78~7.75 (m, 1H), 7.74~7.65 (m, 2H), 7.05~6.94 (m, 6H), 6.92~6.85 (m, 1H), 6.78~6.75 (m, 1H), 6.65~6.67 (m, 1H), 6.35 (d, $J=1.2$ Hz, 1H), 4.73~4.68 (m, 1H), 3.91 (td, $J=9.6, 7.2$ Hz, 1H), 3.45 (d, $J=9.6$ Hz, 1H), 3.39 (s, 3H), 2.39 (dd, $J=13.2, 9.6$ Hz, 1H), 2.26 (dd, $J=13.2, 9.6$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 208.8, 202.9, 200.5, 176.4, 159.2, 141.5, 137.0, 135.8, 135.7, 134.1, 129.2, 128.5, 128.3, 127.4, 126.4, 123.1, 120.6, 117.6, 112.3, 69.4, 57.7, 54.9, 52.6, 51.8, 34.4; HRMS calcd. for $\text{C}_{29}\text{H}_{22}\text{O}_4\text{Na}$ $[\text{M}+\text{Na}]^+$ 457.1410, found 457.1411.

6'-(4-Methoxyphenyl)-1'-phenyl-1',3',3a',6a'-tetrahydro-

4'*H*-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3r**): White semi-solid, 21.5 mg, 50% yield; ¹H NMR (400 MHz, CDCl₃) δ: 7.84~7.80 (m, 1H), 7.79~7.75 (m, 1H), 7.73~7.64 (m, 2H), 7.15~7.09 (m, 2H), 7.05~6.95 (m, 5H), 6.59~6.49 (m, 2H), 6.27 (d, *J*=1.2 Hz, 1H), 4.72~4.63 (m, 1H), 3.93~3.83 (m, 1H), 3.70 (s, 3H), 3.41 (d, *J*=9.6 Hz, 1H), 2.39 (dd, *J*=13.4, 9.6 Hz, 1H), 2.27 (dd, *J*=13.4, 9.6 Hz, 1H); ¹³C NMR (150 MHz, CDCl₃) δ: 208.8, 203.1, 200.6, 176.0, 161.6, 141.5, 137.0, 135.8, 135.6, 129.9, 128.6, 128.5, 128.4, 127.4, 125.2, 124.0, 123.1, 123.1, 113.5, 69.3, 58.1, 55.2, 52.6, 51.7, 34.5; HRMS calcd for C₂₆H₁₈O₄Na [M+Na]⁺ 417.1097, found 417.1094.

6'-(Naphthalen-2-yl)-1'-phenyl-1',3',3a',6a'-tetrahydro-4'*H*-spiro[indene-2,2'-pentalene]-1,3,4'-trione (**3s**): White solid, 25.1 mg, 55% yield (with **C5**, 30.6 mg, 79 % yield). m.p. 205~207 °C; ¹H NMR (400 MHz, CDCl₃) δ: 7.86~7.82 (m, 1H), 7.77~7.74 (m, 1H), 7.73~7.62 (m, 4H), 7.54 (dd, *J*=8.8, 2.0 Hz, 1H), 7.44~7.40 (m, 1H), 7.32~7.28 (m, 2H), 7.09~7.05 (m, 3H), 7.03~7.03 (m, 3H), 6.52 (d, *J*=1.2 Hz, 1H), 4.84~4.80 (m, 1H), 3.96 (td, *J*=9.6, 7.2 Hz, 1H), 3.50 (d, *J*=9.6 Hz, 1H), 2.44 (dd, *J*=13.6, 9.6 Hz, 1H), 2.31 (dd, *J*=13.6, 9.6 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 208.9, 203.1, 200.6, 175.7, 141.6, 141.5, 137.4, 135.9, 135.7, 134.2, 132.4, 129.7, 129.5, 128.8, 128.7, 128.6, 128.0, 127.6, 127.6, 127.4, 126.4, 126.1, 124.6, 123.2, 123.1, 69.4, 57.9, 52.8, 51.9, 34.7; HRMS calcd for C₃₂H₂₂O₃Na [M+Na]⁺ 477.1461, found 477.1473.

1-Benzoyl-1'-benzyl-6-phenyl-1,3,3a,6a-tetrahydro-4*H*-spiro[pentalene-2,3'-pyrrolo[2,3-*b*]pyridine]-2',4'(1'*H*)-dione (**5**): White solid, 24.8 mg, 50% yield. m.p. 210~213 °C; 3 : 1 *dr*; ¹H NMR (400 MHz, CDCl₃) δ (major): 8.09 (dd, *J*=5.4, 1.6 Hz, 1H), 7.56~7.49 (m, 3H), 7.39~7.28 (m, 3H), 7.24~7.18 (m, 2H), 7.17~7.03 (m, 8H), 6.97 (dd, *J*=7.4, 5.3 Hz, 1H), 6.44 (d, *J*=1.7 Hz, 1H), 5.17 (td, *J*=7.2, 1.6 Hz, 1H), 4.55 (d, *J*=14.8 Hz, 1H), 4.43 (d, *J*=14.8 Hz, 1H), 4.31 (d, *J*=7.2 Hz, 1H), 3.58 (m, 1H), 2.46 (dd, *J*=13.0, 10.6 Hz, 1H), 2.27 (dd, *J*=13.0, 8.8 Hz, 1H); δ (minor): 8.02 (dd, *J*=5.4, 1.6 Hz, 1H), 7.48~7.39 (m, 3H), 7.39~7.28 (m, 3H), 7.24~7.18 (m, 2H), 7.17~7.03 (m, 8H), 6.69~6.62 (m, 1H), 6.53 (d, *J*=1.5 Hz, 1H), 5.06~4.98 (m, 3H), 4.47 (d, *J*=8.8 Hz, 1H), 3.80 (dt, *J*=10.0, 7.8 Hz, 1H), 2.80 (dd, *J*=13.5, 8.0 Hz, 1H), 2.39 (dd, *J*=13.6, 9.8 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ (major): 207.1, 198.2, 176.2, 175.8, 155.0, 147.6, 136.3, 135.8, 133.44, 132.9, 132.5, 131.3, 129.00, 128.4, 128.3, 127.9, 127.9, 127.8, 127.4, 127.2, 126.37, 123.8, 118.3, 61.3, 56.7, 51.4, 48.9, 42.7, 39.9; δ (minor): 208.8, 199.1, 179.3, 171.1, 156.0, 147.6, 137.8, 136.6, 133.5, 133.3, 132.8, 130.9, 128.8, 128.7, 128.6, 128.1, 128.1, 127.7, 127.5, 127.2, 121.8, 118.5, 60.6, 53.4, 52.0, 51.8, 50.6, 34.7; HRMS calcd for C₃₄H₂₆N₂O₃Na [M+Na]⁺ 533.1836, found 533.1837.

Supporting Information More condition screenings, NMR spectra of compounds **3a**~**3s** and **5** as well as the X-ray data for racemic product **3a**. The supporting information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Moyano, A.; Rios, R. *Chem. Rev.* **2011**, *111*, 4703.
(b) Volla, C. M. R.; Atodiresei, I.; Rueping, M. *Chem. Rev.* **2014**, *114*, 2390.
(c) Ishihara, K.; Nakano, K. *J. Am. Chem. Soc.* **2007**, *129*, 8930.
(d) Davis, R. L.; Jørgensen, K. A. *J. Am. Chem. Soc.* **2012**, *134*, 2543.
(e) Talavera, G.; Reyes, E.; Vicario, J. L.; Carrillo, L. *Angew. Chem., Int. Ed.* **2012**, *51*, 4104.
(f) Akula, P. S.; Hong, B.-C.; Lee, G.-H. *Org. Lett.* **2018**, *20*, 7835.
(g) Lou, Y.-P.; Zheng, C.-W.; Pan, R.-M.; Jin, Q.-W.; Zhao, G.; Li, Z. *Org. Lett.* **2015**, *17*, 688.
(h) Yuan, Z.; Wei, W.; Lin, A.; Yao, H. *Org. Lett.* **2016**, *18*, 3370.
(i) Sun, X.-X.; Li, C.; He, Y.-Y.; Zhu, Z.-Q.; Mei, G.-J.; Shi, F. *Adv. Synth. Catal.* **2017**, *359*, 2660.
(j) Yang, S.; Lu, D.-F.; Huo, H.-R.; Luo, F.; Gong, Y.-F. *Org. Lett.* **2018**, *20*, 6943.
(k) Huo, H.-R.; A, R.-N.; Gong, Y.-F. *J. Org. Chem.* **2019**, *84*, 2093.
(l) Yao, W.; Dou, X.; Lu, Y. *J. Am. Chem. Soc.* **2015**, *137*, 54.
(m) Zhou, R.; Xiao, W.; Yin, X.; Chen, Y.-C. *Acta Chim. Sinica* **2014**, *72*, 862 (in Chinese).
(周容, 肖微, 尹祥, 陈应春, 化学学报, **2014**, *72*, 862.)
- [2] (a) Wu, T.-C.; Houk, K. N. *J. Am. Chem. Soc.* **1985**, *107*, 5308.
(b) Hong, B.-C.; Shr, Y.-J.; Wu, J.-L.; Gupta, A. K.; Lin, K.-J. *Org. Lett.* **2002**, *4*, 2249.
(c) Hayashi, Y.; Gotoh, H.; Honma, M.; Sankar, K.; Kumar, I.; Ishikawa, H.; Konno, K.; Yui, H.; Tsuzuki, S.; Uchimaru, T. *J. Am. Chem. Soc.* **2011**, *133*, 20175.
(d) Bertuzzi, G.; Thøgersen, M. K.; Giardinetti, M.; Vidal-Albalat, A.; Simon, A.; Houk, K. N.; Jørgensen, K. A. *J. Am. Chem. Soc.* **2019**, *141*, 3288.
- [3] Zhou, Z.; Wang, Z.-X.; Zhou, Y.-C.; Xiao, W.; Ouyang, Q.; Du, W.; Chen, Y.-C. *Nat. Chem.* **2017**, *9*, 590.
- [4] (a) Carrie, E. A.; Alpay, D.; Scott, J. M. *Tetrahedron* **2009**, *65*, 4069.
(b) Kishor C. B. *RSC Adv.* **2015**, *5*, 75923.
(c) Xie, P.-Z.; Huang, Y. *Eur. J. Org. Chem.* **2013**, 6213.
(d) Ni, H.-Z.; Chan, W.-L.; Lu, Y.-X. *Chem. Rev.* **2018**, *118*, 9344.
(e) Fan, Y.-C.; Ohyun K. *Chem. Commun.* **2013**, *49*, 11588.
(f) Methot, J. L.; Roush, W. R. *Adv. Synth. Catal.* **2004**, *346*, 1035.
(g) Gong, J.-J.; Li, T.-Z.; Pan, K.; Wu, X.-Y. *Chem. Commun.* **2011**, *47*, 1491.
(h) Takizawa, S.; Nguyen, T. M.; Grossmann, A.; Enders, D.; Sasai, H. *Angew. Chem., Int. Ed.* **2012**, *51*, 5423.
(i) Su, X.; Zhou, W.; Li, Y.-Y.; Zhang, J.-L. *Angew. Chem., Int. Ed.* **2015**, *54*, 6874.
(j) Zhang, X.-Z.; Gan, K.-J.; Liu, X.-X.; Deng, Y.-H.; Wang, F.-X.; Yu, K.-Y.; Zhang, J.; Fan, C.-A. *Org. Lett.* **2017**, *19*, 3207.
(k) Jin, Z.; Yang, R.; Du, Y.; Tivari, B.; Ganguly, R.; Chi, Y. R. *Org. Lett.* **2012**, *14*, 3226.
(l) Shi, Z.; Yu, P.; Loh, T. P.; Zhong, G.-F. *Angew. Chem., Int. Ed.* **2012**, *51*, 7825.
(m) Dong, X.; Liang, L.; Li, E.; Huang, Y. *Angew. Chem., Int. Ed.* **2015**, *54*, 1621.
(n) Zhou, W.; Su, X.; Tao, M.; Zhu, C.; Zhao, Q.; Zhang, J.-L. *Angew. Chem., Int. Ed.* **2015**, *54*, 14853.
(o) Li, S.; Liu, Y.; Huang, B.; Zhou, T.; Tao, H.; Xiao, Y.; Liu, L.; Zhang, J.-L. *ACS Catal.* **2017**, *7*, 2805.
(p) Qin, C.; Liu, Y.; Yu, Y.; Fu, Y.; Li, H.; Wang, W. *Org. Lett.* **2018**, *20*, 1304.
(q) Wu, X.; Zhou, L.; Maiti, R.; Mou, C.; Pan, L.; Chi, Y. R. *Angew. Chem., Int. Ed.* **2019**, *58*, 477.
(r) Zhou, W.; Gao, L.-H.; Tao, M.-N.; Su, X.; Zhao, Q.-J.; Zhang, J.-L. *Acta Chim. Sinica* **2016**, *74*, 800 (in Chinese).
(周伟, 高利华, 陶梦娜, 张俊良, 化学学报, **2016**, *74*, 800.)
- [5] McDougal, N. T.; Schaus, S. E. *Angew. Chem., Int. Ed.* **2006**, *45*, 3117.

- [6] (a) Gu, J.; Xiao, B.-X.; Chen, Y.-R.; Li, Q.-Z.; Ouyang, Q.; Du, W.; Chen, Y.-C. *Org. Lett.* **2018**, *20*, 2088.
(b) Gu, J.; Xiao, B.-X.; Ouyang, Q.; Du, W.; Chen, Y.-C. *Chin. J. Chem.* **2019**, *37*, 155.
- [7] Thalji, R. K.; Roush, W. R. *J. Am. Chem. Soc.* **2005**, *127*, 16778.
- [8] (a) Johnston, C. P.; Kothari, A.; Sergeieva, T.; Okovytyy, S. I.; Jackson, K. E.; Paton, R. S.; Smith, M. D. *Nat. Chem.* **2015**, *7*, 171.
(b) Chatgililoglu, C.; Ferreri, C.; Guerra, M.; Timokhin, V.; Froudakis, G.; Gimisis, T. *J. Am. Chem. Soc.* **2002**, *124*, 10765.
(c) Caldwell, J. J.; Craig, D. *Angew. Chem., Int. Ed.* **2007**, *46*, 2631.
(d) Auvray, P.; Knochel, P.; Normant, J. F. *Tetrahedron Lett.* **1985**, *26*, 4455.
- [9] (a) Brenninger, C.; Pöthig, A.; Bach, T. *Angew. Chem., Int. Ed.* **2017**, *56*, 4337.
(b) Wenkert, E.; Schorp, M. K. *J. Org. Chem.* **1994**, *59*, 1943.
(c) Fu, X.; Zhang, S.; Yin, J.; McAllister, T. L.; Jiang, S. A.; Tann, C.-H.; Thiruvengadam, T. K.; Zhang, F. *Tetrahedron Lett.* **2002**, *43*, 573.
(d) Bugarin, A.; Jones, K. D.; Connell, B. T. *Chem. Commun.* **2010**, *46*, 1715.
- [10] (a) Lee, C.-J.; Sheu, C.-C.; Tasi, C.-C.; Wu, Z.-Z.; Lin, W. *Chem. Commun.* **2014**, *50*, 5304.
(b) Hassanein, A. Z. *Synth. Commun.* **2000**, *30*, 3883.

(Cheng, F.)