高烯丙基酰肼氧化/卤胺环化串联反应合成吡唑啉类化合物

李鹏飞 黄丹凤 杨天宇 邓周斌 王克虎
王君姣 苏瀛鹏 胡雨来*
(西北师范大学化学化工学院 兰州 730070)

摘要 探索了以 *N*-碘代丁二酰亚胺(NIS)作为卤化试剂, 通过高烯丙基酰肼化合物的串联氧化/卤胺环化反应合成吡唑啉类化合物. 该方法反应条件温和, 不需要添加剂, 以良好的收率得到吡唑啉类化合物.

关键词 氧化/卤胺环化; 吡唑啉; 高烯丙基酰肼; *N*-碘代丁二酰亚胺

Synthesis of Pyrazolines by Cascade Oxidation/Halogenoamino-cyclization Reaction of Homoallylhydrazides

Li, Pengfei Huang, Danfeng Yang, Tianyu Deng, Zhoubin Wang, Kehu
Wang, Junjiao Su, Yingpeng Hu, Yulai*
(College of Chemistry and Chemical Engineering, Northwest Normal University, Lanzhou 730070)

Abstract An efficient cascade oxidation/halogenoaminocyclization reaction of homoallylhydrazides with *N*-halogensuccinimide was developed without addition of other additives under mild reaction conditions to provide pyrazolines. Under the optimized conditions, a variety of highly substituted pyrazoline derivatives were obtained.

Keywords oxidation/halogenoaminocyclization; pyrazolines; homoallylhydrazides; *N*-halogensuccinimide

1 Introduction

Nitrogen-containing heterocycles are vastly distributed in natural products, drugs and agricultural chemicals.^[1] As a class of important heterocycle, pyrazoline derivatives show broad bioactivities such as anticancer,^[2] antidepressant,^[3] anticonvulsant,^[4] antibacterial^[5] and antimalarial activities.^[6] Furthermore, they also attracted attentions in material science because of their luminescent and nonlinear optical properties.^[7] In view of their broad applications, many efforts have been devoted to the synthesis of the pyrazoline derivatives over the past decades. Typically, pyrazolines were synthesized by aza-Michael addition/cyclocondensation reactions of hydrazines with α,β -unsaturated aldehydes or ketones,^[8] 1,3-dipolar cycloaddition of diazoalkanes or nitrilimines with alkenes^[9] and cyclocondensation of hydrazines with dihalides or ditosylates.^[10] As an easy available 1,2-dinitro building blocks, hydrazones were also used to construct the pyrazolines.^[11~14] For instance, [4+1]-annulations of α -chloro acylhydrazone with maleimides, diazo esters or sulfur ylides have been devel-

oped to synthesize pyrazolines.^[12] A visible-light photocatalytic transformation of β,γ -unsaturated hydrazones into pyrazolines was also reported recently.^[13] In our previous works on synthesis of trifluoromethyl compounds, we found that trifluoromethylated homoallylic *N*-acylhydrazines could be conveniently transformed into trifluoromethyl pyrazolines by halogenation reagents such as *N*-iodosuccinimide (NIS), *N*-bromosuccinimide (NBS) and *N*-chlorosuccinimide (NCS).^[15] Herein, we report that this process could also be applied to non-trifluoromethylated homoallylhydrazides to construct pyrazolines in good to excellent yields (Scheme 1).

2 Results and discussion

Based on our previous reaction conditions,^[15] *N*-(1-phenylbut-3-en-1-yl)benzohydrazide (**1a**) was taken as substrate to react with NIS (**2a**) in tetrahydrofuran (THF) at room temperature. After 4 h, *N*-benzoyl-5-iodomethyl-3-phenylpyrazoline (**3a**) was obtained in 91% yield (Table 1). In order to investigate the generalities of the substrates,

* Corresponding author. E-mail: huyul@nwnu.edu.cn

Received March 29, 2019; revised April 26, 2019; published online May 21, 2019.

Project supported by the National Natural Science Foundation of China (Nos. 21662030, 21861033).

国家自然科学基金(Nos. 21662030, 21861033)资助项目.

Scheme 1 Methods for synthesis of pyrazolines from homoallylhydrazides

Table 1 Substrate scope of the reaction^a

^a All reactions were carried out using 0.15 mmol of **1** and 4 equiv. of NIS in 5 mL of THF at r.t. for 4 h. Isolated yields.

the effects of different *N*-acyl groups in homoallylic acylhydrazides **1** were examined. The results in Table 1 showed that both of the aromatic and aliphatic acyl groups were suitable for the reaction, but the substrate with aromatic acyl groups gave the corresponding products in higher yield than that with aliphatic acyl groups.

Next, the influence of *R*² group of homoallylhydrazides **1** on the reaction was studied (Table 2). The homoallylhydrazides **1** with different *R*² groups were taken as substrates to react with NIS under the standard reaction conditions. When *R*² groups were aromatic groups, all the corresponding products obtained in high yields whenever the substitutes on phenyl rings were electron-withdrawing groups or electron-donating groups (**3aa**~**3ai**). However, the yields of the products decreased when *R*² groups were aliphatic groups (**3aj**~**3am**).

After the investigation of the effects of *R*¹ and *R*² groups on the reaction separately, their combined influences on the reaction were checked further. As shown in Table 3, when both *R*¹ and *R*² groups were aromatic groups, the corresponding products were obtained in good to excellent yields (**3ba**~**3fa**). Especially, when *R*¹ groups were electron-poor aromatic groups, the reactions gave the corresponding products in higher yields than those with elec-

Table 2 Substrate scope of the reaction^a

3aa: *R*² = 2-FC₆H₄, 87%; **3ab:** *R*² = 3-FC₆H₄, 92%
3ac: *R*² = 4-FC₆H₄, 95%; **3ad:** *R*² = 2-CH₃C₆H₄, 83%
3ae: *R*² = 3-CH₃C₆H₄, 86%; **3af:** *R*² = 4-CH₃C₆H₄, 90%
3ag: *R*² = 4-ClC₆H₄, 93%; **3ah:** *R*² = 4-CF₃C₆H₄, 96%
3ai: *R*² = 4-CH₃OC₆H₄, 86%; **3aj:** *R*² = Methyl, 49%
3ak: *R*² = *n*-Heptyl, 77%; **3al:** *R*² = Cyclohexyl, 53%
3am: *R*² = Cinnamyl, 50%

^a All reactions were carried out using 0.15 mmol of **1** and 4 equiv. of NIS in 5 mL of THF at r.t. for 4 h. Isolated yields.

tron-rich aromatic groups. If *R*¹ groups were electron-rich aromatic groups and *R*² groups were electron-poor aromatic groups, the yields of products were lower (**3ea** and **3fa**). When one or both of *R*¹ or *R*² were aliphatic groups, the yields of the corresponding products decreased. As an extreme example, there was no product formed when *R*¹ and *R*² groups were sterically hindered isopropyl groups (**3la**).

Next, the influences of halogen sources such as NIS (**2a**), NBS (**2b**) and NCS (**2c**) on the reaction were investigated. The results were collected in Table 4. Among the three halogen sources, NIS gave the products in the highest yields and NCS gave the products in the lowest yields when the same substrates were tested.

To demonstrate the synthetic utility of this methodology, a gram-scale reaction was performed by using homoallylhydrazide **1a** as substrate under the standard reaction conditions (Eq. 1). The reaction proceeded very well to give the product **3a** in 85% yield. More importantly, these dihydropyrazole products could be easily converted into pyrazoles through a simple operation. For example, *N*-free pyrazole **4** was obtained from **3a** in 88% yield in the presence of KOH under refluxing in ethanol for 30 min (Eq. 2).

The mechanism for the oxidative halocyclization of homoallylhydrazides with *N*-halogensuccinimide can refer to our previous work.^[15]

3 Conclusions

In conclusion, a novel and efficient cascade oxidative/iodoaminocyclization reaction of homoallylic *N*-acylhydrazides has been developed, which provides a concise method for the synthesis of potentially bioactive pyrazoline derivatives. This process tolerates a wide range of substrates to produce the products in good to excellent yields. Especially, the products possess a halogen group, which can be used for further transformation into more complex pyrazoline derivatives. Further investigations on this reaction for the synthesis of other heterocycles are underway in our laboratory.

Table 3 Substrate scope of the reaction^a

^a All reactions were carried out using 0.15 mmol of **1** and 4 equiv of NIS in 5 mL of THF at r.t. for 4 h. Isolated yields

Table 4 Substrate scope of the reaction^a

^a All reactions were carried out using 0.15 mmol of **1** and 4 equiv. of NXs in 5 mL of THF at r.t. for 4 h. Isolated yields

4 Experimental section

4.1 Materials and methods

All of the commercially available reagents and solvents were used without further purification. Homoallylhydrazides were prepared through condensation of hydrazides with aldehydes and allyl bromide according to the literature procedure.^[16] Flash chromatography was performed using silica gel 60 (230–400 mesh). Analytical thin layer chromatography (TLC) was conducted using silica gel GF254. TLC plates were analysed by an exposure to ultra-violet (UV) light and/or submersion in a phosphomolybdic acid solution or in Iodine vapour. ¹H NMR spectra were measured on 600 and 400 MHz spectrometers. ¹³C NMR spectra were measured on 150 and 100 MHz spectrometers.

4.2 General procedure for the synthesis of 3

Homoallylhydrazides **1** (0.15 mmol, 1 equiv.), NIS, NBS or NCS **2** (0.6 mmol, 4 equiv.) were placed in a dried round-bottom flask (50 mL) fitted with a magnetic bar. Tetrahydrofuran (5 mL) was then added. The mixture was stirred at room temperature. When the starting materials were consumed completely as monitored by TLC, the saturated Na₂S₂O₃ solution (10 mL) was poured into the mixture and stirred for 10 min. Tetrahydrofuran was removed under vacuum and the mixture was extracted with ethyl acetate (10 mL × 3). The combined organic phases were dried (MgSO₄) and concentrated. Purification of the residue by silica gel column chromatography (230 ~ 400 mesh) using ethyl acetate and petroleum ether as eluent furnished the pure products **3**.

(5-(Iodomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)(phenyl)methanone (**3a**): 53 mg, yield 91%; *R*_f=0.6 [*V*(ethylacetate) : *V*(petroleum ether)=1 : 3]; yellowish solid, m.p. 106~107 °C; ¹H NMR (600 MHz, CDCl₃) δ: 7.98 (d, *J*=7.2 Hz, 2H), 7.69 (dd, *J*=7.2, 1.2 Hz, 2H), 7.51~7.48 (m, 1H), 7.46~7.43 (m, 2H), 7.42~7.38 (m, 3H), 4.93~4.88 (m, 1H), 3.76 (dd, *J*=10.2, 6.6 Hz, 1H), 3.69 (dd, *J*=9.6, 2.4 Hz, 1H), 3.48 (dd, *J*=17.4, 10.8 Hz, 1H), 3.10 (dd, *J*=17.4, 5.4 Hz, 1H); ¹³C NMR (150 MHz, CDCl₃) δ: 167.1, 154.3, 134.1, 131.0, 131.0, 130.4, 130.0, 128.7, 127.6, 126.7, 57.9, 39.2, 9.6; HRMS (ESI) calcd for C₁₇H₁₆IN₂O [M+H]⁺ 391.0302; found 391.0298.

(2-Fluorophenyl)(5-(iodomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3b**): 54 mg, yield 88%; *R*_f=0.5 [*V*(ethylacetate) : *V*(petroleum ether)=1 : 10]; yellow oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.62~7.59 (m, 3H), 7.48~7.34 (m, 4H), 7.21 (t, *J*=7.6 Hz, 1H), 7.13 (t, *J*=8.8 Hz, 1H), 4.93~4.86 (m, 1H), 3.75~3.69 (m, 2H), 3.53 (dd, *J*=18.0, 11.2 Hz, 1H), 3.14 (dd, *J*=18.0, 4.8 Hz, 1H); ¹³C NMR (150 MHz, CDCl₃) δ: 164.2, 159.3 (d, *J*_{C-F}=250.1 Hz), 154.8, 131.9 (d, *J*_{C-F}=8.3 Hz), 130.8, 130.6, 130.0 (d, *J*_{C-F}=3.2 Hz), 128.6, 126.7, 124.0 (d, *J*_{C-F}=15.8 Hz), 123.7 (d, *J*_{C-F}=3.5 Hz), 115.6 (d, *J*_{C-F}=21.8 Hz), 57.8, 39.9, 8.7; HRMS (ESI) calcd for C₁₇H₁₄FI-N₂NaO [M+Na]⁺ 431.0027; found 431.0012.

(3-Fluorophenyl)(5-(iodomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3c**): 54 mg, yield 89%; *R*_f=0.5 [*V*(ethylacetate) : *V*(petroleum ether)=1 : 3]; white solid, m.p. 115~116 °C; ¹H NMR (600 MHz, CDCl₃) δ: 7.78 (d, *J*=7.2 Hz, 1H), 7.72 (d, *J*=9.6 Hz, 1H), 7.69 (d, *J*=7.8 Hz, 2H), 7.45~7.40 (m, 4H), 7.20 (t, *J*=8.4 Hz, 1H), 4.92~4.85 (m, 1H), 3.78~3.76 (m, 1H), 3.67 (d, *J*=9.0 Hz, 1H), 3.50 (dd, *J*=17.4, 10.8 Hz, 1H), 3.11 (dd, *J*=17.4, 4.8 Hz, 1H); ¹³C NMR (150 MHz, CDCl₃) δ: 165.5, 162.0 (d, *J*_{C-F}=247.2 Hz), 154.9, 136.1 (d, *J*_{C-F}=7.4 Hz), 130.8, 130.7, 129.3 (d, *J*_{C-F}=7.8 Hz), 128.8, 126.8, 125.8 (d, *J*_{C-F}=3.0 Hz), 118.1 (d, *J*_{C-F}=21.2 Hz), 117.2 (d, *J*_{C-F}=23.6 Hz), 57.9, 39.2, 9.3; HRMS (ESI) calcd for C₁₇H₁₄FIN₂NaO [M+Na]⁺ 431.0027; found 431.0021.

(4-Fluorophenyl)(5-(iodomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3d**): 58 mg, yield 94%; *R*_f=0.55 [*V*(ethylacetate) : *V*(petroleum ether)=1 : 5]; white solid, m.p. 114~115 °C; ¹H NMR (600 MHz, CDCl₃) δ: 8.04 (dd, *J*=8.4, 5.4 Hz, 2H), 7.68 (d, *J*=7.8 Hz, 2H), 7.45~7.40 (m, 3H), 7.12 (t, *J*=8.4 Hz, 2H), 4.91~4.87 (m, 1H), 3.76 (dd, *J*=10.2, 7.2 Hz, 1H), 3.68 (dd, *J*=10.2, 2.4 Hz, 1H), 3.49 (dd, *J*=18.0, 11.4 Hz, 1H), 3.10 (dd, *J*=18.0, 5.4 Hz, 1H); ¹³C NMR (150 MHz, CDCl₃) δ: 165.8, 164.4 (d, *J*_{C-F}=250.4 Hz), 154.6, 132.6 (d, *J*_{C-F}=8.7 Hz), 130.9, 130.6, 130.1 (d, *J*_{C-F}=3.2 Hz), 128.7, 126.7, 114.7 (d, *J*_{C-F}=21.6 Hz), 57.9, 39.2, 9.5; HRMS (ESI) calcd for C₁₇H₁₅FIN₂O [M+H]⁺ 409.0208; found 409.0195.

(5-(Iodomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)(*o*-tolyl)methanone (**3e**): 47 mg, yield 78%; *R*_f=0.4 [*V*(ethylacetate) : *V*(petroleum ether)=1 : 10]; yellow oil. ¹H NMR (400 MHz, CDCl₃) δ: 7.57 (d, *J*=6.4 Hz, 2H), 7.43 (d, *J*=6.8 Hz, 1H), 7.38~7.31 (m, 4H), 7.23 (t, *J*=5.6 Hz, 2H), 4.92~4.85 (m, 1H), 3.84 (dd, *J*=10.0, 6.4 Hz, 1H), 3.70 (dd, *J*=10.0, 2.4 Hz, 1H), 3.50 (dd, *J*=18.0, 11.2 Hz, 1H), 3.11 (dd, *J*=18.0, 5.2 Hz, 1H), 2.41 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 168.9, 154.5, 135.6, 135.5, 130.8, 130.4, 130.1, 129.3, 128.6, 128.0, 126.6, 125.1, 57.3, 39.7, 20.0, 9.4; HRMS (ESI) calcd for C₁₈H₁₇IN₂NaO [M+Na]⁺ 427.0278; found 427.0286.

(5-(Iodomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)(*m*-tolyl)methanone (**3f**): 48 mg, yield 80%; *R*_f=0.4 [*V*(ethylacetate) : *V*(petroleum ether)=1 : 10]; yellow oil. ¹H NMR (600 MHz, CDCl₃) δ: 7.78 (d, *J*=8.4 Hz, 2H), 7.69~7.67 (m, 2H), 7.43~7.38 (m, 3H), 7.35~7.30 (m, 2H), 4.93~4.88 (m, 1H), 3.75 (dd, *J*=10.2, 7.2 Hz, 1H), 3.70 (dd, *J*=10.2, 3.0 Hz, 1H), 3.49 (dd, *J*=17.4, 10.8 Hz, 1H), 3.11 (dd, *J*=17.4, 4.8 Hz, 1H), 2.42 (s, 3H); ¹³C NMR (150 MHz, CDCl₃) δ: 167.3, 154.2, 137.4, 134.0, 131.8, 131.0, 130.5, 130.4, 128.7, 127.5, 127.2, 126.7, 57.9, 39.2, 21.4, 9.6; HRMS (ESI) calcd for C₁₈H₁₇IN₂NaO [M+Na]⁺ 427.0278; found 427.0279.

(5-(Iodomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)(*p*-tolyl)methanone (**3g**): 51 mg, yield 84%; *R*_f=0.45 [*V*(ethylacetate) : *V*(petroleum ether)=1 : 10]; yellow oil. ¹H NMR (600 MHz, CDCl₃) δ: 7.91 (d, *J*=8.4 Hz, 2H), 7.69 (dd, *J*=7.2, 3.0 Hz, 2H), 7.42~7.39 (m, 3H), 7.25 (d, *J*=7.2 Hz, 2H), 4.92~4.88 (m, 1H), 3.73 (dd, *J*=10.2,

6.6 Hz, 1H), 3.70 (dd, $J=9.6$, 3.0 Hz, 1H), 3.47 (dd, $J=17.4$, 10.8 Hz, 1H), 3.09 (dd, $J=18.0$, 5.4 Hz, 1H), 2.42 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 167.0, 154.1, 141.5, 131.1, 131.0, 130.4, 130.2, 128.7, 128.4, 126.7, 58.0, 39.1, 21.5, 9.7; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{17}\text{IN}_2\text{NaO}$ [$\text{M}+\text{Na}$] $^{+}$ 427.0278; found 427.0266.

(4-Chlorophenyl)(5-(iodomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3h**): 55 mg, yield 86%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 6$]; yellow oil. ^1H NMR (600 MHz, CDCl_3) δ : 7.94 (d, $J=8.4$ Hz, 2H), 7.68 (d, $J=7.2$ Hz, 2H), 7.45~7.38 (m, 5H), 4.91~4.85 (m, 1H), 3.76 (dd, $J=10.2$, 6.6 Hz, 1H), 3.68~3.66 (m, 1H), 3.51~3.45 (m, 1H), 3.13~3.08 (m, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 165.9, 154.8, 137.2, 132.4, 131.5, 130.8, 130.6, 128.7, 127.9, 126.7, 57.9, 39.2, 9.4; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{14}\text{ClIN}_2\text{NaO}$ [$\text{M}+\text{Na}$] $^{+}$ 446.9732; found 446.9751.

Furan-2-yl(5-(iodomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3i**): 49 mg, yield 86%; $R_f=0.6$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 5$]; yellow solid, m.p. 168~169 $^{\circ}\text{C}$; ^1H NMR (600 MHz, CDCl_3) δ : 7.78 (t, $J=9.6$ Hz, 2H), 7.65~7.63 (m, 2H), 7.47~7.46 (m, 3H), 6.57 (t, $J=1.8$ Hz, 1H), 4.92~4.89 (m, 1H), 3.71 (dd, $J=9.6$, 2.4 Hz, 1H), 3.63 (dd, $J=9.6$, 7.8 Hz, 1H), 3.48 (dd, $J=18.0$, 10.8 Hz, 1H), 3.11 (dd, $J=18.0$, 3.6 Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 156.3, 155.1, 145.8, 145.6, 130.9, 130.7, 128.8, 126.7, 119.5, 111.7, 58.2, 39.0, 9.0; HRMS (ESI) calcd for $\text{C}_{15}\text{H}_{13}\text{IN}_2\text{NaO}_2$ [$\text{M}+\text{Na}$] $^{+}$ 402.9914; found 402.9932.

(5-(Iodomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)(naphthalen-2-yl)methanone (**3j**): 57 mg, yield 86%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 5$]; yellow solid, m.p. 165~166 $^{\circ}\text{C}$; ^1H NMR (600 MHz, CDCl_3) δ : 8.55 (s, 1H), 8.05 (dd, $J=8.4$, 1.2 Hz, 1H), 7.96 (d, $J=8.4$ Hz, 1H), 7.90~7.87 (m, 2H), 7.69 (dd, $J=7.2$, 1.2 Hz, 2H), 7.59~7.53 (m, 2H), 7.43~7.39 (m, 3H), 4.99~4.94 (m, 1H), 3.75 (dd, $J=10.2$, 6.6 Hz, 1H), 3.70 (dd, $J=10.2$, 3.0 Hz, 1H), 3.49 (dd, $J=17.4$, 10.8 Hz, 1H), 3.11 (dd, $J=17.4$, 5.4 Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 165.8, 165.2, 163.5, 154.6, 132.6, 132.5, 130.8, 130.6, 130.1, 130.0, 128.7, 126.7, 114.8, 114.6, 57.9, 39.2, 9.5; HRMS (ESI) calcd for $\text{C}_{21}\text{H}_{17}\text{IN}_2\text{NaO}$ [$\text{M}+\text{Na}$] $^{+}$ 463.0278; found 463.0255.

1-(5-(Iodomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)ethan-1-one (**3k**): 37 mg, yield 75%; $R_f=0.6$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 5$]; colorless oil. ^1H NMR (600 MHz, CDCl_3) δ : 7.73~7.71 (m, 2H), 7.42~7.40 (m, 3H), 4.70~4.66 (m, 1H), 3.60 (dd, $J=10.2$, 3.0 Hz, 1H), 3.52 (dd, $J=10.2$, 7.2 Hz, 1H), 3.43 (dd, $J=18.0$, 10.8 Hz, 1H), 3.04 (dd, $J=18.0$, 4.8 Hz, 1H), 2.39 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 169.5, 153.6, 131.0, 130.4, 128.7, 126.6, 57.0, 39.9, 21.9, 9.4; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{14}\text{IN}_2\text{O}$ [$\text{M}+\text{H}$] $^{+}$ 329.0145; found 329.0142.

1-(5-(Iodomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)-2-methylpropan-1-one (**3l**): 36 mg, yield 68%; $R_f=0.6$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 5$]; yellowish oil. ^1H NMR (600 MHz, CDCl_3) δ : 7.73~7.71 (m, 2H),

7.42~7.40 (m, 3H), 4.67~4.63 (m, 1H), 3.61 (dd, $J=10.2$, 6.6 Hz, 1H), 3.56 (dd, $J=10.2$, 3.0 Hz, 1H), 3.49~3.45 (m, 1H), 3.41 (dd, $J=17.4$, 10.8 Hz, 1H), 3.02 (dd, $J=17.4$, 4.8 Hz, 1H), 1.25 (d, $J=6.6$ Hz, 3H), 1.18 (d, $J=6.6$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 176.0, 153.3, 131.2, 130.3, 128.7, 126.5, 56.9, 39.5, 32.0, 19.4, 18.5, 9.7; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{18}\text{IN}_2\text{O}$ [$\text{M}+\text{H}$] $^{+}$ 357.0458; found 357.0458.

Cyclohexyl(5-(iodomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3m**): 46 mg, yield 78%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 4$]; white solid, m.p. 116~117 $^{\circ}\text{C}$; ^1H NMR (600 MHz, CDCl_3) δ : 7.74~7.72 (m, 2H), 7.45~7.41 (m, 3H), 4.66~4.62 (m, 1H), 3.62 (dd, $J=10.2$, 7.2 Hz, 1H), 3.55 (dd, $J=10.2$, 2.4 Hz, 1H), 3.41 (dd, $J=17.4$, 10.8 Hz, 1H), 3.22~3.17 (m, 1H), 3.01 (dd, $J=17.4$, 4.8 Hz, 1H), 2.00 (d, $J=14.4$ Hz, 1H), 1.84~1.80 (m, 3H), 1.71 (d, $J=13.8$ Hz, 1H), 1.57~1.48 (m, 2H), 1.43~1.23 (m, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 175.2, 153.2, 131.3, 130.3, 128.7, 126.5, 56.8, 41.8, 39.5, 29.3, 28.4, 26.0, 25.9, 25.7, 9.9; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{21}\text{IN}_2\text{NaO}$ [$\text{M}+\text{Na}$] $^{+}$ 419.0591; found 419.0586.

1-(5-(Iodomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)dodecan-1-one (**3n**): 51 mg, yield 73%; $R_f=0.6$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 4$]; white solid, m.p. 120~121 $^{\circ}\text{C}$; ^1H NMR (600 MHz, CDCl_3) δ : 7.74~7.71 (m, 2H), 7.44~7.40 (m, 3H), 4.70~4.66 (m, 1H), 3.61~3.55 (m, 2H), 3.43 (dd, $J=18.0$, 11.4 Hz, 1H), 3.04 (dd, $J=18.0$, 5.4 Hz, 1H), 2.83~2.78 (m, 1H), 2.70~2.65 (m, 1H), 1.74~1.69 (m, 2H), 1.41~1.37 (m, 2H), 1.34~1.25 (m, 15H), 0.87 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 172.5, 153.3, 131.2, 130.3, 128.7, 126.5, 57.0, 39.7, 34.2, 31.9, 30.9, 29.7, 29.6, 29.5, 29.4, 29.3, 25.0, 22.7, 14.1, 9.6; HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{34}\text{IN}_2\text{O}$ [$\text{M}+\text{H}$] $^{+}$ 469.1710; found 469.1690.

1-(5-(Iodomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)octadecan-1-one (**3o**): 60 mg, yield 72%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; white solid, m.p. 118~119 $^{\circ}\text{C}$; ^1H NMR (600 MHz, CDCl_3) δ : 7.73~7.72 (m, 2H), 7.43~7.40 (m, 3H), 4.70~4.66 (m, 1H), 3.61~3.55 (m, 2H), 3.42 (dd, $J=18.0$, 11.4 Hz, 1H), 3.04 (dd, $J=18.0$, 5.4 Hz, 1H), 2.83~2.78 (m, 1H), 2.70~2.65 (m, 1H), 1.74~1.69 (m, 2H), 1.36~1.25 (m, 28H), 0.87 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 172.4, 153.2, 131.2, 130.3, 128.7, 126.5, 57.0, 39.7, 34.2, 31.9, 29.7, 29.7, 29.7, 29.6, 29.5, 29.4, 29.3, 25.0, 22.7, 14.1, 9.6; HRMS (ESI) calcd for $\text{C}_{28}\text{H}_{46}\text{IN}_2\text{O}$ [$\text{M}+\text{H}$] $^{+}$ 553.2649; found 553.2634.

(4-Fluorophenyl)(3-(2-fluorophenyl)-5-(iodomethyl)-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3aa**): 55 mg, yield 87%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 4$]; colorless oil. ^1H NMR (600 MHz, CDCl_3) δ : 8.04 (dd, $J=8.4$, 5.4 Hz, 2H), 7.80 (t, $J=7.2$ Hz, 1H), 7.40 (dd, $J=13.8$, 7.2 Hz, 1H), 7.18 (t, $J=7.8$ Hz, 1H), 7.14~7.11 (m, 3H), 4.88~4.84 (m, 1H), 3.81 (dd, $J=10.2$, 6.6 Hz, 1H), 3.63~3.61 (m, 1H), 3.60~3.57 (m, 1H), 3.22~3.17 (m, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ :

165.9, 164.4 (d, $J_{\text{C-F}}=250.5$ Hz), 161.2 (d, $J_{\text{C-F}}=252.5$ Hz), 151.5 (d, $J_{\text{C-F}}=2.9$ Hz), 132.6 (d, $J_{\text{C-F}}=8.9$ Hz), 132.1 (d, $J_{\text{C-F}}=8.6$ Hz), 130.0 (d, $J_{\text{C-F}}=3.2$ Hz), 128.9 (d, $J_{\text{C-F}}=3.0$ Hz), 124.5 (d, $J_{\text{C-F}}=3.5$ Hz), 119.0 (d, $J_{\text{C-F}}=10.9$ Hz), 116.6 (d, $J_{\text{C-F}}=22.0$ Hz), 114.7 (d, $J_{\text{C-F}}=21.6$ Hz), 57.8 (d, $J_{\text{C-F}}=2.7$ Hz), 41.5 (d, $J_{\text{C-F}}=7.8$ Hz), 9.7; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{14}\text{F}_2\text{IN}_2\text{O} [\text{M}+\text{H}]^+$ 427.0113; found 427.0113.

(4-Fluorophenyl)(3-(3-fluorophenyl)-5-(iodomethyl)-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3ab**): 59 mg, yield 92%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 4$]; yellow oil. ^1H NMR (600 MHz, CDCl_3) δ : 8.02 (dd, $J=11.4, 6.0$ Hz, 2H), 7.43 (d, $J=7.8$ Hz, 1H), 7.39 (dd, $J=12.6, 7.8$ Hz, 2H), 7.14~7.11 (m, 3H), 4.91~4.87 (m, 1H), 3.79 (dd, $J=10.2, 6.6$ Hz, 1H), 3.66 (dd, $J=10.2, 2.4$ Hz, 1H), 3.47 (dd, $J=18.0, 11.4$ Hz, 1H), 3.07 (dd, $J=18.0, 5.4$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 166.0, 164.4 (d, $J_{\text{C-F}}=250.5$ Hz), 162.8 (d, $J_{\text{C-F}}=245.4$ Hz), 153.5 (d, $J_{\text{C-F}}=3.0$ Hz), 133.1 (d, $J_{\text{C-F}}=7.9$ Hz), 132.5 (d, $J_{\text{C-F}}=8.7$ Hz), 130.4 (d, $J_{\text{C-F}}=8.5$ Hz), 129.9 (d, $J_{\text{C-F}}=3.3$ Hz), 122.5 (d, $J_{\text{C-F}}=2.9$ Hz), 117.5 (d, $J_{\text{C-F}}=21.3$ Hz), 114.8 (d, $J_{\text{C-F}}=21.6$ Hz), 113.4 (d, $J_{\text{C-F}}=22.8$ Hz), 58.0, 39.2, 9.5; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{13}\text{F}_2\text{IN}_2\text{NaO} [\text{M}+\text{Na}]^+$ 448.9933; found 448.9932.

(4-Fluorophenyl)(3-(4-fluorophenyl)-5-(iodomethyl)-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3ac**): 61 mg, yield 95%; $R_f=0.5$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 4$]; colorless oil. ^1H NMR (600 MHz, CDCl_3) δ : 8.02 (dd, $J=10.2, 5.4$ Hz, 2H), 7.68 (dd, $J=9.0, 4.8$ Hz, 2H), 7.14~7.09 (m, 4H), 4.92~4.87 (m, 1H), 3.77 (dd, $J=10.2, 6.6$ Hz, 1H), 3.67 (dd, $J=10.2, 2.4$ Hz, 1H), 3.47 (dd, $J=18.0, 11.4$ Hz, 1H), 3.08 (dd, $J=18.0, 5.4$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 165.9, 164.4 (d, $J_{\text{C-F}}=250.5$ Hz), 164.1 (d, $J_{\text{C-F}}=250.2$ Hz), 153.6, 132.5 (d, $J_{\text{C-F}}=8.7$ Hz), 130.1 (d, $J_{\text{C-F}}=3.2$ Hz), 128.6 (d, $J_{\text{C-F}}=8.6$ Hz), 127.2 (d, $J_{\text{C-F}}=3.3$ Hz), 116.0 (d, $J_{\text{C-F}}=21.8$ Hz), 114.8 (d, $J_{\text{C-F}}=20.1$ Hz), 58.0, 39.3, 9.6; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{13}\text{F}_2\text{IN}_2\text{NaO} [\text{M}+\text{Na}]^+$ 448.9933; found 448.9925.

(4-Fluorophenyl)(5-(iodomethyl)-3-(*o*-tolyl)-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3ad**): 52 mg, yield 83%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; colorless oil. ^1H NMR (600 MHz, CDCl_3) δ : 8.00 (dd, $J=9.0, 6.6$ Hz, 2H), 7.41 (d, $J=7.8$ Hz, 1H), 7.30 (t, $J=6.6$ Hz, 1H), 7.26 (t, $J=7.2$ Hz, 2H), 7.08 (t, $J=8.4$ Hz, 2H), 4.84~4.79 (m, 1H), 3.83 (dd, $J=10.2, 6.6$ Hz, 1H), 3.64~3.62 (m, 1H), 3.59~3.53 (m, 1H), 3.15~3.10 (m, 1H), 2.49 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 166.2, 164.3 (d, $J_{\text{C-F}}=250.2$ Hz), 155.4, 138.0, 132.3 (d, $J_{\text{C-F}}=8.7$ Hz), 131.9, 130.2 (d, $J_{\text{C-F}}=3.3$ Hz), 129.8, 129.7, 129.0, 126.0, 114.6 (d, $J_{\text{C-F}}=21.6$ Hz), 56.6, 41.4, 23.3, 9.9; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{16}\text{FIN}_2\text{NaO} [\text{M}+\text{Na}]^+$ 445.0184; found 445.0182.

(4-Fluorophenyl)(5-(iodomethyl)-3-(*m*-tolyl)-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3ae**): 54 mg, yield 86%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; colorless oil. ^1H NMR (600 MHz, CDCl_3) δ : 8.04 (dd, $J=9.0, 5.4$ Hz, 2H), 7.49 (d, $J=10.2$ Hz, 2H), 7.30 (t, $J=7.8$

Hz, 1H), 7.25 (s, 1H), 7.13 (d, $J=8.4$ Hz, 2H), 4.90~4.86 (m, 1H), 3.75 (dd, $J=10.2, 6.6$ Hz, 1H), 3.67 (dd, $J=10.2, 2.4$ Hz, 1H), 3.47 (dd, $J=18.0, 11.4$ Hz, 1H), 3.09 (dd, $J=18.0, 5.4$ Hz, 1H), 2.38 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 165.8, 164.4 (d, $J_{\text{C-F}}=251.4$ Hz), 154.9, 138.5, 132.6 (d, $J_{\text{C-F}}=8.7$ Hz), 131.4, 130.8, 130.1 (d, $J_{\text{C-F}}=3.4$ Hz), 128.7, 127.3, 123.9, 114.7 (d, $J_{\text{C-F}}=21.6$ Hz), 57.8, 39.3, 21.4, 9.6; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{16}\text{FIN}_2\text{NaO} [\text{M}+\text{Na}]^+$ 445.0184; found 445.0164.

(4-Fluorophenyl)(5-(iodomethyl)-3-(*p*-tolyl)-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3af**): 57 mg, yield 90%; $R_f=0.6$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 4$]; colorless oil. ^1H NMR (600 MHz, CDCl_3) δ : 8.04 (dd, $J=7.8, 6.0$ Hz, 2H), 7.57 (d, $J=7.8$ Hz, 2H), 7.21 (d, $J=7.8$ Hz, 2H), 7.12 (d, $J=8.0$ Hz, 2H), 4.90~4.85 (m, 1H), 3.73 (dd, $J=9.6, 7.2$ Hz, 1H), 3.67 (dd, $J=9.6, 2.4$ Hz, 1H), 3.46 (dd, $J=18.0, 11.4$ Hz, 1H), 3.08 (dd, $J=18.0, 5.4$ Hz, 1H), 2.38 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 165.7, 164.4 (d, $J_{\text{C-F}}=250.4$ Hz), 154.7, 141.0, 132.6 (d, $J_{\text{C-F}}=8.7$ Hz), 130.2 (d, $J_{\text{C-F}}=3.4$ Hz), 129.5, 128.1, 126.7, 114.7 (d, $J_{\text{C-F}}=21.6$ Hz), 57.9, 39.2, 21.5, 9.6; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{16}\text{FIN}_2\text{NaO} [\text{M}+\text{Na}]^+$ 445.0184; found 445.0177.

(3-(4-Chlorophenyl)-5-(iodomethyl)-4,5-dihydro-1H-pyrazol-1-yl)(4-fluorophenyl)methanone (**3ag**): 61 mg, yield 93%; $R_f=0.5$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; yellow solid, m.p. 113~114 °C; ^1H NMR (600 MHz, CDCl_3) δ : 8.01 (dd, $J=9.0, 6.6$ Hz, 2H), 7.61 (d, $J=8.4$ Hz, 2H), 7.38 (d, $J=8.4$ Hz, 2H), 7.13 (t, $J=8.4$ Hz, 2H), 4.91~4.88 (m, 1H), 3.78 (dd, $J=10.2, 7.2$ Hz, 1H), 3.66 (dd, $J=10.2, 4.2$ Hz, 1H), 3.46 (dd, $J=17.4, 11.4$ Hz, 1H), 3.08 (dd, $J=18.0, 4.2$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 165.9, 164.4 (d, $J_{\text{C-F}}=250.5$ Hz), 153.5, 136.6, 132.5 (d, $J_{\text{C-F}}=8.9$ Hz), 130.0 (d, $J_{\text{C-F}}=3.3$ Hz), 129.4, 129.0, 127.9, 114.8 (d, $J_{\text{C-F}}=21.6$ Hz), 58.0, 39.2, 9.5; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{13}\text{ClFIN}_2\text{NaO} [\text{M}+\text{Na}]^+$ 464.9637; found 464.9617.

(4-Fluorophenyl)(5-(iodomethyl)-3-(4-(trifluoromethyl)phenyl)-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3ah**): 68 mg, yield 96%; $R_f=0.5$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; white solid, m.p. 111~112 °C; ^1H NMR (400 MHz, CDCl_3) δ : 8.02 (dd, $J=9.2, 5.6$ Hz, 2H), 7.79 (d, $J=8.0$ Hz, 2H), 7.67 (d, $J=8.4$ Hz, 2H), 7.13 (t, $J=8.8$ Hz, 2H), 4.95~4.89 (m, 1H), 3.83 (dd, $J=10.4, 6.6$ Hz, 1H), 3.65 (dd, $J=10.4, 2.4$ Hz, 1H), 3.51 (dd, $J=18.0, 11.2$ Hz, 1H), 3.12 (dd, $J=18.0, 5.2$ Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 166.1, 164.5 (d, $J_{\text{C-F}}=250.7$ Hz), 153.1, 134.3, 132.5 (d, $J_{\text{C-F}}=8.9$ Hz), 132.1 (q, $J_{\text{C-F}}=32.7$ Hz), 129.8 (d, $J_{\text{C-F}}=3.2$ Hz), 126.9, 125.7 (q, $J_{\text{C-F}}=3.8$ Hz), 123.7 (q, $J_{\text{C-F}}=270.8$ Hz), 114.8 (d, $J_{\text{C-F}}=21.6$ Hz), 58.1, 39.1, 9.6; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{13}\text{F}_4\text{IN}_2\text{NaO} [\text{M}+\text{Na}]^+$ 498.9901; found 498.9902.

(4-Fluorophenyl)(5-(iodomethyl)-3-(4-methoxyphenyl)-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3ai**): 56 mg, yield 86%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; yellow oil. ^1H NMR (600 MHz, CDCl_3) δ : 8.04 (dd, $J=9.0, 5.4$ Hz, 2H), 7.63 (d, $J=9.0$ Hz, 2H),

7.12 (t, $J=8.4$ Hz, 2H), 6.92 (d, $J=8.4$ Hz, 2H), 4.90~4.85 (m, 1H), 3.84 (s, 3H), 3.73 (dd, $J=10.2$, 7.8 Hz, 1H), 3.68 (dd, $J=10.2$, 2.4 Hz, 1H), 3.45 (dd, $J=17.4$, 10.8 Hz, 1H), 3.07 (dd, $J=17.4$, 4.8 Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 165.6, 164.2 (d, $J_{\text{C-F}}=250$ Hz), 161.5, 154.3, 132.5 (d, $J_{\text{C-F}}=8.7$ Hz), 130.2 (d, $J_{\text{C-F}}=3.1$ Hz), 128.3, 123.4, 114.6 (d, $J_{\text{C-F}}=21.5$ Hz), 114.1, 57.8, 55.3, 39.2, 9.5; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{16}\text{FIN}_2\text{NaO}_2$ $[\text{M}+\text{Na}]^+$ 461.0133; found 461.0148.

(4-Fluorophenyl)(5-(iodomethyl)-3-methyl-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3aj**): 25 mg, yield 49%; $R_f=0.6$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; yellow oil. ^1H NMR (600 MHz, CDCl_3) δ : 7.92 (dd, $J=9.0$, 5.4 Hz, 2H), 7.08 (t, $J=6.4$ Hz, 2H), 4.73~4.69 (m, 1H), 3.74 (dd, $J=10.2$, 6.6 Hz, 1H), 3.55 (dd, $J=10.2$, 2.4 Hz, 1H), 3.08 (dd, $J=18.6$, 10.8 Hz, 1H), 2.65 (dd, $J=18.6$, 5.4 Hz, 1H), 2.04 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 165.7, 164.3 (d, $J_{\text{C-F}}=250.1$ Hz), 156.3, 132.2 (d, $J_{\text{C-F}}=8.7$ Hz), 130.3 (q, $J_{\text{C-F}}=3.3$ Hz), 114.7 (d, $J_{\text{C-F}}=21.6$ Hz), 57.2, 43.2, 16.0, 9.9; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{12}\text{FIN}_2\text{NaO}$ $[\text{M}+\text{Na}]^+$ 368.9871; found 368.9866.

(4-Fluorophenyl)(3-heptyl-5-(iodomethyl)-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3ak**): 50 mg, yield 77%; $R_f=0.6$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 2$]; colorless oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.94 (dd, $J=8.4$, 5.6 Hz, 2H), 7.08 (t, $J=8.4$ Hz, 2H), 4.71~4.65 (m, 1H), 3.76 (dd, $J=10.0$, 6.4 Hz, 1H), 3.53 (dd, $J=10.0$, 2.0 Hz, 1H), 3.05 (dd, $J=18.4$, 11.2 Hz, 1H), 2.66 (dd, $J=18.4$, 5.2 Hz, 1H), 2.37~2.32 (m, 2H), 1.63~1.55 (m, 2H), 1.31~1.26 (m, 8H), 0.89 (t, $J=7.8$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 165.6, 164.2 (d, $J_{\text{C-F}}=249.9$ Hz), 163.3, 132.5 (d, $J_{\text{C-F}}=8.7$ Hz), 130.2 (d, $J_{\text{C-F}}=3.2$ Hz), 114.6 (d, $J_{\text{C-F}}=21.5$ Hz), 56.9, 41.6, 31.7, 30.1, 29.2, 28.9, 26.2, 22.6, 14.0, 10.1; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{24}\text{FIN}_2\text{NaO}$ $[\text{M}+\text{Na}]^+$ 453.0810; found 453.0822.

(3-Cyclohexyl-5-(iodomethyl)-4,5-dihydro-1H-pyrazol-1-yl)(4-fluorophenyl)methanone (**3al**): 34 mg, yield 53%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; colorless oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.97 (t, $J=6.8$ Hz, 2H), 7.07 (t, $J=8.8$ Hz, 2H), 4.71~4.65 (m, 1H), 3.76 (dd, $J=10.0$, 5.4 Hz, 1H), 3.53 (dd, $J=10.0$, 2.4 Hz, 1H), 3.05 (dd, $J=18.4$, 11.2 Hz, 1H), 2.67 (dd, $J=18.4$, 4.8 Hz, 1H), 2.39~2.35 (m, 1H), 1.88 (d, $J=10.4$ Hz, 2H), 1.80 (d, $J=5.6$ Hz, 2H), 1.70 (d, $J=12.8$ Hz, 1H), 1.41~1.17 (m, 5H); ^{13}C NMR (150 MHz, CDCl_3) δ : 165.5, 164.2 (d, $J_{\text{C-F}}=249.9$ Hz), 163.3, 132.5 (d, $J_{\text{C-F}}=8.7$ Hz), 130.2 (d, $J_{\text{C-F}}=3.2$ Hz), 114.6 (d, $J_{\text{C-F}}=21.5$ Hz), 56.6, 39.6, 39.2, 30.2, 25.8, 25.6, 10.2; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{20}\text{FIN}_2\text{NaO}$ $[\text{M}+\text{Na}]^+$ 437.0497; found 437.0507.

(*E*)-(4-Fluorophenyl)(5-(iodomethyl)-3-styryl-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3am**): 33 mg, yield 50%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; colorless oil. ^1H NMR (600 MHz, CDCl_3) δ : 7.97 (dd, $J=6.4$, 4.2 Hz, 2H), 7.48 (d, $J=7.2$ Hz, 2H), 7.37 (t, $J=7.2$ Hz, 2H), 7.32 (t, $J=7.2$ Hz, 1H), 7.11 (t, $J=9.0$ Hz, 2H), 7.02 (d, $J=16.2$ Hz, 1H), 6.86 (d, $J=16.2$ Hz, 1H), 4.84~

4.80 (m, 1H), 3.75 (dd, $J=10.2$, 6.6 Hz, 1H), 3.65 (dd, $J=10.2$, 2.4 Hz, 1H), 3.33 (dd, $J=17.4$, 11.4 Hz, 1H), 2.96 (dd, $J=17.4$, 7.2 Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 165.8, 164.4 (d, $J_{\text{C-F}}=250.4$ Hz), 155.6, 137.9, 135.5, 132.4 (d, $J_{\text{C-F}}=8.9$ Hz), 130.0 (d, $J_{\text{C-F}}=3.2$ Hz), 129.2, 128.9, 127.1, 120.3, 114.8 (d, $J_{\text{C-F}}=21.6$ Hz), 57.7, 38.0, 9.6; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{16}\text{FIN}_2\text{NaO}$ $[\text{M}+\text{Na}]^+$ 457.0184; found 457.0184.

(5-(Iodomethyl)-3-(*p*-tolyl)-4,5-dihydro-1H-pyrazol-1-yl)(*p*-tolyl)methanone (**3ba**): 52 mg, yield 84%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; colorless oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.91 (d, $J=8.0$ Hz, 2H), 7.58 (d, $J=8.4$ Hz, 2H), 7.23 (d, $J=8.0$ Hz, 2H), 7.20 (d, $J=8.4$ Hz, 2H), 4.92~4.85 (m, 1H), 3.73~3.68 (m, 2H), 3.44 (dd, $J=17.6$, 11.2 Hz, 1H), 3.06 (dd, $J=17.6$, 4.8 Hz, 1H), 2.41 (s, 3H), 2.38 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 166.9, 154.2, 141.4, 140.7, 131.2, 130.2, 129.3, 128.3, 128.2, 126.6, 57.9, 39.1, 21.5, 21.5, 9.7; HRMS (ESI) calcd for $\text{C}_{19}\text{H}_{20}\text{IN}_2\text{O}$ $[\text{M}+\text{H}]^+$ 419.0615; found 419.0624.

(4-Chlorophenyl)(3-(4-chlorophenyl)-5-(iodomethyl)-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3ca**): 58 mg, yield 85%; $R_f=0.5$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; yellow solid, m.p. 146~147 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.92 (d, $J=8.4$ Hz, 2H), 7.60 (d, $J=8.4$ Hz, 2H), 7.42 (d, $J=8.4$ Hz, 2H), 7.38 (d, $J=8.4$ Hz, 2H), 4.91~4.86 (m, 1H), 3.78 (dd, $J=10.2$, 6.6 Hz, 1H), 3.66 (dd, $J=10.2$, 2.4 Hz, 1H), 3.46 (dd, $J=18.0$, 11.4 Hz, 1H), 3.07 (dd, $J=17.4$, 5.4 Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 166.0, 153.6, 137.3, 136.6, 132.3, 131.5, 129.3, 129.0, 128.0, 127.9, 58.0, 39.2, 9.4; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{13}\text{Cl}_2\text{IN}_2\text{NaO}$ $[\text{M}+\text{Na}]^+$ 480.9342; found 480.9345.

(4-Chlorophenyl)(3-(4-fluorophenyl)-5-(iodomethyl)-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3da**): 59 mg, yield 88%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; yellow oil. ^1H NMR (400 MHz, CDCl_3) δ : 7.92 (d, $J=8.4$ Hz, 2H), 7.60 (dd, $J=8.8$, 5.6 Hz, 2H), 7.42 (d, $J=8.8$ Hz, 2H), 7.10 (t, $J=8.8$ Hz, 2H), 4.91~4.85 (m, 1H), 3.77 (dd, $J=10.4$, 6.8 Hz, 1H), 3.66 (dd, $J=10.4$, 2.8 Hz, 1H), 3.47 (dd, $J=18.0$, 11.2 Hz, 1H), 3.08 (dd, $J=18.0$, 5.6 Hz, 1H); ^{13}C NMR (150 MHz, CDCl_3) δ : 165.9, 164.1 (d, $J_{\text{C-F}}=250.5$ Hz), 153.7, 137.2, 132.4, 131.4, 128.7 (d, $J_{\text{C-F}}=8.7$ Hz), 127.9, 127.1 (d, $J_{\text{C-F}}=3.2$ Hz), 116.0 (d, $J_{\text{C-F}}=21.9$ Hz), 57.9, 39.3, 9.4; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{13}\text{ClFIN}_2\text{NaO}$ $[\text{M}+\text{Na}]^+$ 464.9637; found 464.9635.

(3-(4-Fluorophenyl)-5-(iodomethyl)-4,5-dihydro-1H-pyrazol-1-yl)(*p*-tolyl)methanone (**3ea**): 42 mg, yield 66%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; yellowish solid, m.p. 125~126 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.88 (d, $J=12.0$ Hz, 2H), 7.67 (dd, $J=12.6$, 7.8 Hz, 2H), 7.25 (d, $J=12.0$ Hz, 2H), 7.08 (t, $J=13.2$ Hz, 2H), 4.92~4.87 (m, 1H), 3.74 (dd, $J=10.2$, 7.2 Hz, 1H), 3.67 (dd, $J=10.2$, 2.4 Hz, 1H), 3.42 (dd, $J=18.0$, 11.4 Hz, 1H), 3.04 (dd, $J=17.4$, 5.4 Hz, 1H), 2.41 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 167.0, 163.9 (d, $J_{\text{C-F}}=250.1$ Hz), 153.0, 141.5, 131.1, 130.1, 128.5 (d, $J_{\text{C-F}}=8.6$ Hz),

128.3, 127.3 (d, $J_{\text{C-F}}=3.3$ Hz), 115.8 (d, $J_{\text{C-F}}=21.9$ Hz), 58.0, 39.2, 21.5, 9.7; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{16}\text{FIN}_2\text{NaO}$ $[\text{M}+\text{Na}]^+$ 445.0184; found 445.0183.

(3-(4-Chlorophenyl)-5-(iodomethyl)-4,5-dihydro-1H-pyrazol-1-yl)(*p*-tolyl)methanone (**3fa**): 43 mg, yield 66%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; yellow solid, m.p. 145~146 °C; ^1H NMR (400 MHz, CDCl_3) δ : 7.88 (d, $J=5.6$ Hz, 2H), 7.60 (d, $J=5.6$ Hz, 2H), 7.35 (d, $J=5.6$ Hz, 2H), 7.25 (d, $J=5.6$ Hz, 2H), 4.90~4.86 (m, 1H), 3.74 (dd, $J=10.4$, 7.2 Hz, 1H), 3.66 (dd, $J=9.8$, 1.6 Hz, 1H), 3.42 (dd, $J=13.0$, 7.6 Hz, 1H), 3.03 (dd, $J=12.6$, 3.6 Hz, 1H), 2.41 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 167.0, 152.9, 141.5, 136.3, 131.0, 130.1, 129.6, 128.9, 128.3, 127.9, 58.0, 39.0, 21.5, 9.6; HRMS (ESI) calcd for $\text{C}_{18}\text{H}_{16}\text{ClIN}_2\text{NaO}$ $[\text{M}+\text{Na}]^+$ 460.9894; found 460.9884.

1-(3-(4-Fluorophenyl)-5-(iodomethyl)-4,5-dihydro-1H-pyrazol-1-yl)ethan-1-one (**3ga**): 22 mg, yield 43%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; yellow oil. ^1H NMR (600 MHz, CDCl_3) δ : 7.73 (dd, $J=9.0$, 5.4 Hz, 2H), 7.12 (t, $J=9.0$ Hz, 2H), 4.72~4.68 (m, 1H), 3.61 (dd, $J=10.2$, 3.0 Hz, 1H), 3.55 (dd, $J=10.2$, 7.8 Hz, 1H), 3.44 (dd, $J=18.0$, 11.4 Hz, 1H), 3.04 (dd, $J=18.0$, 5.4 Hz, 1H), 2.39 (s, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 169.4, 163.9 (d, $J_{\text{C-F}}=250.1$ Hz), 152.6, 128.5 (d, $J_{\text{C-F}}=8.6$ Hz), 127.3 (d, $J_{\text{C-F}}=3.3$ Hz), 115.8 (d, $J_{\text{C-F}}=21.9$ Hz), 57.1, 40.0, 21.8, 9.3; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{13}\text{FIN}_2\text{O}$ $[\text{M}+\text{H}]^+$ 347.0057; found 347.0045.

1-(3-(4-Fluorophenyl)-5-(iodomethyl)-4,5-dihydro-1H-pyrazol-1-yl)octadecan-1-one (**3ha**): 64 mg, yield 76%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; yellowish solid, m.p. 132~134 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.72 (dd, $J=9.0$, 5.4 Hz, 2H), 7.12 (t, $J=9.0$ Hz, 2H), 4.71~4.6 (m, 1H), 3.58 (d, $J=4.8$ Hz, 2H), 3.41 (dd, $J=18.0$, 11.4 Hz, 1H), 3.01 (dd, $J=17.4$, 4.8 Hz, 1H), 2.82~2.77 (m, 1H), 2.69~2.64 (m, 1H), 2.34 (t, $J=7.8$ Hz, 1H), 1.74~1.68 (m, 2H), 1.63 (t, $J=7.2$ Hz, 1H), 1.30~1.25 (m, 26H), 0.87 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (150 MHz, CDCl_3) δ : 172.5, 164.0 (d, $J_{\text{C-F}}=249.9$ Hz), 152.3, 128.5 (d, $J_{\text{C-F}}=8.6$ Hz), 127.5 (d, $J_{\text{C-F}}=3.3$ Hz), 115.9 (d, $J_{\text{C-F}}=21.9$ Hz), 57.1, 39.8, 34.2, 31.9, 29.7, 29.7, 29.7, 29.6, 29.6, 29.5, 29.4, 29.4, 29.3, 29.2, 29.1, 25.0, 24.7, 22.7, 14.1, 9.5; HRMS (ESI) calcd for $\text{C}_{28}\text{H}_{44}\text{FIN}_2\text{NaO}$ $[\text{M}+\text{Na}]^+$ 593.2380; found 593.2367.

1-(5-(Iodomethyl)-3-methyl-4,5-dihydro-1H-pyrazol-1-yl)-2-methylpropan-1-one (**3ia**): 27 mg, yield 62%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 1$]; yellow oil. ^1H NMR (600 MHz, CDCl_3) δ : 4.42~4.38 (m, 1H), 3.52 (dd, $J=10.2$, 6.6 Hz, 1H), 3.38 (dd, $J=10.2$, 2.4 Hz, 1H), 3.24~3.20 (m, 1H), 2.96 (dd, $J=18.0$, 10.8 Hz, 1H), 2.50 (dd, $J=18.0$, 4.8 Hz, 1H), 1.96 (s, 3H), 1.12 (d, $J=7.2$ Hz, 3H), 1.05 (d, $J=6.6$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 175.5, 154.9, 56.2, 43.5, 31.7, 19.4, 18.4, 15.9, 10.1; HRMS (ESI) calcd for $\text{C}_9\text{H}_{15}\text{IN}_2\text{NaO}$ $[\text{M}+\text{Na}]^+$ 317.0127; found 317.0128.

Cyclohexyl(5-(iodomethyl)-3-methyl-4,5-dihydro-1H-pyrazol-1-yl)methanone (**3ja**): 37 mg, yield 73%; $R_f=0.55$

[$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 3$]; colorless oil. ^1H NMR (600 MHz, CDCl_3) δ : 4.48~4.44 (m, 1H), 3.59 (dd, $J=10.2$, 6.6 Hz, 1H), 3.43 (dd, $J=10.2$, 2.4 Hz, 1H), 3.05~2.98 (m, 2H), 2.56 (dd, $J=18.6$, 4.8 Hz, 1H), 2.03 (s, 3H), 1.91 (d, $J=15.6$ Hz, 1H), 1.80~1.77 (m, 3H), 1.68 (d, $J=10.8$ Hz, 1H), 1.52~1.44 (m, 2H), 1.35~1.26 (m, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 174.6, 154.7, 56.1, 43.4, 41.5, 29.3, 28.3, 25.9, 25.8, 25.6, 15.9, 10.3; HRMS (ESI) calcd for $\text{C}_{12}\text{H}_{19}\text{IN}_2\text{NaO}$ $[\text{M}+\text{Na}]^+$ 357.0440; found 357.0432.

1-(5-(Iodomethyl)-3-methyl-4,5-dihydro-1H-pyrazol-1-yl)octadecan-1-one (**3ka**): 44 mg, yield 60%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 2$]; colorless oil. ^1H NMR (600 MHz, CDCl_3) δ : 4.52~4.48 (m, 1H), 3.54 (dd, $J=10.2$, 7.2 Hz, 1H), 3.48 (dd, $J=10.2$, 3.0 Hz, 1H), 3.02 (dd, $J=18.6$, 10.2 Hz, 1H), 2.69~2.64 (m, 1H), 2.60~2.51 (m, 2H), 2.03 (s, 3H), 1.68~1.63 (m, 3H), 1.31~1.25 (m, 27H), 0.88 (t, $J=7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 171.8, 154.8, 56.4, 43.7, 34.2, 31.9, 29.8, 29.7, 29.6, 29.5, 29.4, 29.4, 29.3, 25.0, 22.7, 15.9, 14.1, 9.9; HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{44}\text{IN}_2\text{O}$ $[\text{M}+\text{H}]^+$ 491.2498; found 491.2488.

(*E*)-1-(5-(Iodomethyl)-3-styryl-4,5-dihydro-1H-pyrazol-1-yl)ethan-1-one (**3ma**): 25 mg, yield 48%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 2$]; yellowish oil. ^1H NMR (600 MHz, CDCl_3) δ : 7.49 (d, $J=7.8$ Hz, 2H), 7.38 (t, $J=7.2$ Hz, 2H), 7.33 (t, $J=7.2$ Hz, 1H), 7.02 (d, $J=16.8$ Hz, 1H), 6.83 (d, $J=17.4$ Hz, 1H), 4.66~4.62 (m, 1H), 3.59 (dd, $J=10.2$, 3.0 Hz, 1H), 3.52 (dd, $J=10.2$, 7.8 Hz, 1H), 3.30 (dd, $J=18.0$, 11.4 Hz, 1H), 2.92 (dd, $J=18.0$, 5.4 Hz, 1H), 2.34 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 169.3, 154.6, 137.5, 135.6, 129.1, 128.9, 127.1, 120.4, 56.9, 38.7, 21.8, 9.3; HRMS (ESI) calcd for $\text{C}_{14}\text{H}_{16}\text{IN}_2\text{O}$ $[\text{M}+\text{H}]^+$ 355.0307; found 355.0306.

(5-(Bromomethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)(phenyl)methanone (**3bb**): 45 mg, yield 87%; $R_f=0.55$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 2$]; yellow oil. ^1H NMR (600 MHz, CDCl_3) δ : 7.99 (d, $J=7.8$ Hz, 2H), 7.69 (d, $J=7.2$ Hz, 2H), 7.49 (t, $J=7.2$ Hz, 1H), 7.44 (t, $J=7.8$ Hz, 2H), 7.41~7.38 (m, 3H), 5.13~5.10 (m, 1H), 3.92 (dd, $J=10.2$, 6.6 Hz, 1H), 3.84 (d, $J=10.8$ Hz, 1H), 3.46 (dd, $J=17.4$, 11.4 Hz, 1H), 3.26 (dd, $J=17.4$, 4.8 Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 167.1, 155.0, 134.0, 131.0, 130.9, 130.4, 130.0, 128.6, 127.6, 126.7, 58.3, 44.7, 35.9; HRMS (ESI) calcd for $\text{C}_{17}\text{H}_{16}\text{BrN}_2\text{O}$ $[\text{M}+\text{H}]^+$ 343.0446; found 343.0436.

(5-(Chloromethyl)-3-phenyl-4,5-dihydro-1H-pyrazol-1-yl)(phenyl)methanone (**3bc**): 27 mg, yield 60%; $R_f=0.5$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 2$]; colorless oil. ^1H NMR (600 MHz, CDCl_3) δ : 7.98 (d, $J=6.6$ Hz, 2H), 7.68 (d, $J=7.8$, 1.8 Hz, 2H), 7.49 (t, $J=7.2$ Hz, 1H), 7.44 (t, $J=7.8$ Hz, 2H), 7.42~7.38 (m, 3H), 5.15~5.11 (m, 1H), 4.07 (dd, $J=10.8$, 6.0 Hz, 1H), 3.95 (dd, $J=10.8$, 2.4 Hz, 1H), 3.46 (dd, $J=18.0$, 11.4 Hz, 1H), 3.31 (dd, $J=18.0$, 5.4 Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : 167.1, 155.0, 134.0, 131.0, 130.9, 130.5, 130.0, 128.7, 127.6, 126.7, 58.3, 44.7, 35.9; HRMS (ESI) calcd for

$C_{17}H_{16}ClN_2NaO$ $[M+Na]^+$ 321.0771; found 321.0781.

(5-(Bromomethyl)-3-(*p*-tolyl)-4,5-dihydro-1*H*-pyrazol-1-yl)(*p*-tolyl)methanone (**3cb**): 43 mg, yield 78%; $R_f=0.5$ [V (ethylacetate) : V (petroleum ether) = 1 : 3]; colorless oil. 1H NMR (600 MHz, $CDCl_3$) δ : 7.91 (d, $J=8.4$ Hz, 2H), 7.58 (d, $J=8.4$ Hz, 2H), 7.25 (d, $J=7.8$ Hz, 2H), 7.20 (d, $J=7.8$ Hz, 2H), 5.12~5.08 (m, 1H), 3.87 (dd, $J=10.2$, 6.0 Hz, 1H), 3.85 (dd, $J=10.8$, 3.6 Hz, 1H), 3.44 (dd, $J=18.0$, 11.4 Hz, 1H), 3.23 (dd, $J=17.4$, 5.4 Hz, 1H), 2.41 (s, 3H), 2.38 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 166.9, 154.6, 141.4, 140.8, 131.2, 130.2, 129.4, 128.3, 128.2, 126.7, 57.9, 37.1, 34.1, 21.5, 21.4; HRMS (ESI) calcd for $C_{19}H_{20}BrN_2O$ $[M+H]^+$ 371.0759; found 371.0761.

(5-(Chloromethyl)-3-(*p*-tolyl)-4,5-dihydro-1*H*-pyrazol-1-yl)(*p*-tolyl)methanone (**3cc**): 30 mg, yield 62%; $R_f=0.55$ [V (ethylacetate) : V (petroleum ether) = 1 : 3]; colorless oil. 1H NMR (400 MHz, $CDCl_3$) δ : 7.91 (d, $J=8.0$ Hz, 2H), 7.58 (d, $J=8.0$ Hz, 2H), 7.24 (d, $J=8.0$ Hz, 2H), 7.19 (d, $J=8.0$ Hz, 2H), 5.13~5.07 (m, 1H), 4.01 (dd, $J=11.2$, 6.0 Hz, 1H), 3.95 (dd, $J=11.2$, 2.8 Hz, 1H), 3.43 (dd, $J=17.6$, 11.2 Hz, 1H), 3.28 (dd, $J=17.6$, 4.2 Hz, 1H), 2.41 (s, 3H), 2.37 (s, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 167.0, 154.9, 141.4, 140.7, 131.2, 130.2, 129.3, 128.4, 128.3, 126.7, 58.3, 44.7, 35.9, 21.5, 21.4; HRMS (ESI) calcd for $C_{19}H_{20}ClN_2O$ $[M+H]^+$ 327.1259; found 327.1262.

(5-(Bromomethyl)-3-(4-fluorophenyl)-4,5-dihydro-1*H*-pyrazol-1-yl)(4-fluorophenyl)methanone (**3db**): 44 mg, yield 78%; $R_f=0.55$ [V (ethylacetate) : V (petroleum ether) = 1 : 3]; yellow oil. 1H NMR (600 MHz, $CDCl_3$) δ : 8.02 (dd, $J=9.0$, 5.4 Hz, 2H), 7.68 (dd, $J=9.0$, 5.4 Hz, 2H), 7.14~7.08 (m, 4H), 5.15~5.11 (m, 1H), 3.95 (dd, $J=10.2$, 6.0 Hz, 1H), 3.82 (dd, $J=10.2$, 2.4 Hz, 1H), 3.47 (dd, $J=18.0$, 11.4 Hz, 1H), 3.24 (dd, $J=18.0$, 5.4 Hz, 1H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 165.9, 164.4 (d, $J_{C-F}=250.5$ Hz), 164.1 (d, $J_{C-F}=250.5$ Hz), 154.0, 132.5 (d, $J_{C-F}=8.9$ Hz), 130.0 (d, $J_{C-F}=3.3$ Hz), 128.7 (d, $J_{C-F}=8.4$ Hz), 127.2 (d, $J_{C-F}=3.2$ Hz), 115.9 (d, $J_{C-F}=21.9$ Hz), 114.5 (d, $J_{C-F}=21.6$ Hz), 58.0, 37.2, 34.0; HRMS (ESI) calcd for $C_{17}H_{14}BrF_2N_2O$ $[M+H]^+$ 379.0252; found: 379.0252.

(5-(Chloromethyl)-3-(4-fluorophenyl)-4,5-dihydro-1*H*-pyrazol-1-yl)(4-fluorophenyl)methanone (**3dc**): 25 mg, yield 50%; $R_f=0.55$ [V (ethylacetate) : V (petroleum ether) = 1 : 3]; yellow solid, m.p. 101~102 °C; 1H NMR (600 MHz, $CDCl_3$) δ : 8.02 (dd, $J=8.4$, 5.4 Hz, 2H), 7.68 (dd, $J=8.4$, 5.4 Hz, 2H), 7.14~7.09 (m, 4H), 5.16~5.12 (m, 1H), 4.09 (dd, $J=11.4$, 6.0 Hz, 1H), 3.94 (dd, $J=11.4$, 2.4 Hz, 1H), 3.47 (dd, $J=18.0$, 11.4 Hz, 1H), 3.30 (dd, $J=18.0$, 5.4 Hz, 1H); ^{13}C NMR (150 MHz, $CDCl_3$) δ : 166.0, 164.4 (d, $J_{C-F}=250.7$ Hz), 164.1 (d, $J_{C-F}=250.4$ Hz), 154.2, 132.5 (d, $J_{C-F}=8.7$ Hz), 130.0 (d, $J_{C-F}=3.3$ Hz), 128.7 (d, $J_{C-F}=8.6$ Hz), 127.2 (d, $J_{C-F}=3.2$ Hz), 116.0 (d, $J_{C-F}=21.9$ Hz), 114.7 (d, $J_{C-F}=21.6$ Hz), 58.4, 44.7, 35.9; HRMS (ESI) calcd for $C_{17}H_{13}ClF_2N_2NaO$ $[M+Na]^+$ 357.0582; found 357.0568.

(5-(Bromomethyl)-3-methyl-4,5-dihydro-1*H*-pyrazol-1-

yl)(cyclohexyl)methanone (**3eb**): 30 mg, yield 69%; $R_f=0.5$ [V (ethylacetate) : V (petroleum ether) = 1 : 2]; yellow oil. 1H NMR (600 MHz, $CDCl_3$) δ : 4.71~4.67 (m, 1H), 3.73 (dd, $J=10.2$, 6.0 Hz, 1H), 3.58 (dd, $J=10.2$, 3.0 Hz, 1H), 3.05~2.97 (m, 2H), 2.71 (dd, $J=18.0$, 5.4 Hz, 1H), 2.02 (s, 3H), 1.87 (d, $J=13.2$ Hz, 1H), 1.77~1.74 (m, 3H), 1.67 (d, $J=11.4$ Hz, 1H), 1.50~1.44 (m, 2H), 1.35~1.19 (m, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 174.7, 155.1, 56.2, 41.5, 41.4, 34.4, 29.2, 28.3, 25.9, 25.8, 25.6, 15.9; HRMS (ESI) calcd for $C_{12}H_{19}BrN_2NaO$ $[M+Na]^+$ 309.0573; found 309.0576.

(5-(Chloromethyl)-3-methyl-4,5-dihydro-1*H*-pyrazol-1-yl)(cyclohexyl)methanone (**3ec**): 18 mg, yield 50%; $R_f=0.6$ [V (ethylacetate) : V (petroleum ether) = 1 : 2]; yellow oil. 1H NMR (600 MHz, $CDCl_3$) δ : 4.73~4.69 (m, 1H), 3.87 (dd, $J=10.8$, 6.0 Hz, 1H), 3.70 (dd, $J=10.8$, 3.0 Hz, 1H), 3.06~2.98 (m, 2H), 2.79 (dd, $J=18.0$, 4.8 Hz, 1H), 2.04 (s, 3H), 1.87 (d, $J=13.2$ Hz, 1H), 1.77 (t, $J=13.2$, 3H), 1.68 (d, $J=13.8$ Hz, 1H), 1.51~1.44 (m, 2H), 1.38~1.21 (m, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 174.7, 155.3, 56.6, 44.8, 41.5, 40.1, 29.1, 28.4, 25.9, 25.7, 25.6, 15.9; HRMS (ESI) calcd for $C_{12}H_{20}ClN_2O$ $[M+H]^+$ 243.1259; found 243.1266.

1-(5-(Bromomethyl)-3-methyl-4,5-dihydro-1*H*-pyrazol-1-yl)octadecan-1-one (**3fb**): 37 mg, yield 56%; $R_f=0.6$ [V (ethylacetate) : V (petroleum ether) = 1 : 2]; colorless oil. 1H NMR (600 MHz, $CDCl_3$) δ : 4.73~4.70 (m, 1H), 3.71 (dd, $J=10.2$, 6.6 Hz, 1H), 3.63 (dd, $J=10.2$, 3.0 Hz, 1H), 3.02 (dd, $J=18.0$, 11.4 Hz, 1H), 2.74 (dd, $J=18.6$, 4.8 Hz, 1H), 2.69~2.634 (m, 1H), 2.58~2.53 (m, 1H), 2.03 (s, 3H), 1.67~1.62 (m, 2H), 1.35~1.25 (m, 28H), 0.88 (t, $J=6.6$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 171.9, 155.2, 56.4, 41.6, 34.2, 34.1, 31.9, 29.7, 29.6, 29.6, 29.6, 29.5, 29.4, 29.3, 29.3, 24.9, 22.7, 15.9, 14.1; HRMS (ESI) calcd for $C_{23}H_{44}BrN_2O$ $[M+H]^+$ 443.2637; found 443.2624.

1-(5-(Chloromethyl)-3-methyl-4,5-dihydro-1*H*-pyrazol-1-yl)octadecan-1-one (**3fc**): 32 mg, yield 53%; $R_f=0.55$ [V (ethylacetate) : V (petroleum ether) = 1 : 2]; yellow solid, m.p. 101~102 °C; 1H NMR (600 MHz, $CDCl_3$) δ : 4.75~4.71 (m, 1H), 3.85 (dd, $J=11.4$, 6.6 Hz, 1H), 3.74 (dd, $J=11.4$, 3.0 Hz, 1H), 3.01 (dd, $J=18.6$, 10.8 Hz, 1H), 2.80 (dd, $J=18.0$, 4.8 Hz, 1H), 2.69~2.64 (m, 1H), 2.59~2.54 (m, 1H), 2.34 (t, $J=7.8$ Hz, 1H), 2.03 (s, 3H), 1.66~1.62 (m, 3H), 1.30~1.25 (m, 26H), 0.88 (t, $J=6.6$ Hz, 3H); ^{13}C NMR (100 MHz, $CDCl_3$) δ : 172.0, 155.5, 56.7, 44.8, 40.3, 34.1, 31.9, 29.7, 29.7, 29.6, 29.6, 29.6, 29.5, 29.4, 29.4, 29.3, 29.3, 29.2, 29.1, 24.9, 24.7, 22.7, 15.9, 14.1; HRMS (ESI) calcd for $C_{23}H_{44}ClN_2O$ $[M+H]^+$ 399.3142; found 399.3141.

4.3 General procedure for the synthesis of 4

3a (0.2 mmol) was dissolved in CH_3CH_2OH (2 mL), KOH (2.0 equiv.) was added into the reaction mixture, and then the reaction mixture was refluxed for 30 min until the reaction was completed monitored by TLC analysis. The solvent was removed under vacuum. The crude product

was purified by flash chromatography on silica gel using a mixture of petroleum ether and ethyl acetate as an eluting solvent to give the desired product methyl-3-phenyl-1*H*-pyrazole (**4**)^[17] 28 mg, yield 88%. $R_f=0.5$ [$V(\text{ethylacetate}) : V(\text{petroleum ether})=1 : 2$]; yellow solid, m.p. 101~102 °C; ^1H NMR (600 MHz, CDCl_3) δ : 7.70 (d, $J=7.8$ Hz, 2H), 7.35 (d, $J=7.8$ Hz, 2H), 7.80 (t, $J=7.8$ Hz, 1H), 6.33 (s, 1H), 2.26 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ : 149.9, 143.2, 132.5, 128.6, 127.7, 125.7, 102.0, 11.6

Supporting Information Copies of ^1H NMR, ^{13}C NMR spectra and the HRMS data of compounds **3** and **4**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn/>.

References

- [1] (a) Vitaku, E.; Smith, D. T.; Njardarson, J. T. *J. Med. Chem.* **2014**, *57*, 10257.
(b) Blair, L. M.; Sperry, J. J. *Nat. Prod.* **2013**, *76*, 794.
- [2] (a) Shaharyar, M.; Abdullah, M. M.; Bakht, M. A.; Majeed, J. *Eur. J. Med. Chem.* **2010**, *45*, 114.
(b) Johnson, M.; Younglove, B.; Lee, L.; LeBlanc, R.; Holt, H., Jr.; Hills, P.; Mackay, H.; Brown, T.; Mooberry, S. L.; Lee, M. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 5897.
(c) Manna, F.; Chimenti, F.; Fioravanti, R.; Bolasco, A.; Secci, D.; Chimenti, P.; Ferlinib, C.; Scambia, G. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 4632.
- [3] Prasad, Y. R.; Rao, A. L.; Prasanna, L.; Murali, K.; Kumar, P. R. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 5030.
- [4] Oezdemir, Z.; Kandilci, H. B.; Guemesel, B.; Calis, U.; Bilgin, A. *Eur. J. Med. Chem.* **2007**, *42*, 373.
- [5] (a) Manojkumar, P.; Ravi, T. K.; Gopalakrishnan, S. *Eur. J. Med. Chem.* **2009**, *44*, 4690.
(b) Chimenti, F.; Bizzarri, B.; Manna, F.; Bolasco, A.; Secci, D.; Chimenti, P.; Granese, A.; Rivanera, D.; Lilli, D.; Scaltrito, M. M.; Brenciaglia, M. I. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 603.
- [6] Acharya, B. N.; Saraswat, D.; Tiwari, M.; Shrivastava, A. K.; Ghorpade, R.; Bapna, S.; Kaushik, M. P. *Eur. J. Med. Chem.* **2010**, *45*, 430.
- [7] (a) Bliznyuk, V. N.; Seliman, A. F.; Ishchenko, A. A.; Derevyanko, N. A.; DeVo, T. A. *ACS Appl. Mater. Interfaces* **2016**, *8*, 12843.
(b) Szukalski, A.; Sznitko, L.; Cypriach, K.; Miniewicz, A.; Mysliwiec, J. *J. Phys. Chem. C* **2014**, *118*, 8102.
- (c) Dwivedi, Y.; de Boni, L.; Goncalves, P. J.; Mairink, L. M.; Menegatti, R.; Fonseca, T. L.; Zilio, S. C. *Spectrochim. Acta, Part A* **2013**, *105*, 483.
- [8] (a) Mahé, O.; Dez, I.; Levacher, V.; Brière, J.-F. *Angew. Chem. Int. Ed.* **2010**, *49*, 7072.
(b) Lévai, A.; Jekő, J. *J. Heterocycl. Chem.* **2006**, *43*, 111.
- [9] (a) Du, T.; Du, F.; Ning, Y.; Peng, Y. *Org. Lett.* **2015**, *17*, 1308.
(b) Simovic, D.; Di, M.; Marks, V.; Chatfield, D. C.; Rein, K. S. *J. Org. Chem.* **2007**, *72*, 650.
(c) Kano, T.; Hashimoto, T.; Maruoka, K. *J. Am. Chem. Soc.* **2006**, *128*, 2174.
(d) Ruano, J. L. G.; De Diego, S. A. A.; Blanco, D.; Castro, A. M. M.; Martín, M. R.; Ramos, J. H. R. *Org. Lett.* **2001**, *3*, 3173.
- [10] Ju, Y.; Varma, R. S. *J. Org. Chem.* **2006**, *71*, 135.
- [11] (a) Cheng, J.; Xu, P.; Li, W.; Cheng, Y.; Zhu, C. *Chem. Commun.* **2016**, *52*, 11901.
(b) Wang, G.; Huang, T.; Lin, L.; Feng, X. *Org. Lett.* **2013**, *15*, 76.
(c) Sibi, M. P.; Stanley, L. M.; Soeta, T. *Adv. Synth. Catal.* **2006**, *348*, 2371.
(d) Sibi, M. P.; Stanley, L. M.; Jasperse, C. P. *J. Am. Chem. Soc.* **2005**, *127*, 8276.
- [12] (a) Huang, R.; Tao, H.-Y.; Wang, C.-J. *Org. Lett.* **2017**, *19*, 1176.
(b) Attanasi, O. A.; Crescentini, L. D.; Favi, G. *J. Org. Chem.* **2014**, *79*, 8331.
(c) Chen, J.-R.; Jörres, M.; Bolm, C. *J. Am. Chem. Soc.* **2012**, *134*, 6924.
- [13] (a) Hu, X.-Q.; Chen, J.-R.; Beauchemin, M.; Xiao, W.-J. *Chem. Eur. J.* **2016**, *22*, 14141.
(b) Hu, X.-Q.; Chen, J.-R.; Beauchemin, M.; Xiao, W.-J. *Angew. Chem. Int. Ed.* **2014**, *53*, 12163.
- [14] (a) Yang, M.-N.; Yan, D.-M.; Zhao, Q.-Q.; Chen, J.-R.; Xiao, W.-J. *Org. Lett.* **2017**, *19*, 5208.
(b) Zhao, Q.-Q.; Chen, J.; Yan, D.-M.; Chen, J.-R.; Xiao, W.-J. *Org. Lett.* **2017**, *19*, 3620.
(c) Wu, X.; Wang, M.; Zhang, G.; Zhao, Y.; Wang, J.; Ge, H. *Chem. Sci.* **2015**, *6*, 5882.
(d) Hu, X.-Q.; Deng, Q.-H.; Zou, Y.-Q.; Xiao, W.-J. *Eur. J. Org. Chem.* **2014**, 3082.
- [15] Wang, Y.; Wang, K.-H.; Su, Y.; Yang, Z.; Wen, L.; Liu, L.; Wang, J.; Huang, D.; Hu, Y. *J. Org. Chem.* **2018**, *83*, 939.
- [16] Lu, A.; Wang, F.; Huang, D.; Wang, K.-H.; Su, Y.; Xu, Y.; Hu, Y. *Chin. J. Org. Chem.* **2014**, *34*, 948 (in Chinese).
(陆爱玲, 王凤娇, 黄丹凤, 王克虎, 苏瀛鹏, 徐艳丽, 胡雨来, 有机化学, **2014**, *34*, 948.)
- [17] Wen, J.; Fu, Y.; Zhang, R.-Y.; Zhang, J.; Chen, S.-Y.; Yu, X.-Q. *Tetrahedron* **2011**, *67*, 9618.

(Li, L.; Fan, Y.)