四丁基溴化铵介导的 *N*-甲基苯胺和色氨酸衍生物的
环化芳香化合成 β -咔啉王震^{†,a} 张玲^{†,b} 张富赓^{*,b} 王彬^{*,a}^(a) 南开大学药学院 天津 300353^(b) 南开大学附属环湖医院 天津 300350

摘要 描述了一种四丁基溴化铵介导的环化芳香化反应用于 β -咔啉的合成, 反应物使用容易得到的色氨酸和 *N*-甲基苯胺。这种不使用金属催化剂的策略是现存 β -咔啉合成方法的有益补充。

关键词 β -咔啉; 四丁基溴化铵; *N*-甲基苯胺; 环化芳香化反应

Synthesis of β -Carbolines through Tetra-*n*-butylammonium Bromide-Mediated Cycloaromatization Reaction of *N*-Methylaniline with Tryptophan DerivativesWang, Zhen^{†,a} Zhang, Ling^{†,b} Zhang, Fugeng^{*,b} Wang, Bin^{*,a}^(a) College of Pharmacy, Nankai University, Tianjin 300353^(b) Department of Pharmacy, Affiliated Huanhu Hospital of Nankai University, Tianjin 300350

Abstract A mild and efficient ⁿBu₄NBr-mediated oxidative cycloaromatization to prepare β -carbolines from readily available tryptophans and *N*-methylaniline is described. The present metal free protocol is complementary to the existing methods for the synthesis of aromatic β -carbolines.

Keywords β -carboline; tetra-*n*-butylammonium bromide; *N*-methylaniline; cycloaromatization

1 Introduction

Aromatic β -carbolines represent one of the most important classes of heterocycles that display important biological activity.^[1] They are involved in the treatment of malaria,^[2] cancer,^[3] acquired immune deficiency syndrome (AIDS),^[4] and other diseases.^[5] In addition, some β -carboline derivatives displayed interesting optical properties and acted as photosensitizers in material science.^[6] Generally, syntheses of aromatic β -carbolines involve utilization of traditional methods such as Pictet-Spengler reaction^[7] and Bischler-Napieralski cyclization,^[8] a protic or Lewis acid is needed to obtain the imine intermediate and the reactions take place under harsh conditions. Moreover, a subsequent oxidation step is required to form the desired aromatic β -carbolines.^[9] More recently, intense efforts are

directed towards to synthesize β -carbolines skeleton by use of transition-metal catalyzed oxidative cycloaromatization reactions, including Pd,^[10] Cu,^[11] Ru,^[12] Rh,^[13] Au,^[14] and bimetallic Cu/Rh.^[15] Although these reported protocols were successfully applied to syntheses of β -carboline derivatives, most of the reactions occurred in the presence of expensive metal catalysts, and in certain cases the starting materials were not easily accessible. In view of the importance of the β -carbolines, particularly in medicinal chemistry, the metal-free and mild methods for the preparation of these compounds are desirable. Recently, an eminent contribution independent of metal- and acid-catalyzed reaction has emerged.^[16] In 2012, we^[17] have reported an efficient ⁿBu₄NI-catalyzed C(3)-formylation of indoles using *N*-methylaniline as the carbonyl source (Scheme 1). As a part of major project aimed at developing

* Corresponding authors. E-mail: clare2006@163.com; wangbin@nankai.edu.cn

Received March 30, 2019; revised May 16, 2019; published online June 3, 2019.

Dedicated to the 100th anniversary of the birth of Professor Ruyi Chen.

Project supported by the National Natural Science Foundation of China ((Nos. 21172120, 21472093).

国家自然科学基金(No. 21172120, 21472093)资助项目.

[†] 共同第一作者(These authors contributed equally to this work).

halide-catalyzed cross-dehydrogenative coupling (CDC) reactions^[18] we herein design a new synthetic strategy for the construction of β -carboline skeleton through the reaction of tryptophan ester with *N*-methylaniline (Scheme 1). We envisioned that this metal free protocol is complementary to the existing methods.

Our previous work (Ref. [17])

Scheme 1 *N*-methylaniline acts as carbon source

2 Results and discussion

To validate our hypothesis in Scheme 1, an initial experiment was performed under our previous formylation conditions using the readily available tryptophan methyl ester hydrochloride (**1a**) and *N*-methylaniline (**2**) as the model substrates. Unfortunately, the reactants failed to undergo cyclization to the corresponding β -carboline under our reported conditions^[17a] (Table 1, Entry 1). Attempts to perform the cyclization using other iodine mediators such as I_2 or KI led to the formation of substantial amounts of side products and only trace of desired **3a** was detected (Entries 2 and 3). The reaction was then conducted with the combination of nBu_4NBr /*tert*-butyl peroxybenzoate (TBPB) in 1,4-dioxane in the absence of PivOH. The result indicated that the desired product **3a** was obtained in 37% yield (Entry 4). When *tert*-butyl hydroperoxide (TBHP) was used instead of TBPB, a higher yield of **3a** was observed (Entry 5). In contrast, the use of H_2O_2 led to no conversion (Entry 6). It was found that additives Ph_3P , isatoic anhydride (IA) and perylenetetracarboxylic dianhydride (PTCDA) promoted the cycloaromatization efficiently to afford β -carboline **3a** in 52%, 63% and 61% yields, respectively (Entries 7–9). The satisfactory yields were observed using the combination of Ph_3P /IA or Ph_3P /PTCDA (Entries 10 and 11).

With the optimal conditions in hand (Table 1, Entry 10), the scope of this transformation was next examined and various tryptophans were subjected to the optimized conditions. As illustrated in Table 2, both ethyl and benzyl esters of tryptophan were tolerated (**3b** and **3c**). The substrates with halide groups afforded the corresponding products **3d**–**3f** in 33%–41% yields. The cyclization underwent smoothly with *N*-methyl, *N*-benzyl and allyl substrates to yield the desired products **3h**–**3j** in 67%, 45% and 65%

Table 1 Optimization of the reaction conditions^a

Entry	Halide	Oxidant	Additive	Solvent	Yield/%
1	nBu_4NI	TBPB	PivOH	DMSO	Trace
2	I_2	TBPB	PivOH	DMSO	Trace
3	KI	TBPB	PivOH	DMSO	Trace
4	nBu_4NBr	TBPB	—	1,4-Dioxane	37
5	nBu_4NBr	TBHP	—	1,4-Dioxane	42
6	nBu_4NBr	H_2O_2	—	1,4-Dioxane	0
7	nBu_4NBr	TBHP	Ph_3P	1,4-Dioxane	52
8	nBu_4NBr	TBHP	IA	1,4-Dioxane	63
9	nBu_4NBr	TBHP	PTCDA	1,4-Dioxane	61
10 ^b	nBu_4NBr	TBHP	Ph_3P /IA	1,4-Dioxane	85
11 ^b	nBu_4NBr	TBHP	Ph_3P /PTCDA	1,4-Dioxane	79

^a Reaction conditions: **1a** (0.25 mmol), **2** (0.50 mmol), halide (1.0 equiv.), oxidant (3.5 equiv.), additive (1.0 equiv.), solvent (2.0 mL) under air; isolated yields. ^b Ph_3P (1.0 equiv.), IA (isatoic anhydride, 1.0 equiv.), PTCDA (perylene-tetracarboxylic dianhydride, 1.0 equiv.), TBPB (*tert*-butyl peroxybenzoate), TBHP (*tert*-butyl hydroperoxide).

Table 2 Scope of the substrates^a

^a Reaction conditions: **1** (0.25 mmol), **2** (0.50 mmol), under air; isolated yields.

yields, respectively. However, when *N,N*-dimethylamine as carbon source was subjected to the present conditions, the desired product was not obtained, and a side product 4,4'-methylenebis (*N,N*-dimethylaniline) through the ho-

mocoupling of *N,N*-dimethyl-amine was observed. The similar result has been reported by us in a previous work.^[17a] Unfortunately, tryptophans with 6-hydroxy, 6-cyano, *N*-acetyl or *N*-benzoyl groups failed to give the corresponding products under the present conditions.

To explore the reaction mechanism, the radical inhibition experiments were carried out. We found that the present reaction was completely suppressed by use of 2,2,6,6-tetramethyl-1-piperidinyloxy (TEMPO) or butylated hydroxytoluene (BHT). Based on our previous report^[17a] and this result, a plausible reaction pathway is shown in Scheme 2. First, the reaction of Br[−] with TBHP produces ^tBuO• via a SET pathway. Then ^tBuO• abstract H atom from methyl group of *N*-methylaniline **2** that further yields imine ion **A** by oxidation. Second, the addition reaction of **A** with **1a** takes place to yield intermediate **B**, followed by the oxidation producing intermediate **C**. Third, intermediate **C** undergoes an intramolecular cyclization to provide **D** that further gives intermediate **E** by deprotonation. Finally, Cope elimination of **E** and the followed oxidation afford the final product **3a**. At this stage, the precise role of additives is unknown. We speculate that the cationic bromonium complex Ph₃PBr⁺ may be formed in the transformation.^[19] It acts as a Lewis acid to promote cyclization just as Lewis acid catalyzed Pictet-Spengler reaction.^[7] As for the additives IA and PTCDA, they might promote the deamination of **E**, like dehydrating agents in Bischler-Napieralski cyclization.^[8]

3 Conclusions

In summary, an efficient method for the synthesis of aromatic β -carbolines from readily available starting materials under mild conditions was developed. This protocol is the useful complementary to the known methods for the synthesis of aromatic β -carbolines.

4 Experimental section

4.1 General remarks

¹H NMR and ¹³C NMR spectra were recorded on a Bruker AVANCE AV400 (400 MHz for ¹H NMR and 101 MHz for ¹³C NMR). All NMR chemical shifts were referenced to residual solvent peaks or to Si(CH₃)₄ as an internal standard. NMR spectra recorded in CDCl₃ were referenced to residual CHCl₃ at δ 7.26 for ¹H NMR or 77.0 for ¹³C NMR. NMR spectra recorded in DMSO-*d*₆ were referenced to residual DMSO at δ 2.49 and 3.33 for ¹H NMR or 39.6 for ¹³C NMR. HRMS were measured using a Q-TOF LC-MS and a ESI-FTICR technique. Reactions were monitored by thin-layer chromatography (TLC) on 0.25 mm silica gel glass plates coated with 60 F254. Column chromatography was performed on silica gel (200~300 mesh) using a mixture of dichloromethane and ethyl acetate as eluant. Commercially available reagents **1a**~**1c** were used as received without further purification. Raw materials **1d**~**1j** were prepared by the known methods^[20] and analytical data matched literature values.^[21]

4.2 General procedure for the preparation of product **3a**

Methyl 2-amino-3-(1*H*-indol-3-yl)propanoate hydrochloride (**1a**, 0.25 mmol, 63.7 mg, 1.0 equiv.) was added to a mixture of TBAB (0.25 mmol, 81.4 mg, 1.0 equiv.), PPh₃ (0.25 mmol, 65.5 mg, 1.0 equiv.), isatoic anhydride (0.25 mmol, 42 mg, 1.0 equiv.) and *N*-methylaniline (0.5 mmol, 55 μ L, 2.0 equiv.) under air, and then *tert*-butyl hydroperoxide (0.875 mmol, 120 μ L, 3.5 equiv.) and 1,4-dioxane (2 mL) were added. After the mixture was stirred at 70 °C for 20 h, the residue was mixed with silica gel and concentrated. The resulting mixture was purified by silica gel column chromatography on silica gel with dichloro-

Scheme 2 Plausible reaction pathway

methane/ethyl acetate ($V:V=3:1$) as eluent to give the desired product **3a**.

Methyl 9H-pyrido[3,4-b]indole-3-carboxylate (**3a**):^[22] White solid (48.1 mg, 85%). m.p. 242~244 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 12.10 (s, 1H), 8.99 (s, 1H), 8.94 (s, 1H), 8.42 (d, $J=7.7$ Hz, 1H), 7.68 (d, $J=8.1$ Hz, 1H), 7.61 (t, $J=7.4$ Hz, 1H), 7.32 (t, $J=7.2$ Hz, 1H), 3.92 (s, 3H); ¹³C NMR (101 MHz, DMSO) δ : 166.5, 141.4, 137.9, 136.9, 134.2, 129.2, 127.9, 122.7, 121.3, 120.7, 118.1, 112.9, 52.4; HRMS (ESI) calcd for C₁₃H₁₁N₂O₂ [M+H]⁺ 227.0815, found 227.0811.

Ethyl 9H-pyrido[3,4-b]indole-3-carboxylate (**3b**):^[23] White solid (49.9 mg, 83%). m.p. 225~227 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 12.07 (s, 1H), 8.98 (s, 1H), 8.93 (s, 1H), 8.42 (d, $J=8.0$ Hz, 1H), 7.68 (d, $J=8.4$ Hz, 1H), 7.61 (t, $J=7.8$ Hz, 1H), 7.33 (t, $J=7.6$ Hz, 1H), 4.38 (tt, $J=7.1, 4.5$ Hz, 2H), 1.38 (td, $J=7.2, 2.3$ Hz, 3H); ¹³C NMR (101 MHz, DMSO-*d*₆) δ : 166.0, 141.4, 137.9, 137.2, 134.1, 129.1, 127.9, 122.7, 121.3, 120.7, 118.0, 112.8, 61.0, 14.8. HRMS (ESI) calcd for C₁₄H₁₃N₂O₂ [M+H]⁺ 241.0972, found 241.0968.

Benzyl 9H-pyrido[3,4-b]indole-3-carboxylate (**3c**): White solid (44.6 mg, 59%). m.p. 233~234 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 12.11 (s, 1H), 8.98 (d, $J=4.5$ Hz, 2H), 8.43 (d, $J=7.8$ Hz, 1H), 7.71~7.66 (m, 1H), 7.63 (d, $J=7.0$ Hz, 1H), 7.54 (d, $J=7.2$ Hz, 2H), 7.43 (d, $J=7.5$ Hz, 2H), 7.41~7.37 (m, 1H), 7.32 (d, $J=7.6$ Hz, 1H), 5.43 (s, 2H); HRMS (ESI) calcd for C₁₉H₁₅N₂O₂ [M+H]⁺ 303.1128, found 303.1132.

Methyl 6-chloro-9H-pyrido[3,4-b]indole-3-carboxylate (**3d**):^[24] White solid (26.7 mg, 41%). m.p. 275~277 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 12.20 (s, 1H), 8.99 (s, 1H), 8.94 (s, 1H), 8.43 (d, $J=8.4$ Hz, 1H), 7.73 (s, 1H), 7.33 (d, $J=8.4$ Hz, 1H), 3.95~3.89 (m, 3H); ¹³C NMR (101 MHz, DMSO) δ : 165.8, 141.4, 137.6, 136.8, 133.9, 132.9, 126.9, 123.7, 120.5, 119.7, 117.6, 112.0, 51.9; HRMS (ESI) calcd for C₁₃H₁₀ClN₂O₂ [M+H]⁺ 261.0425, found 261.0429.

Methyl 7-chloro-9H-pyrido[3,4-b]indole-3-carboxylate (**3e**): White solid (26.1 mg, 40%). m.p. 288~289 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 12.18 (s, 1H), 8.99 (s, 1H), 8.94 (s, 1H), 8.44 (d, $J=8.4$ Hz, 1H), 7.73 (s, 1H), 7.34 (d, $J=8.4$ Hz, 1H), 3.92 (s, 3H); ¹³C NMR (101 MHz, DMSO) δ : 166.4, 142.0, 138.2, 137.5, 134.5, 133.5, 127.5, 124.3, 121.0, 120.3, 118.2, 112.6, 52.5; HRMS (ESI) calcd for C₁₃H₁₀ClN₂O₂ [M+H]⁺ 261.0425, found 261.0426.

Methyl 6-bromo-9H-pyrido[3,4-b]indole-3-carboxylate (**3f**):^[25] White solid (25.2 mg, 33%). m.p. 278~280 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 12.25 (s, 1H), 9.00 (s, 2H), 8.71 (d, $J=2.0$ Hz, 1H), 7.73 (dd, $J=8.6, 2.0$ Hz, 1H), 7.65 (d, $J=8.8$ Hz, 1H), 3.92 (s, 3H); ¹³C NMR (101 MHz, DMSO) δ : 166.4, 140.1, 138.1, 137.2, 134.6, 131.7, 127.0, 125.4, 123.2, 118.7, 114.9, 112.8, 52.5; HRMS (ESI) calcd for C₁₃H₁₀BrN₂O₂ [M+H]⁺ 304.9920, 306.9905, 305.9959, 307.9939, found 304.9918, 306.9902, 305.9953, 307.9937.

Methyl 6-methyl-9H-pyrido[3,4-b]indole-3-carboxylate

(**3g**): White solid (37.2 mg, 62%). m.p. 283~284 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 11.97 (s, 1H), 8.94 (s, 1H), 8.88 (s, 1H), 8.19 (s, 1H), 7.57 (d, $J=8.2$ Hz, 1H), 7.44 (d, $J=8.4$ Hz, 1H), 2.50 (s, 3H); ¹³C NMR (101 MHz, DMSO) δ : 166.5, 139.7, 138.1, 136.7, 134.1, 130.6, 129.6, 127.6, 122.2, 121.5, 118.0, 112.6, 52.4, 21.5; HRMS (ESI) calcd for C₁₄H₁₃N₂O₂ [M+H]⁺ 241.0972, found 241.0975.

Methyl 9-methyl-9H-pyrido[3,4-b]indole-3-carboxylate (**3h**):^[26] White solid (40.2 mg, 67%). m.p. 215~216 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 9.12 (s, 1H), 8.93 (s, 1H), 8.44 (d, $J=7.9$ Hz, 1H), 7.77 (d, $J=8.3$ Hz, 1H), 7.69 (t, $J=7.7$ Hz, 1H), 7.37 (t, $J=7.5$ Hz, 1H), 4.05 (s, 3H), 3.93 (s, 3H); ¹³C NMR (101 MHz, DMSO) δ : 166.4, 142.3, 138.3, 137.0, 132.9, 129.3, 127.5, 122.8, 121.0, 120.9, 117.8, 111.0, 52.5, 30.1; HRMS (ESI) calcd for C₁₄H₁₃N₂O₂ [M+H]⁺ 241.0972, found 241.0975.

Methyl 9-benzyl-9H-pyrido[3,4-b]indole-3-carboxylate (**3i**):^[26] White solid (32.4 mg, 41%). m.p. 187~188 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 9.20 (s, 1H), 8.98 (s, 1H), 8.48 (d, $J=7.9$ Hz, 1H), 7.82 (d, $J=8.4$ Hz, 1H), 7.66 (t, $J=7.7$ Hz, 1H), 7.37 (t, $J=7.5$ Hz, 1H), 7.27 (dt, $J=23.3, 7.5$ Hz, 5H), 5.87 (s, 2H), 3.92 (s, 3H); ¹³C NMR (101 MHz, DMSO) δ : 166.4, 141.8, 138.0, 137.5, 133.2, 129.5, 129.2, 128.1, 128.0, 127.3, 123.0, 121.3, 121.2, 118.0, 111.4, 52.5, 46.6; HRMS (ESI) calcd for C₂₀H₁₇N₂O₂ [M+H]⁺ 317.1285, found 317.1290.

Methyl 9-allyl-9H-pyrido[3,4-b]indole-3-carboxylate (**3j**): White solid (43.3 mg, 65%). m.p. 209~211 °C; ¹H NMR (400 MHz, DMSO-*d*₆) δ : 9.11 (s, 1H), 8.97 (s, 1H), 8.47 (d, $J=7.9$ Hz, 1H), 7.77 (d, $J=8.3$ Hz, 1H), 7.67 (t, $J=7.7$ Hz, 1H), 7.37 (t, $J=7.5$ Hz, 1H), 6.06 (ddt, $J=15.8, 10.2, 5.1$ Hz, 1H), 5.25 (d, $J=5.1$ Hz, 2H), 5.16 (d, $J=10.3$ Hz, 1H), 5.01 (d, $J=17.1$ Hz, 1H), 3.93 (s, 3H); HRMS (ESI) calcd for C₁₆H₁₅N₂O₂ [M+H]⁺ 267.1128, found 267.1133.

Supporting Information ¹H NMR, ¹³C NMR and HRMS spectra of compounds **3a**~**3j**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- (a) Cao, R. H.; Peng, W. L.; Wang, Z. H.; Xu, A. L. *Curr. Med. Chem.* **2007**, *14*, 479.
- (b) Lancianesi, S.; Palmieri, A.; Petrini, M. *Chem. Rev.* **2014**, *114*, 7108.
- (c) Nagle, A. S.; Khare, S.; Kumar, A. B.; Supek, F.; Buchynskyy, A.; Mathison, C. J. N.; Chennamaneni, N. K.; Pendem, N.; Buckner, F. S.; Gelb, M. H.; Molteni, V. *Chem. Rev.* **2014**, *114*, 11305.
- (d) Gema, D.; Javier, P. C. *Eur. J. Org. Chem.* **2011**, *2011*, 7243.
- (e) Zhang, G.-J.; Hu, F.; Jiang, H.; Dai, L.-M.; Liao, H.-B.; Li, N.; Wang, H.-S.; Pan, Y.-M.; Liang, D. *Phytochemistry* **2018**, *145*, 68.
- (f) Venkataramana, R. P. O.; Hridhay, M.; Nikhil, K.; Khan, S.; Jha, P. N.; Shah, K.; Kumar, D. *Bioorg. Med. Chem. Lett.* **2018**, *28*, 1278.
- (g) Devi, N.; Kumar, S.; Pandey, S. K.; Singh, V. *Asian J. Org. Chem.* **2018**, *7*, 6.

- [2] (a) Shilabin, A. G.; Kasanah, N.; Tekwani, B. L.; Hamann, M. T. *J. Nat. Prod.* **2008**, *71*, 1218.
(b) Winkler, J. D.; Londregan, A. T.; Hamann, M. T. *Org. Lett.* **2006**, *8*, 2591.
- [3] (a) Boursereau, Y.; Coldham, I. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 5841.
(b) Guan, H.; Chen, H.; Peng, W.; Ma, Y.; Cao, R.; Liu, X.; Xu, A. *Eur. J. Med. Chem.* **2006**, *41*, 1167.
(c) Rashid, M. A.; Gustafson, K. R.; Boyd, M. R. *J. Nat. Prod.* **2001**, *64*, 1454.
(d) Prinsep, M. R.; Blunt, J. W.; Munro, M. H. G. *J. Nat. Prod.* **1991**, *54*, 1068.
(e) Yang, J.-M.; Zhu, Y.-H.; Chen, S.; Lu, X.; Wu, Y.-M.; Ma, F.-E.; Li, L.-P.; Yang, Y.; Shi, Z.-H.; Huang, K.-Y.; Hong, X.; Jiang, P.; Peng, Y. *MedChemComm* **2018**, *9*, 100.
(f) Dighe, S. U.; Khan, S.; Soni, I.; Jain, P.; Shukla, S.; Yadav, R.; Sen, P.; Meeran, S. M.; Batra, S. J. *Med. Chem.* **2015**, *58*, 3485.
(g) Kamal, A.; Srinivasulu, V.; Nayak, V. L.; Sathish, M.; Shankaraiah, N.; Bagul, C.; Reddy, N. V. S.; Rangaraj, N.; Nagesh, N. *ChemMedChem* **2014**, *9*, 2084.
- [4] (a) Tang, J.; Wang, Y.; Wang, R.; Dong, Z.; Yang, L.; Zheng, Y.; Liu, J. *Chem. Biodiversity* **2008**, *5*, 447.
(b) Wang, Y.-H.; Tang, J.-G.; Wang, R.-R.; Yang, L.-M.; Dong, Z.-J.; Du, L.; Shen, X.; Liu, J.-K.; Zheng, Y.-T. *Biochem. Biophys. Res. Commun.* **2007**, *355*, 1091.
(c) Yu, X.; Lin, W.; Li, J.; Yang, M. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 3127.
- [5] (a) Hagen, T. J.; Skolnick, P.; Cook, J. M. *J. Med. Chem.* **1987**, *30*, 750.
(b) Müller, W. E.; Fehske, K. J.; Borbe, H. O.; Wollert, U.; Nanz, C.; Rommelspacher, H. *Pharmacol., Biochem. Behav.* **1981**, *14*, 693.
- [6] (a) Im, Y.; Lee, J. Y. *Chem. Commun.* **2013**, *49*, 5948.
(b) Paul, B. K.; Ghosh, N.; Mukherjee, S. *RSC Adv.* **2016**, *6*, 9984.
(c) Swami, S.; Behera, D.; Agarwala, A.; Verma, V. P.; Shrivastava, R. *New J. Chem.* **2018**, Ahead of Print.
- [7] (a) Whaley, W. M.; Govindachari, T. R. In *Organic Reactions*, John Wiley & Sons, Inc., **2004**.
(b) Cox, E. D.; Cook, J. M. *Chem. Rev.* **1995**, *95*, 1797.
(c) Rao, R. N.; Maiti, B.; Chanda, K. *ACS Comb. Sci.* **2017**, *19*, 199.
(d) Gobe, V.; Gandon, V.; Guinchard, X. *Adv. Synth. Catal.* **2018**, *360*, 1280.
(e) Glenn, C. C.; Jan, B. *Eur. J. Org. Chem.* **2004**, *2004*, 1286.
- [8] Bischler, A.; Napieralski, B. *Ber. Dtsch. Chem. Ges.* **1893**, *26*, 1903.
- [9] (a) Singh, D.; Sharma, P.; Kumar, R.; Pandey, S. K.; Malakar, C. C.; Singh, V. *Asian J. Org. Chem.* **2018**, *7*, 383.
(b) Singh, D.; Hazra, C. K.; Malakar, C. C.; Pandey, S. K.; Kaith, B. S.; Singh, V. *ChemistrySelect* **2018**, *3*, 4859.
(c) Kovvuri, J.; Nagaraju, B.; Nayak, V. L.; Akunuri, R.; Rao, M. P. N.; Ajitha, A.; Nagesh, N.; Kamal, A. *Eur. J. Med. Chem.* **2018**, *143*, 1563.
(d) Singh, D.; Kumar, V.; Devi, N.; Malakar, C. C.; Shankar, R.; Singh, V. *Adv. Synth. Catal.* **2017**, *359*, 1213.
(e) Manasa, K. L.; Tangella, Y.; Ramu, G.; Babu, B. N. *ChemistrySelect* **2017**, *2*, 9162.
(f) Galvis, C. E. P.; Kouznetsov, V. V. *Synthesis* **2017**, *49*, 4535.
(g) Hati, S.; Sen, S. *Tetrahedron Lett.* **2016**, *57*, 1040.
(h) Meesala, R.; Arshad, A. S. M.; Mordí, M. N.; Mansor, S. M. *Tetrahedron* **2016**, *72*, 8537.
(i) Kamal, A.; Tangella, Y.; Manasa, K. L.; Sathish, M.; Srinivasulu, V.; Chetna, J.; Alarifi, A. *Org. Biomol. Chem.* **2015**, *13*, 8652.
(j) Kamal, A.; Sathish, M.; Prasanthi, A. V. G.; Chetna, J.; Tangella, Y.; Srinivasulu, V.; Shankaraiah, N.; Alarifi, A. *RSC Adv.* **2015**, *5*, 90121.
(k) Kamal, A.; Narasimha Rao, M. P.; Swapna, P.; Srinivasulu, V.; Bagul, C.; Shaik, A. B.; Mullagiri, K.; Kovvuri, J.; Reddy, V. S.; Vidyasagar, K.; Nagesh, N. *Org. Biomol. Chem.* **2014**, *12*, 2370.
- [10] (a) Ding, S.; Shi, Z.; Jiao, N. *Org. Lett.* **2010**, *12*, 1540.
(b) Tang, S.; Wang, J.; Xiong, Z.; Xie, Z.; Li, D.; Huang, J.; Zhu, Q. *Org. Lett.* **2017**, *19*, 5577.
(c) Mulcahy, S. P.; Varelas, J. G. *Tetrahedron Lett.* **2013**, *54*, 6599.
(d) Dhara, S.; Singha, R.; Ahmed, A.; Mandal, H.; Ghosh, M.; Nuree, Y.; Ray, J. K. *RSC Adv.* **2014**, *4*, 45163.
(e) Yan, Q.; Gin, E.; Banwell, M. G.; Willis, A. C.; Carr, P. D. *J. Org. Chem.* **2017**, *82*, 4328.
(f) Dhiman, S.; Mishra, U. K.; Ramasastry, S. S. V. *Angew. Chem., Int. Ed.* **2016**, *55*, 7737.
(g) Pan, X.; Bannister, T. D. *Org. Lett.* **2014**, *16*, 6124.
(h) Wang, T.-T.; Zhang, D.; Liao, W.-W. *Chem. Commun.* **2018**, *54*, 2048.
- [11] Dalvi, B. A.; Lokhande, P. D. *Tetrahedron Lett.* **2018**, *59*, 2145.
- [12] Nissen, F.; Richard, V.; Alayrac, C.; Witulski, B. *Chem. Commun.* **2011**, *47*, 6656.
- [13] (a) Varelas, J. G.; Khanal, S.; O'Donnell, M. A.; Mulcahy, S. P. *Org. Lett.* **2015**, *17*, 5512.
(b) Webb, N. J.; Marsden, S. P.; Raw, S. A. *Org. Lett.* **2014**, *16*, 4718.
- [14] (a) Verniest, G.; England, D.; De Kimpe, N.; Padwa, A. *Tetrahedron* **2010**, *66*, 1496.
(b) Subba Reddy, B. V.; Rajashekhar Reddy, M.; Yarlagadda, S.; Ravikumar Reddy, C.; Ravi Kumar, G.; Yadav, J. S.; Sridhar, B. *J. Org. Chem.* **2015**, *80*, 8807.
- [15] Too, P. C.; Chua, S. H.; Wong, S. H.; Chiba, S. *J. Org. Chem.* **2011**, *76*, 6159.
- [16] Zhu, Y.; Liu, M.; Cai, Q.; Jia, F.; Wu, A. *Chem.-Eur. J.* **2013**, *19*, 10132.
- [17] (a) Li, L. T.; Huang, J.; Li, H. Y.; Wen, L. J.; Wang, P.; Wang, B. *Chem. Commun.* **2012**, *48*, 5187.
(b) Li, L. T.; Li, H. Y.; Xing, L. J.; Wen, L. J.; Wang, P.; Wang, B. *Org. Biomol. Chem.* **2012**, *10*, 9519.
- [18] Wang, B.; Wong, H. N. C. *Bull. Chem. Soc. Jpn.* **2018**, *91*, 710.
- [19] Guo, F.; Wang, L.; Mao, S.; Zhang, C.; Yu, J.; Han, J. *Tetrahedron* **2012**, *68*, 8367.
- [20] (a) Benson, S. C.; Lee, L.; Yang, L.; Snyder, J. K. *Tetrahedron* **2000**, *56*, 1165.
(b) Chiotellis, A.; Muller, A.; Mu, L.; Keller, C.; Schibli, R.; Krämer, S. D.; Ametamey, S. M. *Mol. Pharmaceutics* **2014**, *11*, 3839.
- [21] (a) Garcia, J. M.; Curzon, S. S.; Watts, K. R.; Konopelski, J. P. *Org. Lett.* **2012**, *14*, 2054.
(b) Dua, R. K.; Phillips, R. S. *Tetrahedron Lett.* **1992**, *33*, 29.
(c) Tang, J.-G.; Wang, Y.-H.; Wang, R.-R.; Dong, Z.-J.; Yang, L.-M.; Zheng, Y.-T.; Liu, J.-K. *Chem. Biodiversity* **2008**, *5*, 447.
(d) De Croos, P. Z.; Sangdee, P.; Stockwell, B. L.; Kar, L.; Thompson, E. B.; Johnson, M. E.; Currie, B. L. *J. Med. Chem.* **1990**, *33*, 3138.
- [22] Xu, W.; Zhao, M.; Wang, Y.; Zhu, H.; Wang, Y.; Zhao, S.; Wu, J.; Peng, S. *MedChemComm* **2016**, *7*, 1730.
- [23] Sun, R.; Liu, R.; Zhou, C.; Ren, Z.; Guo, L.; Ma, Q.; Fan, W.; Qiu, L.; Yu, H.; Shao, G.; Cao, R. *MedChemComm* **2015**, *6*, 2170.
- [24] Ikeda, R.; Kimura, T.; Tsutsumi, T.; Tamura, S.; Sakai, N.; Konakahara, T. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 3506.
- [25] Primas, N.; El-Kashef, H.; Lancelot, J.-C.; Lesnard, A.; Dodd, R. H.; Raulta, S. *ARKIVOC* **2010**, *2010*, 14.
- [26] Cao, R.; Chen, Q.; Hou, X.; Chen, H.; Guan, H.; Ma, Y.; Peng, W.; Xu, A. *Bioorg. Med. Chem.* **2004**, *12*, 4613.

(Zhao, C.)