基于芳基糠醛功能化巴比妥酸衍生物识别和分离 $\text{Hg}^{2+}/\text{Fe}^{3+}$ 的新型荧光传感器

朱文博 朱伟 丁金东 马小强 姚红
张有明 林奇* 魏太保*

(西北师范大学化学化工学院 生态环境相关高分子材料教育部重点实验室
甘肃省高分子材料重点实验室 兰州 730070)

摘要 合成了一种新型的基于 5-(3-硝基苯基)-呋喃-2-甲醛功能化的巴比妥酸衍生物传感器 **QS**。通过 ^1H NMR, ^{13}C NMR, MS 等方法对 **QS** 进行了表征。**QS** 的荧光光谱结果表明 **QS** 的二甲亚砜(DMSO)溶液在 498 nm 有最大荧光发射, 在 365 nm 紫外灯下照射发出绿色荧光。**QS** 对 Hg^{2+} 和 Fe^{3+} 的水溶液有不同的荧光检测能力, Hg^{2+} 能使探针 **QS** 的荧光增强为橙色, Fe^{3+} 使其荧光猝灭, 实现了实时检测。传感器 **QS** 对 Hg^{2+} 和 Fe^{3+} 的荧光检测限分别为 3.25×10^{-8} 和 $4.0 \times 10^{-8} \text{ mol} \cdot \text{L}^{-1}$ 。根据 Job 曲线和质谱研究得到 **QS** 与汞离子和铁离子化学计量比均为 1:1, 提出了 **QS** 与 Hg^{2+} 和 Fe^{3+} 的可能结合模式。加入 H_2PO_4^- 能使含有 Fe^{3+} 的 **QS** 溶液(**QS-Fe**)荧光恢复, 可用于 Fe^{3+} 的循环检测。此外, 固体 **QS** 可从水溶液中吸附 Hg^{2+} 和 Fe^{3+} (吸附率分别为 92.0% 和 91.8%), 具有较好的吸附能力。

关键词 巴比妥酸衍生物; 荧光传感器; 铁离子; 汞离子; 离子吸附

A Novel Fluorescent Sensor Based on Aryl-furfural Functionalized Barbituric Acid for Recognition and Separation of $\text{Hg}^{2+}/\text{Fe}^{3+}$

Zhu, Wenbo Zhu, Wei Ding, Jindong Ma, Xiaoqiang
Yao, Hong Zhang, Youming Lin, Qi* Wei, Taibao*

(Key Laboratory of Polymer Materials of Gansu Province, College of Chemistry and Chemical Engineering,
Northwest Normal University, Lanzhou 730070)

Abstract A novel fluorescent sensor based on 5-(3-nitrophenyl)-furan-2-carbaldehyde functionalized barbituric acid derivative (**QS**) was successfully synthesized. **QS** was characterized by ^1H NMR, ^{13}C NMR and MS. The maximum fluorescence emission wavelength of **QS** in dimethyl sulfoxide (DMSO) solution was determined to be 498 nm and green fluorescence was emitted under 365 nm ultraviolet lamp. **QS** showed different fluorescence identification ability for aqueous solution of Hg^{2+} and Fe^{3+} . Hg^{2+} can enhance the fluorescence of probe **QS** to orange, and Fe^{3+} quenches its fluorescence, thus realizing real-time detection. The limits of detection of **QS** for Hg^{2+} and Fe^{3+} are 3.25×10^{-8} and $4.0 \times 10^{-8} \text{ mol} \cdot \text{L}^{-1}$, respectively. The fluorescence measurements and MS studies suggest that the binding stoichiometry ratios of **QS** with Hg^{2+} and Fe^{3+} are recognized as 1:1, respectively. The possible modes of **QS** with Hg^{2+} and Fe^{3+} are proposed. The addition of H_2PO_4^- caused the fluorescence of **QS-Fe** recover, indicating that **QS** can be used for cyclic detection of Fe^{3+} . More importantly, the separation percentages of the solid **QS** for Hg^{2+} and Fe^{3+} are 92.0% and 91.8% in aqueous solution, respectively, indicating that it has excellent ingestion capacity.

Keywords barbituric acid derivatives; fluorescence sensor; iron ion; mercury ion; ion absorption

1 Introduction

With rapid development of international globalization

and industrialization, heavy-metal pollution has attracted much more attention in the field of environment and human health.^[1,2] Due to the non-biodegradation nature of

* Corresponding authors. E-mail: weitaibao@126.com, linqi2004@126.com

Received March 24, 2019; revised May 13, 2019; published online June 6, 2019.

Project supported by the National Natural Science Foundation of China (Nos. 21662031, 21661028, 21574104, 21262032).

国家自然科学基金(Nos. 21662031, 21661028, 21574104, 21262032)资助项目。

mercury ion (Hg^{2+}), it can only be diluted but not to be destroyed.^[3] Long term accumulation of this metal in organisms can destroy DNA, inhibit ligand-receptor interactions, disable normal functions of liver and kidney, disrupt the homeostasis of immune system, and even lead to death.^[4–8] Consequently, the search for an effective detection method of Hg^{2+} has attracted increasing attention in the environmental and medical field. Ferric ion (Fe^{3+}), as an essential trace element in plants and animal body, plays important roles in cell metabolism, enzymatic catalysis, oxygen transport in hemoglobin, and acts as a cofactor in enzyme-based reactions.^[9–13] But abnormal levels of ferric iron are signs of diseases, such as anemia, hemochromatosis, Parkinson's disease, arthritis, diabetes, hepatitis and cancer.^[14,15] The Fe^{3+} is also one of important sources of the environmental pollution.^[16,17] Therefore, many studies have tried to develop a simple, sensitive and reliable method to detect and distinguish Hg^{2+} and Fe^{3+} .

Multifunctional fluorescent sensors^[18–23] have attracted significant interest in recent years because of their simplicity, sensitivity, low-cost and real-time performance compared to traditional methods.^[24–35] It has been reported that many fluorescent sensors are used for monitoring Hg^{2+} and Fe^{3+} , respectively.^[36–39] However, a single probe capable of detecting and absorbing Hg^{2+} and Fe^{3+} is still rare. Therefore, search of a fluorescent sensor with high selectivity and sensitivity, rapid response, recyclable and absorption performance is a very challenging issue.

Barbituric acid (BA) as an important compound precursor has been utilized for organic functional materials. BA derivatives can be readily obtained from reaction of BA with chromophore substrates through Knoevenagel reactions under mild conditions.^[2,40,41] A few studies have reported two thymine-like imide-nitrogen atoms in the structure of BA, which have the ability to coordinate with Hg^{2+} via the formation of a mercury(II)-BA coordination complex.^[42] However, BA-based fluorescent sensors that can be used for monitoring and separating Hg^{2+} and Fe^{3+} have not been reported so far.

Based on the above considerations and as part of our ongoing interests in fluorescent sensors,^[43–47] BA was chosen as a receptor moiety for designing fluorescent sensor **QS**, which has simple structure and can be readily synthesized by Knoevenagel reaction of BA with 5-(4-nitrophenyl)-furan-2-carbaldehyde (Scheme 1). Interestingly, sensor **QS** showed highly sensitivity and selectivity for the “individual recognition” of Hg^{2+} and Fe^{3+} with different fluorescence performance. Upon addition of Hg^{2+} and Fe^{3+} into the **QS** solution, respectively, Hg^{2+} induced the fluorescence of **QS** to increase and change to orange fluorescence, Fe^{3+} induced the fluorescence quenching. The limits of detection of **QS** for Hg^{2+} and Fe^{3+} were 3.25×10^{-8} and $4.0 \times 10^{-8} \text{ mol} \cdot \text{L}^{-1}$, respectively. It is worth mentioning that the properties of sensor **QS** based on BA derivatives for Fe^{3+} detection were first reported. Moreover, the fluorescence of **QS-Fe** recovered upon addition of H_2PO_4^- into the solution, which indicate that **QS**

can be used for recyclable detection test for Fe^{3+} . More importantly, sensor **QS** also has good absorption capacity for Hg^{2+} and Fe^{3+} (absorption percentages were up to 92.0% and 91.8%, respectively). Hence, sensor **QS** shows potential application prospect in ion detection and separation of Hg^{2+} and Fe^{3+} .

Scheme 1 Synthesis of the sensor **QS**

2 Results and discussion

To investigate the real-time recognition ability of sensor **QS** towards metal ions (such as Fe^{3+} , Hg^{2+} , Ni^{2+} , Zn^{2+} , Ca^{2+} , Cu^{2+} , Cr^{3+} , Ba^{2+} , Co^{2+} , Ag^+ , Pb^{2+} , Cd^{2+} , Al^{3+} and Mg^{2+}), changes in fluorescence spectra were measured immediately by adding above metal ions to the solution of **QS**. Sensor **QS** exhibited green fluorescence ($\lambda_{\text{ex}} = 355 \text{ nm}$). After addition of Hg^{2+} and Fe^{3+} , an obvious fluorescent color changes from green to orange and dim could be observed, respectively. As depicted in Figure 1, after addition of various metal ions (Fe^{3+} , Hg^{2+} , Ni^{2+} , Zn^{2+} , Ca^{2+} , Cu^{2+} , Cr^{3+} , Ba^{2+} , Co^{2+} , Ag^+ , Pb^{2+} , Cd^{2+} , Al^{3+} and Mg^{2+}) into **QS**, only Hg^{2+} remarkably enhanced fluorescence and accompanied by red shifted from 498 to 526 nm, meanwhile only Fe^{3+} caused significantly quenching at 498 nm. These results indicated that sensor **QS** had excellent selectivity for Hg^{2+} and Fe^{3+} .

Figure 1 Fluorescence spectra of **QS** ($1.0 \times 10^{-4} \text{ mol} \cdot \text{L}^{-1}$) in DMSO with metal ions (Fe^{3+} , Hg^{2+} , Ni^{2+} , Zn^{2+} , Ca^{2+} , Cu^{2+} , Cr^{3+} , Ba^{2+} , Co^{2+} , Ag^+ , Pb^{2+} , Hg^{2+} , Al^{3+} and Mg^{2+}) in response to Hg^{2+} and Fe^{3+} (10 equiv.) upon excitation at 355 nm

Inset: color change of **QS** with metal ions

To further explore the practical applicability of sensor **QS** for Hg^{2+} and Fe^{3+} , respectively, competitive experi-

ments were carried out toward other metal ions (Fe^{3+} , Ni^{2+} , Zn^{2+} , Ca^{2+} , Cu^{2+} , Cr^{3+} , Ba^{2+} , Co^{2+} , Ag^{+} , Pb^{2+} , Hg^{2+} , Al^{3+} and Mg^{2+}). As a result, the fluorescence emission spectra of sensor **QS** with Hg^{2+} was not be influenced by other metal ions (Figure 2a). Similarly, other metal ions did not induce any similar fluorescence changes towards **QS-Fe** solution. Therefore, the sensor **QS** has good selectivity for the detection of Hg^{2+} and Fe^{3+} with different fluorescence performance.

Figure 2 Fluorescence emission data for **QS** ($1.0 \times 10^{-4} \text{ mol} \cdot \text{L}^{-1}$) with different perchlorate salts of cations (10 equiv.) and the previous mixture with (a) Hg^{2+} (10 equiv.) and (b) Fe^{3+} (10 equiv.) in DMSO solutions at room temperature (excitation wavelength = 355 nm)

The fluorescent titration experiments were further carried out to investigate the fluorescent sensitivity of **QS** toward Hg^{2+} and Fe^{3+} , respectively. As shown in Figures 3a and 3b, upon addition of Hg^{2+} (0~11 equiv.) to solution of **QS**, the **QS** exhibited an emission enhancement at $\lambda_{\text{em}} = 498 \text{ nm}$ and lightly red-shifted from 498 to 526 nm, which directly led to the change of fluorescence (changed from green to light orange). However, the addition of Fe^{3+} (0~5.5 equiv.) induced an emission decreasing at $\lambda_{\text{em}} = 498 \text{ nm}$. The above process of fluorescent changing may contribute to the intramolecular charge transfer (ICT) effect induced by Hg^{2+} and Fe^{3+} . Furthermore, the association constants for Hg^{2+} and Fe^{3+} were determined to be 1.42×10^5 and $2.25 \times 10^5 \text{ L} \cdot \text{mol}^{-1}$, respectively. The limits of detection of fluorescence spectral change for Hg^{2+} and Fe^{3+} on the basis of $3\delta/s$ method^[48] are 3.25×10^{-8} and $4.0 \times 10^{-8} \text{ mol} \cdot \text{L}^{-1}$, respectively, which indicated that **QS** has high sensitivity for Hg^{2+} and Fe^{3+} .

Interestingly, the fluorescence of **QS-Fe** reversed to its initial optical state after adding H_2PO_4^- (Figure 4). Because Fe^{3+} has binding ability with H_2PO_4^- to form a complex, these performances make the **QS** act as Fe^{3+} and H_2PO_4^- controlled “OFF-ON-OFF” fluorescent switch. The switch can be reversibly operated for at least three

Figure 3 Fluorescence emission spectra of **QS** ($0.1 \text{ mmol} \cdot \text{L}^{-1}$) excited at 355 nm with addition of increasing concentrations of (a) Hg^{2+} (0~11.0 equiv.) and (b) Fe^{3+} (0~5.5 equiv.) in DMSO solution

Inset: a plot of fluorescence intensity depending on the concentration of (a) Hg^{2+} in the range from 0 to 11 equiv. and (b) Fe^{3+} in the range from 0 to 5.5 equiv.

Figure 4 Fluorescence emission spectra of **QS-Fe** ($0.1 \text{ mmol} \cdot \text{L}^{-1}$) excited at 355 nm with addition of increasing concentrations of H_2PO_4^- in DMSO solution

Inset: fluorescence changes under 365 nm UV lamp

cycles with a slight fluorescent efficiency loss.

Moreover, ingestion capacities of the powder **QS** in aqueous solution for $\text{Hg}^{2+}/\text{Fe}^{3+}$ were assessed by inductively coupled plasma (ICP) analysis. The powder **QS** (0.4 mmol) was added into the aqueous solution of Hg^{2+} and

Fe^{3+} (0.5 equiv.), respectively, and the mixture as stirred at room temperature in water for 30 min, the precipitates were separated by centrifugation (1000 r/min, 10 min) and the supernatant retained. The absorption percentages of powder **QS** for Hg^{2+} and Fe^{3+} were up to 92.0% and 91.8%, respectively. It indicated that the powder **QS** had nice absorption ability for Hg^{2+} and Fe^{3+} in aqueous solution.

Figure 5 FT-IR spectra of powder **QS** and **QS-Hg**

The possible mechanism of **QS** for Hg^{2+} and Fe^{3+} was carefully investigated by Job's method, FT-IR, MS and scanning electron microscope (SEM). The Job's method demonstrated that **QS** was combined with Hg^{2+} and Fe^{3+} by 1 : 1 stoichiometry, respectively. The formation of a 1 : 1 complex was further confirmed by the appearance of peaks at $m/z = 561.2607$ and 415.9336 , corresponding to $[\text{QS} + \text{Hg} - 2\text{H} + \text{Cl}]^-$ and $[\text{QS} + \text{Fe} - 2\text{H} + \text{Cl}]^-$. In FT-IR spectra (Figure 5), the typical stretching vibrations of amides (N—H) at *ca.* 3198 cm^{-1} , which correspond to

stretching vibrations in the spectra of **QS**,^[49–51] were disappeared significantly in the spectrum of **QS-Hg** and **QS-Fe**, respectively. This indicates that the amides (N—H) are involved in the coordination with Hg^{2+} and Fe^{3+} , respectively. In addition, the stretching vibrations of the C=O peaks in the amides [at *ca.* $1730\text{ (}\nu_{\text{C=O, NHCONH}}\text{)}$, $1691\text{ (}\nu_{\text{C=O, NHCOC}}\text{)}\text{ cm}^{-1}$] were obviously weakened in the spectrum of **QS-Hg**, while the corresponding peaks show blue-shift and red-shift respectively [at *ca.* $1759\text{ (}\nu_{\text{C=O, NHCONH}}\text{)}$, $1649\text{ (}\nu_{\text{C=O, NHCOC}}\text{)}\text{ cm}^{-1}$] in the spectrum of **QS-Fe**. This further provides evidence for the binding between **QS** and Hg^{2+} , Fe^{3+} via M—N and M—O bonds ($\text{M} = \text{Hg}^{2+}$, Fe^{3+}). Moreover, the addition of H_2PO_4^- into **QS-Fe** solution caused the stretching vibrations of the C=O peaks in the amides recovered [at *ca.* $1730\text{ (}\nu_{\text{C=O, NHCONH}}\text{)}$, $1658\text{ (}\nu_{\text{C=O, NHCOC}}\text{)}\text{ cm}^{-1}$, respectively. These results indicate that the fluorescent changes were caused by the strong binding affinity of Hg^{2+} , Fe^{3+} with N, O atom in BA ring,^[42,49] respectively. Hence, the enhanced fluorescence caused by Hg^{2+} can be attributed to the strong binding affinity of Hg^{2+} with N atom in BA moiety and the subsequent enhancement of the intramolecular charge transfer (ICT) process.^[49] The quenched fluorescence of **QS** likely was induced by the formation of paramagnetic complex between Fe^{3+} with **QS**,^[18,21] which subsequently enhanced the intramolecular charge transfer (ICT) process.^[18,21,49] Moreover, scanning electron microscope (SEM) was used to study this conjecture. The regular rod-like structure of powder **QS** transformed into an irregular block structure of **QS-Hg**, which further proves the existing of coordination interaction between **QS** and Hg^{2+} . The result is in agreement with previous report.^[49] According to above-mentioned evidences, the possible binding modes of **QS** with Hg^{2+} and Fe^{3+} were shown in Scheme 2.

Scheme 2 Proposed combination of **QS** with Hg^{2+} and Fe^{3+}

3 Conclusions

A novel fluorescent sensor based on 5-(3-nitrophenyl)-furan-2-carbaldehyde functionalized barbituric acid (**QS**) has been successfully constructed. Sensor **QS** showed high sensitivity and selectivity for monitor Hg^{2+} and Fe^{3+} , accompanied by different fluorescent performance, respectively. The limits of detection for Hg^{2+} and Fe^{3+} were determined to be 3.25×10^{-8} and $4.0 \times 10^{-8} \text{ mol} \cdot \text{L}^{-1}$, respectively. Moreover, the quenched fluorescence of **QS-Fe** can be recovered by addition of H_2PO_4^- , which could be performed reversibly for Fe^{3+} detection. More importantly, sensor **QS** showed excellent absorption capacity for Hg^{2+} and Fe^{3+} (absorption percentages were up to 92.0% and 91.8%, respectively). In short, sensor **QS** has the advantages of recoverability, cost effective, high sensitivity, convenient synthesis and would be highly desirable from the view point of practical application.

4 Experimental section

4.1 Synthesis of sensor 5-((5-(3-nitrophenyl)furan-2-yl)methylene)pyrimidine-2,4,6(1H,3H,5H)-trione (**QS**)

5-(3-Nitrophenyl)-furan-2-carbaldehyde (1 mmol, 0.215 g), barbituric acid (1 mmol, 0.128 g) and acetic acid (0.1 mL, as a catalyst) were added into ethanol (10 mL). The mixture was stirred at 78 °C for 8 h. After cooling to room temperature, precipitate of **QS** was obtained as a crude product. A yellow solid of **QS** (0.268 g, yield 82%) was got after washed with ethanol for three times. m.p. > 300 °C; ^1H NMR ($\text{DMSO}-d_6$, 600 MHz) δ : 11.36 (s, 1H), 11.28 (s, 1H), 8.49 (d, $J=3.9$ Hz, 1H), 8.32 (d, $J=8.8$ Hz, 2H), 8.17 (d, $J=8.8$ Hz, 2H), 8.09 (s, 1H), 7.66 (d, $J=3.9$ Hz, 1H); ^{13}C NMR ($\text{DMSO}-d_6$, 101 MHz) δ : 163.18, 162.01, 156.90, 151.15, 150.20, 147.41, 135.77, 134.05, 28.46, 126.02, 124.52, 114.42, 114.05; ESI-MS calcd for $\text{C}_{15}\text{H}_{10}\text{N}_3\text{O}_6 [\text{M}+\text{H}]^+$ 328.2600, found 328.2576.

Supporting Information Characterization of compounds, the fluorescence response of the sensor, FT-IR spectrum, SEM, absorption percentage and so on. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] Wang, J.; Li, Y.; Patel, N. G.; Zhang, G.; Zhou, D.; Pang, Y. *Chem. Commun.* **2014**, 50, 12258.
- [2] Zhuang, P.; McBride, M. B.; Xia, H.; Li, N.; Li, Z. *Sci. Total Environ.* **2009**, 407, 1551.
- [3] Yan, F.; Zou, Y.; Wang, M.; Mu, X.; Yang, N.; Chen, L. *Sens. Actuators, B* **2014**, 192, 488.
- [4] Wu, X. F.; Ma, Q. J.; Wei, X. J.; Hou, Y. M.; Zhu, X. *Sens. Actuators, B* **2013**, 183, 565.
- [5] Yue, Q.; Shen, T.; Wang, J.; Wang, L.; Xu, S.; Li, H.; Liu, J. *Chem. Commun.* **2013**, 49, 1750.
- [6] Kim, H. N.; Ren, W. X.; Kim, J. S.; Yoon, J. S. *Chem. Soc. Rev.* **2012**, 41, 3210.
- [7] López-García, I.; Rivas, R. E.; Hernández-Córdoba, M. *Anal. Chim. Acta* **2012**, 743, 69.
- [8] Sun, Z.; Du, J.; Jing, C. *J. Environ. Sci.* **2016**, 39, 134.
- [9] Yan, Z.; Hu, L.; You, J. *Anal. Methods* **2016**, 8, 5738.
- [10] Wu, J.; Liu, W.; Ge, J.; Zhang, H.; Wang, P. *Chem. Soc. Rev.* **2011**, 40, 3483.
- [11] Sahoo, S. K.; Sharma, D.; Bera, R. K.; Crispin, G.; Callan, J. F. *Chem. Soc. Rev.* **2012**, 41, 7195.
- [12] Liu, Y.; Duan, W.; Song, W.; Liu, J.; Ren, C.; Wu, J.; Liu, D.; Chen, H. *ACS Appl. Mater. Interfaces* **2017**, 9, 12663.
- [13] Rauchfus, T. B. *Acc. Chem. Res.* **2015**, 48, 2107.
- [14] Zhang, S.; Li, J.; Zeng, M.; Xu, J.; Wang, X.; Hu, W. *Nanoscale* **2014**, 6, 4157.
- [15] Shi, B.; Su, Y.; Zhang, L.; Huang, M.; Liu, R.; Zhao, S. *ACS Appl. Mater. Interfaces* **2016**, 8, 10717.
- [16] Han, Y.; Wu, X.; Zhang, X.; Zhou, Z.; Lu, C. *ACS Sustainable Chem. Eng.* **2016**, 4, 5667.
- [17] Khadgi, N.; Upreti, A. R.; Li, Y. *RSC Adv.* **2017**, 7, 27007.
- [18] Zhang, Y.-Y.; Chen, X.-Z.; Liu, X.-Y.; Wang, M.; Liu, J.-J.; Gao, G.; Zhang, X.-Y.; Sun, R.-Z.; Hou, S.-C.; Wang, H.-M. *Sens. Actuators, B* **2018**, 273, 1077.
- [19] Sakunkaewkasem, S.; Petdum, A.; Panchan, W.; Sirirak, J.; Charoenpanich, A.; Sooksimuang, T.; Wanichacheva, N. *ACS Sens.* **2018**, 3, 1016.
- [20] Xia, Q.; Chen, Z.; Zhang, Z.; Liu, R. *Chin. J. Org. Chem.* **2018**, 38, 2700 (in Chinese). (夏琦, 陈子康, 张志德, 刘瑞源, 有机化学, **2018**, 38, 2700.)
- [21] Wu, Y.-C.; Jiang, K.; Luo, S.-H.; Cao, L.; Wu, H.-Q.; Wang, Z.-Y. *Spectrochim. Acta, Part A* **2019**, 206, 632.
- [22] Xu, Z.-H.; Wang, Y.; Wang, Y.; Li, J.-Y.; Luo, W.-F.; Wu, W.-N.; Fan, Y.-C. *Spectrochim. Acta, Part A* **2019**, 212, 146.
- [23] Wang, Z.; Yang, J.; Yang, Y.; Xu, X.; Li, M.; Zhang, Y.; Fang, H.; Xu, H.; Wang, S. *Chin. J. Org. Chem.* **2018**, 38, 1401 (in Chinese). (王忠龙, 杨金来, 杨益琴, 徐徐, 李明新, 张燕, 方华, 徐海军, 王石发, 有机化学, **2018**, 38, 1401.)
- [24] Shekari, Z.; Younesi, H.; Heydari, A.; Tajbakhsh, M.; Chaichi, M.; Shahbazi, J. A.; Saberi, D. *Chemosensors* **2017**, 5, 26.
- [25] Shi, B. B.; Zhang, P.; Wei, T.-B.; Yao, H.; Qi, L.; Liu, J.; Zhang, Y.-M. *Tetrahedron* **2013**, 16, 7981.
- [26] Wang, L.-Y.; Fang, G.-P.; Cao, D.-R. *Sens. Actuators, B* **2015**, 207, 849.
- [27] Abrego, Z.; Unceta, N.; Sanchez, A.; Gomez-Caballero, A.; Berrio-Ochoa, L. M.; Goicolea, M. A.; Barrio, J. *Analyst* **2017**, 142, 1157.
- [28] Asadpour-Zeynali, K.; Amini, R. *Sens. Actuators, B* **2017**, 246, 961.
- [29] Kallithrakas-Kontos, N.; Foteinis, S. C. *Anal. Chem.* **2016**, 12, 22.
- [30] Song, C. Y.; Yang, B. Y.; Yang, Y. J.; Wang, L. H. *Sci. China: Chem.* **2016**, 59, 16.
- [31] Sun, Z. L.; Du, J. J.; Jing, C. Y. *J. Environ. Sci.* **2016**, 39, 134.
- [32] West, M. Ellis, A. T.; Potts, P. J.; Strelci, C.; Vanhoof, C.; Wobruschek, P. *J. Anal. At. Spectrom.* **2016**, 31, 1706.
- [33] van den Berg, C. M. G. *Anal. Chem.* **2006**, 78, 156.
- [34] Lunvongsa, S.; Oshima, M.; Motomizu, S. *Talanta* **2006**, 68, 969.
- [35] Graser, C. H.; Banik, N. I.; Bender, K. A.; Lagos, M.; Marquardt, C. M.; Marsac, R.; Montoya, V.; Geckeis, H. *Anal. Chem.* **2015**, 87, 9786.
- [36] Liu, R.; Zeng, J. *Chin. J. Org. Chem.* **2017**, 37, 3274 (in Chinese). (刘瑞姣, 曾竟, 有机化学, **2017**, 37, 3274.)
- [37] Zhang, M.; Xiao, H.; Han, Z.; Yang, L.; Wu, X. *Chin. J. Org. Chem.* **2018**, 38, 926 (in Chinese). (张敏, 肖慧丰, 韩志湘, 仰榴青, 吴向阳, 有机化学, **2018**, 38, 926.)
- [38] Zhang, H.; Nie, C.; Wang, J.; Guan, R.; Cao, D. *Talanta* **2019**, 195, 713.
- [39] Behera, K. C.; Bag, B. *Dyes Pigm.* **2016**, 135, 143.
- [40] Zhang, X.; Man Lai, E. S.; Martin-Aranda R.; Yeung, K. L. *Appl. Catal. A* **2004**, 261, 109.
- [41] Chen, Z.; Cai, D.; Mou, D.; Yan, Q.; Sun, Y.; Pan, W.; Wan, Y.; Song, H.; Yi, W. *Bioorg. Med. Chem.* **2014**, 22, 3279.
- [42] Wang, J.; Zhang, L.; Qi, Q.; Li, S.; Jiang, Y. *Anal. Methods* **2013**, 5,

- 608.
- [43] Zhang, Y.-M.; Zhu, W.; Qu, W.-J.; Zhong, K.-P.; Chen, X.-P.; Yao, H.; Wei, T.-B.; Lin, Q. *Chem. Commun.* **2018**, 54, 4549.
- [44] Qi, L.; Fan, Y.-Q.; Gong, G.-F.; Mao, P.-P.; Wang, J.; Guan, X.-W.; Liu, J.; Zhang, Y.-M.; Yao, H.; Wei, T.-B. *ACS Sustainable Chem. Eng.* **2018**, 6, 8775.
- [45] Li, X.; Lin, Q.; Qu, W.-J.; Li, Q.; Chen, X.-B.; Li, W.-T.; Zhang, Y.-M.; Yao, H.; Wei, T.-B. *Chin. J. Org. Chem.* **2017**, 37, 889 (in Chinese).
(李翔, 林奇, 曲文娟, 李乔, 程晓斌, 李文婷, 张有明, 姚虹, 魏太保, 有机化学, **2017**, 37, 889.)
- [46] Li, W.-T.; Qu, W.-J.; Zhang, H.-L.; Li, X.; Lin, Q.; Yao, H.; Zhang, Y.-M.; Wei, T.-B. *Chin. J. Org. Chem.* **2017**, 37, 2619 (in Chinese).
(李文婷, 曲文娟, 张海丽, 李翔, 林奇, 姚虹, 张有明, 魏太保, 有机化学, **2017**, 37, 2619.)
- [47] Ma, X.-Q.; Wang, Y.; Wei, T.-B.; Qi, L.-H.; Jiang, X.-M.; Ding, J.-D.; Zhu, W.-B.; Yao, H.; Zhang, Y.-M.; Lin, Q. *Dyes Pigm.* **2019**, 164, 279.
- [48] Kim, H. N.; Ren, W. X.; Kim, J. S.; Yoon, J. *Chem. Soc. Rev.* **2012**, 41, 3210.
- [49] Shi, W.; Zhao, S.; Su, Y.; Hui, Y.; Xie, Z. F. *New J. Chem.* **2016**, 40, 7814.
- [50] Lin, L.; Lv, J.; Ji, Y.; Feng, J.; Liu, Y.; Wang, Z.; Zhang, W. *Anal. Lett.* **2013**, 46, 2890.
- [51] Wang, L.; Zeng, R.; Li, C.; Qiao, R. *Colloids Surf., B* **2009**, 74, 284.

(Zhao, C.)