铜催化的有氧氧化合成靛红类衍生物的方法

Ahmad, Muhammad Siddique 主亚敏 郭云龙 张赛赛 沈增明*

(上海交通大学化学化工学院 上海 200240)

摘要 研究了铜催化的脱羰环化生成靛红的方法, 使用氧气作为最终的氧化剂. 这个互补的 $C(sp^3)-H$ 官能团化方法为合成靛红提供了一种新方案. $Cu/O_2/Co$ 体系显示出良好的反应性和兼容性, 富电子和缺电子的官能团都可以很好地兼容. 并基于机理研究和以前的文献报道, 提出了一个可能的假设机理.

关键词 铜催化; 碳氢键活化; 靛红; 脱羰; 氧气

Copper-Catalyzed Aerobic Oxidation Strategy:
A Concise Route to Isatin

Ahmad, Muhammad Siddique Zhu, Yamin Guo, Yunlong

Zhang, Saisai Shen, Zengming*

(School of Chemistry and Chemical Engineering, Shanghai Jiao Tong University, Shanghai 200240)

Abstract A copper-catalyzed decarbonylation cyclization to form isatins using oxygen as terminal oxidant is developed. This complementary way offers a new protocol for the synthesis of isatins through $C(sp^3)-H$ bond functionalization in $Cu/O_2/Co$ system. This system shows good reactivity and compatibility. Both electron-rich and electron-deficient functional groups can be tolerated. A postulated mechanism is proposed based on mechanistic studies and previous reports.

Keywords Cu catalyst; $C-H$ bond activation; isatin; decarbonylation; oxygen

1 Introduction

Isatins and their derivatives are a group of important structural units which have been found in many bioactive molecules.^[1] Isatins are also powerful building blocks for the construction of heterocyclic systems and spirocyclic compounds.^[2] Therefore, many methods have been developed for the synthesis of isatins. Traditionally, the synthesis of isatins was achieved through the methods developed by Sandmeyer,^[3] Stolle,^[4] and Martinet.^[5] However, these methods suffer from harsh conditions, poor yields and limited substrate choices. In view of this, new synthetic methods to overcome these drawbacks are still highly required. Inspiringly, some efficient methods have been found, such as aryne-based methods,^[6] transition-metal catalyzed oxidation,^[7] oxidative cleavage of carbon-carbon triple bond,^[8] sulfur ylide-mediated carbonyl homologation,^[9] iodine-mediated oxidation,^[10] and $C-H$ amination (Scheme 1).^[11] Although these methods showed some im-

provements, there are still many problems like the requirement of heavy metal catalysts and excess amount of bases or strong oxidants and relatively low yields of products. Thus, the development of a concise and environmentally benign process to access isatins remains highly desirable. In 2010, Li and co-workers^[7c] reported a significant and elegant example concerning copper-catalyzed preparation of isatins through intramolecular aromatic $C-H$ and aldehyde $C-H$ dual activation to construct $C-C$ bonds (Scheme 1). After that, Cheng and co-workers^[7g] documented that 2-aminoacetophenones can undergo intramolecular cyclization to form isatins in the presence of a catalytic amount of copper salt under a balloon of O_2 (Scheme 1). Inspired by these works, we envisioned a complementary way of $C(sp^3)-H$ bond functionalization catalyzed by copper to form β -ketoamide core structure. Herein, a copper-catalyzed decarbonylation cyclization to form isatins using oxygen as terminal oxidant is reported.

* Corresponding author. E-mail: shenzengming@sjtu.edu.cn

Received March 14, 2019; revised May 28, 2019; published online June 12, 2019.

Project supported by the National Natural Sciences Foundation of China (No. 21672144).

国家自然科学基金(No. 21672144)资助项目.

Scheme 1 Synthetic approaches for isatins

1 Result and discussion

To optimize reaction conditions, *N*-(2-acetylphenyl)-*N*-methylformamide (**1a**) derived from readily available 1-(2-aminophenyl)ethanone was chosen as a model substrate. Pleasingly, by using catalytic amounts of CuCl_2 in the presence of THF and oxygen at 100°C , the conversion of **1a** into the desired *N*-methyl isatine **2a** was smoothly performed (34% yield of **2a**, Table 1, Entry 1). Next, a series of other copper catalysts was screened, such as CuCl , CuBr , CuI , CuSCN , CuCN , Cu_2O and $\text{Cu}(\text{OAc})_2$. The results exhibited that activity of these copper salts is inferior to CuCl_2 or even ineffective (Table 1, Entries 2~8). Using CuCl_2 as the catalyst, a variety of Lewis acids were examined. A significant Lewis acid effect was observed. CoCl_2 was the most effective Lewis acid for pro-

moting formation of α -ketoamide structure, affording **2a** in 62% isolated yield (Table 1, Entry 16). ZnCl_2 and CoSO_4 slightly increased yield to 40% and 45%, respectively (Table 1, Entries 9 and 15). Other Lewis acids, such as AlCl_3 , $\text{BF}_3\cdot\text{Et}_2\text{O}$, FeCl_3 , PdCl_2 , $\text{Cu}(\text{ClO}_4)_2\cdot 6\text{H}_2\text{O}$, $\text{Yb}(\text{OTf})_3$, $\text{Ni}(\text{OAc})_2\cdot 4\text{H}_2\text{O}$, $\text{Co}(\text{NO}_3)_2\cdot 6\text{H}_2\text{O}$ and $\text{Cu}(\text{OAc})_2\cdot 4\text{H}_2\text{O}$ suppressed reaction and even shut down the process (Table 1, Entries 10~14 and 17~20). Interestingly, when 1 equiv. of protonic acid, PhCOOH , combined with CoCl_2 (1 equiv.) was added into the reaction, the yield of **2a** was increased to 69% (Table 1, Entry 24). To our surprise, when the loading of CoCl_2 and PhCOOH was decreased to 20 mol% catalytic amount, the efficiency of reaction was further improved and 78% yield of **2a** was obtained (Table 1, Entry 25). Notably, control experiment showed that CoCl_2 could not promote this transformation in the absence

of CuCl₂ (Entry 26). Consequently, the optimal conditions involved the use of CuCl₂ (20 mol%), CoCl₂ (20 mol%), PhCOOH (20 mol%) in O₂ and THF at 100 °C.

Table 1 Optimization of reaction conditions for the synthesis of isatins^{a,b}

Entry	Cat.	Additive	Time/h	Yield ^b /%
1	CuCl ₂	None	49	34
2	CuCl	None	110	23
3	CuBr	None	110	17
4	CuI	None	82	N.R.
5	CuSCN	None	82	N.R.
6	CuCN	None	82	N.R.
7	Cu ₂ O	None	110	Trace
8	Cu(OAc) ₂	None	66	N.R.
9	CuCl ₂	ZnCl ₂ (1 equiv.)	49	40
10	CuCl ₂	AlCl ₃ (1 equiv.)	49	N.R.
11	CuCl ₂	BF ₃ ·Et ₂ O (1 equiv.)	67	N.R.
12	CuCl ₂	FeCl ₃ (1 equiv.)	67	N.R.
13	CuCl ₂	PdCl ₂ (1 equiv.)	67	N.R.
14	CuCl ₂	Cu(ClO ₄)·6H ₂ O (1 equiv.)	67	N.R.
15	CuI	CoSO ₄ (1 equiv.)	59	45
16	CuSCN	CoCl ₂ (1 equiv.)	59	62
17	CuCN	Yb(OTf) ₃ (1 equiv.)	48	N.R.
18	Cu ₂ O	Ni(OAc) ₂ ·4H ₂ O (1 equiv.)	65	16
19	Cu(OAc) ₂	Co(NO ₃) ₂ ·6H ₂ O (1 equiv.)	65	Trace
20	CuCl ₂	Cu(OAc) ₂ ·4H ₂ O (1 equiv.)	65	N.R.
21	CuCl ₂	LiCl (1 equiv.)	49	N.R.
22	CuCl ₂	CaCl ₂ (1 equiv.)	49	23
23	CuCl ₂	CoCl ₂ (1 equiv.) + PivOH (1 equiv.)	58	47
24	CuCl ₂	CoCl ₂ (1 equiv.) + PhCOOH (1 equiv.)	57	69
25	CuCl ₂	CoCl ₂ (20 mol%) + PhCOOH (20 mol%)	84	78
26	None	CoCl ₂ (1 equiv.) + PhCOOH (20 mol%)	77	N.R.

^a Conditions: **1a** (50 mg, 0.28 mmol), CuCl₂ (20 mol%), additive (1 equiv.), THF (1.2 mL), O₂, 100 °C. ^b Isolated yield.

With the optimized reaction conditions in hand, the generality and limitations of this Cu-catalyzed decarbonylation cyclization for forming isatins were explored (Table 2). A series of electron-donating and electron-withdrawing substituents on the aromatic ring was examined. It was found that these substrates could be successfully converted to the corresponding isatins in moderate to good yields. Isatin derivatives **2f** and **2g** containing Br and F atoms were smoothly obtained in 71% and 56% yields, respectively,

which had potential application for further functionalization into some bioactive compounds (Table 2). It is worth noting that electron-withdrawing CF₃ and COOMe substituents at the *meta* position on the aromatic rings were tolerated well to furnish the products **2h** and **2i** in moderate yields. To the best of our knowledge, the previous approach reported by Cheng *et al.*^[7g] could not synthesize these isatins. Accordingly, a complementary access to isatin derivatives with electron-withdrawing groups was offered. In addition to examining the substituents on the aromatic ring, the substrates with different size chain on nitrogen atom were also investigated. As expected, substrates with Et, Bn, *n*-Bu on nitrogen efficiently underwent the Cu-catalyzed annulation to give the corresponding isatins in 54% ~ 69% yields (Table 2). To our surprise, *N*-(2-acetylphenyl)formamide without methyl group on the nitrogen also worked well under the standard conditions and provided the desired product **2m** in 48% yield (Table 2).

Table 2 Scope of substrates for the synthesis of isatins^a

2b 62%	2c 44%	2d 61%

2e , 65%	2f , 71%	2g , 56%

2h , 45%	2i , 50%	2j , 69%

2k , 54%	2l , 62%	2m , 48%

^a Conditions: **1** (0.3 mmol), CuCl₂ (20 mol%), CoCl₂ (20 mol%), PhCOOH (20 mol%), THF (1.2 mL), O₂, 100 °C. Isolated yield.

Substrates **1p** and **1q** were also subjected to the standard conditions. Substrate **1p** did not work. Substrate **1q** could give the desired product **2a** in 33% yield after 131 h. These substrates did not show good reactivity (Scheme 2).

To gain insight into the reaction mechanism, some mechanistic studies were conducted. Compound **1n** bearing

Scheme 2 Reaction of compounds **1p** and **1q** in standard conditions

^{13}C isotope on aldehyde group was prepared, and **1n** was subjected to the standard conditions (Eq. 1). It was found that the carbon of amide in **2a** came from methyl group, not aldehyde. In addition, it was found that the substrate **1o** without aldehyde on nitrogen was performed sluggishly under Cu catalytic conditions to furnish the corresponding product **2a** with low 29% yield (Eq. 2). It was worth noting that an intermediate was discovered during the course of reaction, which was determined as a dichloronated compound **3a** by the analysis of $^1\text{H}/^{13}\text{C}$ NMR and HRMS. To our surprise, compound **3a** smoothly converted into the deserted product **2a** under the standard conditions (75% yield, Eq. 3), which revealed that compound **3a** was a key intermediate in this reaction. Consequently, the reaction profile using **1a** as the model substrate was obtained. It was found that the desired product **2a** was produced with the formation of intermediate **3a** (Figure 1). It further supported dichloronated compound **3a** as a key intermediate in this reaction. When a radical trapping agent such as 2,2,6,6-tetramethyl-1-piperidinyloxy free radical (TEMPO) (20 mol%) was added into the reaction, no reaction occurred (Eq. 4). However, although the mechanism for the present catalytic reaction is not clear, a radical process for this reaction can be possible.

Figure 1 Reaction profile using **1a** as the model substrate

On the base of the above mechanistic studies and previous reports,^[7] a tentative reaction mechanism was proposed (Scheme 3). First, substrate reacts with CuCl_2 under the acidic conditions to undergo the halogenation of ketone, forming a dichloronated intermediate **A**,^[12] followed by decarbonylation in Cu/Co system to give a secondary amine radical intermediate **C**.^[7i,7j] The secondary amine radical attacks one of C—Cl bonds to yield compound **D** which undergo substitution reaction with H_2O to provide the cyclic α -keto alcohol **E**. Finally, intermediate **E** is oxidized by $\text{Cu(II)}/\text{O}_2$ to form isatin **2**.

Scheme 3 Postulated mechanism for the formation of isatins

2 Conclusion

In summary, a copper-catalyzed decarbonylation cyclization to form isatins using oxygen as a terminal oxidant was developed. This complementary way offers a new

protocol for the synthesis of isatins through $C(sp^3)$ —H bond functionalization in Cu/O₂/Co system. This system shows good reactivity and compatibility. Both electron-rich and electron-deficient functional groups can be tolerated. Further exploration of the generality and application of this approach is ongoing in our laboratory.

3 Experimental section

3.1 General considerations

All experiments were carried out under an oxygen atmosphere unless otherwise noted. Reactions were monitored using thin-layer chromatography (TLC). ¹H NMR and ¹³C NMR spectra were obtained at 400 and 100 MHz, respectively on a Mercury Plus-400. NMR spectra were run in a solution of deuterated chloroform (CDCl₃). High-resolution mass spectra (HRMS) were recorded using electrospray ionization (ESI) with a Q-TOF MS Spectrometer. The substrates were prepared according to the reported literatures.^[13,14]

3.2 General procedure for Cu-catalyzed decarbonylation cyclization for the synthesis of isatins

An oven-dried Schlenk tube equipped with a magnetic stir bar was evacuated and backfilled with oxygen three times. Under oxygen, substrates **1** (0.3 mmol), CuCl₂ (20 mol%, 0.06 mmol, 8.1 mg), CoCl₂ (20 mol%, 0.06 mmol, 7.8 mg), PhCOOH (20 mol%, 0.06 mmol, 7.3 mg) and tetrahydrofuran (THF) (1.2 mL) were added into the tube. Then the reaction was stirred at 100 °C for specific time. THF was removed under reduced pressure. Then, the mixture was extracted with EtOAc (10 mL × 3). The combined organic layers were dried over Na₂SO₄, filtered and concentrated *in vacuo*. The residue was purified by column chromatography to give the corresponding isatin products. All of products are known compounds.^[14–18]

1-Methylindoline-2,3-dione (**2a**):^[15] 78% yield. ¹H NMR (400 MHz, CDCl₃) δ: 3.26 (s, 3H), 6.91 (d, *J*=8.0 Hz, 1H), 7.16 (td, *J*=1.2, 7.6 Hz, 1H), 7.62~7.59 (m, 2H); HRMS (ESI) calcd for C₉H₈NO₂ 162.0555, found 162.0544.

1,6-Dimethylindoline-2,3-dione (**2b**):^[15] 62% yield. ¹H NMR (400 MHz, CDCl₃) δ: 2.45 (s, 3H), 3.23 (s, 3H), 6.69 (s, 1H), 6.93 (d, *J*=7.6 Hz, 1H), 7.50 (d, *J*=7.6 Hz, 1H); HRMS (ESI) calcd for C₁₀H₁₀NO₂ 176.0712, found 176.0704.

6-Methoxy-1-methylindoline-2,3-dione (**2c**):^[16] 44% yield. ¹H NMR (400 MHz, CDCl₃) δ: 3.22 (s, 3H), 3.93 (s, 3H), 6.37 (d, *J*=2.4 Hz, 1H), 6.57 (dd, *J*=8.4, 2.0 Hz, 1H), 7.59 (d, *J*=8.4 Hz, 1H); HRMS (ESI) calcd for C₁₀H₁₀NO₃ 192.0661, found 192.0647.

1,5-Dimethylindoline-2,3-dione (**2d**):^[15] 61% yield; ¹H NMR (400 MHz, CDCl₃) δ: 2.33 (s, 3H), 3.22 (s, 3H), 6.80 (d, *J*=7.6 Hz, 1H), 7.41~7.38 (m, 2H); HRMS (ESI) calcd for C₁₀H₁₀NO₂ 176.0712, found 176.0700.

5-Methoxy-1-methylindoline-2,3-dione (**2e**):^[15] 65% yield; ¹H NMR (400 MHz, CDCl₃) δ: 3.24 (s, 3H), 3.82 (s, 3H), 6.84 (d, *J*=8.4 Hz, 1H), 7.19~7.15 (m, 2H); HRMS

(ESI) calcd for C₁₀H₁₀NO₃ 192.0661, found 192.0651.

5-Bromo-1-methylindoline-2,3-dione (**2f**):^[15] 71% yield; ¹H NMR (400 MHz, CDCl₃) δ: 3.25 (s, 3H), 6.82 (dd, *J*=7.2, 1.6 Hz, 1H), 7.71 (s, 1H), 7.74 (dd, *J*=7.6, 2.0 Hz, 1H); HRMS (ESI) calcd for C₉H₇NO₃Br 239.9660, found 239.9640.

5-Fluoro-1-methylindoline-2,3-dione (**2g**):^[15] Orange solid, 56% yield. m.p. 150.4~151.3 °C (lit. 150.7~152.5 °C); ¹H NMR (400 MHz, CDCl₃) δ: 3.26 (s, 3H), 6.88~6.85 (m, 1H), 7.36~7.31 (m, 2H); HRMS (ESI) calcd for C₉H₇NO₃F 180.0461, found 180.0454.

1-Methyl-6-(trifluoromethyl)indoline-2,3-dione (**2h**):^[17] 45% yield. ¹H NMR (400 MHz, CDCl₃) δ: 3.32 (s, 3H), 7.11 (s, 1H), 7.43 (d, *J*=8.0 Hz, 1H), 7.73 (d, *J*=8.0 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 26.4, 106.9 (d, *J*=3.5 Hz, 1C), 119.4, 120.8 (d, *J*=3.7 Hz, 1C), 124.1, 125.5, 139.2 (d, *J*=33.3 Hz, 1C), 151.5, 157.4, 182.4; HRMS (ESI) calcd for C₁₀H₇NO₂F₃ 230.0429, found 230.0425.

1-Methyl-2,3-dioxindoline-6-carboxylate (**2i**):^[18] 50% yield. ¹H NMR (400 MHz, CDCl₃) δ: 3.32 (s, 3H), 3.98 (s, 3H), 7.54 (d, *J*=1.2 Hz, 1H), 7.68 (dd, *J*=8.0, 0.8 Hz, 1H), 7.84 (dd, *J*=7.6, 1.2 Hz, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 26.4, 52.9, 110.6, 120.0, 125.2, 138.5, 151.1, 157.6, 165.2, 183.1; HRMS (ESI) calcd for C₁₁H₁₀NO₄ 220.0610, found 220.0604.

1-Ethylindoline-2,3-dione (**2j**):^[19] 69% yield. ¹H NMR (400 MHz, CDCl₃) δ: 1.34 (t, *J*=7.2 Hz, 1H), 3.82 (q, *J*=7.2 Hz, 2H), 6.92 (d, *J*=7.6 Hz, 1H), 7.13 (td, *J*=0.8, 7.2 Hz, 1H), 7.62~7.57 (m, 2H); HRMS (ESI) calcd for C₁₀H₁₀NO₂ 176.0712, found 176.0700.

1-Butylindoline-2,3-dione (**2k**):^[20] 54% yield. ¹H NMR (400 MHz, CDCl₃) δ: 0.99 (t, *J*=7.2 Hz, 3H), 1.46~1.37 (m, 2H), 1.72~1.65 (m, 2H), 3.74 (t, *J*=7.2 Hz, 2H), 6.91 (d, *J*=8.0 Hz, 1H), 7.13 (dt, *J*=7.6, 0.8 Hz, 1H), 7.61~7.56 (m, 2H); HRMS (ESI) calcd for C₁₂H₁₄NO₂ 204.1025, found 204.1011.

1-Benzylindoline-2,3-dione (**2l**):^[15] 62% yield. ¹H NMR (400 MHz, CDCl₃) δ: 4.93 (s, 2H), 6.78 (d, *J*=8.0 Hz, 1H), 7.11 (t, *J*=7.6 Hz, 1H), 7.37~7.30 (m, 5H), 7.50 (dt, *J*=7.6, 0.8 Hz, 1H), 7.62 (d, *J*=7.6 Hz, 1H); HRMS (ESI) calcd for C₁₅H₁₂NO₂ 238.0868, found 238.0858.

Indoline-2,3-dione (**2m**):^[15] 48% yield; ¹H NMR (400 MHz, CDCl₃) δ: 6.93 (d, *J*=8.0 Hz, 1H), 7.15 (td, *J*=0.8, 7.6 Hz, 1H), 7.59 (td, *J*=1.2, 7.6 Hz, 1H), 7.63 (d, *J*=7.6 Hz, 1H), 8.09 (br s, 1H); HRMS (ESI) calcd for C₈H₆NO₂ 148.0399, found 148.0386.

N-(2-Acetylphenyl)-*N*-methylformamide-¹³C (**1n**): Major : Minor = 3 : 1; ¹H NMR (400 MHz, CDCl₃) δ: Major: 2.51 (s, 3H), 3.26 (s, 3H), 7.25 (dd, *J*=1.2, 7.2 Hz, 1H), 7.43 (td, *J*=1.2, 7.6 Hz, 1H), 7.57 (td, *J*=1.6, 7.6 Hz, 1H), 7.70 (dd, *J*=7.6, 1.6 Hz, 1H), 8.38 (s, 1H); Minor: 2.56 (s, 3H), 3.37 (s, 3H), 7.24 (d, *J*=7.2 Hz, 1H), 7.38 (td, *J*=1.2, 7.6 Hz, 1H), 7.53 (td, *J*=1.6, 7.6 Hz, 1H), 7.65 (dd, *J*=1.2, 7.6 Hz, 1H), 8.40 (s, 1H); ¹³C NMR (100 MHz, CDCl₃) δ: 29.5, 33.8, 128.0, 129.8, 132.7, 136.4, 162.6, 199.6; HRMS (ESI) calcd for C₁₀H₁₂NO₂ 178.0902, found 178.0891.

N-(2-(2,2-Dichloroacetyl)phenyl)-*N*-methylformamide (**3a**): Major : Minor = 3 : 1; ^1H NMR (400 MHz, CDCl_3) δ : Major: 3.25 (s, 3H), 6.48 (s, 1H), 7.33 (d, $J=7.6$ Hz, 1H), 7.52 (t, $J=7.6$ Hz, 1H), 7.67 (t, $J=7.6$ Hz, 1H), 7.84 (d, $J=7.6$ Hz, 1H), 8.19 (s, 1H); Minor: 3.39 (s, 3H), 6.51 (s, 1H), 7.30 (d, $J=8.0$ Hz, 1H), 7.44 (t, $J=8.0$ Hz, 1H), 7.64 (t, $J=8.0$ Hz, 1H), 7.91 (d, $J=8.0$ Hz, 1H), 8.18 (s, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ : Major: 34.6, 69.3, 128.7, 129.5, 130.4, 132.1, 134.5, 141.8, 162.8, 188.1; Minor: 38.0, 69.0, 127.9, 128.1, 129.3, 132.7, 134.0, 139.1, 163.1, 188.3; HRMS (ESI) calcd for $\text{C}_{10}\text{H}_9\text{Cl}_2\text{NO}_2\text{Na}$ ($\text{M}+\text{Na}^+$) 267.9908, found 267.9903.

Supporting Information ^1H NMR spectra of all products and data of intermediate **3a**. The Supporting Information is available free of charge via the Internet at <http://sioc-journal.cn>.

References

- [1] (a) Sumpter, W. C. *Chem. Rev.* **1944**, *34*, 393.
(b) da Silva, J. F. M.; Garden, S. J.; Pinto, A. C. *J. Braz. Chem. Soc.* **2001**, *12*, 273.
- [2] (a) Ratan, B. T.; Anand, B.; Yogeewari, P.; Sriram, D. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 4451.
(b) Raj, A.; Raghunathan, R.; Sridevikumaria, M. R.; Raman, N. *Bioorg. Med. Chem.* **2003**, *11*, 407.
(c) Verma, M.; Pandeya, S. N.; Singh, K. N.; Stables, J. P. *Acta Pharm.* **2004**, *54*, 49.
(d) Jiang, T.; Kuhen, K. L.; Wolff, K.; Yin, H.; Bieza, K.; Caldwell, J.; Bursulaya, B.; Tuntlad, T.; Zhang, K.; Karanewsky, D.; He, Y. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 2109.
(e) Aboul-Fadl, T.; Bin-Jubair, F. A. S. *Int. J. Res. Pharm. Sci.* **2010**, *1*, 113.
(f) Sharma, S.; Gupta, M. K.; Saxena, A. K.; Bedi, P. M. S. *Bioorg. Med. Chem.* **2015**, *23*, 7165.
(g) Harbinder, S.; Jatinder, V. S.; Gupta, M. K.; Sharma, S.; Nepali, K.; Bedi, P. M. S. *Bioorg. Med. Chem. Lett.* **2017**, *27*, 3974.
- [3] Sandmeyer, T. *Helv. Chim. Acta* **1919**, *2*, 234.
- [4] (a) Stollé, R. *Ber. Dtsch. Chem. Ges.* **1913**, *46*, 3915.
(b) Stollé, R. *J. Prakt. Chem.* **1922**, *106*, 137.
- [5] (a) Martinet, J. *Compt. Rend.* **1918**, *166*, 85.
(b) Bonnefoy, J.; Martinet, J. *Compt. Rend.* **1921**, *172*, 220.
- [6] Xie, Y. *Chem. Commun.* **2016**, *52*, 12372.
- [7] (a) Sun, J.; Liu, B.; Xu, B. *RSC Adv.* **2013**, *3*, 5824.
(b) Liu, T.; Yang, H.; Jiang, Y.; Fu, H. *Adv. Synth. Catal.* **2013**, *355*, 1169.
- (c) Tang, B.-X.; Song, R.-J.; Wu, C.-Y.; Liu, Y.; Zhou, M.-B.; Wei, W.-T.; Deng, G.-B.; Yin, D.-L.; Li, J.-H. *J. Am. Chem. Soc.* **2010**, *132*, 8900.
- (d) Liu, T.; Yang, H.; Jiang, Y.; Fu, H. *Adv. Synth. Catal.* **2013**, *355*, 1169.
- (e) Sun, J.; Liu, B.-X.; Xu, B. *RSC Adv.* **2013**, *3*, 5824.
- (f) Ilangoan, A.; Satish, G. *Org. Lett.* **2013**, *15*, 5726.
- (g) Huang, P. C.; Gandeepan, P.; Cheng, C. H. *Chem. Commun.* **2013**, *49*, 8540.
- (h) Li, J.; Zheng, Y.; Yu, X. L.; Lv, S. Y.; Wang, Q. T.; Hai, L.; Wu, Y. *RSC Adv.* **2015**, *5*, 103280.
- (i) Liu, Z.; Zhang, J.; Chen, S.; Shi, E.; Xu, Y.; Wan, X. *Angew. Chem., Int. Ed.* **2012**, *51*, 3231.
- (j) Meng, Q.; Wang, F.; Li, M. *J. Mol. Model.* **2013**, *19*, 2225.
- [8] Liao, Y.-Y.; Gao, Y.-C.; Zheng, W.; Tang, R.-Y. *Adv. Synth. Catal.* **2018**, *360*, 3391.
- [9] Lollar, C. T.; Krennek, K. M.; Bruemmer, K. J.; Lippert, A. R. *Org. Biomol. Chem.* **2014**, *12*, 406.
- [10] (a) Satish, G.; Polu, A.; Ramar, T.; Ilangoan, A. *J. Org. Chem.* **2015**, *80*, 5167.
(b) Reddy, M. R.; Rao, N. N.; Ramakrishna, K.; Meshram, H. M. *Tetrahedron Lett.* **2014**, *55*, 4758.
(c) Ilangoan, A.; Satish, G. *J. Org. Chem.* **2014**, *79*, 4984.
(d) Gao, F. F.; Xue, W. J.; Wang, J. G.; Wu, A. X. *Tetrahedron* **2014**, *70*, 4331.
- [11] (a) Huang, P. C.; Gandeepan, P.; Cheng, C. H. *Chem. Commun.* **2013**, *49*, 8540.
(b) Wu, H.; Zhang, Z. G.; Liu, Q. F.; Liu, T. X.; Ma, N. N.; Zhang, G. S. *Org. Lett.* **2018**, *20*, 2897.
(c) Salvanna, N.; Reddy, L. M.; Kumar, R. A.; Das, B. *ChemistrySelect* **2018**, *3*, 8019.
- [12] Nobrega, J. A.; Goncalves, S. M. C.; Peppe, C. *Synth. Commun.* **2002**, *32*, 3711.
- [13] Shekhar, A. C.; Kumar, A. R.; Sathaiiah, G.; Paul, V. L.; Sridhar, M.; Rao, P. S. *Tetrahedron Lett.* **2009**, *50*, 7099.
- [14] Kirincich, S. J.; Xiang, J.; Green, N.; Tam, S.; Yang, H. Y.; Shim, J.; Clark, J. D.; McKew, J. C. *Bioorg. Med. Chem.* **2009**, *17*, 4383.
- [15] Luo, J. F.; Gao, S. S.; Ma, Y. R.; Ge, G. P. *Synlett* **2018**, *29*, 969.
- [16] Ji, H. H.; Zhu, Y. Z.; Shao, Y.; Liu, J.; Yuan, Y.; Jia, X. D. *J. Org. Chem.* **2017**, *82*, 9859.
- [17] Wang, H. Y.; Wang, K. Y.; Man, Y. Q.; Gao, X. N.; Yang, L. M.; Ren, Y. F.; Li, N.; Tang, B.; Zhao, G. *Adv. Synth. Catal.* **2017**, *359*, 3934.
- [18] Bredenkampa, A.; Mohrb, F.; Kirsch, S. F. *Synthesis* **2015**, *47*, 1937.
- [19] Zhang, C.; Li, S.; Filip, B.; Richmond, L.; Ye, X.; Jiang, Z. *ACS Catal.* **2016**, *6*, 10, 6853.
- [20] Salvanna, N.; Ramesh, P.; Kumarc, K. S.; Das, B. *New J. Chem.* **2017**, *41*, 13754.

(Zhao, C.)